

1 **Bisaccate pollen from the Early Permian OSPZ3a Sub-Biozone of the**
2 **Lower Gharif Member, Oman**

3
4 Michael H. Stephenson *

5
6 *British Geological Survey, Keyworth, Nottingham, NG12 5GG, UK*

7 * Corresponding author. *E-mail address:* mhste@bgs.ac.uk.
8

9
10 **Abstract**

11
12 The OSPZ3a Sub-Biozone, associated with the lowest part of the Lower Gharif Member, is
13 part of biozonal scheme that was intended to unify the palynological schemes across Arabia.
14 This paper describes and illustrates the main bisaccate pollen taxa from the OSPZ3a Sub-
15 Biozone of the Well A cored well, Oman, between 2842.69 and 2852.82 m, where they are
16 unusually well preserved. *Pteruchipollenites indarraensis* which is the most common
17 bisaccate pollen taxon, reaching 40 to 50% of assemblages, is here placed in synonymy with
18 *Alisporites tenuicarpus* Balme, 1970, the latter being its junior synonym. *Striatopodocarpites*
19 *cancellatus* consistently first occurs in the OSPZ3a Sub-Biozone, and well-preserved
20 specimens are present in Well A, but Arabian specimens appear to have a wider range of
21 morphology, mainly in the arrangement of taeniae, than the type material. The relationship of
22 the genus *Striatopodocarpites* to *Verticipollenites* Bharadwaj, 1962, *Lahirites* Bharadwaj,
23 1962 and *Hindipollenites* Bharadwaj, 1962 is also examined with the result that
24 *Striatopodocarpites* is asserted as the senior synonym. The taeniata bisaccate pollen
25 *Hamiapollenites fusiformis* Marques-Toigo, 1974 is unusually common in the Well A

26 assemblages and its morphology is found to be distinct from the similar multi-taeniate
27 bisaccate taxon *Striatoabieites multistriatus* (Balme and Hennelly) Hart, 1964, with which it
28 is sometimes placed in synonymy.

29

30 **Key words:**

31

32

33 **1. Introduction**

34

35 The Gharif Formation (the upper formation of the Haushi Group, Hughes Clarke, 1988)
36 overlies the Al Khlata Formation both disconformably and conformably and is in turn
37 overlain conformably by the marine carbonates and marginal marine to non-marine 'red-bed'
38 clastics of the Khuff Formation (Osterloff et al., 2004). The formation is subdivided into 3
39 members: the Lower, Middle and Upper Gharif members, using subsurface sections (Forbes
40 et al., 2010; Fig. 1). In South Oman, the lower part of the Lower Gharif Member is a complex
41 of fluvial and fluviodeltaic clastics succeeded by marginal marine clastics toward the top;
42 while in North Oman similar lower clastics give way to bioclastic limestone, known locally
43 as the Haushi limestone (Fig. 1). The surface equivalent of the Lower Gharif Member was
44 termed the Saiwan Formation by the Bureau de Recherches Géologiques et Minières
45 (BRGM) (Dubreuilh et al., 1992; Platel et al., 1992; Roger et al., 1992). The Middle Gharif
46 Member is a sequence of marginal marine clastics overlain by lacustrine and fluvial units,
47 capped by stacked palaeosols (the 'Playa Shale' sensu Guit et al., 1995), deposited in a semi-
48 arid climate. Lying unconformably above the Middle Gharif Member is the Upper Gharif
49 Member. The upper part of this clastic unit contains abundant plant remains (Broutin et al.,
50 1995) in the outcrops of the Northern Huqf area.

51

52 Age determinations for the Lower Gharif member rely mainly on macropaleontological dates
53 from the Haushi Limestone. Miller and Furnish (1957) and Hudson and Sudbury (1959)
54 suggested Mid Permian and Sakmarian-Artinskian ages respectively for fauna from the
55 Haushi limestone. More recent foraminiferal evidence (Angiolini et al., 2006) suggests a
56 Sakmarian age for the Haushi limestone, providing an upper age limit for the Lower Gharif
57 Member.

58

59

60 **2. OSPZ3a**

61

62 The OSPZ3a Sub-Biozone, is part of biozonal scheme that was intended to unify the
63 palynological schemes across Arabia (Stephenson et al., 2003). The OSPZ3a Sub-Biozone is
64 succeeded by the OSPZ3b and OSPZ3c sub-biozones. Compared with the other biozones of
65 the OSPZ scheme, the three sub-biozones are smaller in scale and somewhat localised in
66 geographical extent, and appear not to be recognisable throughout Arabia either due to
67 palaeophytogeographical variation or hiatus.

68

69 The base of OSPZ3a Sub-Biozone is marked by the most distinct palynological discontinuity
70 in the Lower Permian section, which corresponds closely to the transition between the Rahab
71 Shale Bed of the Al Khlata Formation and the Lower Gharif Member, and may also be linked
72 with post-glacial climatic change (Stephenson and Osterloff, 2002; Stephenson et al., 2005).

73 The base is defined by the abrupt increase of the small non-taeniate bisaccate pollen

74 *Pteruchipollenites indarraensis* from approximately 10 to 50 or 60% of assemblages. This

75 increase is accompanied by an increase in coarsely ornamented forms of *Cristatisporites* (to a

76 maximum of approximately 4% of assemblages). Other taxa that occur first consistently in
77 OSPZ3a Sub-Biozone are the taeniate bisaccate pollen *Striatopodocarpites cancellatus* and *S.*
78 *fusus*, and the taeniate 'circumstriate' pollen taxa such as *Circumstriatites talchirensis* and
79 *Striasulcites tectus*. The colpate pollen *Kingiacolpites subcircularis* is common throughout,
80 occasionally reaching 50% of assemblages, but more typically 5-10% of assemblages. The
81 base of OSPZ3b Sub-Biozone is defined by the first uphole appearance of the algal cyst
82 *Ulanisphaeridium omanensis* Stephenson and Osterloff, 2002.

83

84 It is difficult to generalise about the relationship between the 2141C Biozone of Penney et al.
85 (2008) and OSPZ3a, and there may be some overlap in character. However 2141C contains
86 more common *Cycadopites cymbatus*, and coarsely ornamented triangular fern spores such as
87 *Converrucosiporites grandegrnulatus* and *Converrucosiporites* sp. A of Stephenson and
88 Osterloff (2002) (see also Stephenson, 2004). 2141C is regarded as an approximate
89 equivalent of the *Converrucosiporites* sp. A – *Microbaculispora grandegrnulata* Biozone of
90 Stephenson and Osterloff (2002).

91

92 This paper describes the main bisaccate pollen taxa from the OSPZ3a Sub-Biozone of the
93 Well A cored well (Fig. 2) between 2842.69 and 2852.82 m (driller's depths) within the
94 Lower Gharif Member. Well A assemblages represent a typical post-glacial Lower Gharif
95 Member flora. The simple bisaccate pollen may have been produced by upland plants, while
96 the lowland may have been populated by colpate pollen-producing cycad-like plants
97 (Stephenson and Osterloff, 2002; Stephenson et al., 2005). Spores of the Lower Permian
98 succession have been described previously (Stephenson, 2004), but pollen have not, mainly
99 because preservation is generally poor amongst the bisaccate pollen of the Lower Gharif
100 Member. However the core of Well A yielded relatively well-preserved bisaccate pollen from

101 the OSPZ3a Sub-Biozone, allowing this part of the palynological succession in Oman to be
102 described and illustrated.

103

104

105 **3. Materials and methods**

106

107 Preparation of strew mounts for palynological analysis involved well-established procedures
108 of crushing followed by hydrochloric and hydrofluoric acid treatments (Wood et al., 1996).

109 Post-hydrofluoric acid organic residues were oxidised with Schulze's Solution and dilute
110 nitric acid. Slides are held in the collection of the British Geological Survey, Keyworth,
111 Nottingham, NG12 5GG.

112

113 The terminology used is that of Punt et al. (1994) and Smith and Butterworth (1967).

114 Maximum equatorial dimensions are given in microns (μm); and the scheme of dimensions as
115 given in Fig 3. Previous records of taxa given here are not meant to be exhaustive, but to
116 focus on Middle Eastern occurrences. Stratigraphical ages for these occurrences are those
117 suggested by the respective authors.

118

119

120 **4. Systematic palynology**

121

122 Genus *Pteruchipollenites* Couper, 1958

123

124 *Type species: Pteruchipollenites thomasii* Couper, 1958.

125

126 *Pteruchipollenites indarraensis* (Segroves) Foster, 1979 (Plate I, 1–18)

127

128 1970 *Alisporites tenuicarpus* Balme; p. 394; pl. 15, figs. 1-4.

129 1988 *Pteruchipollenites* sp. 1 MacRae; p. 56-57; pl. 22, figs. 7-8, 11-13, 15-16, 18-19.

130

131 *Description:* Pollen bilaterally symmetrical, bisaccate, alete; amb haploxyelonoid oval to

132 slightly irregular due to flaccid sacci. Corpus latitudinally oval, circular or rarely

133 longitudinally oval; intexine thin ($<0.5\mu\text{m}$); exoexine over corpus intexine infrareticulate.

134 Cappula usually indistinct; when distinct, oval; margins sometimes imperceptible due to the

135 similarity of corpus and saccus exine. Saccus inclination distal; proximal saccus detachment

136 equatorial; distal saccus detachment $5\text{-}8\mu\text{m}$ in from the distal outer margin of the corpus.

137 Sacci variable in size, usually with width 30% of the corpus width; crescentic in outline;

138 flaccid; infrareticulation fine (brochi size less than $1\mu\text{m}$ in diameter).

139 *Mean dimensions:* (37 specimens): corpus width $36.8\mu\text{m}$; total length $34.26\mu\text{m}$; maximum

140 offlap $7.05\mu\text{m}$; maximum onlap $10.06\mu\text{m}$; total width $51.11\mu\text{m}$.

141 *Remarks:* The rapid uphole increase in the abundance of *P. indarraensis* is not recorded in the

142 cored section of Well A and thus the base of the OSPZ3a Sub-Biozone is not present, but

143 Figure 4 serves to indicate its abundance. Figure 5 shows the base of the OSPZ3a Sub-

144 Biozone across the Rahab-2, Thuleilat-42 and 16, and Marmul-151 wells (data from

145 Stephenson and Osterloff, 2002). Due to its delicacy, it is often poorly preserved. The

146 preservation is poorer in the upper part of the Well A core, between 2842.69 and 2852.82 m

147 (Fig. 4), with the result that many likely specimens of *P. indarraensis* were recorded as

148 ‘bisaccate pollen indeterminate’ (Fig. 4). Poorly preserved specimens of *P. indarraensis*

149 usually occur as detached sacci or corpi.

150 *Pteruchipollenites indarraensis* is similar to *Alisporites tenuicarpus* Balme, 1970, as noted by
151 Foster (1979). *Alisporites tenuicarpus* is diagnosed as having minor distal saccus inclination
152 and an oval cappula of width about half that of the corpus. *P. indarraensis* is diagnosed as
153 having a parallel-sided cappula of width 1/5-2/3 the width of the corpus. Segroves'
154 description of the cappula as parallel-sided is inconsistent, however, with the cappulae
155 figured (Segroves 1969; pl. 6, figs. A-E), which are oval in outline.

156 A careful study of a large number of the present specimens has shown that the distally-
157 inclined, flaccid sacci are compressed variably; sometimes inward toward the cappula,
158 sometimes outward to expose the cappula. When the sacci are pushed inward, folding
159 obscures the distal saccus roots and the cappula may appear to be narrow. When the sacci are
160 pushed outward by compression the cappula is exposed and stretched sideways so that it may
161 appear artificially wide. Furthermore, the similarity of saccus and corpus exine often makes
162 the determination of saccus onlap and cappula shape difficult to discern.

163 In view of the fact that *P. indarraensis* and *A. tenuicarpus* are separated on minor size
164 difference and cappula width (which may be influenced by preservation), it is suggested that
165 the two species be placed in synonymy with *P. indarraensis* as the senior synonym.

166 *Pteruchipollenites* sp. 1 MacRae, 1988 is also synonymous with *P. indarraensis*.

167 *Previous records*: Iran, Permian (Chateauneuf and Stampfli, 1979); Africa, Permian
168 (MacRae, 1988); Pakistan, Permian (Balme, 1970); Australia, Permian (Foster, 1979;
169 Segroves, 1969); Middle East, Permian (Stephenson and Osterloff, 2002; Stephenson et al.,
170 2003, 2005).

171

172 *Genus* ***Hamiapollenites*** Wilson emend. Tshudy and Kosanke, 1966

173

174 *Type species*: *Hamiapollenites saccatus* Wilson, 1962.

175

176 *Hamiapollenites fusiformis* Marques-Toigo emend. Archangelsky and Gamarro, 1979 (Plate
177 II, 1–10)

178

179 *Description:* Pollen bilaterally symmetrical, bisaccate, monolete, dilete or trilete, taeniate;
180 amb oval or diploxytonoid with very small sacci. Corpus oval, dark in colour; intexine thin,
181 punctate. Cappa has approximately 9 taeniae of dense, dark-coloured exoexine separated by
182 narrow (0.5µm wide) clefts which are floored by intexine. Taeniae parallel. Central striation
183 deepened in the polar part of the cappa to form distinct monolete, dilete or rarely trilete mark.
184 Sacci small (width <20% of the width of the corpus); situated at the latitudinal extremities of
185 the corpus only; sacci distally inclined. Occasionally thin unexpanded exoexine occurs at the
186 margin of the corpus in the longitudinal positions; unexpanded exoexine 1-3µm thick. Distal
187 exine of the corpus ?laevigate, usually with 2 indistinct longitudinal folds or ?thickenings;
188 thickenings masked by the thick cappa.

189 *Mean dimensions:* (7 specimens): total length 35.57 µm; maximum offlap 5.57 µm; saccus
190 length 10.71 µm; total width 46.14 µm.

191 *Remarks:* Marques-Toigo (1974) did not refer to haptotypic features in her diagnosis, but the
192 figured holotype and a paratype (Marques-Toigo 1974; pl. 3, fig. 9 (holotype), fig. 7
193 (paratype) clearly have a monolete mark.

194 The present species is very similar to *Hamiapollenites karrooensis* (Hart, 1963) Hart, 1964.

195 The latter species differs, however, in having a smaller number of wider proximal taeniae and
196 in lacking a haptotypic mark (see Stephenson, 2008). *H. bullaeformis* differs from the present
197 species in having a single distal, longitudinal keel-like thickening (see Samoilovich, 1953).

198 Foster and Waterhouse (1988) tentatively considered *Hamiapollenites fusiformis* to be

199 synonymous with *Striatoabieites multistriatus* (Balme and Hennelly) Hart, 1964. Marques-

200 Toigo (1974) did not compare her species with *S. multistriatus*. Via the respective diagnosis
201 and description (Marques-Toigo, 1974, p. 611; Balme and Hennelly, 1955, p. 93)
202 comparisons are difficult to make. A visual comparison of the respective figured specimens
203 (Marques-Toigo, 1974; pl. 3, figs. 7-10 (holotype fig. 9); Balme and Hennelly, 1955; pl. 2,
204 figs. 16-20 (lectotype fig. 17, designated by Hart (1964) however, show that the corpus of *S.*
205 *multistriatus* bears a larger number of narrower taeniae (approximately 20 taeniae in each of
206 the 5 figured specimens) than does that of *H. fusiformis*. Balme and Hennelly (1955) did not
207 specify the number of taeniae in their ‘description’, but Hart (1964) in his later ‘diagnosis’
208 specified 12-16 taeniae. Marques-Toigo (1974) diagnosed 9-12 taeniae for *H. fusiformis*.
209 In addition, the figures appear to indicate that *H. fusiformis* has a generally darker and more
210 clearly oval corpus than *S. multistriatus*, and bears longitudinal distal thickenings which are
211 absent in *S. multistriatus*. *Hamiapollenites fusiformis* bears a haptotypic mark (usually
212 monolete) which appears to be absent in the figured specimens of *S. multistriatus*.
213 *Previous records*: Uruguay, Early Permian (Marques-Toigo, 1974); Argentina, Permo
214 Carboniferous (Vergel, 1987; Archangelsky and Gamarro, 1979, Césari et al., 1995); Middle
215 East, Permian (Stephenson and Osterloff, 2002).

216

217 Genus *Striatopodocarpites* Sedova emend. Hart, 1964

218

219 *Type species*: *Striatopodocarpites tojmensis* Sedova, 1956.

220

221 1962 *Verticipollenites* Bharadwaj: p. 90-91; pl. 9, figs. 126-127, 129-136; pl. 10, figs. 137-
222 139, 143-146; pl. 11, figs. 158-159; pl. 12, figs. 160, 162-165, 168-171, 173; pl. 13, figs. 177-
223 178, 180, 186.

224 1962 *Hindipollenites* Bharadwaj: p. 92-93; pl. 10, figs. 141-142.

225 1962 *Lahirites* Bharadwaj: p. 91-92; pl. 11, figs. 152-153; pl. 12, fig. 172; pl. 13, figs. 181,
226 183, 188.

227

228 *Remarks:* The present author concurs with Hart (1964) in placing *Verticipollenites* Bharadwaj
229 1962, *Lahirites* Bharadwaj, 1962 and *Hindipollenites* Bharadwaj, 1962 in synonymy with the
230 present genus. Hart (1964) did not give reasons for his synonymisation. An attempt will be
231 made here to justify synonymisation.

232 Bharadwaj (1962) did not compare *Verticipollenites* with *Striatopodocarpites* but his
233 comparison with other diploxylonoid taeniate pollen indicates that he believed

234 *Verticipollenites* to be distinct because of the outline of the saccus which is 'pitcher' or flask-
235 shaped (see Bharadwaj, 1962, text-fig. 9). The figure of the holotype of the type species, *V.*

236 *secretus* (Bharadwaj, 1962; pl. 12, fig. 160), does not clearly show this feature and the sacci

237 outlines in other figured specimens are similarly obscure. In the absence of definite evidence

238 to suggest that saccate arrangement in *Verticipollenites* is distinct, the latter genus, which has

239 no other distinguishing features, is considered a junior synonym of the present genus.

240 Bharadwaj (1962) asserted that *Hindipollenites* is distinct from *Verticipollenites* only in its

241 intrapunctate ?corpus exine because it also has 'pitcher' or flask-shaped sacci. Generic

242 differentiation based solely on minor corpus exine- taeniae exine differences is considered

243 unsafe because these may be of secondary origin. As in the figured specimens of

244 *Verticipollenites*, the figured specimens of *Hindipollenites* (Bharadwaj, 1962; pl. 10, figs.

245 141-142) do not clearly show "pitcher" or flask-shaped sacci and so *Hindipollenites* is

246 similarly considered to be a junior synonym of *Striatopodocarpites*. *Lahirites* purportedly

247 differs from *Striatopodocarpites* in lacking the latter's structured corpus exine. As has been

248 suggested above, the present author considers such differences as possibly secondary in

249 origin and for this reason considers *Lahirites* to be synonymous with *Striatopodocarpites*.

250

251 *Striatopodocarpites cancellatus* (Balme and Hennelly) Hart, 1964 (Plate III, 1–8; Plate IV,
252 1–3)

253

254 *Description:* Pollen, bisaccate, bilaterally symmetrical taeniate; amb diploxytonoid. Corpus
255 distinct, circular, dark in colour; cappa 1–2 μ m thick, distinctly taeniate. Cappula distinct,
256 narrow (20% of the corpus width), parallel sided; delineated by narrow distal intexinal folds.
257 Cappa with approximately 8 taeniae; taeniae narrow (2–3 μ m wide), parallel or sub-parallel;
258 extend the width of the corpus; striations between <1 μ m wide. Proximal saccus detachment
259 equatorial; distal saccus detachment close to distal pole; sacci distally inclined. Sacci outline
260 is semi circular; sacci coarsely infrareticulate (brochi 1–2 μ m wide); sacci flaccid.

261 *Mean dimensions* (5 specimens): total width 40 μ m, total length 23 μ m, corpus width 16 μ m,
262 saccus offlap 12 μ m, saccus onlap 7 μ m, cappula width 4 μ m.

263 *Remarks:* *Striatopodocarpites cancellatus* is the most common diploxytonoid multitaeniate
264 bisaccate pollen occurring in OSPZ3a. Though most specimens fit well within the concept of
265 the species as described by, for example, Balme and Hennelly (1955) and Foster (1979),
266 some vary from that concept in two ways: in the form of taeniae, and in the development of
267 rudimentary haptotypic marks. Although this is not discussed by Balme and Hennelly (1955)
268 and Foster (1979) and other authors, Australian specimens of *S. cancellatus* tend to show
269 rather regular, parallel taeniae. In a small proportion of Oman specimens, and also in Saudi
270 Arabian and Yemeni specimens, the taeniae are discontinuous or only sub-parallel. Some
271 Pakistan specimens of *S. cancellatus* from the Salt Range also show non-parallel taeniae (see
272 Balme, 1970). There is a continuum between such specimens and those with regular, parallel
273 taeniae, and thus it was not thought judicious to separate the two groups taxonomically.
274 Arabian specimens of *S. cancellatus* also sometimes bear a disruption of the taeniae in the

275 central part of the cappa, suggesting a rudimentary haptotypic mark, usually a monolete or
276 dilete mark. Again such structures are not shown in illustrations of Australian specimens of *S.*
277 *cancellatus*, nor are they mentioned in descriptions. In cases where monolete marks are clear
278 (e.g. Plate IV, 4–5), such specimens are assigned to *Strotersporites* (see below).

279 *Previous records:* Israel, Late Permian (Eshet, 1990); Saudi Arabia, Late Permian (Hemer,
280 1965); Iran, Late Permian (Chateauneuf and Stampfli, 1979, Ghavidel-syooki, 1997); Oman,
281 Permian (Love, 1994; Broutin et al., 1995). Middle East, Permian (Stephenson and Osterloff,
282 2002; Stephenson et al., 2003).

283

284

285 Genus *Strotersporites* Wilson emend. Klaus, 1963

286

287 *Type species:* *Strotersporites communis* Wilson, 1962.

288

289 *Remarks:* Klaus' (1963) emendation of *Strotersporites* Wilson created a useful category for
290 monolete/dilete, taeniate, bisaccate grains otherwise similar to *Striatopodocarpites* Sedova
291 emend. Hart, 1964.

292

293 *Strotersporites indicus* Tiwari, 1965 (Plate IV, 4–5)

294

295 *Description:* Pollen bilaterally symmetrical, bisaccate, monolete or dilete, taeniate; amb
296 latitudinally elongate haploxyloid. Corpus oval or barrel-shaped. Longitudinal margins of
297 the corpus often flat and concordant with the longitudinal extremities of the sacci so that the
298 grain overall has flat parallel longitudinal extremities. Intexine relatively thick <1µm;
299 expanded infrareticulate exoexine occurs on the taeniae and in irregular patches on the

300 corpus. Cappula distinct, parallel-sided or rarely fusiform; bounded by two distal intexinal
301 folds marking distal saccus detachment zones. Intexinal folds lunate in shape ~10 μm wide in
302 the central part of the corpus. Width of cappula about 40% of the corpus width. 7- 11
303 proximal taeniae occur, 3-7 μm wide, separated by narrow (1-2 μm wide) striations.
304 Longitudinal extremities of corpus have narrower, convergent taeniae. Proximal saccus
305 detachment equatorial; distal saccus detachment close to distal pole. Sacci distally inclined;
306 outline greater than semi circular; robust, joined at the longitudinal extremities of the corpus
307 by thin strips of expanded exoexine. Saccus infrareticulation coarse (brochi 1-2 μm , elongate,
308 radially arranged on distal side close to the corpus edge). Monolete mark large, distinct,
309 straight or geniculate; situated between the central two taeniae; length 50-80% of the corpus
310 width; intexine of corpus visible along the commissures. Rarely a dilete mark or
311 asymmetrical trilete mark is present.

312 *Mean dimensions:* (14 specimens): corpus width 44.78 μm ; total length 51.64 μm ; maximum
313 offlap 17.36 μm ; maximum onlap 16.64 μm ; total width 78.36 μm .

314 *Remarks:* The present specimens show a wider range of variation than that permitted by
315 Tiwari (1965). Tiwari allows 4-8 striations (=5-9 taeniae) whereas the present specimens
316 have between 7 and 11 (mean 8) taeniae. The specimens of Tiwari (1965) are also
317 considerably larger. These differences however are not considered to justify further
318 separation.

319 Rare specimens have a large dilete or asymmetrical trilete mark which is similar to the "type
320 3" branching striation described by Jizba (1962) in specimens of *Complexisporites*
321 *polymorphus* Jizba, 1962. The mark in the present specimens, however, is never associated
322 with a circumpolar striation as in the latter species. Small specimens of the present species
323 with poorly preserved corpi are however difficult to distinguish from *C. polymorphus*.

324 *Previous records:* Libya, Ghzelian-Early Asselian (Loboziak and Clayton, 1988); India,
325 Early Permian (Tiwari, 1965); Middle East, Permian (Stephenson and Osterloff, 2002).

326

327 Genus *Protohaploxylinus* Samoilovich emend. Morbey, 1975

328

329 *Type species:* *Protohaploxylinus latissimus* (Luber and Valts) Samoilovich 1953.

330 1962 *Faunipollenites* Bharadwaj: p. 95, text-fig. 12; pl. 17, figs. 220-228; pl. 18, figs. 229-
331 234.

332

333 *Remarks:* Bharadwaj (1962) erected *Faunipollenites* to include taeniate, haploxytonoid pollen
334 grains similar to *Protohaploxylinus* but with an ill-defined corpus and infrareticulate cappa.

335 As defined, therefore, it appears to be similar to the type species of *Protohaploxylinus*

336 Samoilovich 1953 (see Luber and Valts, 1941; pl. 13, fig. 221) and Samoilovich (1953; pl. 4,
337 fig. 4) and presumably on this basis was rejected by Hart (1964).

338 Species of *Protohaploxylinus* in OSPZ3a consistently have thin, poorly defined corpi that

339 make them very distinct from the diploxytonoid bisaccate pollen such as *Striatopodocarpites*
340 which have smaller, darker, more distinct corpi.

341

342 *Protohaploxylinus amplus* (Balme and Hennesly) Hart, 1964 (Plate IV, 8)

343

344 *Description:* Pollen bilaterally symmetrical, bisaccate, alete, taeniate; amb oval to sub-
345 rectangular, haploxytonoid. Corpus slightly elongate oval or circular; intexine thin. Cappa
346 exoexine partly expanded, infrareticulate on the taeniae. Cappula parallel-sided to boat
347 shaped, width about 50% of the corpus; delineated by a pair of distal intexinal folds.

348 Approximately 8-10 latitudinal proximal taeniae occur; taeniae exoexine infrareticulate. Sacci

349 distally inclined; proximal saccus detachment equatorial; distal saccus detachment close to
350 distal pole. Sacci hemispherical in outline; appear to join adjacent to the longitudinal
351 extremities of the corpus. Sacci robust; infrareticulation coarse, brochi 1-2 μ m in diameter.
352 *Mean dimensions:* (8 specimens): corpus width 72.25 μ m; corpus length 72 μ m; maximum
353 offlap 22.37 μ m; maximum onlap 24.87 μ m; total width 109.25 μ m.

354 *Remarks:* *Protohaploxypinus limpidus* (Balme and Hennelly) Balme and Playford, 1967 was
355 considered by Balme and Hennelly (1955) to be distinct from *P. amplus* because of its
356 smaller size and thinner, finely granulate body exine. Later workers (e.g. Powis, 1979;
357 unpublished PhD thesis) have shown that the latter also has a larger number of taeniae.

358 *Previous records:* Iran, Permian (Ghavidel-syooki, 1997; Chateauneuf and Stampfli, 1979);
359 Libya, Ghzelian-Artinskian (Brugman et al., 1985 (as *Striatoabietites amplus sic*);
360 Gondwana, Permian (e.g. Balme and Hennelly, 1955; Bose and Kar, 1966; Balme and
361 Playford, 1967; Foster, 1975; Backhouse, 1991; MacRae, 1988; Césari et al., 1995;
362 Stephenson and Osterloff, 2002).

363

364 *Protohaploxypinus limpidus* (Balme and Hennelly) Balme and Playford, 1967 (Plate IV, 9)

365

366 *Description:* Pollen bilaterally symmetrical, bisaccate, taeniate, alete; amb oval
367 haploxytonoid. Corpus latitudinally oval; intexine thin; taeniae exoexine partially
368 infrareticulate. Cappula distinct, 5-10 μ m wide (approximately 10% of the width of the
369 corpus); parallel-sided, extends the length of the corpus. Cappa with 5-8 taeniae; usually
370 convergent, 3-14 μ m in width. Sacci strongly distally inclined with very narrow sacci
371 connections at the longitudinal margins of the corpus. Sacci detached equatorially on
372 proximal side of corpus, cappula margins mark the distal saccus detachment; saccus offlap
373 dimension usually approximately equal to saccus onlap dimension. Sacci semicircular in

374 outline, roughly the same size as the corpus; infrareticulation fine to coarse (0.5-2 μm brochi
375 diameter).

376 *Mean dimensions:* (12 specimens): total width 69.33 μm ; saccus offlap 14.83 μm ; saccus
377 onlap 14.42 μm ; corpus width 43.67 μm ; corpus length 50.83 μm ; total length 51.67 μm .

378 *Previous records:* Iran, Permian (Ghavidel-syooki, 1997, Chateauneuf and Stampfli, 1979);
379 Gondwana (e.g. Powis, 1979, Lindström, 1996, Backhouse, 1991; Stephenson and Osterloff,
380 2002; Stephenson et al., 2003).

381

382

383 **Acknowledgements**

384

385 The author is grateful to the Ministry of Oil and Gas and Petroleum Development, Oman for
386 permission to publish this paper. The author acknowledges constructive reviews by Stewart
387 G. Molyneux, James B. Riding and Gordon Forbes. Michael H. Stephenson publishes with
388 the permission of the Executive Director, British Geological Survey (NERC).

389

390

391 **References**

392

393 Angiolini, L., M.H. Stephenson and E.J. Leven 2006. Correlation of the Lower Permian
394 surface Saiwan Formation and subsurface Haushi limestone, Central Oman.

395 *GeoArabia*, v. 11, no. 3, p. 17-38.

396 Archangelsky, S. and J.C. Gamero 1979. *Palinología del Palaeozoico Superior en el*

397 *Subsuelo de la Cuenca Chacoparanense, República Argentina 1. Estudio sistemático*

398 de los palinomorfos de tres perforaciones de la Provincia de Córdoba. Revista
399 Española de Micropaleontología, v. 11, p. 417-478.

400 Backhouse, J. 1991. Permian palynostratigraphy of the Collie Basin, western Australia.
401 Review of Palaeobotany and Palynology, v. 67, p. 237-314.

402 Balme, B.E. 1970. Palynology of Permian and Triassic strata in the Salt Range and Surghar
403 Range, west Pakistan. In, B. Kummel and C. Teichert (Eds.), Stratigraphic Boundary
404 Problems: Permian and Triassic of West Pakistan. University Press of Kansas,
405 Department of Geology Special Publication 4, p. 306-453.

406 Balme, B.E. and G. Playford 1967. Late Permian plant microfossils from the Prince Charles
407 Mountains Antarctica. Revue de Micropaléontologie, v. 10, p. 179-192.

408 Balme, B.E. and J.P. Hennelly 1955. Bisaccate sporomorphs from Australian Permian coals.
409 Australian Journal of Botany, v. 3, p. 89-98.

410 Bharadwaj, D.C. 1962. The miospore genera in the coals of the Raniganj Stage, Upper
411 Permian, India. Palaeobotanist, v. 9, p. 68-106.

412 Bose, M.N. and R.K Kar 1966. Palaeozoic spora dispersae from Zaire (Congo) I.
413 Kindu\Kalima and Walikale regions. Annales Musee Royal de L'Afrique Centrale,
414 Sér. 8vo, v. 53, p. 1-168.

415 Broutin, J., J. Roger, J.-P. Platel, L. Angiolini, A. Baud, H. Bucher, J. Marcoux and H. Al
416 Hasmi 1995. The Permian Pangea. Phytogeographic implications of new
417 palaeontological discoveries in Oman (Arabian Peninsula). Compte Rendus de
418 l'Academie des Sciences de Paris, Series IIa, v. 321, p. 1069-1086.

419 Brugman, W.A., J.W. Eggink, S. Loboziak and H. Visscher 1985. Late Carboniferous - Early
420 Permian (Ghzelian - Artinskian) palynomorphs. Journal of Micropalaeontology, v. 4,
421 p. 93-106.

422 Césari, S., S. Archangelsky and L. de Seoane 1995. Palinología del Palaeozoico Superior de
423 la perforación Las Mochas, Provincia de Santa Fe, Argentina. *Ameghiniana*, v. 32, p.
424 73-106.

425 Chateauneuf, J.J. and G. Stampfli 1979. Preliminary report on Permian palynology of Iran.
426 IV International Palynological Conference, Lucknow, v. 2, p. 186-198.

427 Dubreuilh, J., J.-P. Platel, J. Le Métour, J. Roger, R. Wyns, F. Béchenec and A. Berthiaux
428 1992. Explanatory notes to the geologic map of the Khaluf Quadrangle, Sultanate of
429 Oman. Geoscience map, scale 1:250,000, sheet NF 40-15. Ministry of Petroleum and
430 Minerals, Directorate General of Minerals, Sultanate of Oman. 92 p.

431 Eshet, Y. 1990. Paleozoic-Mesozoic palynology of Israel, I. Palynological aspects of the
432 Permo-Triassic succession in the subsurface. *Geological Survey of Israel Bulletin* 81,
433 p. 1-20.

434 Forbes, G.A., H.S.M. Jansen and J. Schreurs 2010. Lexicon of Oman subsurface stratigraphy:
435 Reference guide to the stratigraphy of Oman's hydrocarbon basins. *GeoArabia*
436 Special Publication 5, Gulf PetroLink, Bahrain, 371 p.

437 Foster, C.B. 1975. Permian plant microfossils from the Blair Athol Coal Measures, Central
438 Queensland, Australia. *Palaeontographica, Abteilung B*, v. 154, p. 121-171.

439 Foster, C.B. 1979. Permian Plant Microfossils of the Blair Atholl Coal Measures, Baralaba
440 Coal Measures and Basal Rewan Formation of Queensland. *Geological Survey of*
441 *Queensland Publication no. 372*, 244 p.

442 Foster, C.B. and J. Waterhouse 1988. The *Granulatisporites confluens* Opper Zone and Early
443 Permian marine faunas from the Grant Formation on the Barbwire Terrace, Canning
444 Basin, Australia. *Australian Journal of Earth Sciences*, v. 35, p. 135-157.

- 445 Ghavidel-syooki, M. 1997. Palynostratigraphy and palaeogeography of Early Permian strata
446 in the Zagros basin, southeast-southwest Iran. *Journal of Science of the Islamic*
447 *Republic of Iran*, v. 8, p. 243-261.
- 448 Guit, F.A, M.H. Al-Lawati and P.J.R. Nederlof 1995. Seeking new potential in the early-Late
449 Permian Gharif play, west central Oman. In, M.I. Al-Husseini (Ed.), *Middle East*
450 *Petroleum Geosciences Conference, GEO'94. Gulf PetroLink, Bahrain*, v. 2, p. 447-
451 462.
- 452 Hart, G.F. 1964. A review of the classification and distribution of the Permian miospore:
453 *Disaccate Striatiti. Cinqième Congrès International de Stratigraphie et de Géologie du*
454 *Carbonifère, Paris 1963. Compte Rendu*, v. 3, p. 1171-1199.
- 455 Hemer, D.O. 1965. Application of palynology in Saudi Arabia. *Fifth Arab Petroleum*
456 *Congress, Cairo, March 16th-23rd*, 31 p.
- 457 Hudson, R.G.S. and M. Sudbury 1959. Permian brachiopoda from south east Asia. *Notes et*
458 *Memoires sur le Moyen-Orient, Museum National d'Histoire Naturelle, Paris* v. 7, p.
459 19-55.
- 460 Hughes Clarke, M.W.H. 1988. Stratigraphy and rock unit nomenclature in the oil-producing
461 area of Interior Oman. *Journal of Petroleum Geology*, v. 11, no. 1, p. 5-60.
- 462 Jizba, K.M.M. 1962. Late Palaeozoic bisaccate pollen from the United States midcontinent
463 area. *Journal of Paleontology*, v. 36, p. 871-887.
- 464 Klaus, W. 1963. Sporen aus dem südalpinen Perm. *Jahrbuch der Geologischen Bundesanstalt,*
465 *Wien*, v. 106, p. 229-363.
- 466 Lindström, S. 1996. Late Permian palynology of Fossilryggen, Vestfjella, Dronning Maud
467 Land, Antarctica. *Palynology*, v. 20, p. 15-48.

- 468 Loboziak, S. and G. Clayton 1988. The Carboniferous palynostratigraphy of northeast Libya.
469 In, A. El-Arnauti, B. Owens and B. Thusu (Eds.), *Subsurface Palynostratigraphy of*
470 *Northeast Libya*. Garyounis University Publications, Benghazi, Libya, p. 129-145.
- 471 Love, C.F. 1994. The palynostratigraphy of the Haushi Group (Westphalian-Artinskian) in
472 Oman. In, M.D. Simmons (Ed.), *Micropalaeontology and Hydrocarbon Exploration in*
473 *the Middle East*. Chapman and Hall, p. 23-39.
- 474 Luber, A.A. and I. E. Waltz 1941. Atlas of microspores and pollen grains of the Palaeozoic of
475 the USSR. *Trudy Vsesoiuznogo Nauchno-Issledovatel'skogo Geologicheskogo*
476 *Instituta, Gosgeolizdat, Moscow/Leningrad*, v. 139, p. 1-107. (In Russian).
- 477 MacRae, C.S. 1988. Palynostratigraphical correlation between the Lower Karoo sequence of
478 the Waterburg and Pafuri coal basins and the Hammanskraal plant macrofossil
479 locality, RSA. *Memoirs of the Geological Survey of South Africa*, v. 75, 217 p.
- 480 Marques-Toigo, M. 1974. Some new species of spores and pollens of Lower Permian age
481 from the San Gregorio Formation in Uruguay. *Anais da Academia Brasileira de*
482 *Ciências*, v. 46, p. 601-616.
- 483 Miller, A.K. and W.M. Furnish 1957. Permian ammonoids from southern Arabia. *Journal of*
484 *Palaeontology*, v. 31, p. 1043-1051.
- 485 Osterloff, P., A. Al-Harthy, R. Penney, P. Spaak, G. Williams, F. Al-Zadjali, N. Jones, R.
486 Knox, M.H. Stephenson, G. Oliver and M.I. Al-Husseini 2004. Depositional sequence
487 of the Gharif and Khuff formations, subsurface Interior Oman. In M.I. Al-Husseini
488 (Editor), *Carboniferous, Permian and Early Triassic Arabian Stratigraphy*. GeoArabia
489 Special Publication 3, Gulf PetroLink, Bahrain, p. 83-147.
- 490 Penney, R.A., I. Al Barram and M.H. Stephenson 2008. A high resolution palynozonation for
491 the Al Khlata Formation (Pennsylvanian to Lower Permian), South Oman.
492 *Palynology*, 32, 213-231.

493 Platel, J.-P., J. Le Métour, A. Berthiaux, M. Beurrier and J. Roger 1992. Explanatory notes to
494 the geologic map of the Juzor Al Halaaniyaat Quadrangle, Sultanate of Oman.
495 Geoscience map, scale 1:250,000, sheet NE 40-10. Ministry of Petroleum and
496 Minerals, Directorate General of Minerals, Sultanate of Oman. Muscat. 56 p.

497 Powis, G.D. 1979. Palynology of the Late Palaeozoic glacial sequence, Canning Basin,
498 Western Australia. Unpublished Ph.D. thesis, University of Western Australia.

499 Punt, W., S. Blackmore, S. Nilsson and A. Le Thomas 1994. Glossary of pollen and spore
500 terminology. LPP Foundation Contributions Series no. 1, 71 p.

501 Roger, J., S. Chevrel, J.-P. Platel, F. Bechenec, J. Dubreuilh, J. Le Metour and R. Wyns
502 1992. Geological map of Khaluf, sheet NF 40-15, scale 1:250,000 and explanatory
503 notes: Directorate General of Minerals, Oman Ministry of Petroleum and Minerals.
504 Muscat. 34 p.

505 Samoilovich, S.R. 1953. Pollen and spores from the Permian deposits of the Cherdyia and
506 Aktyubinsk areas, Cis-Urals. Oklahoma Geological Survey Circular 56, University of
507 Oklahoma Press, p. 1-103. (originally in Russian, translated by Elias, M. K.).

508 Segroves, K.L. 1969. Saccate plant microfossils from the Perth Basin of Western Australia.
509 *Grana Palynologica*, v. 9, p. 174-227.

510 Smith, A.H.V. and M.A. Butterworth 1967. Miospores in the coal seams of the Carboniferous
511 of Great Britain. *Special Papers of the Palaeontological Association*, no. 1, p. 1-324.

512 Stephenson, M.H. 2004. Early Permian Spores from Oman and Saudi Arabia. *GeoArabia*
513 *Special Publication 3*, Gulf PetroLink, Bahrain, p. 185-215.

514 Stephenson, M. H. 2008. Spores and Pollen from the Middle and Upper Gharif members
515 (Permian) of Oman. *Palynology*, v. 32, p. 157-183.

516 Stephenson, M.H. and P.L. Osterloff 2002. Palynology of the deglaciation sequence
517 represented by the Lower Permian Rahab and Lower Gharif members, Oman.

518 American Association of Stratigraphic Palynologists, Contribution Series no. 40, p. 1-
519 41.

520 Stephenson, M.H., P.L. Osterloff and J. Filatoff 2003. Palynological biozonation of the
521 Permian of Oman and Saudi Arabia: Progress and challenges. *GeoArabia*, v. 8, no. 3,
522 p. 467-496.

523 Stephenson, M.H., M.J. Leng, C.H. Vane, P.L. Osterloff and C. Arrowsmith 2005.
524 Investigating the record of Permian climate change from argillaceous sediments,
525 Oman. *Journal of the Geological Society, London*, v. 162, p. 641–651.

526 Tiwari, R.S. 1965. Miospore assemblage in some coals of the Barakar Stage (Lower
527 Gondwana) of India. *Palaeobotanist*, v. 13, p. 168-214.

528 Vergel, M. del M. 1987. Consideraciones sobre el contenido microflorístico de al perforación
529 YPF SE AB (Palaeozoico Superior), Aabol Blanco, Provincia de Santiago del Estero,
530 Argentina. VII Simposio Argentino de Paleobotánica y Palinología, Actas. Buenos
531 Aires, Abril, 1987, p. 75-76.

532 Wood, G.D., A.M. Gabriel and J.C. Lawson 1996. Chapter 3. Palynological techniques -
533 Processing and microscopy. In, J. Jansonius and D.C McGregor (Eds.), *Palynology:
534 Principles and Applications*. American Association of Stratigraphic Palynologists
535 Foundation, v. 1, p. 29-50.

536 **Figure captions**

537

538 Fig. 1. Summary stratigraphy of the Carboniferous-Permian of Oman.

539 Fig. 2. Location of wells discussed in the text.

540 Fig. 3. Measurement and orientation scheme used for bisaccate pollen in this study, based on

541 Segroves (1969).

542 Fig. 4. Quantitative character of bisaccate pollen assemblages in Well A, Oman.

543 Fig. 5. The base of the OSPZ3a Sub-Biozone across the Rahab-2, Thuleilat-42 and 16, and

544 Marmul-151 wells (data from Stephenson and Osterloff, 2002).

545

546

547 **Plate captions**

548

549 Plate I. All figures with differential interference contrast (DIC) unless noted. Scale bar

550 indicates 10 μm .

551

552 1–18. *Pteruchipollenites indarraensis*.

553 1. R36, MPA 51778 showing variable form of the cappula.

554 2. R36, MPA 51778 non-DIC.

555 3. Q35, MPA 51778.

556 4. P33/4, MPA 51778 non-DIC.

557 5. R33/2, MPA 51778 non-DIC.

558 6. Q29/2, MPA 51778 non-DIC, slightly oblique compression.

559 7. Q29, MPA 51778 non-DIC.

560 8. Q28/2, MPA 51778 non-DIC.

- 561 9. S29/1, MPA 51778 non-DIC, this specimen shows evidence of weak taeniae and may
562 be transitional to *Protohaploxypinus limpidus*.
- 563 10. N28/3, MPA 51778 lateral compression non-DIC.
- 564 11. N28/2, MPA 51778 lateral compression non-DIC.
- 565 12. N28/2, MPA 51778 showing dense cappa.
- 566 13. M29/3, MPA 51778 non-DIC.
- 567 14. M29/3, MPA 51778.
- 568 15. K16, MPA 51778 non-DIC.
- 569 16. P8/1, MPA 51778 non-DIC.
- 570 17. F21/2, MPA 51778 non-DIC.
- 571 18. F21/2, MPA 51778.

572

573 Plate II. All figures with differential interference contrast (DIC) unless noted. Scale bar
574 indicates 10 μm .

575

576 1–10. *Hamiapollenites fusiformis*.

- 577 1. J21/3, MPA 51777 proximal focus.
- 578 2. J21/3, MPA 51777 focus on distal saccus roots.
- 579 3. Q16, MPA 51777 proximal focus.
- 580 4. Q16, MPA 51777 focus on distal saccus roots.
- 581 5. Q21/1, MPA 51784 proximal focus.
- 582 6. Q21/1, MPA 51784 focus on distal saccus roots.
- 583 7. N22, MPA 51774 proximal focus.
- 584 8. N22, MPA 51774 focus on distal saccus roots.
- 585 9. P38, MPA 51795 non-DIC.

586 10. P38, MPA 51795.

587

588 Plate III. All figures with differential interference contrast (DIC) unless noted. Scale bar

589 indicates 10 μm .

590

591 1–8. *Striatopodocarpites cancellatus*.

592 1. N27, MPA 51787 proximal focus.

593 2. N27, MPA 51787 focus on saccus.

594 3. S16, MPA 51787 proximal focus.

595 4. S16, MPA 51787 focus on saccus.

596 5. S12, MPA 51770 non-DIC.

597 6. S12, MPA 51770 focus on saccus.

598 7. M19/4, MPA 51790 proximal focus.

599 8. M19/4, MPA 51790 focus on distal saccus roots.

600

601 Plate IV. All figures with differential interference contrast (DIC) unless noted. Scale bar

602 indicates 10 μm .

603

604 1–3. *Striatopodocarpites cancellatus*.

605 1. H9/1, MPA 51779 proximal focus.

606 2. H9/1, MPA 51779 focus on cappula.


607 3. H9/1, MPA 51779 focus on distal saccus roots.

608 4–5. *Strotersporites indicus*.

609 4. G35/2, MPA 51779.


610 5. G35/2, MPA 51779.

- 611 6–7. *Strotersporites cf. indicus*.
- 612 6. G35/1, MPA 51779 proximal focus.
- 613 7. K23, MPA 51779 focus on folds at distal saccus roots.
- 614 8. *Protohaploxypinus amplus*. F35/4, MPA 51779.
- 615 9. *Protohaploxypinus limpidus*. E27/4, MPA 51779.




Diploxyelonoid Bisaccate Grain

Proximodistal view




Haploxyelonoid Bisaccate Grain

Proximodistal view




Orientation

Proximodistal view



Longitudinal
meridian



Key


- 1 Total Width
- 2 Saccus Width
- 3 Corpus Width
- 4 Cappula Width
- 5 Saccus Offlap
- 6 Saccus Length
- 7 Corpus Length
- 8 Saccus Onlap

Saccus detachment

P= proximal pole, D= distal pole

Asymmetrical
saccus
detachment:
distal inclination

Symmetrical
saccus
detachment



Proximodistal
view

Cross
section along
latitudinal
meridian

Samples (m)

Selected bisaccate pollen (no. specimens counted)

Pteruchipollenites indarraensis
% of total assemblage

Bisaccates %
of total assemblage

Pteruchipollenites indarraensis

Bisaccate indet

Protohaploxypinus limpidus

Striatopodocarpites spp.


Strotersporites indicus

Striatopodocarpites cancellatus


Hamiapollenites fusiformis

Protohaploxypinus spp.


Striatopodocarpites fusus




Rahab-2




Thuleilat-42



Thuleilat-16



Marmul-151



Base of the OSPZ3a
Sub-Biozone

