



Article (refereed) - postprint

Maes, Dirk; Isaac, Nick J.B.; Harrower, Colin A.; Collen, Ben; van Strien, Arco J.; Roy, David B. 2015. **The use of opportunistic data for IUCN Red List assessments** [in special issue: Fifty years of the Biological Records Centre] *Biological Journal of the Linnean Society*, 115 (3). 690-706. [10.1111/bij.12530](https://doi.org/10.1111/bij.12530)

© 2015 The Linnean Society of London

This version available <http://nora.nerc.ac.uk/511086/>

NERC has developed NORA to enable users to access research outputs wholly or partially funded by NERC. Copyright and other rights for material on this site are retained by the rights owners. Users should read the terms and conditions of use of this material at <http://nora.nerc.ac.uk/policies.html#access>

This document is the author's final manuscript version of the journal article, incorporating any revisions agreed during the peer review process. There may be differences between this and the publisher's version. You are advised to consult the publisher's version if you wish to cite from this article.

The definitive version is available at <http://onlinelibrary.wiley.com/>

Contact CEH NORA team at
noraceh@ceh.ac.uk

1 **JOURNAL**

2 Biological Journal of the Linnean Society

3

4 **TITLE**

5 The use of opportunistic data for IUCN Red List assessments

6

7 **RUNNING HEAD**

8 IUCN Red Listing using opportunistic data

9

10 **AUTHORS**

11 Dirk Maes^a

12 Nick J.B. Isaac^b

13 Colin A. Harrower^b

14 Ben Collen^c

15 Arco J. van Strien^d

16 David B. Roy^b

17

18 **AFFILIATIONS**

19 ^aResearch Institute for Nature and Forest (INBO), Kliniekstraat 25, B-1070 Brussels, Belgium;

20 dirk.maes@inbo.be

21 ^bBiological Records Centre, CEH Wallingford, Maclean Building, Crowmarsh Gifford,

22 Wallingford, Oxfordshire, OX10 8BB, England; njbi@ceh.ac.uk, corr@ceh.ac.uk, dbr@ceh.ac.uk

23 ^cCentre for Biodiversity & Environment Research, Department of Genetics, Evolution &

24 Environment, University College London, Gower Street, London WC1E 6BT, UK;

25 b.collen@ucl.ac.uk

26 ^dStatistics Netherlands, PO Box 24500, NL-2490 HA Den Haag, The Netherlands; asin@cbs.nl

27

28 ***FULL ADDRESS FOR CORRESPONDENCE**

29 Dirk Maes, Research Institute for Nature and Forest (INBO), Kliniekstraat 25, B-1070 Brussels,

30 Belgium, e-mail: dirk.maes@inbo.be

31

32 Version

33 29/06/2015

34

35 Word count (Title – Discussion): 5 956 (title – discussion, without references)

36

37 **The use of opportunistic data for IUCN Red List assessments**

38 DIRK MAES, NICK J.B. ISAAC, COLIN A. HARROWER, BEN COLLEN, ARCO J. VAN STRIEN and
39 DAVID B. ROY

40

41 *IUCN Red Lists are recognized worldwide as powerful instruments for the conservation of*
42 *species. Quantitative criteria to standardise approaches for estimating population trends,*
43 *geographic ranges and population sizes have been developed at global and sub-global levels.*
44 *Little attention has been given to the data needed to estimate species trends and range sizes*
45 *for IUCN Red List assessments. Few regions collect monitoring data in a structured way and*
46 *usually only for a limited number of taxa. Therefore, opportunistic data are increasingly used*
47 *for estimating trends and geographic range sizes. Trend calculations use a range of proxies: i)*
48 *monitoring sentinel populations, ii) estimating changes in available habitat or iii) statistical*
49 *models of change based on opportunistic records. Geographic ranges have been determined*
50 *using: i) marginal occurrences, ii) habitat distributions, iii) range-wide occurrences, iv) species*
51 *distribution modelling (including site-occupancy models) and v) process-based modelling. Red*
52 *List assessments differ strongly among regions (Europe, Britain and Flanders, north Belgium).*
53 *Across different taxonomic groups, in European Red Lists IUCN criterion B and D resulted in the*
54 *highest level of threat. In Britain, this was the case for criterion D and criterion A, while in*
55 *Flanders criterion B and criterion A resulted in the highest threat level. Among taxonomic*
56 *groups, however, large differences in the use of IUCN criteria were revealed. We give examples*
57 *from Europe, Britain and Flemish Red List assessments using opportunistic data and give*
58 *recommendations for a more uniform use of IUCN criteria among regions and among*
59 *taxonomic groups.*

60 **ADDITIONAL KEYWORDS:** Britain – citizen science – Europe – Flanders (north Belgium) –
61 geographic range size – threatened species – trend calculations

62

63

INTRODUCTION

64 IUCN Red Lists are recognized worldwide as very powerful instruments for the conservation of
65 threatened species (Lamoreux *et al.*, 2003; Rodrigues *et al.*, 2006). Although theoretically Red
66 Lists are designed for estimating the extinction risk of species, they are used in conjunction
67 with other information for setting priorities in the compilation of species action plans (e.g.,
68 Keller & Bollmann, 2004; Fitzpatrick *et al.*, 2007), reserve selection and management (e.g.,
69 Simaika & Samways, 2009) and as indicators for the state of the environment (Butchart *et al.*,
70 2006). The compilation of IUCN Red Lists has a long history (Scott, Burton & Fitter, 1987): the
71 first assessments based on (subjective) expert opinion were produced in the 1970's for
72 mammals (IUCN, 1972), followed by fish (IUCN, 1977), birds (IUCN, 1978), plants (Lucas &
73 Synge, 1978), amphibians and reptiles (IUCN, 1979) and invertebrates (IUCN, 1983). Following
74 recognition of the need to standardise approaches to avoid issues such as severity of threat
75 and likelihood of extinction, more objective and quantitative criteria were developed in the
76 1990's (Mace & Lande, 1991; Mace *et al.*, 1993). These criteria have become widely
77 implemented at the global (Mace *et al.*, 2008), national and regional level (Gärdenfors *et al.*,
78 2001; Miller *et al.*, 2007) as a means of classifying the relative risk of extinction of species.

79 As well as on the global level, Red Lists can also be compiled on continental (e.g., European,
80 African), national (e.g., Eaton *et al.*, 2005; Keller *et al.*, 2005; Rodríguez, 2008; Brito *et al.*,
81 2010; Collen *et al.*, 2013; Juslén, Hyvärinen & Virtanen, 2013; Stojanovic *et al.*, 2013) or
82 regional (sub-national) scales (e.g., Maes *et al.*, 2012; Verreycken *et al.*, 2014). Research has
83 mainly focused on the implementation of the IUCN criteria at sub-global levels (Gärdenfors *et al.*
84 *et al.*, 2001), but far less attention has been given to the data needed and/or used to estimate
85 species trends and rarity. The number of species assessed at the global (76 000 species in the
86 latest IUCN update) and sub-global level is large and increasing, and consequently greater
87 scrutiny has been brought to bear on the types of data available to conduct such assessments
88 (e.g., the latest update of the National Red List database contains 135 000 species
89 assessments; www.nationalredlist.org).

90 Only few regions in the world collect data on trends, geographic range size and population
91 sizes in a structured way (e.g., statistically sound monitoring networks – Thomas, 2005),
92 usually for a limited number of taxa (e.g., birds – Baillie, 1990; butterflies – van Swaay *et al.*,
93 2008). Such data collection is often done with a network of volunteer experts (i.e., citizen
94 science) under the co-ordination of professionals (e.g., Jiguet *et al.*, 2012; Pescott *et al.*, 2015).
95 Monitoring data collected in a structured way allow for the use of most of the IUCN criteria,
96 but require sustained funding (Hermoso, Kennard & Linke, 2014). Increasingly, opportunistic
97 data (i.e., distribution records collected by volunteers in a non-structured way) are used for
98 regional Red List assessments (e.g., Fox *et al.*, 2011; Maes *et al.*, 2012). Especially in NW
99 Europe (Britain, the Netherlands, Belgium), the number of volunteers contributing to
100 distribution and monitoring data is increasing yearly (Pocock *et al.*, 2015). In Flanders, for
101 example, the online data portal www.waarnemingen.be of the volunteer nature NGO
102 Natuurpunt started in 2008 and now has almost 20 000 active distribution record providers.
103 The total number of records in the data portal at present amounts to more than 15 million, of
104 which almost 2 million are accompanied by a picture to check identifications. Birds are by far
105 the most recorded taxonomic group in Flanders (51%), followed by plants (26%), moths (8%),
106 butterflies (5%), mushrooms, mammals (both 2%), dragonflies, beetles, flies, bees and wasps,
107 amphibians and reptiles and grasshoppers (all 1%). Whilst the number of records collated is
108 impressive, it is less clear how suitable these opportunistic data are for Red Listing.

109 Opportunistic data are often biased, both in time (e.g., recent periods are usually much
110 better surveyed than ‘historical’ ones), in space (e.g., not all areas are surveyed with an equal
111 intensity – Dennis, Sparks & Hardy, 1999), but also in volunteer preferences for taxonomic
112 groups (e.g., birds, mammals, butterflies) and in differences in observation volunteer skills
113 (e.g., identification errors, detectability - Dennis *et al.*, 2006). A growing diversity of
114 approaches, however, has been developed to take these biases in opportunistic data into
115 account when calculating trends in both abundance and in distribution and geographic ranges
116 (Isaac *et al.*, 2014).

117 Here, we focus on opportunistic citizen science data used to classify species into IUCN Red
118 List categories at sub-global levels. We review the assessment of IUCN criteria in Europe,
119 Britain and Flanders (north Belgium) and give examples of how they were applied in the
120 different regions. Specifically, we examine the role of opportunistic data and compare them
121 with data that have been collected in a standardized way, mainly for the estimation of
122 population trends (IUCN criterion A) and for species' geographic range sizes (IUCN criterion B).

123

124 HOW RED LIST ASSESSMENTS WORK: IUCN CRITERIA AND CATEGORIES

125 Red List categories provide an approximate measure of species' extinction risk in a given
126 region, by quantitatively evaluating some of the key symptoms of risk: 1) a trend in population
127 size or distribution, 2) rarity (abundance) and/or restriction (geographic range) and 3)
128 population size (number of reproductive individuals). These measures reflect the major
129 determinants of risk identified by conservation biology (Caughley, 1994): species are at
130 greatest risk of extinction when population sizes are small, decline rate is high and fluctuations
131 are high relative to population growth. Very small populations are also more susceptible to
132 negative genetic, demographic and environmental effects. At relatively large scales (e.g.,
133 global, continental), data are often very patchy (e.g., GBIF – Beck *et al.*, 2014), but this can also
134 be the case on national or regional levels when survey intensity is low. The over-riding
135 philosophy is to 'make do' with the available data, since the conservation problem is too
136 pressing to wait for more robust data (Hermoso, Kennard & Linke, 2014). IUCN criteria are,
137 therefore, designed to be used with different types of data (Mace, 1994).

138 The IUCN applies five main criteria to classify species in Red List categories:

- 139 A. Population size reduction
- 140 B. Geographic range size
- 141 C. Small population size and decline
- 142 D. Very small population or restricted distribution
- 143 E. Quantitative analysis of extinction risk.

144 Eleven IUCN categories are used for listing species in sub-global Red Lists (Fig. 1 –
145 Gärdenfors *et al.*, 2001). These use the same quantitative criteria as those applied to global
146 Red Lists, but with an additional criterion of downgrading the risk category when rescue
147 effects, across national or regional borders can occur (Gärdenfors *et al.*, 2001). During a Red
148 List assessment, all taxa are assessed against as many IUCN criteria as possible and the Red List
149 category that results in the highest level of extinction risk is assigned to a taxon. Opportunistic
150 data are most often used for assessing IUCN criteria A (population trends) and B (geographic
151 range sizes). But, by making use of expert opinion and when the focal region is well-surveyed,
152 criterion C (population sizes) and D (very small AOO or very limited number of populations) can
153 also be assessed with opportunistic data.

154

155 IUCN CRITERION USE IN EUROPE, BRITAIN AND FLANDERS

156 Many countries and regions make use of the IUCN Red List criteria to estimate species'
157 extinction risks at sub-global levels. Here, we review the use of the different IUCN criteria for
158 Red List assessments in three 'regions': Europe (continental), Britain (national) and Flanders
159 (north Belgium – regional; Table 1). We also give examples of appropriate methods to estimate
160 trends and geographic range sizes for regional Red List assessments.

161 The proportions of the different criteria assessed over all taxonomic groups in Europe,
162 Britain and Flanders are given in Fig. 2. For the European Red Lists, the criteria that resulted in
163 the highest threat level were B (57%) and D (32%). In Britain, this applies to criterion D (47%)
164 and criterion A (27%), while in Flanders; this was the case for criterion B (57%) and criterion A
165 (25%). Among taxonomic groups, however, large differences in the use of the different IUCN
166 criteria were revealed (Fig. 3). In Europe, criterion A resulted in the highest threat level for
167 mammals (44%) and butterflies (43%), criterion B for saproxylic beetles (85%), amphibians
168 (68%) and reptiles (63%), criterion C for dragonflies (21%) and criterion D for terrestrial (51%)
169 and freshwater molluscs (39% – Fig. 3). In Britain, criterion A resulted in the highest threat
170 levels for butterflies (67%) and plants (44%), criterion B for dragonflies (100%) and water

171 beetles (80%), criterion C for flies (30%) and criterion D for boletes (100%) and lichens (68% –
172 Fig. 3). In Flanders, criterion A lead to the highest threat level in water bugs (50%), freshwater
173 fishes (29%) and ladybirds (27%), criterion B for reptiles (100%) and amphibians (83%),
174 criterion C for mammals (18%) and amphibians (17%) and criterion D for mammals only (44% –
175 Fig. 3).

176

177

POPULATION TREND ESTIMATES

178 Few species globally have their entire population monitored regularly in order to accurately
179 assess trends in population size. One of several shortcuts is, therefore, typically employed. A
180 first possible shortcut is to use a small number of sentinel populations that are monitored
181 regularly, either at long-term research sites or as part of co-ordinated schemes such as the UK
182 or Dutch Butterfly Monitoring Scheme (Botham *et al.*, 2013; van Swaay *et al.*, 2013) or the
183 Breeding Bird Survey in the UK or Flanders (Harris *et al.*, 2014; Vermeersch & Onkelinx, 2014).
184 This approach can deliver precise trend estimates, but in most cases the populations are a
185 biased subset and may not be representative of the wider species' population (Brereton *et al.*,
186 2011). A second and coarser tool is to estimate changes in the amount of available habitat,
187 typically from polygon maps, but problems with this approach (commission and omission
188 errors, see further) have been documented and discussed (Boitani *et al.*, 2011). The approach
189 is appealing, as remote sensed data on change in habitat extent can be cost-effectively applied
190 to a range of species. However, even if changes in habitat can be captured accurately, it is
191 unclear how trends reflect actual trends in abundance (Van Dyck *et al.*, 2009). Thus, both these
192 proxies rely on a large number of untested assumptions. A third proxy is to construct a
193 statistical model of change based on opportunistic biological records. Often, measures of
194 change from biological records have been derived from simple 'grid cell counts' between atlas
195 periods (e.g., Maes & van Swaay, 1997; Maes & Van Dyck, 2001; Thomas *et al.*, 2004; Maes *et*
196 *al.*, 2012), which is conceptually similar to the use of habitat extent maps described above.
197 Estimating change from biological records is complicated, because the intensity of recording

198 varies in space and time (Prendergast *et al.*, 1993; Isaac & Pocock, 2015) and can be difficult to
199 estimate from the records alone (Hill, 2012). The development of methods for estimating
200 trends from biological records has recently been the subject of considerable research effort
201 and several robust approaches are increasingly being used. Abundance data is generally
202 considered superior to distributional data for trend estimation (Isaac *et al.*, 2014) and
203 statistical methods are starting to be developed which derive composite trends using models
204 that combine information from both data types (Pagel *et al.*, 2014).

205 Using the IUCN criteria, a population trend (criterion A) can be assessed in five different
206 ways. Criterion Aa (direct observation of population decline) is only rarely used: in the
207 European Red List, eight freshwater fishes, six freshwater molluscs, two terrestrial molluscs
208 and one mammal, plant, reptile and saproxylic beetle were assessed against this criterion. In
209 the UK, criterion Aa was only applied to four vascular plant species, while in Flanders this
210 criterion is not yet used in Red List assessments. The use of criterion Ab (an index of
211 abundance) depends strongly on the taxonomic group (e.g., for British butterflies, an index of
212 abundance (criterion Ab) is available for 49 out of 62 resident species (79%), Fox *et al.*, 2011 –
213 Box 1). Criterion Ac (a decline in geographic range or in habitat quality – Box 2), is the most
214 often used criterion in Britain (93%), in Flanders (91%) and Europe (50% – Fig. 4). Criterion Ad
215 (actual or potential levels of exploitation) is mainly used in European Red List assessments for
216 freshwater fishes (13 species) and mammals (four species). Finally, criterion Ae (effects of
217 introduced taxa, hybridization, pathogens, pollutants, competitors or parasites) is used in 22%
218 of the cases (Fig. 4). Criterion Ae was used mainly for freshwater organisms such as fishes and
219 molluscs where invasive species are a major problem (Strayer, 2010; Roy *et al.*, 2015b). In
220 Flanders, this criterion was also used for the negative effect of the Harlequin ladybird on native
221 ladybirds (Roy *et al.*, 2012a).

222

223 **Box 1 – Trend calculations using abundance data from standardized citizen science**
224 **monitoring data (IUCN criterion Ab)**

225 There is a wide spectrum of citizen science approaches which contribute to monitoring
226 biodiversity, ranging from simple protocols with wide participation to structured approaches
227 which often include elements of professional support and co-ordination (Schmeller *et al.*,
228 2009; Roy *et al.*, 2012b; Isaac & Pocock, 2015; Pescott *et al.*, 2015). Structured, participatory
229 monitoring schemes such as those established for birds, butterflies and mammals in Europe
230 and North America (Devictor, Whittaker & Beltrame, 2010) typically comprise counts of target
231 species throughout the year, repeated annually at fixed locations across a region. For example,
232 the UK Butterfly Monitoring Scheme (UKBMS) provides a standardised annual measure (index)
233 of butterfly populations at line-transect sites (Rothery & Roy, 2001).

234 The UKBMS was initiated in 1976 with 34 sites, rising to more than 100 sites per year from
235 1979 onwards and currently comprises 2000 sites recorded annually. The UKBMS also
236 incorporates a Wider Countryside Butterfly Scheme component to improve the spatial
237 coverage of the scheme (Roy *et al.*, 2015a) Indices from different UKBMS sites over years are
238 combined to derive regional and national collated indices, which can be used to assess long-
239 and short-term population trends (Pannekoek & van Strien, 2003). The UKBMS has been used
240 to assess threat status of 49 out of 62 species (79%) over two time periods: (i) 10 years (1995–
241 2004) and (ii) long-term (typically 1976–2004) for the Red List of British Butterflies (Fox *et al.*,
242 2011). Other examples of the use of structured monitoring schemes are the bird scheme in the
243 UK where 22 out of 74 species (30%) were classified as threatened on the basis of trends in
244 abundances (Eaton *et al.*, 2005).

245 One advantage of a volunteer-based, structured monitoring scheme is good statistical
246 power for measuring trends (e.g. Roy, Rothery & Brereton, 2007) and the capacity to generate
247 time series with comprehensive spatial coverage of a region. They have also provided a rich
248 resource for scientific research, investigating large-scale pattern and processes (Thomas,
249 2005). Although there has been a growth in the number of such schemes in some regions (e.g.,
250 N America, NW Europe) during the current century (Nature Editorials, 2009), there remains a
251 paucity for many species groups in most parts of the world. Successful schemes often rely on

252 institutional support and funding, as well as having a large pool of potential contributors.
253 Although we recommend adopting best practice from established schemes to further their
254 value for future Red List criterion Ab assessments, distribution data (criteria Ac) is typically
255 available for a wider set of species groups and for more regions of the world (see Box 2).

256

257

258 **Box 2 – Trend calculations using opportunistic distribution data (IUCN criterion Ac)**

259 Citizen science data are a potentially valuable source of information of changes in
260 distributions, but they suffer from uneven and unstandardized observation effort (Isaac &
261 Pocock, 2015). Changes in observation efforts across years may easily lead to artificial trends
262 or mask existing trends in species' distributions.

263 In the past, researchers used broad time periods in their comparisons of distribution to
264 ensure sufficient effort and spatial coverage in each time period (van Swaay, 1990). Other
265 authors have filtered their data and used thresholds of completeness of sampling per grid cell
266 (cf. Soberón *et al.*, 2007) for estimating trends (e.g., Maes *et al.*, 2012). Recently, the methods
267 available for trend estimations have developed substantially (Powney & Isaac, 2015). Isaac *et*
268 *al.* (2014) tested a number of approaches for estimating trends from noisy data. Using
269 simulations, they found that simple methods may easily produce biased trend estimates,
270 and/or had low power to detect genuine trends in distribution. Two sophisticated methods
271 known as Frescalo and site-occupancy models emerged as especially promising.

272 Frescalo uses information about sites' similarity to neighbouring sites to assign local
273 benchmark species (Hill, 2012). These benchmarks provide a measure of local observation
274 effort that can be statistically corrected. Frescalo was used to assess changes in plant species
275 distributions for the recent vascular plant Red List for England (Stroh *et al.*, 2014).

276 Site-occupancy models have a special mechanism to adjust for observation effort. They
277 separate occupancy (the *presence* of a species in a site) from detection (the *observation* of the
278 species in that site) when analysing field survey data (MacKenzie *et al.*, 2006). The models

279 require that species are recorded as an assemblage, such that observations of one species can
280 be used to infer non-detection of others (Isaac & Pocock, 2015). Detection can be estimated
281 from sites that were surveyed multiple times in any given time period (e.g., a year). If
282 observation effort increases over time, a species will be observed during more visits, which
283 leads to a higher detection probability, but not to a higher occupancy probability (van Strien,
284 van Swaay & Termaat, 2013). Site-occupancy models have been successfully used in status
285 assessments of butterflies and dragonflies in the Netherlands (van Strien *et al.*, 2010; van
286 Strien, van Swaay & Termaat, 2013).

287

288 METHODS FOR ESTIMATING GEOGRAPHIC RANGE SIZE

289 The IUCN Red List criteria embrace two different measures of geographic range: Extent of
290 Occurrence (EOO) and Area of Occupancy (AOO). The EOO (criterion B1) is defined as the area
291 contained within the shortest continuous imaginary boundary which can encompass all the
292 known, inferred or projected sites of present occurrence of a taxon, excluding cases of
293 vagrancy. The AOO (criterion B2) is intended to represent the total amount of occupied habitat
294 (excluding cases of vagrancy). IUCN guidelines advocate the use of 2 x 2 km² grid cells to
295 estimate AOO (IUCN, 2013), so it is generally used for species with restricted geographic
296 ranges.

297 Different approaches can be applied to estimate geographic range sizes: marginal
298 occurrences, habitat distributions, range-wide occurrences, species distribution modelling
299 (including site-occupancy models) and process-based modelling (Gaston & Fuller, 2009). i)
300 marginal occurrences, i.e., mapping the outer boundaries of species and subsequently
301 interpolating the area in between (Boitani *et al.*, 2011). Such maps are often displayed in field
302 guides to illustrate the possible species distribution range in a usually large region (e.g., world,
303 continent – Graham & Hijmans, 2006). ii) habitat and/or associations with environmental
304 variables as a proxy (Boitani *et al.*, 2011). iii) when range-wide occurrences are available for a
305 focal region (country), records are often assigned to a grid cell projection (e.g., Universal

306 Transverse Mercator – UTM) to produce local or regional distribution atlases. At fine resolution
307 (e.g., 1 x 1 km² or 5 x 5 km²), these data are sufficient to capture a species' distribution, so long
308 as sampling intensity is relatively equally spread over the region (Gaston & Fuller, 2009).
309 Coarse grid cells (e.g., 10 x 10 km² or even 50 x 50 km²) are seldom useful for regional
310 conservation purposes, because they include too much unsuitable habitat (Rondinini *et al.*,
311 2006), but recently, downscaling methods have been proposed to estimate local occupancy
312 from coarse-grain distribution atlas data (Barwell *et al.*, 2014). iv) species distribution
313 modelling is a helpful tool to determine species geographic ranges (Pena *et al.*, 2014).
314 Typically, presence/absence or presence-only data are used in different modelling techniques
315 (Guisan *et al.*, 2013) to 'predict' where suitable environmental conditions occur in a given
316 region for a given species (e.g., Thomaes, Kervyn & Maes, 2008; Cassini, 2011; Syfert *et al.*,
317 2014). v) processed-based modelling using small-scale environmental variables (e.g.,
318 microclimate) can be applied to estimate the possible geographic range of species (e.g.,
319 Kearney, 2006; Kearney *et al.*, 2014; Tomlinson *et al.*, 2014; Panzacchi *et al.*, 2015). Range-
320 wide occurrences tend to underestimate the geographic range of species due to incomplete
321 sampling (omission errors), while the other approaches tend to overestimate the distribution
322 range of species (commission errors) because it incorporates large areas in which the species
323 cannot occur (Gaston & Fuller, 2009).

324

325 ESTIMATING GEOGRAPHIC RANGE SIZES WITH OPPORTUNISTIC DATA

326 EOO and AOO reflect two different processes (spread of extinction risk and vulnerability due to
327 a restricted range, respectively) and it is, therefore, useful to estimate both criteria in Red List
328 assessments. All three regions assessed taxa against both EOO and AOO (Fig. 4). In Europe, the
329 joint use of both EOO and AOO (50%) and AOO alone (50%) resulted equally often in the
330 highest threat level for criterion B, probably depending on individual species' data availability.
331 In Britain, the combined use of EOO and AOO resulted in the highest Red List category (76%),
332 while in Flanders this was the case for AOO (86% - Fig. 4).

333

334 **Box 3 Estimating geographic range sizes (criterion B)**

335 ***EOO (criterion B1): Minimum Convex Polygons for plants and bees in the UK***

336 One of the simplest methods to estimate a species' EOO is to calculate the Minimum Convex
337 Polygon (MCP), the smallest polygon that will contain all the points and in which no internal
338 angle is greater than 180 degrees (Fig. 5b). The MCP has, however, been criticised as being
339 sensitive to errors in location, being derived from the most extreme points (Burgmann & Fox,
340 2003) and for incorporating large areas of unsuitable habitat. Two alternative methods to
341 calculate species ranges that are less susceptible to these issues are: 1) the α -hull (Burgmann &
342 Fox, 2003) and 2) the Localised Convex Hulls (LoCoH) (Getz & Wilmers, 2004). It should be
343 noted that the IUCN guidelines recommend such methods, designed to exclude discontinuous
344 or outlying areas, only when comparing changes in EOO over time discouraging their use when
345 estimating the EOO itself for assessment via criterion B1, as these outlying areas are important
346 in determining the risk associated with geographic range. Both of these methods have recently
347 been applied to Red List assessments in the UK for vascular plants (Stroh *et al.*, 2014) and
348 aculeate Hymenoptera (www.bwars.com; Edwards *et al.*, in prep). The α -hull is derived from a
349 mathematical algorithm for converting points (the locations of records) into triangles based on
350 a threshold parameter α (Burgmann & Fox, 2003). The hull produced becomes more inclusive
351 and approaches the MCP as α increases (Fig. 5c).

352 The Localised Convex Hull (LoCoH) is an adaptation of the MCP but rather than fitting one
353 hull to the entire dataset, the LoCoH is the result of the union of a set of 'localised' MCPs
354 created by fitting the MCP to subsets of the data (Getz & Wilmers, 2004). There are several
355 ways in which these local subsets can be determined (Getz *et al.*, 2007): 1) fixed number of
356 points (k -LoCoH) in which subsets consist of $k-1$ closest points to each root point, 2) fixed
357 sphere-of-influence (r -LoCoH) in which subsets consist of all points within a radius r of each
358 root point, and 3) adaptive sphere-of-influence (a -LoCoH) in which subsets consist of the root
359 point and the closest points where the sum of the distances between the points in the subset

360 and root is less than α . In the UK Red Listing exercises for vascular plants and aculeate
361 Hymenoptera, the fixed sphere-of-influence method (r -LoCoH) was used as it facilitated the
362 data review for the taxonomic exports and because it gave a visual understanding of the final
363 Red Listing decisions (Fig. 6d). This variant of LoCoH is also fairly insensitive to sporadic but
364 spatially clustered recording which is relatively common in opportunistic citizen science data.

365 In both the α -hull and LoCoH, the resulting area is dependent on the value of a control
366 parameter (α for α -hull and k , r , or a for the LoCoH variants). The selection of this parameter is
367 a non-trivial process as it has a marked impact on the EOO estimates. Conceptually, there is no
368 'correct' value. Rather, the most suitable value depends upon i) the aims of the study, i.e., a
369 trade-off between being as inclusive as possible at the cost of including some unsuitable areas
370 (commission errors) or being cautious at the cost of excluding of some suitable areas (omission
371 errors), ii) the degree of spatial coverage in the data (with poorly sampled data requiring
372 higher parameter values) and iii) the properties of the taxa being investigated (e.g., for highly
373 mobile taxa, the most appropriate value is larger than for sedentary ones while large values for
374 linearly distributed taxa (e.g., coastal species) can result in the incorporation of large areas of
375 unsuitable habitat). In the UK Red Listing exercises mentioned above, the parameter values
376 were selected to match the IUCN guidelines and previous Red Listing exercises (i.e., vascular
377 plants – Cheffings *et al.*, 2005) on the one hand or through expert opinion based on the
378 outputs produced using a series of parameter values on the other.

379

380 ***AOO (criterion B2): Ecological ecodistricts for ladybirds in Flanders (north Belgium)***

381 For some regions and for particular taxonomic groups, opportunistic data are available on a
382 high resolution and covering a large part or even the entire region (e.g., birds in the UK –
383 Balmer *et al.*, 2013; butterflies in Flanders – Maes *et al.*, 2012). In such cases, the AOO can be
384 estimated by summing the area of these high resolution grid cells in which a species was
385 observed in a recent period (e.g., 1 x 1 km² – Maes *et al.*, 2012 or 2 x 2 km² – Fox *et al.*, 2011).
386 In regions where mapping coverage for taxonomic groups is fairly incomplete (e.g., ladybirds in

387 Flanders), AOO can be strongly underestimated by using the sum of the area of high resolution
388 grid cells (Sheth *et al.*, 2012). On the other hand, EOO is much less likely to be biased by
389 incomplete sampling, as it uses only the outer boundaries of the distribution. As EOO for
390 ladybirds in Flanders, we, therefore, used the sum of the areas of the ecological districts (i.e.,
391 relatively small and geographical units with a very similar climatology, geology, relief,
392 geomorphology, landscape, etc. – n = 36, Fig. 6) when the species was observed in at least
393 three 1 x 1 km² grid cells in the period 2006-2013. The minimum number of three grid cells per
394 ecological district was applied to exclude single observations of vagrant or erratic individuals.
395 (Adriaens *et al.*, 2015).

396

397

DISCUSSION

398 IUCN enables the use of five different criteria to estimate the extinction risk of species: A)
399 population size reduction, B) geographic range size, C) small population size and decline, D)
400 very small population and/or restricted distribution and/or E) quantitative analysis of
401 extinction risk. In the ideal case, the presence of a statistically sound monitoring scheme in a
402 focal region would allow the use of all IUCN criteria to assess the Red List status of species.
403 With opportunistic data, IUCN criteria A and B can be assessed applying different statistical
404 techniques. But, when mapping intensity is sufficiently high, opportunistic data can also serve
405 to estimate population size classes (criterion C) of some relatively well-known taxonomic
406 groups (e.g., mammals, birds) and for determining species with very small AOO's or a very
407 small number of populations (criterion D).

408 Before assessing taxa against IUCN criteria, it would be desirable to assess whether a focal
409 region has the appropriate data to calculate 'reliable' trends and geographic ranges for a given
410 taxonomic group. In Flanders, prior to the compilation of an IUCN Red List, the institute co-
411 ordinating all regional Red List assessments (i.e., the Research Institute for Nature and Forest –
412 INBO) applies a quantitative and simple procedure to judge whether a dataset is sufficiently
413 good to reliably estimate trends and range sizes. First, the Red List compilers determine which

414 periods will be compared to calculate population trends. Here, IUCN recommends a recent
415 period of 10 year or three generations, whichever is the longer (IUCN, 2003), but many Red List
416 compilers use historical periods that are longer than 10 years usually to compensate for the
417 lower number of historical records in many data sets (e.g., the English Red List of plants – Stroh
418 *et al.*, 2014). Second, for these periods, the grid cells (e.g., 1 x 1 km² or 5 x 5 km²) that have
419 been sufficiently well mapped in common in both periods are located. Mapping intensity can
420 be estimated using species completeness measures (Soberón *et al.*, 2007), rarefaction
421 measures (Carvalho *et al.*, 2013), reference species (Maes & van Swaay, 1997) etc. In a third
422 step, the sufficiently well-surveyed grid cells are attributed to the twelve ecological regions in
423 Flanders (i.e., regions with similar biotopes, soil types and landscapes – Couvreur *et al.*, 2004).
424 To make a representative Red List for a focal region, the recommendation for Flanders is that
425 distribution data should be available in a minimum number of the grid cells (e.g., 10%) in all
426 the (relevant) ecological regions for the given taxonomic group. If a data set of a taxonomic
427 group does not fulfil these criteria, it is considered as currently insufficient for the compilation
428 of an IUCN Red List in Flanders. Fig. 7 visualizes this procedure for dolichopodid flies and
429 butterflies. The first group failed to pass, while the latter fulfilled the criteria (Maes *et al.*,
430 2012).

431 Even in data-rich regions or countries, the estimated trends and geographic ranges, as well
432 as the Red List categories are subject to a degree of uncertainty (Akçakaya *et al.*, 2000). To
433 inform users of Red Lists about this, the IUCN Red List Categories and Criteria (IUCN, 2013)
434 suggests the inclusion of metadata about this uncertainty, including a range of plausible values
435 for the Red List assessment. These will be affected by how well a species has been surveyed in
436 time and space. This approach adds transparency to the Red Listing process, and helps defining
437 the Data Deficient category more objectively (e.g., when the range of uncertainty ranges from
438 Least Concern to Critically Endangered).

439 On larger scales (e.g., world, continental, European Union), it would be biologically more
440 meaningful to make Red Lists per ecological and/or biogeographical regions as, for example,

441 for the global biodiversity hotspot of the Mediterranean region (Myers *et al.*, 2000). In this
442 region, such lists have been compiled for mammals (Temple & Cuttelod, 2009), dragonflies
443 (Riservato *et al.*, 2009), freshwater fishes (Smith & Darwall, 2006), cartilaginous fishes
444 (Cavanagh & Gibson, 2007) and amphibians and reptiles (Cox, Chanson & Stuart, 2006). On the
445 other hand, conservation planning is usually the responsibility of national governments, which
446 makes biogeographical Red Lists difficult to apply in the field.

447 Due to differences in scale requirements and longevity among species (e.g., short-lived
448 invertebrates versus long-lived vertebrates or trees), but also because of differences in data
449 availability, some have argued that IUCN criteria should be differentiated for taxonomic groups
450 (e.g., invertebrates – Cardoso *et al.*, 2011; Cardoso *et al.*, 2012) and/or for spatial scales (Brito
451 *et al.*, 2010). Some countries continue to use national Red List criteria and categories instead
452 of those of the IUCN criteria because they judge them unusable in smaller regions (e.g., the
453 Netherlands – de longh & Bal, 2007). If applied correctly and even with the use of
454 opportunistic and/or data, we are convinced that the present-day IUCN criteria can be applied
455 to a wide variety of taxa, including invertebrates (Collen & Böhm, 2012) and at many different
456 spatial scales (from global to regional). The key point is that such data should be scrutinised
457 and not used blindly. IUCN Red Lists are useful to countries or regions since they need to
458 understand and track the fate of species within their borders. Legislation such as the
459 Convention on Biological Diversity encourages countries to do this at a national level (Zamin *et*
460 *al.*, 2010). For example, should Britain care about a butterfly species that is at the edge of its
461 northern range in a restricted area within the south of the region? From a global or continental
462 extinction risk perspective, probably not. The vast population in the rest of mainland Europe
463 means that the potential loss of the species in Britain is no threat to its overall survival. Since
464 the butterfly is part of Britain's biodiversity and is considered nationally threatened, however,
465 it should be protected and conserved. This clearly demonstrates the difference between a Red
466 List which 'only' estimates the extinction risk of a given species in a focal region on the one
467 hand and a national or regional list of conservation priorities on the other (Lamoreux *et al.*,

468 2003). Red Lists should, therefore, be considered as decision *support* tools and not as decision
469 *making* tools (Possingham *et al.*, 2002).

470 To conclude, we give some recommendations that may help to apply IUCN criteria more
471 uniformly across taxa and across regions from an organisational point of view but also for
472 peers that compile Red List in other parts of the world. Documenting a Red List assessment is
473 of vital importance to understand trend analyses and geographic range size estimates.
474 Therefore, it is important to document spatial and temporal mapping intensity in the focal
475 region, to give detailed information on how trends, distribution ranges and population sizes
476 were calculated and which assumptions were made in the analyses. Important organisational
477 aspects that can improve Red List assessments are, among others, the assignment of a Red List
478 co-ordinator in a region to have consistency among Red Lists of different taxonomic groups
479 (e.g., BRC in Britain, the Research Institute for Nature and Forest (INBO) in Flanders), the
480 availability of the dataset used for the Red List assessment for peers (open access data, e.g.,
481 GBIF, National Red List database; www.nationalredlist.org), and the motivation and
482 documentation of expert-judgement when using subcriteria such as fragmentation,
483 fluctuations and rescue effects or for the estimation of population sizes.

484

485

ACKNOWLEDGEMENTS

486 We are indebted to all the many volunteers who collect data on the occurrence of species and
487 made them available for research and conservation. The Biological Records Centre is a
488 partnership of the Centre for Ecology & Hydrology (CEH) and the Joint Nature Conservation
489 Committee. CH, NI and DR were funded by the CEH National Capability funding from the
490 Natural Environmental Research Council (Project NEC04932). We thank Marc Pollet for making
491 the dolichopodid flies data available for checking the procedure for Red List assessments in
492 Flanders. We would also like to thank Pete Stroh, Marc Pollet and an anonymous reviewer for
493 very useful comments on a previous version of the manuscript.

494

- 496 **Adriaens T, San Martin y Gomez G, Bogaert J, Crevecoeur L, Beuckx JP, Maes D. 2015.** Testing
 497 the applicability of regional IUCN Red List criteria on ladybirds (Coleoptera,
 498 Coccinellidae) in Flanders (north Belgium): opportunities for conservation. *Insect*
 499 *Conservation and Diversity* in press.
- 500 **Ainsworth AM, Smith JH, Boddy L, Dentinger BTM, Jordan M, Parfitt D, Rogers HJ, Skeates SJ.**
 501 **2013.** *Red List of Fungi for Great Britain: Boletaceae; A pilot conservation assessment*
 502 *based on national database records, fruit body morphology and DNA barcoding.* Joint
 503 Nature Conservation Committee: Peterborough.
- 504 **Akçakaya HR, Ferson S, Burgman MA, Keith DA, Mace GM, Todd CR. 2000.** Making consistent
 505 IUCN classifications under uncertainty. *Conservation Biology* **14**: 1001-1013.
- 506 **Baillie SR. 1990.** Integrated-Population Monitoring of Breeding Birds in Britain and Ireland. *Ibis*
 507 **132**: 151-166.
- 508 **Balmer DE, Gillings S, Caffrey BJ, Swann RL, Downie IS, Fuller RJ. 2013.** *Bird Atlas 2007-11:*
 509 *The Breeding and Wintering Birds of Britain and Ireland.* BTO Books: Thetford.
- 510 **Barwell LJ, Azaele S, Kunin WE, Isaac NJB. 2014.** Can coarse-grain patterns in insect atlas data
 511 predict local occupancy? *Diversity and Distributions* **20**: 895-907.
- 512 **Beck J, Böller M, Erhardt A, Schwanghart W. 2014.** Spatial bias in the GBIF database and its
 513 effect on modeling species' geographic distributions. *Ecological Informatics* **19**: 10-15.
- 514 **Bilz M, Kell SP, Maxted N, Lansdown RV. 2011.** *European Red List of vascular plants.*
 515 Publications Office of the European Union: Luxembourg.
- 516 **Boitani L, Maiorano L, Baisero D, Falcucci A, Visconti P, Rondinini C. 2011.** What spatial data
 517 do we need to develop global mammal conservation strategies? *Philosophical*
 518 *Transactions of the Royal Society of London. Series B, Biological Sciences* **366**: 2623–
 519 2632.
- 520 **Botham MS, Brereton TM, Middlebrook I, Randle Z, Roy DB. 2013.** *United Kingdom Butterfly*
 521 *Monitoring Scheme report for 2012.* Centre for Ecology & Hydrology: Wallingford.
- 522 **Brereton T, Roy DB, Middlebrook I, Botham M, Warren M. 2011.** The development of
 523 butterfly indicators in the United Kingdom and assessments in 2010. *Journal of Insect*
 524 *Conservation* **15**: 139-151.
- 525 **Brito D, Ambal RG, Brooks T, De Silva N, Foster M, Hao W, Hilton-Taylor C, Paglia A,**
 526 **Rodríguez JP, Rodríguez JV. 2010.** How similar are national red lists and the IUCN Red
 527 List? *Biological Conservation* **143**: 1154-1158.
- 528 **Burgmann MA, Fox JC. 2003.** Bias in species range estimates from minimum convex polygons:
 529 implications for conservation and options for improved planning. *Animal Conservation*
 530 **6**: 19–28.
- 531 **Butchart SHM, Akçakaya HR, Kennedy E, Hilton-Taylor C. 2006.** Biodiversity Indicators Based
 532 on Trends in Conservation Status: Strengths of the IUCN Red List Index. *Conservation*
 533 *Biology* **20**: 579-581.
- 534 **Cardoso P, Borges PAV, Triantis KA, Ferrández MA, Martín JL. 2011.** Adapting the IUCN Red
 535 List criteria for invertebrates. *Biological Conservation* **144**: 2432-2440.
- 536 **Cardoso P, Borges PAV, Triantis KA, Ferrández MA, Martín JL. 2012.** The underrepresentation
 537 and misinterpretation of invertebrates in the IUCN Red List. *Biological Conservation*
 538 **149**: 147-148.
- 539 **Carvalho LG, Kunin WE, Keil P, Aguirre-Gutiérrez J, Ellis WN, Fox R, Groom QJ, Hennekens**
 540 **SM, Van Landuyt W, Maes D, Van de Meutter F, Michez D, Rasmont P, Odé B, Potts**
 541 **SG, Reemer M, Masson Roberts SP, Schaminée JHJ, WallisdeVries MF, Biesmeijer JC.**
 542 **2013.** Biodiversity declines and biotic homogenization have slowed for NW Europe
 543 pollinators and plants. *Ecology Letters* **16**: 870-878.
- 544 **Cassini MH. 2011.** Ranking threats using species distribution models in the IUCN Red List
 545 assessment process. *Biodiversity and Conservation* **20**: 3689-3692.
- 546 **Caughley G. 1994.** Directions in conservation biology. *Journal of Animal Ecology* **63**: 215-244.

- 547 **Cavanagh RD, Gibson C. 2007.** *Overview of the Conservation Status of Cartilaginous Fishes*
548 *(Chondrichthyans) in the Mediterranean Sea: Gland, Switzerland and Malaga, Spain.*
- 549 **Cheffings CM, Farrell LE, Dines TD, Jones RA, Leach SJ, McKean DR, Pearman DA, Preston CD,**
550 **Rumsey FJ, Taylor I. 2005.** *The Vascular Plant Red Data List for Great Britain.* Joint
551 Nature Conservation Committee: Peterborough.
- 552 **Collen B, Böhm M. 2012.** The growing availability of invertebrate extinction risk assessments -
553 A response to Cardoso et al. (October 2012): Adapting the IUCN Red List criteria to
554 invertebrates. *Biological Conservation* **149**: 145-146.
- 555 **Collen B, Griffiths J, Friedmann Y, Rodriguez JP, Rojas-Suàrez F, Baillie JEM. 2013.** Tracking
556 Change in National-Level Conservation Status: National Red Lists *Biodiversity*
557 *Monitoring and Conservation: Wiley-Blackwell.* 17-44.
- 558 **Couvreur M, Menschaert J, Sevenant M, Ronse A, Van Landuyt W, De Blust G, Antrop M,**
559 **Hermy H. 2004.** Ecodistricten en ecoregio's als instrument voor natuurstudie en
560 milieubeleid. *Natuur.focus* **3**: 51-58.
- 561 **Cox NA, Chanson JS, Stuart SN. 2006.** *The Status and Distribution of Reptiles and Amphibians*
562 *of the Mediterranean Basin.* IUCN: Gland, Switzerland and Cambridge, UK.
- 563 **Cox NA, Temple HJ. 2009.** *European Red List of reptiles.* Publications Office of the European
564 Union: Luxembourg.
- 565 **Cuttelod A, Seddon M, Neubert E. 2011.** *European Red List of non-marine molluscs.*
566 Publications Office of the European Union: Luxembourg.
- 567 **Daguet CA, French GC, Taylor P. 2008.** *The Odonata Red Data List for Great Britain.* Joint
568 Nature Conservation Committee: Peterborough.
- 569 **de longh HH, Bal D. 2007.** Harmonization of Red Lists in Europe: some lessons learned in the
570 Netherlands when applying the new IUCN Red List Categories and Criteria version 3.1.
571 *Endangered Species Research* **3**: 53-60.
- 572 **Dennis RLH, Shreeve TG, Isaac NB, Roy DB, Hardy PB, Fox R, Asher J. 2006.** The effects of
573 visual apparency on bias in butterfly recording and monitoring. *Biological Conservation*
574 **128**: 486-492.
- 575 **Dennis RLH, Sparks TH, Hardy PB. 1999.** Bias in butterfly distribution maps: the effects of
576 sampling effort. *Journal of Insect Conservation* **3**: 33-42.
- 577 **Devictor V, Whittaker RJ, Beltrame C. 2010.** Beyond scarcity: citizen science programmes as
578 useful tools for conservation biogeography. *Diversity and Distributions* **16**: 354-362.
- 579 **Eaton MA, Gregory RD, Noble DG, Robinson JA, Hughes J, Procter D, Brown AF, Gibbons DW.**
580 **2005.** Regional IUCN red listing: the process as applied to birds in the United Kingdom.
581 *Conservation Biology* **19**: 1557-1570.
- 582 **Falk SJ, Chandler PJ. 2005.** *A review of the scarce and threatened flies of Great Britain. Part 2:*
583 *Nematocera and Aschiza not dealt with by Falk (1991).* Joint Nature Conservation
584 Committee: Peterborough.
- 585 **Falk SJ, Crossley R. 2005.** *A review of the scarce and threatened flies of Great Britain. Part 3:*
586 *Empidoidea.* Joint Nature Conservation Committee: Peterborough.
- 587 **Fitzpatrick U, Murray TE, Paxton RJ, Brown MJF. 2007.** Building on IUCN regional red lists to
588 produce lists of species of conservation priority: a model with Irish bees. *Conservation*
589 *Biology* **21**: 1324-1332.
- 590 **Foster GN. 2010.** *A review of the scarce and threatened Coleoptera of Great Britain Part (3):*
591 *Water beetles of Great Britain.* Joint Nature Conservation Committee: Peterborough.
- 592 **Fox R, Warren MS, Brereton TM. 2010.** *A new Red List of British Butterflies.* Joint Nature
593 Conservation Committee: Peterborough.
- 594 **Fox R, Warren MS, Brereton TM, Roy DB, Robinson A. 2011.** A new Red List of British
595 butterflies. *Insect Conservation and Diversity* **4**: 159-172.
- 596 **Freyhof J, Brooks E. 2011.** *European Red List of freshwater fishes.* Publications Office of the
597 European Union: Luxembourg.
- 598 **Gärdenfors U, Hilton-Taylor C, Mace GM, Rodríguez JP. 2001.** The application of IUCN Red List
599 criteria at regional levels. *Conservation Biology* **15**: 1206-1212.

- 600 **Gaston KJ, Fuller RA. 2009.** The sizes of species' geographic ranges. *Journal of Applied Ecology*
601 **46:** 1-9.
- 602 **Getz WM, Fortmann-Roe S, Cross PC, Lyons AJ, Ryan SJ, Wilmers CC. 2007.** LoCoH:
603 Nonparametric Kernel Methods for Constructing Home Ranges and Utilization
604 Distributions. *Plos One* **2:** e207.
- 605 **Getz WM, Wilmers CC. 2004.** A local nearest-neighbor convex-hull construction of home
606 ranges and utilization distributions. *Ecography* **27:** 489-505.
- 607 **Graham CH, Hijmans RJ. 2006.** A comparison of methods for mapping species ranges and
608 species richness. *Global Ecology and Biogeography* **15:** 578-587.
- 609 **Guisan A, Tingley R, Baumgartner JB, Naujokaitis-Lewis I, Sutcliffe PR, Tulloch AIT, Regan TJ,
610 Brotons L, McDonald-Madden E, Mantyka-Pringle C, Martin TG, Rhodes JR, Maggini
611 R, Setterfield SA, Elith J, Schwartz MW, Wintle BA, Broennimann O, Austin M, Ferrier
612 S, Kearney MR, Possingham HP, Buckley YM. 2013.** Predicting species distributions for
613 conservation decisions. *Ecology Letters* **16:** 1424-1435.
- 614 **Harris SJ, Risely K, Massimino D, Newson SE, Eaton MA, Musgrove AJ, Noble DG, Procter D,
615 Baillie SR. 2014.** *The Breeding Bird Survey 2013*. British Trust for Ornithology: Thetford.
- 616 **Hermoso V, Kennard MJ, Linke S. 2014.** Evaluating the costs and benefits of systematic data
617 acquisition for conservation assessments. *Ecography* in press.
- 618 **Hill MO. 2012.** Local frequency as a key to interpreting species occurrence data when
619 recording effort is not known. *Methods in Ecology and Evolution* **3:** 195-205.
- 620 **Isaac NJB, Pocock MJ. 2015.** Bias and information in biological records. *Biological Journal of*
621 *the Linnean Society* **115:** 522-531.
- 622 **Isaac NJB, van Strien AJ, August TA, de Zeeuw MP, Roy DB. 2014.** Statistics for citizen science:
623 extracting signals of change from noisy ecological data. *Methods in Ecology and*
624 *Evolution* **5:** 1052-1060.
- 625 **IUCN. 1972.** *Red data book. I. Mammals*. IUCN: Cambridge.
- 626 **IUCN. 1977.** *Red data book. IV. Fish*. IUCN: Cambridge.
- 627 **IUCN. 1978.** *Red data book. II. Birds*. IUCN: Cambridge.
- 628 **IUCN. 1979.** *Red data book. III. Amphibia and reptiles*. IUCN: Cambridge.
- 629 **IUCN. 1983.** *The IUCN invertebrate red data book*. IUCN: Gland.
- 630 **IUCN. 2003.** *Guidelines for Application of IUCN Red List Criteria at Regional Levels: Version 3.0*.
631 IUCN Species Survival Commission, IUCN: Gland, Switzerland and Cambridge, UK.
- 632 **IUCN. 2013.** *Guidelines for Using the IUCN Red List Categories and Criteria. Version 10*.
633 *Prepared by the Standards and Petitions Subcommittee*. IUCN: Gland, Switzerland and
634 Cambridge, UK.
- 635 **Jiguet F, Devictor V, Julliard R, Couvet D. 2012.** French citizens monitoring ordinary birds
636 provide tools for conservation and ecological sciences. *Acta Oecologica-International*
637 *Journal of Ecology* **44:** 58-66.
- 638 **Jooris R, Engelen P, Speybroeck J, Lewylle I, Louette G, Bauwens D, Maes D. 2012.** *De IUCN*
639 *Rode Lijst van de amfibieën en reptielen in Vlaanderen*. Instituut voor Natuur- en
640 Bosonderzoek: Brussel.
- 641 **Juslén A, Hyvärinen E, Virtanen LK. 2013.** Application of the Red-List Index at a National Level
642 for Multiple Species Groups. *Conservation Biology* **27:** 398-406.
- 643 **Kalkman VJ, Boudot JP, Bernard R, Conze KJ, De Knijf G, Dyatlova E, Ferreria S, Jovic M, Ott J,
644 Riservato E, Sahl G. 2010.** *European Red List of Dragonflies*. Publications Office of the
645 European Union: Luxembourg.
- 646 **Kearney MR. 2006.** Habitat, environment and niche: what are we modelling? *Oikos* **115:** 186-
647 191.
- 648 **Kearney MR, Shamakhy A, Tingley R, Karoly DJ, Hoffmann AA, Briggs PR, Porter WP. 2014.**
649 Microclimate modelling at macro scales: a test of a general microclimate model
650 integrated with gridded continental-scale soil and weather data. *Methods in Ecology*
651 *and Evolution* **5:** 273-286.
- 652 **Keller V, Bollmann K. 2004.** From red lists to species of conservation concern. *Conservation*
653 *Biology* **18:** 1636-1644.

- 654 **Keller V, Zbinden N, Schmid H, Volet B. 2005.** A case study in applying the IUCN regional
655 guidelines for national red lists and justifications for their modification. *Conservation*
656 *Biology* **19**: 1827-1834.
- 657 **Lamoreux J, Akçakaya HR, Bennun L, Collar NJ, Boitani L, Brackett D, Brautigam A, Brooks**
658 **TM, de Fonseca GAB, Mittermeier RA, Rylands AB, Gärdenfors U, Hilton-Taylor C,**
659 **Mace G, Stein BA, Stuart S. 2003.** Value of the IUCN Red List. *Trends in Ecology &*
660 *Evolution* **18**: 214-215.
- 661 **Lock K, Stoffelen E, Vercouteren T, Bosmans R, Adriaens T. 2013.** Updated Red List of the
662 water bugs of Flanders (Belgium) (Hemiptera : Gerromorpha & Nepomorpha). *Bulletin*
663 *de la Société royale belge d'Entomologie/Bulletin van de Koninklijke Belgische*
664 *Vereniging voor Entomologie* **149**: 57-63.
- 665 **Lucas G, Syngé H. 1978.** *The IUCN Plant Red Data Book*. IUCN: Morges.
- 666 **Mace GM. 1994.** Classifying threatend species: means and ends. *Philosophical Transactions of*
667 *the Royal Society of London, B* **344**: 91-97.
- 668 **Mace GM, Collar NJ, Cooke J, Gaston KJ, Ginsberg J, Leader Williams N, Maunder M, Milner-**
669 **Gulland EJ. 1993.** The development of new criteria for listing species on the IUCN Red
670 List. *Species* **19**: 16-22.
- 671 **Mace GM, Collar NJ, Gaston KJ, Hilton-Taylor C, Akçakaya HR, Leader-Williams N, Milner-**
672 **Gulland EJ, Stuart SN. 2008.** Quantification of Extinction Risk: IUCN's System for
673 Classifying Threatened Species. *Conservation Biology* **22**: 1424-1442.
- 674 **Mace GM, Lande R. 1991.** Assessing extinction threats: toward a re-evaluation of IUCN
675 threatened species categories. *Conservation Biology* **5**: 148-157.
- 676 **MacKenzie DI, Nichols JD, Royle JA, Pollock KH, Hines JE, Bailey LL. 2006.** *Occupancy*
677 *estimation and modeling: Inferring patterns and dynamics of species occurrence*.
678 Elsevier: San Diego.
- 679 **Maes D, Baert K, Boers K, Casaer J, Crevecoeur L, Criel D, Dekeukeleire D, Gouwy J, Gyselings**
680 **R, Haelters J, Herman D, Herremans M, Lefebvre, J., Lefevre A, Onkelinx T, Stuyck J,**
681 **Thomaes A, Van Den Berge K, Vandendriessche B, Verbeylen G, Vercayie D. 2014.** *De*
682 *IUCN Rode Lijst van de zoogdieren in Vlaanderen*. Instituut voor Natuur- en
683 Bosonderzoek: Brussel.
- 684 **Maes D, Van Dyck H. 2001.** Butterfly diversity loss in Flanders (north Belgium): Europe's worst
685 case scenario? *Biological Conservation* **99**: 263-276.
- 686 **Maes D, van Swaay CAM. 1997.** A new methodology for compiling national Red Lists applied
687 on butterflies (Lepidoptera, Rhopalocera) in Flanders (N.-Belgium) and in The
688 Netherlands. *Journal of Insect Conservation* **1**: 113-124.
- 689 **Maes D, Vanreusel W, Jacobs I, Berwaerts K, Van Dyck H. 2012.** Applying IUCN Red List
690 criteria at a small regional level: A test case with butterflies in Flanders (north
691 Belgium). *Biological Conservation* **145**: 258-266.
- 692 **Miller RM, Rodríguez JP, Aniskowicz-Fowler T, Bambaradeniya C, Boles R, Eaton MA,**
693 **Gärdenfors U, Keller V, Molur S, Walker S, Pollock C. 2007.** National threatened
694 species listing based on IUCN criteria and regional guidelines: Current status and future
695 perspectives. *Conservation Biology* **21**: 684-696.
- 696 **Myers N, Mittermeier RA, Mittermeier CG, daFonseca GAB, Kent J. 2000.** Biodiversity
697 hotspots for conservation priorities. *Nature* **403**: 853-858.
- 698 **Nature Editorials. 2009.** A public service. The Christmas bird count is a model to be emulated
699 in distributed, volunteer science. *Nature* **457**: 8.
- 700 **Nieto A, Alexander KNA. 2010.** *European Red List of saproxylic beetles*. Publications Office of
701 the European Union: Luxembourg.
- 702 **Pagel J, Anderson BJ, O'Hara RB, Cramer W, Fox R, Jeltsch F, Roy DB, Thomas CD, Schurr FM.**
703 **2014.** Quantifying range-wide variation in population trends from local abundance
704 surveys and widespread opportunistic occurrence records. *Methods in Ecology and*
705 *Evolution* **5**: 751-760.
- 706 **Pannekoek J, van Strien A. 2003.** *TRIM 3 Manual. Trends and indices for monitoring data*.
707 Centraal Bureau voor de Statistiek: Voorburg.


- 708 **Panzacchi M, Van Moorter B, Strand O, Egil Loe L, Reimers E. 2015.** Searching for the
709 fundamental niche using individual-based habitat selection modelling across
710 populations *Ecography* **in press**.
- 711 **Pena JCD, Kamino LHY, Rodrigues M, Mariano-Neto E, de Siqueira MF. 2014.** Assessing the
712 conservation status of species with limited available data and disjunct distribution.
713 *Biological Conservation* **170**: 130-136.
- 714 **Pescott OL, Walker KJ, Pocock MJ, Jitlal M, Outhwaite CW, Cheffings CM, Harris F, Roy DB.**
715 **2015.** Ecological monitoring for citizen science: the history, design and implementation
716 of schemes for plants in Britain and Ireland. *Biological Journal of the Linnean Society*
717 **115**: 505-521.
- 718 **Pocock MJ, Roy HE, Preston CD, Roy DB. 2015.** The Biological Records Centre in the United
719 Kingdom: a pioneer of citizen science. *Biological Journal of the Linnean Society* **115**:
720 475-493.
- 721 **Possingham HP, Andelman SJ, Burgman MA, Medellín RA, Master LL, Keith DA. 2002.** Limits to
722 the use of threatened species lists. *Trends in Ecology & Evolution* **17**: 503-507.
- 723 **Powney GD, Isaac NJB. 2015.** Beyond maps: a review of the applications of biological records.
724 *Biological Journal of the Linnean Society* **115**: 535-542.
- 725 **Prendergast JR, Wood SN, Lawton JH, Eversham BC. 1993.** Correcting for variation in
726 recording effort in analyses of diversity hotspots. *Biodiversity Letters* **1**: 39-53.
- 727 **Riservato E, Boudot JP, Ferreira S, Jović M, Kalkman VJ, Schneider W, Samraoui B, Cuttelod**
728 **A. 2009.** *The Status and Distribution of Dragonflies of the Mediterranean Basin*: Gland,
729 Switzerland and Malaga, Spain.
- 730 **Rodrigues ASL, Pilgrim JD, Lamoreux JF, Hoffmann M, Brooks TM. 2006.** The value of the
731 IUCN Red List for conservation. *Trends in Ecology & Evolution* **21**: 71-76.
- 732 **Rodríguez JP. 2008.** National Red Lists: the largest global market for IUCN Red List Categories
733 and Criteria. *Endangered Species Research* **6**: 193-198.
- 734 **Rondinini C, Wilson KA, Boitani L, Grantham H, Possingham HP. 2006.** Tradeoffs of different
735 types of species occurrence data for use in systematic conservation planning. *Ecology*
736 *Letters* **9**: 1136-1145.
- 737 **Rothery P, Roy DB. 2001.** Application of generalized additive models to butterfly transect
738 count data. *Journal of Applied Statistics* **28**: 897-909.
- 739 **Roy DB, Ploquin EF, Randle Z, Risely K, Botham MS, Middlebrook I, Noble D, Cruickshanks K,**
740 **Freeman SN, Brereton TM. 2015a.** Comparison of trends in butterfly populations
741 between monitoring schemes. *Journal of Insect Conservation* **in press**.
- 742 **Roy DB, Rothery P, Brereton T. 2007.** Reduced-effort schemes for monitoring butterfly
743 populations. *Journal of Applied Ecology* **44**: 993-1000.
- 744 **Roy HE, Adriaens T, Isaac NB, Kenis M, San Martín y Gómez G, Onkelinx T, Brown PMJ,**
745 **Poland R, Ravn HP, Roy DB, Comont R, Grégoire JC, de Biseau JC, Hautier L, Eschen R,**
746 **Frost R, Zindel R, Van Vlaenderen J, Nedved O, Maes D. 2012a.** Invasive alien
747 predator causes rapid declines of native European ladybirds. *Diversity and*
748 *Distributions* **18**: 717-725.
- 749 **Roy HE, Pocock MJO, Preston CD, Roy DB, Savage J, Tweddle JC, Robinson LD. 2012b.**
750 *Understanding citizen science and environmental monitoring: Final report on behalf of*
751 *UK-EOF*. NERC Centre for Ecology & Hydrology and Natural History Museum:
752 Wallingford.
- 753 **Roy HE, Rorke SL, Beckmann B, Booy O, Botham MS, Brown PMJ, Harrower C, Noble D,**
754 **Sewell J, Walker KJ. 2015b.** The contribution of volunteer recorders to our
755 understanding of biological invasions. *Biological Journal of the Linnean Society* **115**:
756 678-689.
- 757 **Schmeller DS, Henry PY, Julliard R, Gruber B, Clobert J, Dziock F, Lengyel S, Nowicki P, Deri E,**
758 **Budrys E, Kull T, Tali K, Bauch B, Settele J, Van Swaay C, Kobler A, Babij V,**
759 **Papastergiadou E, Henle K. 2009.** Advantages of Volunteer-Based Biodiversity
760 Monitoring in Europe. *Conservation Biology* **23**: 307-316.

- 761 **Scott P, Burton JA, Fitter R. 1987.** Red Data Books: the historical background. In: Fitter R and
762 Fitter M, eds. *The road to extinction*. Gland: IUCN. 1-5.
- 763 **Sheth SN, Lohmann LG, Distler T, Jimenez I. 2012.** Understanding bias in geographic range size
764 estimates. *Global Ecology and Biogeography* **21**: 732-742.
- 765 **Simaika JP, Samways MJ. 2009.** Reserve selection using Red Listed taxa in three global
766 biodiversity hotspots: Dragonflies in South Africa. *Biological Conservation* **142**: 638-
767 651.
- 768 **Smith KG, Darwall WRT. 2006.** *The Status and Distribution of Freshwater Fish Endemic to the*
769 *Mediterranean Basin*. IUCN: Gland, Switzerland and Cambridge, UK.
- 770 **Soberón J, Jiménez R, Golubov J, Koleff P. 2007.** Assessing completeness of biodiversity
771 databases at different spatial scales. *Ecography* **30**: 152-160.
- 772 **Stojanovic DV, Curcic SB, Curcic BPM, Makarov SE. 2013.** The application of IUCN Red List
773 criteria to assess the conservation status of moths at the regional level: a case of
774 provisional Red List of Noctuidae (Lepidoptera) in Serbia. *Journal of Insect*
775 *Conservation* **17**: 451-464.
- 776 **Strayer DL. 2010.** Alien species in fresh waters: ecological effects, interactions with other
777 stressors, and prospects for the future. *Freshwater Biology* **55**: 152-174.
- 778 **Stroh PA, Leach SJ, August TA, Walker KJ, Pearman DA, Rumsey FJ, Harrower CA, Fay MF,**
779 **Martin JP, Pankhurst T, Preston CD, Taylor I. 2014.** *A Vascular Plant Red List for*
780 *England*. Botanical Society of Britain and Ireland. : Bristol.
- 781 **Syfert MM, Joppa L, Smith MJ, Coomes DA, Bachman SP, Brummitt NA. 2014.** Using species
782 distribution models to inform IUCN Red List assessments. *Biological Conservation* **177**:
783 174-184.
- 784 **Temple HJ, Cox NA. 2009.** *European Red List of amphibians*. Publications Office of the
785 European Union: Luxembourg.
- 786 **Temple HJ, Cuttelod A. 2009.** *The Status and Distribution of Mediterranean Mammals*. IUCN:
787 Gland, Switzerland and Cambridge, UK
- 788 **Temple HJ, Terry A. 2007.** *The status and distribution of European mammals*. Office for Official
789 Publications of the European Communities: Luxembourg.
- 790 **Thomaes A, Kervyn T, Maes D. 2008.** Applying species distribution modelling for the
791 conservation of the threatened saproxylic Stag Beetle (*Lucanus cervus*). *Biological*
792 *Conservation* **141**: 1400-1410.
- 793 **Thomaes A, Maes D. 2014.** *Rode-Lijststatus van het Vliegend hert (Lucanus cervus)*. Instituut
794 voor Natuur- en Bosonderzoek: Geraardsbergen.
- 795 **Thomas JA. 2005.** Monitoring change in the abundance and distribution of insects using
796 butterflies and other indicator groups. *Philosophical Transactions of the Royal Society*
797 *of London B* **360**: 339-357.
- 798 **Thomas JA, Telfer MG, Roy DB, Preston CD, Fox R, Clarke RT, Lawton JH. 2004.** Comparative
799 losses in British butterflies, birds, and plants and the global extinction crisis. *Science*
800 **303**: 1879-1881.
- 801 **Tomlinson S, Arnall SG, Munn A, Bradshaw SD, Maloney SK, Dixon KW, Didham RK. 2014.**
802 Applications and implications of ecological energetics. *Trends in Ecology & Evolution*
803 **29**: 280-290.
- 804 **Van Dyck H, van Strien AJ, Maes D, van Swaay CAM. 2009.** Declines in common, widespread
805 butterflies in a landscape under intense human use. *Conservation Biology* **23**: 957-965.
- 806 **van Strien AJ, Termaat T, Groenendijk D, Mensing V, Kéry M. 2010.** Site-occupancy models
807 may offer new opportunities for dragonfly monitoring based on daily species lists.
808 *Basic and Applied Ecology* **11**: 495-503.
- 809 **van Strien AJ, van Swaay CAM, Termaat T. 2013.** Opportunistic citizen science data of animal
810 species produce reliable estimates of distribution trends if analysed with occupancy
811 models. *Journal of Applied Ecology* **50**: 1450-1458.
- 812 **van Swaay CAM. 1990.** An assessment of the changes in butterfly abundance in the
813 Netherlands during the 20th century. *Biological Conservation* **52**: 287-302.


- 814 **van Swaay CAM, Cuttelod A, Collins S, Maes D, Munguira ML, Šašić M, Settele J, Verovnik R,**
815 **Verstrael T, Warren MS, Wiemers M, Wynhoff I. 2010.** *European Red List of*
816 *Butterflies*. Publications Office of the European Union: Luxembourg.
- 817 **van Swaay CAM, Nowicki P, Settele J, van Strien AJ. 2008.** Butterfly monitoring in Europe:
818 methods, applications and perspectives. *Biodiversity and Conservation* **17**: 3455-3469.
- 819 **van Swaay CAM, Veling K, Termaat T, Huskens K, Plate CL. 2013.** Vlinders en libellen geteld.
820 Jaarverslag 2012. Wageningen: De Vlinderstichting.
- 821 **Vermeersch G, Onkelinx T. 2014.** ABV-project: trends na de tweede volledige telcyclus.
822 *Vogelnieuws* **19**: 29-31.
- 823 **Verreycken H, Belpaire C, Van Thuyne G, Breine J, Buysse D, Coeck J, Mouton A, Stevens M,**
824 **Vandenneucker T, De Bruyn L, Maes D. 2014.** An IUCN Red List of lampreys and
825 freshwater fishes in Flanders (north Belgium). *Fisheries Management and Ecology* **21**:
826 122-132.
- 827 **Woods RG, Coppins BJ. 2012.** *A Conservation Evaluation of British Lichens and Lichenicolous*
828 *Fungi*. Joint Nature Conservation Committee: Peterborough.
- 829 **Zamin TJ, Baillie JEM, Miller RM, Rodriguez JP, Ardid A, Collen B. 2010.** National Red Listing
830 Beyond the 2010 Target. *Conservation Biology* **24**: 1012-1020.
- 831
- 832

833	Table 1. IUCN Red Lists in Europe, Britain and Flanders that were screened on the use of the different IUCN criteria.
834	
835	
836	Europe (ec.europa.eu/environment/nature/conservation/species/redlist/)
837	
838	Amphibians (Temple & Cox, 2009); Butterflies (van Swaay <i>et al.</i> , 2010); Dragonflies (Kalkman <i>et al.</i> ,
839	2010); Freshwater fishes (Freyhof & Brooks, 2011); Freshwater molluscs (Cuttelod, Seddon & Neubert,
840	2011); Mammals (Temple & Terry, 2007); Reptiles (Cox & Temple, 2009); Saproxyllic beetles (Nieto &
841	Alexander, 2010); Terrestrial molluscs (Cuttelod, Seddon & Neubert, 2011); Vascular plants, partim (Bilz
842	<i>et al.</i> , 2011)
843	
844	Britain (jncc.defra.gov.uk/page-3352)
845	
846	Boletes (Ainsworth <i>et al.</i> , 2013); Butterflies (Fox, Warren & Brereton, 2010); Dragonflies (Daguet, French
847	& Taylor, 2008); Flies (Falk & Crossley, 2005; Falk & Chandler, 2005); Lichens and lichenicolous fungi
848	(Woods & Coppins, 2012); Vascular plants (Cheffings <i>et al.</i> , 2005); Water beetles (Foster, 2010)
849	
850	Flanders (http://wwwl.inbo.be/nl/rode-lijsten-vlaanderen)
851	
852	Amphibians (Jooris <i>et al.</i> , 2012); Butterflies (Maes <i>et al.</i> , 2012); Freshwater fishes (Verreycken <i>et al.</i> ,
853	2014); Ladybirds (Adriaens <i>et al.</i> , 2015); Mammals (Maes <i>et al.</i> , 2014); Reptiles (Jooris <i>et al.</i> , 2012); Stag
854	beetle (Thomaes & Maes, 2014); Water bugs (Lock <i>et al.</i> , 2013)


855
856




857
858 **Figure 1.** IUCN categories at the regional level (IUCN, 2003).
859



860
861 **Figure 2.** Overall criterion use for species in Britain (total number of threatened species = 1569), Europe (n = 714) and
862 Flanders (n = 125). Criterion A = Population size reduction, Criterion B = Geographic range size, Criterion C = Small
863 population size and decline, Criterion D = Very small or restricted population, Criterion E = Quantitative analysis of
864 extinction risk.
865




866
867 **Figure 3.** Criterion use per taxonomic group in Britain (left), Europe (middle) and Flanders (right).
868




869
 870 **Figure 4.** Use of approaches in IUCN criterion A (population size reduction, left) and IUCN criterion B (geographic
 871 range size, right) in Red List assessments in Britain, Europe and Flanders. Criterion A: Aa = direct observation, Ab = an
 872 index of abundance appropriate to the taxon, Ac = a decline in AOO, EOO and/or habitat quality, Ad = actual or
 873 potential level of exploitation, Ae = effects of introduced taxa, hybridization, pathogens, pollutants, competitors or
 874 parasites; Criterion B: B1 = EOO, B2 = AOO, B1+B2 = EOO + AOO.
 875


876



877



878 **Figure 5.** Maps showing the EOO estimates for *Andrena bicolor* in the UK between 1996-2010 using a) observed 10 x
879 10 km² grid squares (total area = 46 100 km²), b) Minimum Convex Polygon (MCP – 324 850 km² for full MCP or 208
880 150 km² for intersection of MCP with land area) c) α -hull (101 895km²) with $\alpha = 40\ 000$ m and d) r-LoCoH (101 919
881 km²) with $r = 40\ 000$ m. These figures were produced for a Red Listing assessment of aculeate Hymenoptera in Great
882 Britain (Edwards *et al.*, in prep) using data collected by the Bees, Wasp & Ants Recording Scheme (BWARS).
883
884





885
886 **Figure 6.** AOO of the ladybird *Coccinella hieroglyphica* using the 36 ecological districts in Flanders (north Belgium) in
887 the period 2006-2013. The distribution of the species is shown using 1 x 1 km² grid cells (black dots). Only ecological
888 districts (in grey) in which the species was observed in at least three grid cells were incorporated in the estimate of the
889 AOO (i.e., 3 087 km² – Adriaens *et al.*, 2015).
890

891



Dolichopodid flies

Butterflies





892

a)




893

b)



894

c)



895

d)

896 **Figure 7.** Visualization of the procedure used in Flanders (north Belgium) to judge whether enough data are available
897 for a Red List assessment. As a background, the 12 ecological regions of Flanders are shown. a) all grid cells (5 x 5 km²)
898 surveyed in the first period for dolichopodid flies (left) and butterflies (right), b) all grid cells surveyed in the second
899 period, c) all grid cells surveyed in common in both periods, d) all grid cells in common in both periods that are
900 considered as sufficiently well surveyed (i.e., ≥ 10 species per grid cell in both periods).