

# Block correlation and the spatial resolution of soil property maps made by kriging.

R.M. Lark\*

*British Geological Survey, Keyworth, Nottinghamshire NG12 5GG, U.K.*

---

## Abstract

1 The block correlation is the correlation between the block kriging prediction of a variable  
2 and the true spatial mean which it estimates, computed for a particular sampling configu-  
3 ration and block size over the stochastic model which underlies the kriging prediction. This  
4 correlation can be computed if the variogram and disposition of sample points are known. It  
5 is also possible to compute the concordance correlation, a modified correlation which mea-  
6 sures the extent to which the block kriging prediction and true block spatial mean conform  
7 to the 1:1 line, and so is sensitive to the tendency of the kriging predictor to over-smooth.  
8 It is proposed that block concordance correlation has two particular advantages over krig-  
9 ing variance for communicating uncertainty in predicted values. First, as a measure on a  
10 bounded scale it is more intuitively understood by the non-specialist data user, particularly  
11 one who is interested in a synoptic overview of soil variation across a region. Second, because  
12 it accounts for the variability of the spatial means and their kriged estimates, as well as the  
13 uncertainty of the latter, it can be more readily compared between blocks of different size  
14 than can a kriging variance.

15 Using the block correlation and concordance correlation it is shown that the uncer-  
16 tainty of block kriged predictions depends on block size, but this effect depends on the  
17 interaction of the autocorrelation of the random variable and the sampling intensity. In  
18 some circumstances (where the dominant component of variation is at a long range relative  
19 to sample spacing) the block correlation and concordance correlation are insensitive to block  
20 size, but if the grid spacing is closer to the range of correlation of a significant component  
21 then block size can have a substantial effect on block correlation. It is proposed that (i)  
22 block concordance correlation is used to communicate the uncertainty in kriged predictions  
23 to a range of audiences (ii) that it is used to explore sensitivity to block size when plan-  
24 ning mapping and (iii) as a general operational rule a block size is selected to give a block  
25 concordance correlation of 0.8 or larger where this can be achieved without extra sampling.

26 *Keywords:* Kriging, Quality measures, Spatial resolution.

---

\**E-mail address:* [mlark@nerc.ac.uk](mailto:mlark@nerc.ac.uk) (R.M. Lark).

## 1. Introduction

In conventional soil survey the scale of the published map plays a tacit role in communicating to the user an understanding of the uncertainty in the information that it conveys. The intensity of field effort affects the uncertainty of predictions made from the resulting soil map in terms of soil classes and soil properties (Beckett and Burrough, 1971). Soil survey organizations have conventionally required that maps be supported by some fixed density of field observations on the published map (Burrough and Beckett, 1971). For example, soil maps in British Columbia should be supported by about 1 observation per  $\text{cm}^2$  of published map, with an acceptable range of 0.2–2 observations  $\text{cm}^{-2}$  (Resources Inventory Committee, 1995). In Australia it has been recommended that the density of observations is in the range 0.25–1  $\text{cm}^{-2}$  (Gunn et al, 1988). The larger the density of observations *on the ground* the larger the cartographic scale of the map which these observations can support. For this reason, the larger the scale ratio of a map the greater confidence a user can have in predicting likely soil conditions at a site. Maps published at different scales are therefore suitable for different purposes. Large scale maps (e.g. 1:10 000–1:25 000) are called ‘detailed’ maps and are recommended for agricultural extension and irrigation planning, whereas maps at 1:120 000 –1:500 000 are called ‘reconnaissance’ maps and are recommended for national-scale land use planning and tentative selection of project locations (Dent and Young, 1981).

The printed map with a particular cartographic scale ratio is increasingly superseded by the raster layer in a Geographical Information System (GIS), a digital soil map. As GIS technology emerged cartographers explored how their traditional questions about scale ratios should be addressed in this new setting. They made a link with the spatial resolution, expressed by the dimension of the raster cell or pixel (I assume square pixels in this paper). Tobler (1988) proposed a heuristic rule by which information that could be communicated effectively on a map of scale ratio no smaller than 1:s requires raster cells length  $r$  such that  $r \leq s/2000$  m. Hengl (2006) reviewed and proposed a similar set of rules, for example, if the

53 effective range of the variogram of a continuous variable is  $a$  units then this variable should  
54 be mapped on pixels of length  $r \leq a/2$ .

55 The question of the resolution of a raster soil map, here I focus on maps of individual  
56 soil properties, has received revived attention with the emergence of the GlobalSoilMap  
57 project (Hempel et al., 2014). The project specification (Arrouays et al., 2014a) requires  
58 that, in addition to a grid of predictions on a notional 2×2-m ‘pedon’ support (first tier  
59 information), the spatial mean of soil properties, to a specified depth, is predicted over a  
60 100×100-m block (second tier information).

61 As Arrouays et al. (2014b) acknowledge, this selection of a cell size was not un-  
62 controversial. In particular they note that the data user is likely make inferences about  
63 uncertainty from block size (resolution), and that a map of global extent on 100-m blocks,  
64 given available data, may seem implausibly ambitious. They observe, however, that in the  
65 context of digital soil mapping the uncertainty of an individual prediction, on a block of any  
66 size, can be quantified directly. Uncertainty, therefore, should not be tacitly communicated  
67 by scale or block size, and one is free to select a block size on the basis of other criteria such  
68 as the resolution of ancillary data to be used in the prediction.

69 This argument is correct, but it is necessary to reflect on how this approach adds to  
70 the challenge of successfully communicating the uncertainty attached to soil information to  
71 the user of this information. Three points are particularly germane.

72 First, if one presents a data user with a map with predictions made on 100-m blocks,  
73 it is reasonable for the user to infer that one is making a tacit claim to be able to identify  
74 differences between those blocks, and that features of variation that can be resolved should  
75 be assumed to be genuine.

76 Second, while it is true, as Heuvelink (2014) points out, that one can create sets of  
77 geostatistical maps in which precision and resolution are decoupled, other things being equal  
78 (the sampling grid, the covariates), decreasing the block size reduces the precision of kriged  
79 predictions as measured by their mean-square error. Heuvelink (2014) created his sets of

80 precise or imprecise and coarse- or fine-resolution maps by working from sparse or dense  
81 data sets. In practice one cannot reduce the grid cell size while maintaining the precision of  
82 predictions unless more data or covariates are used for prediction.

83 Third, one may state to the data user that the block size is selected on purely op-  
84 erational grounds, and that they should examine the corresponding uncertainty map to  
85 understand the claims made for the information. However, while uncertainty measures such  
86 as a kriging variance have considerable value, and are understood by statisticians and sci-  
87 entists, it is not clear that they always succeed in conveying to the general data user a clear  
88 understanding of uncertainty. Even if one changes the kriging variance to a standard error,  
89 so that it is on the scale of the original measurement, the user, particularly one who is  
90 interested in the overall variation across a region rather than decision making at one or a  
91 few locations, may be challenged to make a judgement as to which features of the mapped  
92 pattern can be interpreted with confidence and which cannot.

93 In this paper I propose a new measure of the uncertainty of block kriging predictions.  
94 This is the block correlation, the expected correlation of the block prediction with the value  
95 that it estimates: the spatial mean of the target variable across the block. The block  
96 concordance correlation (after Lin, 1989) can also be calculated, and may be more useful for  
97 communicating uncertainty of kriged predictions. This is because it measures not simply the  
98 linear correlation between variables but their conformity to the 1:1 line. In the case of the  
99 correlation between the block spatial mean and the kriging predictor systematic deviations  
100 from the 1:1 line are due to over-smoothing by kriging, as the procedure is unbiased.

101 The block correlation has two particular advantages over kriging variance, which make  
102 it pertinent to the questions above. First, it can be compared between blocks of different sizes  
103 more readily than the prediction error variance, because changing block size also changes  
104 the variance of the block means. This makes the block correlation conceptually useful  
105 for investigating the effects of block size on prediction uncertainty. Second, as a bounded  
106 and dimensionless quantity, correlations may be more readily interpreted by the general

107 user as a measure of information quality than a kriging variance, although this remains a  
 108 key uncertainty measure for quantitative assessments of uncertainty either through error  
 109 propagation or the comparison of prediction distributions to threshold values. For reasons  
 110 explained in the previous paragraph, the block concordance correlation is particularly useful  
 111 for communication. The user can understand that a concordance correlation of 1 implies  
 112 that the block prediction is perfect, and zero that the prediction includes no information,  
 113 with intermediate values implying some degree of deviation from the 1:1 line.

114 In this paper I develop the concept of the block correlation and concordance correla-  
 115 tion and show how they can be computed for block kriging. I then use them to examine the  
 116 extent to which, *ceteris paribus*, the block length for the kriged map serves as a proxy for  
 117 information quality. In particular I consider how the block concordance correlation might be  
 118 used to select both block size and sample density for a soil map where there is some freedom  
 119 to select a block size. In principle the block correlation could be fixed at some value as a  
 120 quality standard and block size adjusted so as to achieve this.

## 121 2. Theory

### 122 2.1. Block correlation

123 Supports are defined in two dimensions, assuming sampling to some fixed depth for  
 124 intensive properties such as concentrations. The region of interest is denoted by  $\mathcal{R} \in \mathbb{R}^2$ .  
 125 The variable of interest, measured at location  $\mathbf{x} \in \mathcal{R}$ , is denoted by  $z(\mathbf{x})$  and is treated  
 126 as a realization of a random variable  $Z(\mathbf{x})$ . I denote some block of support  $\mathcal{B}$  centred  
 127 at location  $\mathbf{x}$  by  $\mathbf{x}_{\mathcal{B}}$ ; so, for example, if  $\mathcal{B}$  is a square support of 200 m  $\times$  200 m then  
 128  $\mathbf{s} \in \mathbf{x}_{\mathcal{B}} \rightarrow \max_{i=1,2} |s_i - x_i| \leq 100$ . The quantity of interest is the spatial mean of  $z$  across  
 129  $\mathbf{x}_{\mathcal{B}}$ ,

$$\bar{z}(\mathbf{x}_{\mathcal{B}}) = \int_{\mathbf{s} \in \mathbf{x}_{\mathcal{B}}} z(\mathbf{s}) \, d\mathbf{s}. \quad (1)$$

130 The ordinary kriging estimate of this quantity is denoted by  $\tilde{Z}(\mathbf{x}_{\mathcal{B}})$ , and the ordinary

131 kriging variance,  $\sigma_K^2(\mathbf{x}_B)$  is obtained with the estimate where

$$\sigma_K^2(\mathbf{x}_B) = \text{E} \left[ \left\{ \tilde{Z}(\mathbf{x}_B) - \bar{z}(\mathbf{x}_B) \right\}^2 \right]. \quad (2)$$

132 By the block correlation we mean

$$\rho_B = \text{Corr} \left[ \tilde{Z}(\mathbf{x}_B), \bar{z}(\mathbf{x}_B) \right], \quad (3)$$

133 where  $\text{Corr}[\cdot, \cdot]$  denotes the correlation of the two variables in the brackets. Note that  
 134 this quantity is defined for a particular block, given the configuration of sampling points  
 135 from which the prediction is derived, over the random model which underlies the kriging  
 136 prediction. It is in this sense a superpopulation statistic, and a measure of confidence for  
 137 the particular prediction. It should not be confused with a population statistic, such as  
 138 the correlation of the spatial means of a set of distinct block with their respective kriging  
 139 predictions from a particular set of data. Bishop et al. (2015) in a recent study attempted  
 140 to estimate population validation statistics for spatial prediction of soil properties on blocks  
 141 of different size.

142 The block correlation can be computed directly from terms computed to solve the  
 143 ordinary block kriging equation. Because the ordinary kriging predictor is unbiased the  
 144 ordinary kriging variance is the variance of the difference between the ordinary kriging  
 145 predictor,  $\tilde{Z}(\mathbf{x}_B)$ , and the true spatial mean of the block which it predicts,  $\bar{z}(\mathbf{x}_B)$ , and  
 146 therefore

$$\text{Cov} \left[ \tilde{Z}(\mathbf{x}_B), \bar{z}(\mathbf{x}_B) \right] = \frac{\text{Var} \left[ \tilde{Z}(\mathbf{x}_B) \right] + \text{Var} \left[ \bar{z}(\mathbf{x}_B) \right] - \sigma_K^2(\mathbf{x}_B)}{2}, \quad (4)$$

147 where  $\text{Cov}[\cdot, \cdot]$  denotes the covariance of the two variables in the brackets and  $\text{Var}[\cdot]$  denotes  
 148 the variance of a variable.

149 The block kriging prediction is obtained from a set of  $N$  observations, and the covari-  
 150 ance matrix of these observations under the model used for kriging is denoted by  $\mathbf{C}$ , which  
 151 is a submatrix of a matrix in the ordinary kriging equation. If  $\boldsymbol{\lambda}$  is the vector of ordinary  
 152 block kriging weights then it follows from the familiar properties of linear combinations of

153 random variables that

$$\text{Var} \left[ \tilde{Z}(\mathbf{x}_B) \right] = \boldsymbol{\lambda}^T \mathbf{C} \boldsymbol{\lambda}. \quad (5)$$

154 We now require the variance of the block spatial mean. This can be obtained by  
 155 applying Krige's relation (Journel and Huijbregts, 1978) to give

$$\text{Var} [\bar{z}(\mathbf{x}_B)] = \text{Var}[Z] - \sigma_{\mathcal{D},\mathcal{B}}^2, \quad (6)$$

156 where the variance of  $Z$  is the a priori variance of the variable and  $\sigma_{\mathcal{D},\mathcal{B}}^2$  is the dispersion  
 157 variance of  $Z$  within the block of support  $\mathcal{B}$  defined as

$$\sigma_{\mathcal{D},\mathcal{B}}^2 = \int_{\mathbf{s}_1 \in \mathbf{x}_B} \int_{\mathbf{s}_2 \in \mathbf{x}_B} \gamma(\mathbf{s}_1 - \mathbf{s}_2) \, d\mathbf{s}_2 \, d\mathbf{s}_1, \quad (7)$$

158 where  $\gamma(\mathbf{h})$  is the variogram function used for kriging. Again, this term is calculated as part  
 159 of the computation for ordinary block kriging.

160 It is therefore possible to calculate all the terms required to find the covariance of the  
 161 block mean with the ordinary kriging predictor, Eq[4], and this can then be standardized to  
 162 obtain the block correlation

$$\rho_B = \frac{\text{Cov} \left[ \tilde{Z}(\mathbf{x}_B), \bar{z}(\mathbf{x}_B) \right]}{\sqrt{\text{Var} \left[ \tilde{Z}(\mathbf{x}_B) \right] \text{Var} [\bar{z}(\mathbf{x}_B)]}}. \quad (8)$$

163 The correlation is a measure of the strength of the linear relationship between two  
 164 variables, and takes values in the interval  $[-1, 1]$ . Because the variances and covariance in  
 165 Eq [8] are derived from a common stochastic model with authorized (negative semi-definite)  
 166 variograms, it follows that  $\rho_B \in [-1, 1]$  for arbitrary real-valued weights in  $\boldsymbol{\lambda}$ . When  $\boldsymbol{\lambda}$  are  
 167 kriging weights the worse-case scenario for spatial prediction is where  $Z$  is a pure nugget  
 168 random variable with no spatial correlation, and in this case  $\rho_B = 0$  (see appendix).

169 One disadvantage of the correlation as a measure of the reproduction of some quan-  
 170 tity by a predictor is that it is simply a measure of strength of linear association, and so  
 171 may take large values even when the predictions are biased, or over- or under-estimate the  
 172 variance. This was recognized by Lin (1989) who developed the concordance correlation as  
 173 an alternative. The concordance correlation measures the extent to which a variable and its

174 associated predictor fall near the 1:1 line. If predictions are perfect (all observations on the  
175 1:1 line) the concordance correlation (and correlation) are 1, if the variable and predictions  
176 are independent the expected concordance correlation (and correlation) are zero. However,  
177 the predictions and variable may be strongly linearly correlated, but differ markedly with  
178 respect to mean, variance or both. In this case the concordance correlation is smaller than  
179 the correlation, and a more useful measure of association. The concordance correlation has  
180 been used elsewhere in soil science (e.g Corstanje et al., 2008).

181 The concordance correlation for two variables can be written as

$$\rho_c = \frac{2 \text{Cov} [x, x']}{\text{Var} [x] + \text{Var} [x'] + \{\text{E} [x] - \text{E} [x']\}^2}. \quad (9)$$

182 Because ordinary kriging is unbiased the difference between the means in the denom-  
183 inator goes to zero, and so we can write the block concordance correlation as

$$\rho_{\mathcal{B},c} = \frac{\text{Cov} [\tilde{Z}(\mathbf{x}_{\mathcal{B}}), \bar{z}(\mathbf{x}_{\mathcal{B}})]}{\text{Var} [\tilde{Z}(\mathbf{x}_{\mathcal{B}})] + \text{Var} [\bar{z}(\mathbf{x}_{\mathcal{B}})]}. \quad (10)$$

184 Substituting the expression for the covariance in Eq [4] gives

$$\rho_{\mathcal{B},c} = 1 - \frac{\sigma_{\text{K}}^2(\mathbf{x}_{\mathcal{B}})}{\text{Var} [\tilde{Z}(\mathbf{x}_{\mathcal{B}})] + \text{Var} [\bar{z}(\mathbf{x}_{\mathcal{B}})]}. \quad (11)$$

185 From Eqs [8] and [10] it can be seen that

$$\rho_{\mathcal{B},c} = 2\rho_{\mathcal{B}} \frac{\sqrt{\text{Var} [\tilde{Z}(\mathbf{x}_{\mathcal{B}})] \text{Var} [\bar{z}(\mathbf{x}_{\mathcal{B}})]}}{\text{Var} [\tilde{Z}(\mathbf{x}_{\mathcal{B}})] + \text{Var} [\bar{z}(\mathbf{x}_{\mathcal{B}})]}, \quad (12)$$

186 from which it follows that the two correlations are identical if and only if  $\text{Var} [\tilde{Z}(\mathbf{x}_{\mathcal{B}})] =$   
187  $\text{Var} [\bar{z}(\mathbf{x}_{\mathcal{B}})]$ , and that otherwise  $\rho_{\mathcal{B},c} < \rho_{\mathcal{B}}$ . If the block concordance correlation is smaller  
188 than the block correlation this is due to differences in the variance because of the smoothing  
189 effect of the kriging predictor, which will be most pronounced when observations are collected  
190 on sampling grids which are coarse relative to the range of spatial correlation.

## 191 2.2. Hypothetical examples

192 The expressions given above were used to compute block correlations and concordance  
193 correlations in different settings. In each case I considered the block correlation and concor-  
194 dance correlation for a cell-centred square block using the nearest 400 observations from a



195 square sampling grid. Note that in this paper ‘grid spacing’ always denotes the spacing of  
196 the sample grid. Grid spacings from 100 to 1000 m were considered, and correlations were  
197 computed for square blocks of length 10 m to 500 m.

198 Three spatial models were considered. In each case there was a nugget component  
199 with variance  $c_0$ , and two nested spherical components one of range 250 m (comparable  
200 to the shorter grid spacings considered) and variance  $c_1$  and one of range 5000 m (longer  
201 than any grid spacings) and variance  $c_2$ . The first model was ‘nugget-dominated’ with  
202  $c_0 = 0.7$ ,  $c_1 = 0.1$ ,  $c_2 = 0.2$ . The second model was dominated by the short range term:  
203  $c_0 = 0.1$ ,  $c_1 = 0.7$ ,  $c_2 = 0.2$ . The third model was long-range-dominated:  $c_0 = 0.1$ ,  $c_1 =$   
204  $0.2$ ,  $c_2 = 0.7$ . Figure 1 shows the block correlations and block concordance correlations  
205 for each model plotted as contours to show the effect of grid spacing and block length, and  
206 Figure 2 shows block correlation and block concordance correlation plotted against block  
207 length for a 300-m sampling grid.

208 The following key properties of the block correlation emerge

- 209 i. For any grid spacing the block correlation increases with block length, but the effect  
210 of block length depends on the dominant scales of variation relative to block size and  
211 grid spacing. In the short-range-dominated case the block correlation increases more  
212 rapidly with block length with smaller grid spacings than with coarser ones. In the  
213 long-range-dominated case the block correlation is insensitive to block length, other  
214 factors remaining constant.
- 215 ii. For any block size the block correlation declines with increasing grid spacing. The  
216 sensitivity to grid spacing depends on the correlation structure of the random variable,  
217 and on the grid spacing in the short-range-dominated case.
- 218 iii. For any fixed grid spacing and block size the block correlation is smallest in the short-  
219 range dominated case and largest in the long-range dominated case.

220 The first point shows that the effect of block length on uncertainty of the prediction  
221 depends strongly on the important scales of spatial variation and the sampling intensity.

222 The effect may be small, in which case there is no strong reason to take uncertainty and its  
223 communication into account when selecting a block length. However, in other conditions (as  
224 in the short-range dominated case), with sampling on a grid of spacing similar in order to  
225 the range of correlation of a component of the variable, the uncertainty of a block prediction  
226 may be very sensitive to block size.

227 Note, in respect of point (iii) that the variance at lags greater than 250 m is the same  
228 in the short-range and nugget-dominated cases. When most of this variance is in the nugget  
229 term then the effect of this on the uncertainty of the block mean is removed by any spatial  
230 aggregation, and the block size has little effect. It is the short-range component of variation  
231 which largely contributes to the increase in block correlation with block length. There is a  
232 practical consequence of this effect. Consider a case in which all information on a variable  
233 with the short-range dominated variability was available from a 400-m grid. In this case  
234 the short-range component would not be resolved, and its variance would all be attributed  
235 to a nugget component. This would result in overestimation of the block correlation. It is  
236 important to base sampling decisions on variograms estimated from sampling schemes that  
237 provide information on short lags relative to potential block sizes. However, a conservative  
238 approach would be to compute block correlations with a variogram function in which the  
239 nugget component is replaced with a nested spherical or other authorised model with vari-  
240 ance equal to the nugget and range equal to the shortest lag distance at which the variogram  
241 is estimated from supporting data. Note that the model substituted for the lag should be  
242 one like the spherical where the autocorrelation goes exactly to zero at the lag. If the mea-  
243 surement error of the variable is known independently then the nugget variance could be  
244 partitioned into a measurement error component and a component of variance correlated at  
245 fine scales. Only the latter component would be treated as correlated up to the shortest lag  
246 interval in the data, and the measurement error component treated as a nugget effect.

247 The block concordance correlations in Figure 1 show similar behaviour to the block  
248 correlations, with similar effects of grid size and block length depending on the underlying

249 spatial dependence. In Figure 2 there is little difference between the block correlations and  
250 block concordance correlations for different block lengths with a sample grid spacing of 300-m  
251 in the long-range-dominated and nugget-dominated cases. However, the block concordance  
252 correlation decreases more markedly with block length than does block correlation in the  
253 short-range-dominated case.

254 Figure 3 shows the ratio of block concordance correlation to block correlation for  
255 different block lengths and grid spacings in the short-range-dominated, nugget-dominated  
256 and long-range-dominated cases. In the latter two cases the ratio is nowhere very far from  
257 1.0, but in the short-range dominated case the minimum value over the cases explored is  
258 a little less than 0.6. The ratio is reduced by predicting means for smaller blocks or by  
259 increasing the grid spacing. This indicates that the tendency for the kriging predictor to  
260 smooth is increased by sparser sampling but also by predicting for smaller blocks. This latter  
261 effect is only made clear by the block concordance correlation. It would not be apparent in  
262 a visual assessment of block kriging on different supports because the absolute variance of  
263 the smaller blocks is larger.

264 The block concordance correlation shows comparable effects of block length on pre-  
265 diction uncertainty to the block correlation. It is more useful than the block correlation as  
266 an absolute measure of uncertainty because it can be understood as a measure of scatter  
267 about the 1:1 line and not just a measure of linear association. I therefore suggest that  
268 the block concordance correlation is preferred for communication of the uncertainty of block  
269 kriging predictions, and the selection of a block size where there is flexibility to adjust this.

### 270 **3. A case study with soil data**

271 Next I present block correlations computed from a variogram for a soil property. The  
272 property is total nickel (Ni) content of topsoil determined from soil samples collected across  
273 the Humber–Trent region of eastern England as part of the Geochemical Baselines Survey  
274 of the Environment (Rawlins et al., 2003; Johnson et al., 2005). The data were obtained  
275 from sample sites at a mean density of about one per 2 km<sup>2</sup> across the region. Each sample

276 was a composite formed from cores collected at the centre and vertices of a 20-m square.  
277 The cores were length 15 cm excluding surface litter. Material was subsequently air-dried,  
278 disaggregated and sieved to pass 2 mm and sub-sampled by coning and quartering. A 50-g  
279 sub-sample was ground in an agate planetary ball mill until 95% of the material was finer  
280 than 53  $\mu\text{m}$ . Total concentration of Ni, along with 25 other elements was determined for  
281 each sample by wavelength dispersive X-Ray Fluorescence Spectrometry.

282 Details of the analysis of these data are provided by Lark and Lapworth (2013). In  
283 summary, exploratory analysis suggested that there was no pronounced anisotropy in this  
284 variable. Isotropic variograms were estimated using the standard estimator of Matheron  
285 (1962) and alternative estimators, including the resistant estimator of Cressie and Hawkins  
286 (1980). Models were fitted to each set of estimates by weighted least squares and then tested  
287 by cross-validation. For each model the standardized squared cross-validation error, the  
288 square error divided by the point ordinary kriging variance, was computed for each datum  
289 and, following Lark (2000) the median value over all data was computed as a validation  
290 statistic. Lark and Lapworth (2013) tabulate the validation statistics. On the basis of  
291 these the model fitted to the empirical variogram obtained with the estimator of Cressie  
292 and Hawkins (1980) was selected. The estimates and fitted model are shown in Figure 4.  
293 Note that the model was fitted to variogram estimates at lag distances from approximately  
294 200 m to 30 000 m. The model is a double spherical with distance parameters of 2 535 and  
295 16 115 m and corresponding variance components of 42.5 and 82.7 respectively. The nugget  
296 variance is 11.6.

297 The procedures described above were used to compute block correlations and block  
298 concordance correlations for Ni on square blocks from 10 m to 1 km in length from square  
299 grids of interval 2 to 5 km. Note that, as discussed in section 2.2, the block correlations  
300 were computed with a variogram model identical to the fitted one shown in Figure 4 except  
301 that the nugget component was replaced by a spherical variogram with range 200 m and  
302 variance equal to the nugget variance in the fitted model. Figure 5 shows contours of the

303 block correlation and concordance correlation over the range of block sizes and grid spacings  
304 considered. Over this space the block correlation varies from 0.60 to 0.86 and the block  
305 concordance correlation varies from 0.55 to 0.85. Figure 6 shows graphs of block correlation  
306 and concordance correlation against block length for 2-km and 5-km sample grids. With  
307 a 2-km square grid, the average sampling intensity of the Humber–Trent survey, the block  
308 concordance correlation is 0.8 for a 350-m block. This block length could be selected for  
309 kriging the variable to provide an overview of variation in nickel concentration when other  
310 operational factors do not determine the block size. However, the contour map for block  
311 correlations in Figure 5 shows that the sensitivity of the block correlation to block length is  
312 not very great for fixed grid spacing over the range considered. For example, if one preferred  
313 to use a 100-m block for practical reasons for prediction from a 2-km sampling grid the block  
314 concordance correlation is 0.76. This is because most of the variance (60%) is spatially  
315 correlated with a range of 2.5-km or more. With a 5-km sample grid — the sample intensity  
316 of the National Soil Inventory in England and Wales, (McGrath and Loveland, 1992) — the  
317 block concordance correlation for a 350-m block is 0.58. As seen in Figures 5 and 6, the  
318 block concordance correlation is not very sensitive to block length, and increasing the block  
319 length to 1000 m increases it to just 0.63. The dominant limitation on the confidence we  
320 can have in kriged results is the sampling density rather than the block size. For comparison  
321 Figure 7 shows block kriging variances for blocks up to 1000 m in length for prediction from  
322 2-km and 5-km sampling grids.

323 Figure 8 shows block-kriged maps of Ni across the region with square blocks of 350-m  
324 length, and the local kriging variance, and Figure 9 shows the block correlations and block  
325 concordance correlations computed for each block over the stochastic model. Note that the  
326 concordance correlations are mostly larger than 0.8 (the median value is 0.88), which is the  
327 block correlation for a worst-case scenario prediction from a 2-km square grid: a cell-centred  
328 block as far as possible from any observations. The kriging variances, block correlations and  
329 block concordance correlations show some variation, since the sampling intensity is smaller

330 than planned in some areas due to problems of access.

#### 331 4. Discussion

332 The key finding of this paper is that one cannot generalize about the relationship  
333 between block size and prediction quality. While increasing the block size increases block  
334 correlation and concordance correlation, the effect may be very small in cases where the  
335 variation is dominated by a component with a range of correlation which is long relative  
336 to grid spacing and block size. Thus, as in the Humber–Trent case, reducing the block  
337 length from 350 m to 100 m reduces block concordance correlation by a small amount (0.80  
338 to 0.76), particularly if one considers the inevitable uncertainty in the variogram estimate.  
339 The block length has no bearing on uncertainty of the predictions and can be selected on  
340 other criteria. If the block correlation is not deemed sufficient then this can be improved  
341 only by increasing the density of sampling, or finding an appropriate covariate.

342 However, in cases where the sample grid spacing is of similar order to the correlation  
343 range of a significant component of the random variable, block correlation can be sensitive  
344 to block length. In the short-range-dominated hypothetical example (Figures 1 and 2) with  
345 a sample grid of 300 m, the block concordance correlation is increased from 0.35 to 0.80  
346 by using a 350-m block rather than a 50-m block. In these circumstances careful attention  
347 should be paid to both sample spacing and block length, and it could be concluded that  
348 making predictions on a 50-m block is not justified without increasing the sample density.  
349 More generally one might adhere to an operational rule that, at least when mapping to  
350 provide a synoptic overview, the block length is selected to achieve a block concordance  
351 correlation of no less than 0.8, and if this is not achievable at the sample density available  
352 then this is flagged for the data user. If the blocks were fixed at 50 m for practical reasons  
353 (management zones, for example), then the data user should be aware of the weak block  
354 concordance correlations and encouraged to examine the kriging variances and assess the  
355 implications of this uncertainty for any decisions made with the data.

356 In addition to these observations it is suggested that the block concordance correla-

357 tion is used as an alternative to the kriging variance as the primary means to communicate  
358 the uncertainty of geostatistical maps, particularly for general users interested in a synoptic  
359 overview of the soil variable rather than those making specific decisions from the predictions  
360 (e.g. on land remediation) where more focussed decision analysis is necessary. In partic-  
361 ular one might use a verbal scale to indicate the strength of correlation such as the one  
362 proposed by Campbell and Swinscow (2011) whereby a correlation in the interval 0–0.19 is  
363 called ‘very weak’, 0.2–0.39 is ‘weak’, 0.40–0.59 is ‘moderate’, 0.6–0.79 is ‘strong’ and >0.8  
364 is ‘very strong’. It would be advisable to include the numerical values of the concordance  
365 correlation along with the verbal labels to improve the consistency of these interpretations  
366 by data users, and to avoid regressive interpretations (Budescu et al., 2009). If the kriging  
367 variances are also provided then the user, having formed an initial impression of uncertainty  
368 from the concordance correlations may be able to consider the implications of the confidence  
369 intervals of the kriging predictions on the scale of measurement of the target variable. This  
370 approach is consistent with the recommended practice of ‘progressive disclosure’ of informa-  
371 tion about uncertainty (e.g. Wardekker et al., 2008). More accessible and general measures  
372 of uncertainty are provided initially as a prelude to more focussed measures, which may  
373 require greater statistical understanding and may not be needed by all data users.

374 In this paper I have introduced the block correlation and concordance correlation in  
375 the context of ordinary univariate kriging. However, these statistics could be computed  
376 in a straightforward way for predictions obtained by other model-based methods. The  
377 formulation of the block concordance correlation in Eq [11], for example, shows that we  
378 require the prediction error variance, variance of the block spatial mean and variance of the  
379 block mean estimate. In the case of block cokriging the first of these terms is the block  
380 cokriging variance, and the second two terms can be obtained from equations in this paper  
381 and the autovariogram for the target variable in the linear model of coregionalization. In the  
382 case of a best linear unbiased prediction with one or more covariates (kriging with external  
383 drift) the block correlation or concordance correlation is conditional on the block mean of the

384 covariate(s) in addition to the spatial configuration of sample points (as is the prediction  
385 error variance), but can be computed in the same way for different values of this mean  
386 using the underlying variance parameters. In some cases block estimates and prediction  
387 error variances are formed by aggregating point estimates, but in these circumstances the  
388 prediction error variances are only approximated and so it cannot be guaranteed that block  
389 correlations or concordance correlations computed from them are in the interval  $[-1, 1]$ .

## 390 **5. Conclusions**

391 This paper presents the block correlation and concordance correlation, measures of  
392 the quality of a block-kriging prediction of a random variable. Using these measures of  
393 uncertainty it is shown that the size of a block on which a variable is predicted is not always  
394 a good proxy of the uncertainty in the information, although in some circumstances it may  
395 be, specifically when the sample grid is of comparable order to the range of correlation of  
396 a significant component of the random variable. Given this, in the context of the Global-  
397 SoilMap project, it is likely that for much of the globe, with relatively sparse soil data, the  
398 communication of uncertainty is unlikely to have much bearing on the choice of block size.  
399 However, it would be advisable to include block correlations along with other uncertainty  
400 measures to aid communication of uncertainty to the general data user, particularly as the  
401 output is likely to be of value as a general overview of soil variation, prior to planning more  
402 detailed sampling to support local projects.

## **Acknowledgements**

The contribution of all British Geological Survey staff involved in the geochemical sampling of the Humber Trent region, and the subsequent soil analysis, is acknowledged. This paper is published with the permission of the Director of the British Geological Survey (NERC). I am grateful for helpful comments by the Editor and two reviewers, one of whom suggested the inclusion of the concordance correlation.



## References

- Arrouays, D., McBratney, A.B., Minasny, B., Hempel, J.W. et al. 2014a. The Global-SoilMap project specifications. In: Arrouays, D., McKenzie, N., Hempel, J., Richer de Forges, A.C., McBratney, A. (Eds.) *GlobalSoilMap, Basis of the global spatial soil information system*. CRC Press, Boca Raton. pp, 9–12.
- Arrouays, D., Grundy, M.G., Hartemink, A.E., Hempel, J.W., et al. 2014b. Global-SoilMap: toward a fine-resolution global grid of soil properties. *Advances in Agronomy* 125, 93–134.
- Beckett, P.H.T., Burrough, P.A. 1971. The relation between cost and utility in soil survey. IV. Comparison of the utilities of soil maps produced by different survey procedures, and to different scales. *Journal of Soil Science* 22, 466–480.
- Bishop, T.F.A., Horta, A., Karunaratne, S.B. 2015. Validation of digital soil maps at different spatial supports. *Geoderma*, 241–242, 238–249.
- Budescu, D.V., Broomell, S., Por, H.-H. 2009. Improving communication of uncertainty in the reports of the Intergovernmental Panel on Climate Change. *Psychological Science* 20, 299–308.
- Burrough, P.A., Beckett, P.H.T. 1971. The relation between cost and utility in soil survey. I. The design of the Experiment. *Journal of Soil Science* 22, 360–368.
- Campbell, M.J., Swinscow, T.D.V. 2011. *Statistics at square one*. John Wiley and Sons, Chichester.
- Corstanje, R., Kirk, G.J.D., Lark, R.M. 2008. The behaviour of soil process models of ammonia volatilization at contrasting spatial scales. *European Journal of Soil Science*, 59, 1271–1283.
- Cressie, N., Hawkins, D. 1980. Robust estimation of the variogram. *Journal of the International Association of Mathematical Geology* 12, 115–125.
- Dent, D., Young, A. 1981. *Soil Survey and Land Evaluation*. George Allen and Unwin, London.
- Gunn, R.H., Beattie, J.A., Riddler, A.M.H., Lawrie, R.A. 1988. Mapping. In: Gunn, R.H., Beattie, J.A., Reid, R.E., van de Graaff, R.H.M. (Eds.), *Australian soil and land survey handbook: guidelines for conducting surveys*. Inkata Press, Melbourne.

- Hempel, J.W., McBratney, A.B., Arrouays, D., McKenzie, N.J., Hartemink, A.E. 2014. GlobalSoilMap project history. In: Arrouays, D., McKenzie, N., Hempel, J., Richer de Forges, A.C., McBratney, A. (Eds.) GlobalSoilMap, Basis of the global spatial soil information system. CRC Press, Boca Raton. pp, 3–8.
- Hengl, T. 2006. Finding the right pixel size. *Computers and Geosciences*, 32, 1283–1298.
- Heuvelink, G.B.M. 2014. It's the uncertainty, stupid. *Pedometron*, 36, 19–21.
- Johnson, C.C., Breward, N., Ander, E. L., Ault, L. 2005. G-BASE: Baseline geochemical mapping of Great Britain and Northern Ireland. *Geochemistry: Exploration, Environment, Analysis* 5, 1–13.
- Journel, A.G., Huijbregts, C.J., 1978, *Mining Geostatistics*. Academic Press, London.
- Lark, R.M. 2000. A comparison of some robust estimators of the variogram for use in soil survey. *European Journal of Soil Science* 51, 137–157.
- Lark, R.M., Lapworth, D.J. 2013. A novel quality measure for planning geochemical surveys of the soil by kriging. *Geoderma*, 197–198, 27–35.
- Lin, L. I-K. 1989. A concordance correlation coefficient to evaluate reproducibility. *Biometrics*, 45, 255–268.
- Matheron, G. 1962. *Traité de Géostatistique Appliqué, Tome 1. Mémoires du Bureau de Recherches Géologiques et Minières*, Paris.
- McGrath, S.P., Loveland, P.J. 1992. *The Soil Geochemical Atlas of England and Wales*. Blackie, London.
- Rawlins, B.G., Webster, R., Lister, T.R. 2003. The influence of parent material on topsoil geochemistry in Eastern England. *Earth Surface Processes and Landforms* 28, 1389–1409.
- Resources Inventory Committee. 1995. *Soil Inventory Methods for British Columbia*, British Columbia Ministry of Environment. [http://www.for.gov.bc.ca/hts/risc/pubs/teecolo/soil/soil\\_inv\\_methodsbc.pdf](http://www.for.gov.bc.ca/hts/risc/pubs/teecolo/soil/soil_inv_methodsbc.pdf)
- Tobler, W. 1988. Resolution, resampling and all that. In: Mounsey, H., Tomlinson, R.F. (Eds.), *Building databases for global science*. Taylor and Francis, London. 129–137.

- Wardekker, J.A., van der Sluijs, J.P., Janssen, P.H.M., Kloprogge, P., Petersen, A.C. 2007. Uncertainty communication in environmental assessments: views from the Dutch science-policy interface. *Environmental Science & Policy*, 11, 627–641.
- Webster, R., Oliver, M.A. 2007. *Geostatistics for Environmental Scientists*. 2nd Edition John Wiley & Sons, Chichester.

**Appendix. For ordinary block kriging with a pure nugget variogram,  $\rho_B$  and  $\rho_{B,c}$  are zero.**

The ordinary block kriging weights for prediction from  $n$  unique observation sites,  $\lambda_i$ ,  $i = 1, 2, \dots, n$ , are found by solution of the equation:

$$\mathbf{A}\boldsymbol{\lambda}_k = \mathbf{b}, \quad (13)$$

where

$$\mathbf{b} = [\bar{\gamma}_1, \bar{\gamma}_2, \dots, \bar{\gamma}_n, 1]^T$$

and  $\bar{\gamma}_i$  denotes the mean semivariance between the  $i$ th observation and the block. Matrix  $\mathbf{A}$  is  $(n+1) \times (n+1)$  with all diagonal elements zero, and off-diagonal elements  $[i, j]$  equal to the semivariance between the  $i$ th and  $j$ th observation if  $i \leq n$  and  $j \leq n$ . All off-diagonal elements in the  $(n+1)$ th row and column are 1. In addition

$$\boldsymbol{\lambda}_k = [\lambda_1, \lambda_2, \dots, \lambda_n, \psi]^T,$$

where  $\psi$  is a Lagrange multiplier (Webster and Oliver, 2007). In the case of the pure nugget variogram all observations are uncorrelated with each other and with the block so all weights are equal, and, because of the Lagrange multiplier required by the unbiasedness condition of ordinary kriging,

$$\lambda_i = \frac{1}{n}, \quad i = 1, 2, \dots, n. \quad (14)$$

In the case of the pure nugget variogram, all off-diagonal elements of  $\mathbf{A}$   $[i, j]$ ,  $i \leq n$  and  $j \leq n$  are equal to  $\text{Var}[Z]$ , as are the first  $n$  elements of  $\mathbf{b}$ . Any of the first  $n$  equations in the system in Eq [13] therefore takes the form

$$\frac{n-1}{n}\text{Var}[Z] + \psi = \text{Var}[Z],$$

and so

$$\psi = \frac{\text{Var}[Z]}{n}. \quad (15)$$

The ordinary block kriging variance is

$$\sigma_K^2(\mathbf{x}_B) = \mathbf{b}^T \boldsymbol{\lambda}_k - \sigma_{D,B}^2, \quad (16)$$

(Webster and Oliver, 2007).

We can therefore write from Eqs [4],[6] and [16]

$$\text{Cov} \left[ \tilde{Z}(\mathbf{x}_B), \bar{z}(\mathbf{x}_B) \right] = \frac{\text{Var} \left[ \tilde{Z}(\mathbf{x}_B) \right] + \text{Var} [Z] - \mathbf{b}^T \boldsymbol{\lambda}_k}{2}. \quad (17)$$

In the pure nugget case

$$\text{Var} \left[ \tilde{Z}(\mathbf{x}_B) \right] = \boldsymbol{\lambda}^T \mathbf{C} \boldsymbol{\lambda} = \frac{\text{Var}[Z]}{n}, \quad (18)$$

as all elements in  $\boldsymbol{\lambda}$  are  $\frac{1}{n}$  and  $\mathbf{C} = \text{diag}\{\text{Var}[Z]\}$ ; it is also the case that

$$\mathbf{b}^T \boldsymbol{\lambda}_k = \text{Var}[Z] + \psi. \quad (19)$$

Substituting Eqs [18] and [19] into Eq [17], and noting Eq [15] gives

$$\begin{aligned} \text{Cov} \left[ \tilde{Z}(\mathbf{x}_B), \bar{z}(\mathbf{x}_B) \right] &= \frac{1}{2} \left( \frac{\text{Var}[Z]}{n} - \psi \right) \\ &= 0, \end{aligned} \quad (20)$$

from which it follows that  $\rho_B$  and  $\rho_{B,c}$  are zero  $\square$

## Figure captions

1. Block correlation (left) and block concordance correlation (right) as a function of grid spacing and block length for nugget-dominated, short-range-dominated and long-range-dominated random variables.
2. Block correlation as a function of block length for nugget-dominated, short-range-dominated and long-range-dominated random variables kriged from a 300-m square grid.
3. The ratio of block concordance correlation to block correlation as a function of grid spacing and block length for nugget-dominated, short-range-dominated and long-range-dominated random variables.
4. Empirical variogram for topsoil Ni content in the Humber–Trent region with fitted double-spherical model.
5. Block correlation (left) and block concordance correlation (right) as a function of grid spacing and block length for Ni in the Humber–Trent region.
6. Block correlation (left) and block concordance correlation (right) as a function of block length for Ni in the Humber–Trent region kriged from a 2-km or 5-km square grid.
7. Block kriging variance as a function of block length for Ni in the Humber–Trent region kriged from a 2-km or 5-km square grid.
8. Kriged estimates of Ni content on discrete 350-m blocks across the Humber–Trent region (Top) and corresponding block kriging variances (Bottom).
9. Block correlations (top) and concordance correlations (bottom) for kriged estimates of Ni content on 350-m blocks across the Humber–Trent region.




Figure 1: Block correlation (left) and block concordance correlation (right) as a function of grid spacing and block length for nugget-dominated, short-range-dominated and long-range-dominated random variables.




Figure 2: Block correlation as a function of block length for nugget-dominated, short-range-dominated and long-range-dominated random variables kriged from a 300-m square grid.



### Block concordance correlation/Block correlation




Figure 3: The ratio of block concordance correlation to block correlation as a function of grid spacing and block length for nugget-dominated, short-range-dominated and long-range-dominated random variables.




Figure 4: Empirical variogram for topsoil Ni content in the Humber–Trent region with fitted double-spherical model.




Figure 5: Block correlation (left) and block concordance correlation (right) as a function of grid spacing and block length for Ni in the Humber–Trent region.




Figure 6: Block correlation (left) and block concordance correlation (right) as a function of block length for Ni in the Humber–Trent region kriged from a 2-km or 5-km square grid.




Figure 7: Block kriging variance as a function of block length for Ni in the Humber-Trent region kriged from a 2-km or 5-km square grid.

