

Accepted Manuscript

Title: Complete genome of *Planococcus rifietoensis* M8^T, a halotolerant and potentially plant growth promoting bacterium

Author: Wah-Seng See-Too Peter Convey David A. Pearce
Yan Lue Lim Robson Ee Wai-Fong Yin Kok-Gan Chan



PII: S0168-1656(16)30029-3
DOI: <http://dx.doi.org/doi:10.1016/j.jbiotec.2016.01.026>
Reference: BIOTEC 7385

To appear in: *Journal of Biotechnology*

Received date: 16-1-2016
Accepted date: 20-1-2016

Please cite this article as: See-Too, Wah-Seng, Convey, Peter, Pearce, David A., Lim, Yan Lue, Ee, Robson, Yin, Wai-Fong, Chan, Kok-Gan, Complete genome of *Planococcus rifietoensis* M8T, a halotolerant and potentially plant growth promoting bacterium. *Journal of Biotechnology* <http://dx.doi.org/10.1016/j.jbiotec.2016.01.026>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Complete genome of *Planococcus rifietoensis* M8^T, a halotolerant and potentially plant growth promoting bacterium

Wah-Seng See-Too^{a,b}, Peter Convey^{b,c}, David A. Pearce^{b,c,d}, Yan Lue Lim^a, Robson Ee^a, Wai-Fong Yin^a, Kok-Gan Chan^{a,e*}

^a Division of Genetics and Molecular Biology, Institute of Biological Sciences, Faculty of Science, University of Malaya, Kuala Lumpur, Malaysia

^b National Antarctic Research Centre (NARC), Institute of Postgraduate Studies, University of Malaya, 50603 Kuala Lumpur, Malaysia

^c British Antarctic Survey, NERC, High Cross, Madingley Road, Cambridge CB3 0ET, UK

^d Faculty of Health and Life Sciences, University of Northumbria, Newcastle Upon Tyne NE1 8ST, UK

^e UM Omics Centre, University of Malaya, Kuala Lumpur

*Corresponding author

Tel.: +603–79675162; E-mail address: kokgan@um.edu.my

Highlights

- *Planococcus rifietoensis* M8^T (=DSM 15069^T =ATCC BAA-790^T) is a halotolerant bacterium with potential plant growth promoting properties
- Various key genes coding for these properties were identified
- This is the first complete genome sequence of *Planococcus rifietoensis*

Abstract

Planococcus rifietoensis M8^T (=DSM 15069^T =ATCC BAA-790^T) is a halotolerant bacterium with potential plant growth promoting properties isolated from an algal mat collected from a sulfurous spring in Campania (Italy). This paper presents the first complete genome of *P. rifietoensis* M8^T. Genes coding for various potentially plant growth promoting properties were identified within its genome.

Keyword: nitrogen fixing, potassium homeostasis, biotechnology, biofertilizer

High salinity in agricultural soils is of global concern, affecting productivity of crops (Mayak et al., 2004). The plant growth promoting (PGP) properties of halotolerant bacteria are highly sought after biotechnological traits for facilitating plant growth in saline soils (Yildirim & Taylor, 2005; Barassi et al., 2006). *Planococcus rifietoensis* is a moderately halotolerant (Romano et al., 2003) bacterium capable of promoting wheat growth by converting ammonia to nitrogen, thereby enabling fertilization of soil under high salinity stress, as well as metabolizing potassium to reduce ion imbalance of plant cells (Rajput et al., 2013). Here, we present the first complete genome of *Planococcus rifietoensis* M8^T, identifying finding of its various key genes for PGP properties.

Genomic DNA of *P. rifietoensis* M8^T was isolated from an overnight cell suspension culture using the MasterPure™ Gram positive DNA purification kit (Epicentre Technologies). The genomic DNA was then constructed into a 20-kb SMRTbell template library. Pacific Biosciences (PacBio) RSII sequencing platform was used to perform whole genome sequencing using C4 chemistry in two single molecule real time (SMRT) cells (Ee et al., 2015). A total of 41,347 reads with a mean read length of 12,808 bp were generated. The reads were *de novo* assembled using hierarchical genome assembly process (HGAP) algorithm version 2 into a circular contig with an average genome coverage of 124-fold (Goh et al., 2015).

Genome annotation was performed using NCBI Prokaryotic Genome Annotation Pipeline (PGAP) version 2.10 and Rapid Annotation using Subsystem Technology (RAST) version 2.0 (Aziz et al., 2008, Overbeek et al., 2014, Brettin et al., 2015). The genome of *P. rifietoensis* M8^T consists of a 3,527,718 bp circular chromosome with a G+C content of 48.4%. A total of 3,514 protein coding genes were predicted along with 6 rRNA and 49 tRNA genes (Table 1).

Functional annotation of the genome revealed genes coding for assimilation of ammonia via both the GDH pathway using NAD-specific glutamate dehydrogenase (NCBI locus tag: AUC31_04065 & AUC31_0581; EC 1.4.1.2 & EC 1.4.1.4) the and glutamine synthetase (GS)-glutamate synthase (GOGAT) pathway using glutamine synthetase (NCBI locus tag: AUC31_00390, EC 6.3.1.2), glutamate synthase (NCBI locus tag: AUC31_12670, AUC31_12675, AUC31_03115; EC 1.4.1.13), serine hydroxymethyltransferase (NCBI locus tag: AUC31_14095, EC 2.1.2.1), glycine dehydrogenase (NCBI locus tag: AUC31_02400, AUC31_02405; EC 1.4.4.2), serine-glyoxylate aminotransferase (EC 2.6.1.45), potassium homeostasis gene (NCBI locus tag: AUC31_11595, AUC31_11650, AUC31_11655, AUC31_16700, AUC31_05695, AUC31_0775, EC: 2.7.3) and phosphate metabolism and transporter gene (NCBI locus tag: AUC31_02645, AUC31_02760, AUC31_02525, EC: 2.7.13.3, TC: 3.A.1.7.1). Furthermore, genes coding for various chemical compounds metabolism processes were found, including genes for degradation of gentisate, glycerol, mannitol, inositol, glycerate, formaldehyde, ribulose and carotenoids, isoprenoids, and isopropanoids. This finding may indicate the application of this strain in bioremediation (Urbieta et al.,

2015). Further study of the full PGP capacity of this strain is required in order to further confirm its potential for biotechnological application.

Nucleotide sequence accession number

The complete chromosome sequence has been deposited in GenBank under the accession number CP013659.

Acknowledgements

This work was supported by the University of Malaya High Impact Research Grants (UM-MOHE HIR Grant UM.C/625/1/HIR/MOHE/CHAN/14/1, Grant No. H-50001-A000027; UM-MOHE HIR Grant UM.C/625/1/HIR/MOHE/CHAN/01, Grant No. A-000001-50001) awarded to Kok-Gan Chan. Peter Convey is supported by NERC core funding to the British Antarctic Survey's 'Biodiversity, Evolution and Adaptation' Team.

References

- Aziz, R.K., Bartels, D., Best, A.A., DeJongh, M., Disz, T., Edwards, R.A., Formsma, K., Gerdes, S., Glass, E.M., Kubal, M., Meyer, F., Olsen, G.J., Olson, R., Osterman, A.L., Overbeek, R.A., McNeil, L.K., Paarmann, D., Paczian, T., Parrello, B., Pusch, G.D., Reich, C., Stevens, R., Vassieva, O., Vonstein, V., Wilke, A., Zagnitko, O., 2008. The RAST Server: rapid annotations using subsystems technology. *BMC Genomics* 9, 75.
- Barassi, C.A., Ayrault, G., Creus, C.M., Sueldo, R.J., Sobrero, M.T. 2006. Seed inoculation with *Azospirillum* mitigates NaCl effects on lettuce. *Sci. Hort.* 109, 8-14.
- Brettin, T., Davis, J.J., Disz, T., Edwards, R.A., Gerdes, S., Olsen, G.J., Olson, R., Overbeek, R., Parrello, B., Pusch, G.D., Shukla, M., Thomason, J.A., Stevens, R., Vonstein, V., Wattam, A.R., Xia, F., 2015. RASTtk: A modular and extensible implementation of the RAST algorithm for building custom annotation pipelines and annotating batches of genomes. *Sci. Rep.* 5, 8365.
- Ee, R., Yong, D., Lim, Y.L., Yin, W.F. and Chan, K.G., 2015. Complete genome sequence of oxalate-degrading bacterium *Pandoraea vervacti* DSM 23571 T. *J. Biotechnol.* 204, 5-6.
- Goh, K.M., Chan, K.G., Yaakop, A.S. and Ee, R., 2015. Complete genome of *Jeotgalibacillus malaysiensis* D5^T consisting of a chromosome and a circular megaplasmid. *J. Biotechnol.* 204, 13-14.
- Mayak, S., Tirosh, T., Glick, B.R. 2004 Plant growth-promoting bacteria confer resistance in tomato plants to salt stress. *Plant Physiol. Biochem.* 42, 565–572.
- Overbeek, R., Olson, R., Pusch, G.D., Olsen, G.J., Davis, J.J., Disz, T., Edwards, R.A., Gerdes, S., Parrello, B., Shukla, M., Vonstein, V., Wattam, A.R., Xia, F., Stevens, R., 2014. The SEED and the Rapid Annotation of microbial genomes using Subsystems Technology (RAST). *Nucleic Acids Res.* 42, D206–D214
- Rajput., L. , Imran, A., Mubeen, F., Hafeez, F. Y. 2013. Salt-tolerant PGPR strain *Planococcus rifietoensis* promotes the growth and yield of wheat (*Triticum aestivum* L.) cultivated in saline soil. *Pak. J. Bot.* 45, 1955–1962.
- Romano, I., Giordano, A., Lama, L., Nicolaus, B., Gambacorta, A. 2003. *Planococcus rifietoensis* sp. nov., isolated from algal mat collected from a sulfurous spring in Campania (Italy). *Syst. Appl. Microbiol.* 26, 357–366
- Urbieta, M.S., Donati, E.R., Chan, K.G., Shahar, S., Sin, L.L. and Goh, K.M., 2015. Thermophiles in the genomic era: Biodiversity, science, and applications. *Biotechnol. Adv.* 33(6), 633-647.
- Yildirim, E., Taylor, A.G. 2005. Effect of biological treatments on growth of bean plants under Salt Stress. *Ann. Rep. Bean Improv. Coop.* 48, 176-177.

Table 1. Genome features of *P. rifietoensis* M8^T.

Attributes	Value
Genome size (bp)	3,527,718
GC content (%)	48.4
Total number of genes	4,141
Protein coding genes	3,901
Contigs	1
rRNA genes	27
tRNA genes	63