


Received: 24 June 2020 | Revised: 27 July 2020 | Accepted: 5 August 2020

DOI: 10.1111/andr.12884

## ORIGINAL ARTICLE

ANDROLOGY   WILEY

# Long-term effect of gender-affirming hormone treatment on depression and anxiety symptoms in transgender people: A prospective cohort study

Zoë Aldridge<sup>1,2</sup> | Shireen Patel<sup>1</sup> | Boliang Guo<sup>1</sup> | Elena Nixon<sup>1</sup> |  
Walter Pierre Bouman<sup>1,3</sup> | Gemma L. Witcomb<sup>3</sup> | Jon Arcelus<sup>1,3</sup> 

<sup>1</sup>Faculty of Medicine and Health Sciences, Institute of Mental Health, University of Nottingham, Nottingham, UK

<sup>2</sup>School of Sport, Exercise and Health Sciences, Loughborough University, Loughborough, UK

<sup>3</sup>Nottingham Centre for Transgender Health, Nottingham, UK

**Correspondence**

Jon Arcelus. Institute of Mental Health, Jubilee Campus, Triumph Road, University of Nottingham, Nottingham NG7 2TU, UK.  
Email: [Jon.Arcelus@nottingham.ac.uk](mailto:Jon.Arcelus@nottingham.ac.uk)

**Funding information**

The data analysis only was funded by the National Institute for Health Research (NIHR) Applied Research Collaboration East Midlands (ARC EM), NIHR Research for Patient Benefit (RfPB) and Nottinghamshire Healthcare NHS Foundation Trust.

**Abstract**

**Background:** Cross-sectional studies show that transgender people are more likely than cisgender people to experience depression and anxiety before gender-affirming hormone treatment (GAHT). However, the effect of GAHT on mental health in transgender people, and the role of other factors that may have a predictive effect, is poorly explored.

**Objectives:** Using a longitudinal methodology, this study investigated the effect of 18-month GAHT on depression and anxiety symptomatology and the predictors on mental health outcomes in a large population of transgender people.

**Materials and methods:** Participants ( $n = 178$ ) completed a socio-demographic questionnaire, the Hospital Anxiety and Depression Scale (HADS), the Multidimensional Scale of Perceived Social Support (MSPSS) and the Autism Spectrum Quotient—Short Version (AQ-Short) at pre-assessment (T0) and at 18 months after initiation of GAHT (T1).

**Results:** From T0 to T1, symptomatology was significantly decreased for depression ( $P < .001$ ) and non-significantly reduced for anxiety ( $P = .37$ ). Scores on the MSPSS predicted reduction in depression, while scores on the AQ-Short predicted reduction in anxiety.

**Discussion:** GAHT reduces symptoms of depression which are predicted by having higher levels of social support. Although anxiety symptoms also reduce, the changes are not significant and high levels of anxiety still remain post-GAHT.

**Conclusions:** These results highlight the important mental health benefits of GAHT. Support services (professional, third sector or peer support) aiming at increasing social support for transgender individuals should be made available.

**KEYWORDS**

autism, hormone therapy, longitudinal, mental health, social support, transgender

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *Andrology* published by Wiley Periodicals LLC on behalf of American Society of Andrology and European Academy of Andrology

## 1 | INTRODUCTION

Treatment-seeking transgender people who are not on hormone treatment have reported high levels of mental health problems, particularly anxiety, depression and self-harm, which are likely caused by a number of internal and external stressors.<sup>1-12</sup> Studies examining mental health in transgender people have primarily focused on individuals attending transgender health services and hence those who are likely to experience a higher level of distress about their assigned sex at birth. These studies have primarily looked cross-sectionally at levels of anxiety,<sup>7,13,14</sup> depression<sup>7,15-17</sup> and self-harm.<sup>18-20</sup>

With regard to anxiety, several studies have demonstrated high levels in transgender people before gender-affirming hormone treatment (GAHT).<sup>21,22</sup> For example, Bouman et al<sup>13</sup> found that levels of anxiety in transgender people were three times higher than those in a matched sample from the general population. This study also found that transgender males were more anxious than transgender females. Interestingly, the high scores on autistic traits found among this population have been suggested to be a product of the high levels of anxiety and low self-esteem often experienced by this group<sup>15</sup> and not autism per se.<sup>23</sup> However, a recent study has demonstrated stability in autistic traits following GAHT.<sup>24</sup>

Similar to anxiety, high levels of depression have also been reported in transgender individuals, prior to GAHT.<sup>16,18,22</sup> Witcomb et al<sup>16</sup> reported that transgender people prior to receiving GAHT had a fourfold increased risk of a probable depressive disorder compared to a matched control sample from the general population. Why this is the case is unclear, but social factors such as lack of general social support,<sup>9,25-27</sup> parental support<sup>28</sup> and peer support have been found to be associated with depressive symptoms among transgender people.<sup>29,30</sup> Experiences of transphobic discrimination are associated with increased odds of suffering from depression<sup>31</sup> independent of other types of discrimination, for example racism. This suggests that transgender people who are ethnic minorities are at even greater risk, because of the intersectional experience of discriminatory events. In addition, while unemployment increases the risk of depression in the general population<sup>32</sup> and transgender people have been found to have a higher unemployment rate than cisgender people<sup>33</sup>—being in employment is associated with higher levels of experienced transphobia and fear of disclosing mental health problems in the transgender population.<sup>34,35</sup>

Another factor that has been associated with mental health problems in treatment-seeking transgender people is age. Younger transgender people report high levels of bullying<sup>36</sup> and very high levels of self-harm,<sup>19,37</sup> which have been associated with increased anxiety as well as effects on self-esteem, family relationships and social life, which all negatively influence mental well-being.

While these studies have provided valuable insight, the use of cross-sectional methodologies to examine the impact of the above factors, particularly the role of GAHT in mental health, is limited.

Therefore, it is critical to explore this on a within-subject basis using a longitudinal design. This is the most effective approach to show the effects of GAHT on mental health as it provides the opportunity to examine individuals prior to and during GAHT.

A small number of longitudinal studies that focus on the effect of GAHT on mental health do exist. Colizzi et al<sup>38</sup> reported significant reductions in mental health symptoms after the initiation of GAHT with anxiety reducing from 50% to 17% and depression from 24% to 11%. Heylens et al<sup>39</sup> also showed significant reductions in symptoms of anxiety and depression after the initiation of GAHT to the point where they resemble those of the general population. These studies are, however, not without limitations. Heylens et al's<sup>39</sup> study has a small sample size ( $n = 57$ ), while Colizzi et al's<sup>38</sup> study is limited by the lack of evaluation of factors that may have impacted on the mental health of their participants, such as social support. Both studies describe the need to replicate their findings. In contrast, Bränström and Pachankis<sup>40</sup> using the Swedish population register showed no significant association between the likelihood of accessing mental health treatment and time since initiation of GAHT. The limitation of their study includes primarily that accessing mental health services does not necessarily reflect actual mental health and there is little additional information about the type of mental health treatment received by their participants. These limitations mean that this study cannot provide reliable evidence regarding the role of GAHT on the mental health symptoms of transgender people, and this information is vital in order to provide an evidence base of GAHT improving overall quality of life of transgender people.

While the available longitudinal studies have provided valuable evidence of the effect of GAHT on transgender people's mental health, there is a requirement to replicate these studies addressing their limitations. With this in mind, the primary aim of this study is to examine the effect of GAHT on anxiety and depression symptoms. The study will focus on those who have been on treatment for over 18 months as this allows for enough time for GAHT to produce physical, bodily changes but before surgical procedures have taken place, which could bias the results. As some physical changes can be quicker in assigned females at birth than in assigned males at birth (eg voice change with testosterone),<sup>2</sup> which can affect mental health outcome following GAHT, the results of GAHT in anxiety and depression for both groups will be presented separately. It is hypothesized that an improvement in mental health will take place in those assigned male and female at birth following GAHT treatment. Unfortunately, because of the long waiting list for gender-affirming surgical treatment in the United Kingdom (UK), it is unlikely that people will have undergone these interventions before this time. The secondary aim of this study is to examine pre-GAHT factors which may be predicting changes in anxiety and depression following GAHT. The predictors selected for this study are based on the literature and include ethnicity, age, assigned sex at birth, civil status, employment, social support, and autistic traits. This study hypothesized that symptoms of depression and anxiety would be significantly decreased after 18 months of GAHT.

## 2 | MATERIALS AND METHODS

### 2.1 | Participants

Participants were invited to take part through a national transgender health service in Nottingham, UK. This service is part of the National Health Service (NHS) and offers assessment for suitability of GAHT as well as chest and genital reconstructive surgery. The service also offers GAHT and speech and language therapy. The service accepts referrals from people aged 17 and over who are seeking, or considering, medical transition.

### 2.2 | Procedures

The sample consisted of individuals who attended an assessment at the transgender health service from November 2014 to March 2018, who agreed participation and who were not on GAHT prior to the assessment. Prior to the clinical assessment, every patient was invited to participate in the study. If agreed, they were invited to complete a baseline questionnaire pack (T0). The pack included a socio-demographics questionnaire (age, sex assigned at birth, gender identity, ethnicity, employment status, relationship status and whether participants were taking cross-sex hormones and/or blockers pre-assessment—as a significant proportion of young people are referred from the only existing child and adolescent transgender health services in the United Kingdom). Validated questionnaires regarding anxiety and depression (HADS), social support (MSPSS) and autistic traits (AQ-Short) were also included in the information pack. Data were only included if participants returned a signed consent form with the study questionnaires.

Participants who consented and returned T0 questionnaires were invited to complete a T1 questionnaire 18 months after commencing GAHT. The T1 questionnaire pack consisted of a HADS questionnaire. This allowed a comparison of changes in depression and anxiety symptoms before and after GAHT. Data were collected in October 2019. Except for the data analysis, the study was primarily unfunded and set up in a busy clinic.

### 2.3 | Tools

#### 2.3.1 | The Hospital Anxiety and Depression Scale (HADS)

The HADS is a 14-item self-report screening scale originally developed to indicate the possible presence of anxiety and depression states in medical non-psychiatric outpatient clinics.<sup>41</sup> The HADS consists of two subscales, HADS—Depression (HADS-D) and HADS—Anxiety (HADS-A). Each subscale has seven items that are rated on a 4-point Likert scale that ranges from 0 to 4 with some items being reverse-scored. A maximum total of 21 can be obtained on each subscale. A score of 0-7 on both scales implies a non-clinical range, while a score of 8-10 suggests the possible presence of a depressive

or anxiety disorder. A score of 11 or higher suggests the probable presence of a depressive or anxiety disorder. Caseness of depression and anxiety has been suggested for scores above 8.<sup>42</sup> The HADS has previously been used with transgender individuals.<sup>13,43</sup> For depression (HADS-D), this gave a specificity of 0.7 and a sensitivity of 0.9. For anxiety (HADS-A), this gave a specificity of 0.78 and a sensitivity of 0.9. In this study; Cronbach's alpha for depression was 0.76 and for anxiety 0.68.

#### 2.3.2 | The multidimensional scale of perceived social support (MSPSS)

The MSPSS is a 12-item self-report scale to record levels of social support from family, friends and significant others.<sup>44</sup> The measure consists of three subscales to measure the three different types of support. Items are rated on a Likert scale that ranges from 1 ('very strongly agree') to 7 ('very strongly disagree'). To calculate subscale scores, items from each subscale are added together and divided by 4. A total score is calculated by adding together all 12 items and dividing by 12. The mean and total scores range from 1 to 7 with a higher score indicating a higher level of perceived social support. A mean total scale score ranging from 1 to 2.9 can be considered low support; a score of 3 to 5 can be considered moderate support; and a score from 5.1 to 7 can be considered high support. The MSPSS has previously been used with transgender individuals.<sup>26,45</sup> In this study, Cronbach's alpha was 0.89.

#### 2.3.3 | Autism spectrum quotient—short version (AQ-Short)

The AQ-Short is a 28-item self-report questionnaire designed to measure autistic traits to give an indication of where the person lies on the continuum of the spectrum, ranging from healthy to autistic.<sup>46,47</sup> It is a shortened version of the validated AQ-50.<sup>48</sup> It consists of two higher order factors related to autistic traits, including numbers and patterns (which assess the extent to which people are fascinated by numbers, dates, patterns and categories) and social behaviours. The AQ-Short is a 4-point Likert scale ranging from 'definitely agree' to 'definitely disagree', with some items being reverse-scored. Total scores range between 28 and 112. The AQ-Short has previously been used with transgender populations.<sup>23,24,49</sup> Higher scores represent higher levels of autistic traits. Although not intended to be a diagnostic tool, a cut-off of  $\geq 70$  was found to have a sensitivity of 0.94 and specificity of 0.91 to discriminate between an autism sample and a community sample. Cronbach's alpha was 0.86.

#### 2.3.4 | Data analysis

Data analyses were performed using the Statistical Software Package Stata 16.<sup>50</sup> Stata 16 was used to conduct power analysis. Only those

participants not on GAHT at assessment (T0) were included in the regression analysis. All missingness was imputed using analytical model with 20 imputed data sets generated for each model. Paired sample t tests were used to determine whether there had been a significant change in the HADS-D and HADS-A subscales from T0 to T1. Multiple regression was conducted to explore ethnicity, employment status, relationship status, age, assigned sex, MSPSS and AQ. The hypothesis regarding whether the specific factors were predictive of changes in anxiety and depression was tested via a moderator analysis, entering only the subscales found to be significant in the linear regressions and a product of their combined centred scores. This was tested via a multiple regression. Bonferroni corrections were used to correct multiplicity issue if needed. Although data are not normally distributed, Allison<sup>51</sup> states that normality is the least important assumption of regression and as data met the assumptions for linearity, homoscedasticity and absence of multicollinearity or extreme outliers, a multiple regression analysis was conducted. The socio-demographic categories were split into two distinct groups for each category as seen in Table 1. Assigned sex at birth instead of gender identity was used in the socio-demographics in view of the many different gendered identities described, as an analysis based on gender identities would have made the analyses too complex to interpret; this followed previous study approaches.<sup>23,24</sup> The MSPSS and AQ-28 Short at T0 were significant factors predicting change in HADS-D and HADS-A at T1. To check the robustness of regression estimates sensitive to missingness, all regression models were re-run on observed data only and the results were examined against the results from imputed data set.

Ethical approval for the study was received from the NHS Ethics Committee (14/EM/0092) and the Research and Development Department at Nottinghamshire Healthcare NHS Foundation Trust in line with Health Research Authority guidance,<sup>52</sup> which included approval for individuals aged 17 and over to sign giving their consent without the need for additional parental consent.

### 3 | RESULTS

#### 3.1 | Socio-demographic characteristics of the participants

A total of 1,271 participants were assessed between November 2014 and March 2018, completed T0 questionnaires and agreed participation in the study. Seventy-one per cent (N = 906) could be included in the analysis as they had not received hormones prior to assessment. Of these, 178 (20%) went on to complete a T1 questionnaire after 18 months of GAHT, indicating a response rate of 20%. Responders did not differ from non-responders in terms of either demographic characteristics or baseline AQ-Short scores, but they were significantly less anxious at baseline than non-responders (median 9 vs 8,  $P = .001$ ;  $z = 3.225$ ) (see Table 1).

The age range from the 178 participants that completed T0 and T1 questionnaires ranged from 17 to 79 years with a median age of 23 years. More than half of the participants (n = 95; 53.3%) were

**TABLE 1** Socio-demographic characteristics (n,%) and mean (SD) scores on HADS, MSPSS and AQ-Short score of all responders assigned male and female at birth

|                                | Responders    |    | Assigned male at birth (2) |    | Assigned female at birth (1) |    |
|--------------------------------|---------------|----|----------------------------|----|------------------------------|----|
|                                | n = 178       | n% | n = 95                     | n% | n = 83                       | n% |
| Age, median (range) years      | 23 (1774)     |    | 28 (1779)                  |    | 21 (1764)                    |    |
| Ethnic origin n (%)            |               |    |                            |    |                              |    |
| White                          | 167           | 94 | 92                         | 97 | 75                           | 91 |
| Other                          | 11            | 6  | 3                          | 3  | 8                            | 9  |
| Not known                      | -             | -  | -                          | -  | -                            | -  |
| Employment status n (%)        |               |    |                            |    |                              |    |
| Employed (1)                   | 75            | 41 | 45                         | 47 | 28                           | 33 |
| Student (2)                    | 58            | 32 | 20                         | 21 | 37                           | 44 |
| Housewife/<br>househusband (4) | -             | -  | -                          | -  | -                            | -  |
| Voluntary work (3)             | 7             | 4  | 5                          | 5  | 2                            | 2  |
| Retired (6)                    | 5             | 3  | 6                          | 6  | -                            | -  |
| Disabled (5)                   | 7             | 5  | 5                          | 5  | 3                            | 3  |
| Unemployed (0)                 | 26            | 14 | 14                         | 14 | 12                           | 14 |
| Not known (9)                  | 1             | 1  |                            |    | 1                            | 1  |
| Civil status n (%)             |               |    |                            |    |                              |    |
| Single (1)                     | 120           | 69 | 55                         | 57 | 65                           | 78 |
| Married (2)                    | 15            | 7  | 12                         | 12 | 1                            | 1  |
| Civil partner (3)              | 5             | 3  | -                          | -  | 5                            | 6  |
| In a relationship (7)          | 4             | 3  | 1                          | 1  | 4                            | 4  |
| Divorced/separated (4)         | 20            | 11 | 18                         | 18 | 2                            | 2  |
| Widowed (5)                    | 2             | 2  | 2                          | 2  | 1                            | 1  |
| Other (6)                      | -             | -  | -                          | -  | -                            | -  |
| Not known (9 + Blanks)         | 12            | 7  | 7                          | 7  | 5                            | 6  |
| HADSD                          | 7.24 (4.03)   |    | 7.03 (4.11)                |    | 7.48 (3.94)                  |    |
| HADSA                          | 8.07 (4.34)   |    | 7.54 (4.31)                |    | 8.69 (4.32)                  |    |
| MSPSS                          | 4.85 (1.29)   |    | 4.64 (1.35)                |    | 5.1 (1.16)                   |    |
| AQ                             | 64.77 (11.86) |    | 62.83 (11.94)              |    | 66.97 (11.44)                |    |

assigned male sex at birth, and 83 (46.7%) were assigned female sex at birth. The large majority of participants classified themselves as white (n = 167; 94%), were single (n = 120; 69%) and were in employment (n = 75; 41%) or students (n = 58; 32%). Participants who were assigned male sex at birth were more likely to be in employment compared to participants assigned female sex at birth (47% vs 33%), while more participants assigned female sex at birth were single at the time of assessment (78% vs 57%) and a higher percentage of participants assigned male sex at birth were divorced/separated (18% vs 2%) (see Table 1).

TABLE 2 Means (SD) of HADS-D and HADS-A scores of responders at T0 and T1

|               | All responders |             |   | Assigned male at birth |             |   | Assigned female at birth |             |   |
|---------------|----------------|-------------|---|------------------------|-------------|---|--------------------------|-------------|---|
|               | N              | Mean (SD)   | Mean change from T0 to T1 (95% CI), P-value | n                      | Mean (SD)   | Mean change from T0 to T1 (95% CI), P-value | n                        | Mean (SD)   | Mean change from T0 to T1 (95% CI), P-value |
| <b>HADS-D</b> |                |             |   |                        |             |   |                          |             |   |
| T0            | 178            | 7.24 (4.03) | -2.05<br>(-2.72 to -1.38)<br>P = .00        | 95                     | 7.03 (4.11) | -1.91<br>(-2.80 to -1.01)<br>P = .00        | 83                       | 7.48 (3.94) | -2.21<br>(to 3.23 to -1.20)<br>P = .00      |
| T1            | 178            | 5.19 (3.73) |   | 95                     | 5.13 (3.92) |   | 83                       | 5.26 (3.52) |   |
| <b>HADS-A</b> |                |             |   |                        |             |   |                          |             |   |
| T0            | 178            | 8.07 (4.34) | -0.31<br>(-0.97 to 0.36),<br>P = .37        | 95                     | 7.54 (4.31) | -1.16<br>(-1.50 to 0.39)<br>P = .25         | 83                       | 8.69 (4.32) | -0.55<br>(-0.97 to 0.92) P = .97            |
| T1            | 178            | 7.77 (3.90) |   | 95                     | 6.98 (3.96) |   | 83                       | 8.66 (3.65) |   |

### 3.2 | Anxiety and depression scores

The mean score for the total group for anxiety was 8.07 (SD: 4.34). It was higher in those assigned female (8.69 (SD: 4.32)) versus those assigned male at birth (7.54 (SD: 4.31)). The mean score for depression was 7.24 (SD: 4.03), also higher in assigned females (7.48 (SD: 3.94)) than assigned males at birth (7.03 (SD: 4.11)).

At T0 (before hormone treatment), 51.13% of participants scored 8 or over on the HADS-A subscale, and in the case of the HADS-D subscale, 47.75% of participants scored above 8 placing these participants within the categories of possible to probable presence of an anxiety or depression disorder. At T1, 47.19% of participants scored 8 or above in the HADS-A subscale showing a reduction of 3.94%, and in the HADS-D subscale, 25.84% of people scored 8 or above showing a reduction of 21.91%.

### 3.3 | Change in anxiety and depression scores between T0 and T1

There was a statistically significant reduction in mean scores of HADS-D from T0 to T1 (mean change difference, -2.05; 95% CI, -2.72 to -1.38;  $P = .00$ ). This indicated a reduction in depression following 18 months of GAHT. There was also a reduction in the HADS-A score from T0 to T1, but this was not statistically significant (mean change difference, -0.31; 95% CI, -0.97 to 0.36,  $P = .37$ ). The same findings (a significant reduction in HADS-D and a non-significant reduction in HADS-A) were found when comparing T0 and T1 according to sex assigned at birth (see Table 2).

### 3.4 | Predictors of anxiety and depression change after hormone treatment

Two multiple regressions with seven predictor variables were conducted to explore the predictors of change, from T0 to T1, in scores on HADS-D and HADS-A. The predictors for each were ethnicity, employment status, relationship status, assigned sex at birth, and age. MSPSS and AQ-Short at T0 were used as independent variables for both regressions.

The results for the first regression showed that overall, the model was significant ( $F(7,152)=2.09$ ,  $P = .04$ ) and explained 8.8% ( $R^2 = 0.088$ ) of the total variance in depression scores. The model also showed that mean MSPSS scores at T0 were the only significant predictor of HADS-D change between T0 and T1 ( $\beta = 0.81$ ,  $P = .006$ ). The second regression showed that overall, the model was significant ( $F(7,152)=2.09$ ,  $P = .048$ ) and explained 8.8% ( $R^2 = 0.088$ ) of the total variance in anxiety scores. The model also showed mean AQ-Short scores at T0 was a significant predictor of HADS-A change between T0 and T1 ( $\beta = -0.069$ ,  $P = .034$ ). The findings suggest that only levels of social support (MSPSS scores) and autistic spectrum traits (AQ scores) were able to predict changes in anxiety and depression following 18 months of GAHT. Having higher levels of social support

(higher scores of MSPSS) predicted a reduction in depression scores following 18 months of GAHT ( $P = .006$ ) and having lower levels of autistic spectrum traits (lower AQ scores) predicted a reduction of anxiety symptoms following 18 months of GAHT ( $P = .03$ ), although this reduction was statistically non-significant (see Table 3).

## 4 | DISCUSSION

This prospective longitudinal study aimed to explore whether 18 months of GAHT reduces symptoms of anxiety and depression in transgender people, while addressing the limitations of previous studies, by recruiting a large sample of participants within the same setting. The study found a significant reduction in symptoms of depression in transgender individuals after 18 months of starting GAHT, with a more than one-fifth decrease in the number of participants who scores reflected a possible or probable depressive disorder. A statistical reduction of anxiety was not found. While reductions in depression, and to a lesser extent anxiety, were seen, a significant proportion of participants still present, post-treatment, with a possible or probable depressive disorder (25.84%) or anxiety disorder (47.19%). Data from previous studies in the field were used to compare these findings with the general population.<sup>13,16</sup> Acknowledging that direct comparison cannot be made, our study showed that the levels of possible and probable anxiety and depressive disorder after GAHT were still significantly higher than those reported in the general population (4.5% for possible or probable depressive disorder<sup>16</sup> and 34.5% for possible or probable anxiety disorder).<sup>13</sup> Whether these elevated levels will reduce further (when a longer use of GAHT +/- surgical interventions) needs to be explored. Thus, future longitudinal studies would benefit from following people for longer in order to track the longer-term impact of interventions.

These findings do confirm, once again, the high levels of possible anxiety and depressive disorders before GAHT and the benefit

that this treatment brings. It highlights the need to facilitate the expedited use of GAHT to aid the reduction of poor mental health symptoms in the transgender population, when possible and appropriate. This conclusion supports the literature which has called for longitudinal studies such as this to replicate the findings from cross-sectional studies.<sup>15,16,22,25,38,45</sup> The large reduction in depression, comparing to anxiety, may indicate that GAHT targets the dysphoria that many people attending transgender health services present with, which is manifested as depression (rather than anxiety). The fact that many transgender people still feel anxious after GAHT may be because of the victimisation, discrimination and social rejection experienced by the transgender population.<sup>3,5,10,53</sup> Unfortunately for some, these experiences do not necessarily stop after initiation of GAHT, and in some cases, they may increase. Clinical and community services should take these findings into consideration and increase the support offered even after gender-affirming medical treatment is over.

Importantly, this study also highlights how levels of pre-treatment social support are predictive of reduced risk post-treatment, as higher levels of social support prior to receiving GAHT significantly predict a greater reduction in depression symptoms after 18 months of receiving GAHT. This indicates the importance of increasing social support in transgender people.

The study also found that those with higher levels of autistic spectrum condition (ASC) traits prior to the receipt of GAHT had lower reductions in anxiety symptoms. This could indicate that those with greater ASC traits have higher anxiety symptoms even after post-treatment or that higher ASC traits are simply reflective of difficulties in social interactions, as a result of being anxious, grounded in an individual's gender identity status. These findings corroborate cross-sectional studies which have shown that interpersonal interactions can have an impact upon transgender people's psychological well-being.<sup>9,15,23,25,27</sup> However, these results need to be interpreted with caution because of the lack of validity of the AQ-Short in this population.

**TABLE 3** Predictive role of ethnicity, employment status, relationship status, assigned gender at birth, age, MSPSS and AQ-Short for change in HADS-D and HADS-A from T0 to T1

|   | HADS - D |                 |      | HADS - A |               |      |
|---|----------|-----------------|------|----------|---------------|------|
|   | Coef.    | 95% CI          | P    | Coef.    | 95% CI        | P    |
| Ethnicity (grouped as White and all other at T0)  | -0.37    | -3.19 to 2.45   | .794 | -0.67    | -3.62 to 2.28 | .652 |
| Employment (grouped as unemployed and disabled and all other at T0)                         | -0.97    | -2.75 to 0.81   | .284 | -0.51    | -2.37 to 1.35 | .591 |
| Relationship (grouped as single, widowed, divorced/separated and other at T0 and all other) | 0.31     | -1.82 to 2.43   | .776 | -0.61    | -2.83 to 1.62 | .590 |
| Assigned sex at T0  | 0.36     | -1.21 to 1.93   | .651 | -0.17    | -1.81 to 1.47 | .841 |
| Age at T0   | 0.06     | -0.50 to 0.61   | .843 | -0.03    | -0.08 to 0.03 | .409 |
| Mean MSPSS at T0  | 0.81     | 0.24 to 1.39    | .006 | 0.56     | -0.04 - 1.16  | .065 |
| Mean AQ - Short at T0   | -0.04    | -0.096 to 0.025 | .250 | -0.07    | -0.13 to 0.05 | .034 |
| Constant  | -1.93    | -4.76 to 0.90   | .179 | 0.41     | -2.34 to 3.16 | .787 |

With social support (from families, friends or significant others) being highlighted as a factor in depression, it is important to consider approaches to aiding in the building of social support networks for transgender people accessing transgender services. Such support does not necessarily need to be based in clinical settings, as online and offline peer support and the work of tertiary services are available, less stigmatizing and have demonstrated a positive impact.<sup>29,30,54</sup> Resources such as peer support workers (PSWs) and online peer-to-peer support may be a valuable tool to provide social support for transgender individuals awaiting and receiving GAHT. Additional support is recommended in particular to those with higher levels of ASC traits because of the lower reductions in anxiety symptoms found in this population. However, PSWs need to be appropriately trained in order to reduce any potential risks related to managing their own stress as well as power imbalances within peer and professional relationship.<sup>30</sup>

Regarding expedited access to GAHT, there are practical issues that must be considered.<sup>55-57</sup> Many countries either lack clinical services specializing in transgender health care or have significant waiting lists to access these services.<sup>13,58-60</sup> These practical issues surrounding accessing treatment can lead to self-prescribing of GAHT, with 23% of individuals referred to transgender health clinics using GAHT prior to their first appointment, 70% of which was sourced online.<sup>61</sup> Self-prescribing without medical oversight presents its own risks, most notably a lack of specialized knowledge required to minimize health risks.<sup>61</sup>

The issues surrounding access to GAHT may be compounded by how services are configured. The role of many mental health workers in transgender health services in the UK is seen as gatekeepers, focused on the assessment of transgender people with limited attention given to psychological support. A shift in roles from gatekeeping to tangible mental health support would allow for a partial addressing of the power imbalance between transgender people and mental health professionals. This would potentially allow for transgender people to feel more able to discuss issues without fear of rejection for treatment. This in turn would provide mental health professionals with the ability to focus more on supporting the mental health of those transgender people who need it. From the evidence provided by this study, it appears that this would be most important for those with low levels of social support and those with autistic traits.

The strengths of this paper are the large sample and the naturalistic prospective longitudinal design within one transgender health service, which allows for within-subject comparisons and so provides a highly valid insight into the impact of GAHT on depression and anxiety of treatment-seeking transgender people. It is important to add that this is a national NHS service, which offers gender-affirming medical interventions and assessment free at the point of access to people from different geographical regions within the UK. This study is one of the few in the literature currently addressing the role of GAHT on mental health with such a methodology.

There are, however, limitations to the study. We must acknowledge that participants may downplay their symptoms of depression and anxiety pre-assessment, for fear of not being treated.

Consequently, participants may have attenuated their mental health symptoms because of concern of not being accepted for GAMT. This could indicate that the changes between T0 and T1 may be more significant than recorded. As the method of recruitment was through a transgender health clinic as part of the NHS, the findings are only generalizable to treatment-seeking transgender individuals. A limitation may also be the response rate. Although this is low, it is in line with other clinical studies and it may be a reflection of the unfunded nature of the study. Another limitation is that the sample consisted of predominantly white participants, which may explain why ethnicity was not a predictive factor and the lack of control group where the intervention (GAHT) is not being offered (eg waiting list), but this will have its limitations too. A full randomized control study within this area will not be ethically possible. Studies using data from a clinical setting must also be aware of the context in which their data are being gathered.

In conclusion, this study shows that the mental health of transgender individuals improves following GAHT, particularly for those who reported high levels of social support prior to receiving GAHT. These results highlight the important mental health benefits of GAHT and emphasises the need for interventions focused on developing social support for transgender individuals.

## ACKNOWLEDGEMENTS

The authors thank all participants in the study for having given their time so generously.

## CONFLICT OF INTEREST

The authors have no other conflicts of interest relevant to this article to disclose. All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

## AUTHORS' CONTRIBUTIONS

JA conceptualized the study and participated in the study design. ZA, SP, JA, WPB and BG handled data collection management. BG carried out the statistical analysis. ZA drafted the initial manuscript. SP drafted the initial methods and results section of the manuscript. All authors contributed to the critical review of the manuscript and approved the final version of the manuscript.

## ORCID

Jon Arcelus  <https://orcid.org/0000-0002-3805-0180>

## REFERENCES

1. Birkett M, Newcomb ME, Mustanski B. Does it get better? A longitudinal analysis of psychological distress and victimization in lesbian, gay, bisexual, transgender, and questioning youth. *J Adolesc Health*. 2015;56(3):280-285.
2. Bouman WP, Arcelus J. *The Transgender Handbook: A Guide for Transgender People, Their Families, and Professionals*. New York: Nova Publishers; 2017.

3. Breslow AS, Brewster ME, Velez BL, Wong S, Geiger E, Soderstrom B. Resilience and collective action: Exploring buffers against minority stress for transgender individuals. *Psychol Sex Orientation Gender Div*. 2015;2(3):253-265.
4. Reisner SL, Vetteser R, Leclerc M, et al. Mental health of transgender youth in care at an adolescent urban community health center: a matched retrospective cohort study. *J Adolesc Health*. 2015;56(3):274-279.
5. Carmel TC, Erickson-Schroth L. Mental health and the transgender population. *J Psychosoc Nurs Ment Health Serv*. 2016;54(12):44-48.
6. Claes L, Bouman WP, Witcomb G, Thurston M, Fernandez-Aranda F, Arcelus J. Non-suicidal self-injury in trans people: associations with psychological symptoms, victimization, interpersonal functioning, and perceived social support. *J Sex Med*. 2015;12(1):168-179.
7. Dhejne C, Van Vlerken R, Heylens G, Arcelus J. Mental health and gender dysphoria: A review of the literature. *Int Rev Psychiatry*. 2016;28(1):44-57.
8. Seelman KL, Colon-Diaz MJP, LeCroix RH, Xavier-Brier M, Kattari L. Transgender noninclusive healthcare and delaying care because of fear: connections to general health and mental health among transgender adults. *Transgender Health*. 2017;2(1):17-28.
9. Simons L, Schrage SM, Clark LF, Belzer M, Olson J. Parental support and mental health among transgender adolescents. *J Adolesc Health*. 2013;53(6):791-793.
10. Herman JL. Gendered Restrooms And Minority Stress: The Public Regulation of Gender and its Impact on Transgender People's Lives. *Journal of Public Management & Social Policy*. 2013;19(1):65-80.
11. Meyer IH. Prejudice, social stress, and mental health in lesbian, gay, and bisexual populations: conceptual issues and research evidence. *Psychol Bull*. 2003;129(5):674-697.
12. Meyer IH, Schwartz S, Frost DM. Social patterning of stress and coping: does disadvantaged social statuses confer more stress and fewer coping resources? *Soc Sci Med*. 2008;67(3):368-379.
13. Bouman WP, Claes L, Brewin N, et al. Transgender and anxiety: a comparative study between transgender people and the general population. *Int J Transgenderism*. 2017;18(1):16-26.
14. Millet N, Longworth J, Arcelus J. Prevalence of anxiety symptoms and disorders in the transgender population: A systematic review of the literature. *Int J Transgenderism*. 2017;18(1):27-38.
15. Bouman WP, Davey A, Meyer C, Witcomb GL, Arcelus J. Predictors of psychological well-being among treatment seeking transgender individuals. *Sex Relat Ther*. 2016;31(3):359-375.
16. Witcomb GL, Bouman WP, Claes L, Brewin N, Crawford JR, Arcelus J. Levels of depression in transgender people and its predictors: results of a large matched control study with transgender people accessing clinical services. *J Affect Disord*. 2018;235:308-315.
17. Veale JF, Watson RJ, Peter T, Saewyc EM. Mental health disparities among Canadian transgender youth. *J Adolesc Health*. 2017;60(1):44-49.
18. Marshall E, Claes L, Bouman WP, Witcomb GL, Arcelus J. Non-suicidal self-injury and suicidality in trans people: A systematic review of the literature. *Int Rev Psychiatry*. 2016;28(1):58-69.
19. Butler C, Joiner R, Bradley R, et al. Self-harm prevalence and ideation in a community sample of cis, trans and other youth. *Int J Transgenderism*. 2019;20(4):447-458.
20. Davey A, Arcelus J, Meyer C, Bouman WP. Self-injury among trans individuals in transition and matched controls: Prevalence and associated factors. *Health Soc Care Community*. 2016;24:485-494.
21. Nobili A, Glazebrook C, Bouman WP, et al. Autistic traits in treatment-seeking transgender adults. *J Autism Dev Dis*. 2018;48(12):3984-3994.
22. Nobili A, Glazebrook C, Bouman WP, Baron-Cohen S, Allison C, Arcelus J. The stability of autistic traits in transgender adults following cross-sex hormone treatment. *Int J Transgender Health*. 2020;48(12):3984-3994.
23. Meier CSL, Fitzgerald KM, Pardo ST, Babcock J. The effects of hormonal gender affirmation treatment on mental health in female-to-male transsexuals. *J Gay Lesbian Mental Health*. 2011;15(3):281-299.
24. Budge SL, Adelson JL, Howard KAS. Anxiety and depression in transgender individuals: the roles of transition status, loss, social support, and coping. *J Consult Clin Psychol*. 2013;81(3):545-557.
25. Davey A, Bouman WP, Arcelus J, Meyer C. Social support and psychological wellbeing: a comparison of patients with gender dysphoria and matched controls. *J Sex Med*. 2014;1(12):2976-2985.
26. Hoffman B. An overview of depression among transgender women. *Depress Res Treat*. 2014;2014:394283.
27. Olson KR, Durwood L, DeMeules M, McLaughlin KA. Mental health of transgender children who are supported in their identities. *Pediatrics*. 2016;137(3):e20153223.
28. Naslund JA, Aschbrenner KA, Marsch LA, Bartels SJ. The future of mental health care: peer-to-peer support and social media. *Epidemiol Psychiatr Sci*. 2016;25(2):113-122.
29. Repper J, Carter T. A review of the literature on peer support in mental health services. *J Ment Health*. 2011;20(4):392-411.
30. Jefferson K, Neilands TB, Sevelius J. Transgender women of color: discrimination and depression symptoms. *Ethn and Inequal Health Soc Care*. 2013;6(4):121-136.
31. Dooley D, Catalano R, Wilson G. Depression and unemployment: panel findings from the Epidemiologic Catchment Area study. *Am J Community Psychol*. 1994;22(6):745-765.
32. Leppel K. The labor force status of transgender men and women. *Int J Transgenderism*. 2016;17(3-4):155-164.
33. Mizock L, Mueser KT. Employment, mental health, internalized stigma, and coping with transphobia among transgender individuals. *Psychol Sex Orientation Gender Div*. 2014;1(2):146-158.
34. Mizock L, Woodrum TD, Riley J, Sotillo EA, Yuen N, Ormerod AJ. Coping with transphobia in employment: strategies used by transgender and gender-diverse people in the United States. *Int J Transgenderism*. 2017;18(3):282-292.
35. Witcomb GL, Claes L, Bouman WP, Nixon E, Motmans J, Arcelus J. Experiences and psychological wellbeing outcomes associated with bullying in treatment-seeking transgender and gender-diverse youth. *LGBT Health*. 2019;6(5):216-226.
36. Arcelus J, Claes L, Witcomb GL, Marshall E, Bouman WP. Risk factors for non-suicidal self-injury among trans youth. *J Sex Med*. 2016;13(3):402-412.
37. Colizzi M, Costa R, Todarello O. Transsexual patients' psychiatric comorbidity and positive effect of cross-sex hormonal treatment on mental health: results from a longitudinal study. *Psychoneuroendocrinology*. 2014;39(1):65-73.
38. Heylens G, Verroken C, De Cock S, T'Sjoen G, De Cuypere G. Effects of different steps in gender reassignment therapy on psychopathology: A prospective study of persons with a gender identity disorder. *J Sex Med*. 2014;11(1):119-126.
39. Bränström R, Pachankis JE. Reduction in mental health treatment utilization among transgender individuals after gender-affirming surgeries: a total population study. *Am J Psychiatry*. 2019.
40. Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand*. 1983;67(6):361-370.
41. Bjelland I, Dahl AA, Haug TT, Neckelmann D. The validity of the hospital anxiety and depression scale. An updated literature review. *J Psychosom Res*. 2002;52(2):69-77.
42. Gómez-Gil E, Zubiaurre-Elorza L, Esteva I, et al. Hormone-treated transsexual report less social distress, anxiety, and depression. *Psychoneuroendocrinology*. 2012;37(5):662-670.
43. Zimet GD, Powell SS, Farley GK, Werkman S, Berkoff KA. Psychometric characteristics of the Multidimensional Scale of Perceived Social Support. *J Pers Assess*. 1990;55:610-617.



44. Bouman WP, Claes L, Marshall E, et al. Socio-demographic variables, clinical features and the role of pre-assessment cross-sex hormones in older trans people. *J Sex Med.* 2016;13(4):711-719.
45. Hoekstra RA, Vinkhuyzen AAE, Wheelwright S, et al. The construction and validation of an abridged version of the Autism-Spectrum Quotient (AQ-Short). *J Autism Dev Disord.* 2011;41(5):589-596.
46. Woodbury-Smith MR, Robinson J, Wheelwright S, Baron-Cohen S. Screening adults for Asperger Syndrome using the AQ: a preliminary study of its diagnostic validity in clinical practice. *J Autism Dev Disord.* 2005;35(3):331-335.
47. Baron-Cohen S, Wheelwright S, Skinner R, Martin J, Clubley E. The Autism-Spectrum Quotient (AQ): Evidence from Asperger Syndrome/High-Functioning Autism, Males and Females, Scientists and Mathematicians. *J Autism Dev Disord.* 2001;31(1):5-17.
48. Pasterski V, Gilligan L, Curtis R. Traits of autism spectrum disorders in adults with gender dysphoria. *Arch Sex Behavior.* 2014;43(2):387-393.
49. StataCorp *Stata Statistical Software: Release 16.* College Station, TX: StataCorp LLC; 2019.
50. Allison PD. Comparing logit and probit coefficients across groups. *Sociol Methods & Res.* 1999;28(2):186-208.
51. Health Research Authority (HRA). *Guidance for NHS Research Studies.* London: Health Research Authority; 2013.
52. Fuller KA, Riggs DW. Family support and discrimination and their relationship to psychological distress and resilience amongst transgender people. A systematic review. *Int J Transgenderism.* 2018;19(4):379-388.
53. Bockting WO, Miner MH, Swinburne Romine RE, Hamilton A, Coleman E. Stigma, mental health, and resilience in an online sample of the US transgender population. *Am J Public Health.* 2013;103(5):943-951.
54. Bouman WP, de Vries ALC, T'Sjoen G. Gender Dysphoria and Gender Incongruence: An evolving inter-disciplinary field. *Int Rev Psychiatry.* 2016;28(1):1-4.
55. Motmans J, Nieder TO, Bouman WP. Transforming the paradigm of nonbinary transgender health: A field in transition. *Int J Transgenderism.* 2019;20(2-3):118-124.
56. Richards C, Arcelus J, Barrett J, et al. Trans is not a disorder – but should still receive funding. *Sex Relats Ther.* 2015;30(3):309-313.
57. Mepham N, Bouman WP, Arcelus J, Hayter M, Wylie KR. People with gender dysphoria who self-prescribe cross-sex hormones: prevalence, sources, and side effects knowledge. *J Sex Med.* 2014;11(12):2995-3001.
58. Bergero-Miguel T, Garcia-Encinas MA, Villena-Jimena A, et al. Gender Dysphoria and social anxiety: An exploratory study in Spain. *J Sex Med.* 2016;13(8):1270-1278.
59. Bouman WP, Richards C. Diagnostic and treatment issues for people with gender dysphoria in the United Kingdom. *Sex Relats Ther.* 2013;28(3):165-171.
60. Davies A, Bouman WP, Richards C, et al. Patient satisfaction with gender identity clinic services in the United Kingdom. *Sex Relats Ther.* 2013;28(4):400-418.
61. Winter S, Diamond M, Green J, et al. Transgender people: health at the margins of society. *Lancet.* 2016;388(10042):390-400.

**How to cite this article:** Aldridge Z, Patel S, Guo B, et al. Long-term effect of gender-affirming hormone treatment on depression and anxiety symptoms in transgender people: A prospective cohort study. *Andrology.* 2020;00:1-9. <https://doi.org/10.1111/andr.12884>