



City Research Online

City, University of London Institutional Repository

Citation: Fring, A. ORCID: 0000-0002-7896-7161 and Taira, T. (2020). Complex BPS solitons with real energies from duality. *Journal of Physics A: Mathematical and Theoretical*, 53, 455701.. doi: 10.1088/1751-8121/abb92a

This is the accepted version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/24987/>

Link to published version: <http://dx.doi.org/10.1088/1751-8121/abb92a>

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

City Research Online:

<http://openaccess.city.ac.uk/>

publications@city.ac.uk

Complex BPS solitons with real energies from duality

Andreas Fring and Takanobu Taira

Department of Mathematics, City, University of London,

Northampton Square, London EC1V 0HB, UK

E-mail: a.fring@city.ac.uk, takanobu.taira@city.ac.uk

ABSTRACT: Following a generic approach that leads to Bogomolny-Prasad-Sommerfield (BPS) soliton solutions by imposing self-duality, we investigate three different types of non-Hermitian field theories. We consider a complex version of a logarithmic potential that possess BPS super-exponential kink and antikink solutions and two different types of complex generalisations of systems of coupled sine-Gordon models with kink and antikink solution of complex versions of arctan type. Despite the fact that all soliton solutions obtained in this manner are complex in the non-Hermitian theories we show that they possess real energies. For the complex extended sine-Gordon model we establish explicitly that the energies are the same as those in an equivalent pair of a non-Hermitian and Hermitian theory obtained from a pseudo-Hermitian approach by means of a Dyson map. We argue that the reality of the energy is due to the topological properties of the complex BPS solutions. These properties result in general from modified versions of antilinear CPT symmetries that relate self-dual and an anti-self-dual theories.

1. Introduction

T'Hooft [1] and Polyakov [2] established more than 45 years ago that gauge theories almost inevitably contain monopole solutions. The corresponding soliton solutions that interpolate between different vacua of the theory are usually constructed explicitly by means of Bogomolny-Prasad-Sommerfield (BPS) [3, 4] multiple scaling limits. The validity of these limits can be justified on physical grounds when assuming that certain mass ratios in the theory are very small. We have recently demonstrated [5] that such type of solutions can also be constructed in certain domains of the parameter space of a non-Hermitian field theory with local non-Abelian $SU(2)$ gauge symmetry and a modified antilinear CPT -symmetry.

Here we investigate the properties of complex soliton solutions resulting in a general setting of BPS theories and in particular show that the reality of their energies are attributed to a modified version of a CPT -symmetry that relates a self-dual to an anti-self-dual theory and governs their topological properties. The underlying reason that ensures the reality of

the energy is slightly different to what has been observed previously for integrable complex nonlinear equations, such as for instance complex versions of the Korteweg-de Vries equation or Calogero-Sutherland-Moser systems [6, 7, 8, 9, 10, 11, 12, 13]. The governing equations to be solved have additional specific structures that need to be respected by the \mathcal{CPT} -symmetry. First of all, the soliton solutions solve the BPS equations, that are by construction of lower order than the equations of motion. Moreover, while the soliton solutions studied in [10, 12, 13] all vanish asymptotically, the soliton solutions that maybe associated to magnetic monopoles are of kink or antikink type with nontrivial asymptotic behaviour. It is this latter topological behaviour that completely governs the energy, which is bounded from below by the topological charge of the theory, the Bogomolny bound. Crucially in our approach is that the BPS equations occur in pairs involving self-dual and anti-self-dual functions of the fields their first order derivatives that have the same energy. The modified versions of the \mathcal{CPT} -symmetry relate the solutions of these two pairs of equations.

One may approach the study of BPS systems in several alternative ways. The original and most direct way is to investigate a concrete full-fledged gauge theory and carry out the appropriate limits, see [5] for a non-Hermitian system. Alternatively one can take the above mentioned general properties as the defining relations for a BPS theory and derive them in a simpler setting as was shown for instance in [14, 15, 16, 17]. While the discussion in [14] is generic for any dimension we restrict our considerations here to complex scalar field theories in two dimensions described by Lagrangians of the general form

$$\mathcal{L} = \frac{1}{2} \eta_{ab} \partial_\mu \phi_a \partial^\mu \phi_b - \mathcal{V}(\phi), \quad (1.1)$$

where η_{ab} is a target space metric, the metric g in space-time is taken to be Lorentzian $\text{diag } g = (1, -1)$ and the potential $V(\phi)$ depends on the complex scalar field components ϕ_a , $a = 1, \dots, \ell$.

We present three different types of systems purposely chosen to illustrate different types of features. For all models we construct the explicit complex BPS kink and antikink soliton solutions, we identify the different versions of the modified antilinear \mathcal{CPT} -symmetry that can be used to argue that the corresponding energy is real in certain regimes of the parameter space. The models exhibit different types of symmetry breaking and appear to possess exceptional points in their energy spectrum. However, we demonstrate that none of these points is a genuine exceptional point [18] in the standard sense of non-Hermitian theories [19, 20]. We study the stability of the vacua and identify the explicit soliton solutions that interpolate between them. Our approach to analyse directly the non-Hermitian system is justified further in section 5 where we present an explicit example of a pair of a non-Hermitian and a Hermitian Hamiltonian that are related by a nontrivial Dyson map and show that the energy of the two systems is identical in the well-defined \mathcal{CPT} -symmetric regime of the parameter space.

Our manuscript is organized as follows: In section 2 we present the general set up for the study of BPS solitons from the requirement that the theory contains self-dual or anti-self-dual functionals of the fields and their derivatives. In section 3, 4 and 5 we investigate three different types of models in the way described above and in section 6 we state our conclusions.

2. BPS solitons from self-duality and anti-self-duality

The authors in [14] take an energy functional E and a topological charge Q of the form

$$E = \frac{1}{2} \int d^2x \left(A_\alpha^2 + \tilde{A}_\alpha^2 \right), \quad \text{and} \quad Q = \int d^2x A_\alpha \tilde{A}_\alpha, \quad (2.1)$$

as a starting point for the setup of a BPS theory, where the quantities $A_\alpha(\phi, \partial_\mu\phi)$, $\tilde{A}_\alpha(\phi, \partial_\mu\phi)$ are functions of the fields ϕ appearing in the Lagrangian \mathcal{L} of the field theory under consideration and at most of first order derivatives thereof. It is clear that the relations in (2.1) ensure that the topological charge is always a lower bound for the energy $E \geq |Q|$. Following [14], one may then use these definitions to derive two equations, one being the Euler-Lagrange equation resulting from varying E and the other from considering infinitesimal changes $\delta\phi$ in Q and demanding $\delta Q = 0$. The latter requirement incorporates that Q is interpreted as a topological charge, which should be a homotopy invariant, i.e. invariant under smooth variations in the fields. The compatibility between these two equations then implies (anti)-self-duality of the quantities A_α , \tilde{A}_α and moreover that Q saturates the Bogomolny bound for the energy E

$$A_\alpha = \pm \tilde{A}_\alpha, \quad \text{and} \quad E = |Q|. \quad (2.2)$$

Evidently the energies of the self-dual and anti-self-dual fields are the same. Assuming next the existence of a pre-potential $U(\phi)$, that is a function of the fields in the theory only, one may write the energy functional and the topological charge for the static solutions as

$$E = \int_{-\infty}^{\infty} dx \left(\frac{1}{2} \eta_{ab} \partial_\mu \phi_a \partial^\mu \phi_b + \mathcal{V}(\phi) \right) = \frac{1}{2} \int_{-\infty}^{\infty} dx \left(\eta_{ab} \partial_\mu \phi_a \partial^\mu \phi_b + \eta_{ab}^{-1} \frac{\partial U}{\partial \phi_a} \frac{\partial U}{\partial \phi_b} \right), \quad (2.3)$$

$$Q = \int_{-\infty}^{\infty} dx \frac{\partial U}{\partial x} = \int_{-\infty}^{\infty} dx \frac{\partial U}{\partial \phi_a} \partial_x \phi_a = \lim_{x \rightarrow \infty} U[\phi(x)] - \lim_{x \rightarrow -\infty} U[\phi(x)]. \quad (2.4)$$

Comparing the general expressions for A_α and \tilde{A}_α in (2.1) with those for $U(\phi)$ in (2.3), (2.4) implies the identifications

$$A_a = \rho_{ab} \partial_x \phi_b, \quad \text{and} \quad \tilde{A}_a = \frac{\partial U}{\partial \phi_b} \rho_{ba}^{-1}, \quad (2.5)$$

where ρ factorizes the target space metric as $\rho^T \rho = \eta$. The (anti)-self-duality relations in (2.2), then become equivalent to the pair of BPS equations in the form

$$\partial_x \phi_b = \pm \eta_{ab}^{-1} \frac{\partial U}{\partial \phi_b}. \quad (2.6)$$

Allowing the scalar fields to be complex and the potential to be non-Hermitian, the reality of the energy could be guaranteed when the Hamiltonian is \mathcal{CPT} -symmetric satisfying $\mathcal{H}[\phi(x)] = \mathcal{H}^\dagger[\phi(-x)]$ by employing the same argument as in [7]

$$E = \int_{-\infty}^{\infty} dx \mathcal{H}[\phi(x)] = - \int_{\infty}^{-\infty} dx \mathcal{H}[\phi(-x)] = \int_{-\infty}^{\infty} dx \mathcal{H}^\dagger[\phi(x)] = E^*. \quad (2.7)$$

Since in the scenario considered here the self-duality imposes the kinetic energy to equal the potential energy, it would suffice therefore to establish that

$$\mathcal{V}[\phi_{\pm}(x)] = \mathcal{V}^{\dagger}[\phi_{\pm}(-x)], \quad \text{or} \quad \mathcal{V}[\phi_{\pm}(x)] = \mathcal{V}^{\dagger}[\phi_{\mp}(-x)] \quad (2.8)$$

in order to ensure the reality of the energy by means of (2.7). We have denoted here by ϕ_{\pm} the solutions of (2.6) corresponding to the two options for the sign in (2.2). Evidently it follows from (2.1) that the energy is the same for either choice. The second option in (2.8) is novel due to the set up involving anti-self-duality and not available in the standard setting of integrable systems [7, 10, 12, 13]. We shall demonstrate below that actually this novel option of the \mathcal{CPT} -symmetry is the guarand for the reality of the energy for the systems considered. Evidently, for a direct analysis it is clear that the energy is real if

$$\lim_{x \rightarrow \infty} \text{Im} \{U[\phi(x)]\} = \lim_{x \rightarrow -\infty} \text{Im} \{U[\phi(x)]\}. \quad (2.9)$$

We shall now analyse several different theories with concrete choices for pre-potential that lead to non-Hermitian scalar field theory with an antilinear symmetry. We shall demonstrate that the first version of the \mathcal{CPT} -symmetry in (2.8) is in fact broken, but the second version can be realised by the various solutions in our examples.

3. A non-Hermitian BPS theory with super-exponential kink solutions

We start by generalizing a Hermitian one field theory that was recently studied by Kumar, Khare and Saxena [21] to one with two component complex fields in a non-Hermitian setting. The original model was motivated in parts by its proximity to a ϕ^6 -type potential and its feature of minimal nonlinearity. A very interesting aspect of this model is that it possesses kink and antikink solutions with a super-exponential profile rather than the more standard arctan type solutions. This feature survives our generalization and moreover the complex BPS solutions interpolating between five out of nine vacua of our model have real energies.

To set up the field theory we choose the target space metric and the pre-potential as

$$\eta = \begin{pmatrix} 1 & -i\lambda \\ -i\lambda & 1 \end{pmatrix}, \quad \text{and} \quad U(\phi_1, \phi_2) = \frac{\mu_1}{2} \phi_1^2 \ln(\phi_1^2) + \frac{\mu_2}{2} \phi_2^2 \ln(\phi_2^2), \quad \lambda, \mu_1, \mu_2 \in \mathbb{R}, \quad (3.1)$$

respectively. Using the relation between the potential and the pre-potential (2.3) we obtain from the Ansatz (3.1) the non-Hermitian potential

$$\mathcal{V}(\phi_1, \phi_2) = \frac{1}{1 + \lambda^2} \sum_{i=1}^2 \frac{\mu_i^2}{2} [\phi_i + \phi_i \ln(\phi_i^2)]^2 + i \frac{\lambda}{1 + \lambda^2} \prod_{i=1}^2 \mu_i [\phi_i + \phi_i \ln(\phi_i^2)]. \quad (3.2)$$

According to the standard pseudo-Hermitian approach to non-Hermitian field theories one may seek a similarity transformation by means of a well defined Dyson map, e.g. [22, 23, 24, 25, 26], to map the theory to a Hermitian theory or introduce non-vanishing surface terms [27, 28, 29, 30, 31, 32] and analyse these systems. However, as we shall demonstrate

below, just as in a standard quantum mechanical setting [19, 20], the energy is preserved in this process so that one may also analyse the solutions of the non-Hermitian theory directly. Our approach is further justified in section 5 where we shall present an explicit system for which a non-Hermitian Hamiltonian is related to a Hermitian Hamiltonian by means of an explicit nontrivial Dyson map.

Using the BPS equations (2.6), the static solutions associated to the potential (3.2) are the two pairs of coupled first order differential equations

$$BPS_1^\pm : \quad \partial_x \phi_1 = \pm \frac{\mu_1 [\phi_1 + \phi_1 \ln(\phi_1^2)]}{\lambda^2 + 1} \pm i\lambda \frac{\mu_2 [\phi_2 + \phi_2 \ln(\phi_2^2)]}{\lambda^2 + 1} =: F_1^\pm, \quad (3.3)$$

$$BPS_2^\pm : \quad \partial_x \phi_2 = \pm i\lambda \frac{\mu_1 [\phi_1 + \phi_1 \ln(\phi_1^2)]}{\lambda^2 + 1} \pm \frac{\mu_2 [\phi_2 + \phi_2 \ln(\phi_2^2)]}{\lambda^2 + 1} =: F_2^\pm. \quad (3.4)$$

We will need both versions in (3.3) and (3.4) to verify the general argument that guarantees the reality of the energy. We observe that these equations are compatible under two types of modified \mathcal{CPT} -transformations

$$\mathcal{CPT}_\pm : \quad \phi_1(x) \rightarrow \pm [\phi_1(-x)]^\dagger, \quad \phi_2(x) \rightarrow \mp [\phi_2(-x)]^\dagger, \quad \Leftrightarrow \quad BPS_i^\pm \rightarrow (BPS_i^\mp)^*. \quad (3.5)$$

Using these symmetries we can derive the second relation in (2.8). We notice that a modified \mathcal{CT} -transformation $\phi_1(x) \rightarrow -[\phi_1(x)]^\dagger$, $\phi_2(x) \rightarrow -[\phi_2(x)]^\dagger$ is achieving the compatibility $BPS_i^\pm \rightarrow (BPS_i^\pm)^*$. However, this symmetry can not be employed in the argument in (2.7) that guarantees the reality of the energy. The introduction of time by means of a standard Lorentz transformation, $x \rightarrow (x - vt)/\sqrt{1 - v^2}$, will not change this feature, so that the reality of the energy is not a consequence of this particular antilinear symmetry. Moreover, we do not find solutions below that posses this kind of \mathcal{CT} -symmetry.

Let us now solve the pair of the two BPS equations (3.3) and (3.4). In the Hermitian limit, when $\lambda = 0$, the equations decouple and the solutions can be obtained in an explicit analytical form as double exponentials

$$\phi_i(x) = \exp\left(-\frac{1}{2} + \frac{1}{2}e^{2(\mu_i x + \kappa_i)}\right), \quad (3.6)$$

with integration constants $\kappa_i \in \mathbb{C}$ and $i = 1, 2$. We fix our constants in such a way that we obtain proper kink and antikink solutions with well-defined asymptotic behaviour. We select our solutions as

$$\phi_i^{a+}(x) = \exp\left(-\frac{1}{2} - \frac{1}{2}e^{2\mu_i x}\right), \quad \phi_i^{k+}(x) = -\exp\left(-\frac{1}{2} - \frac{1}{2}e^{2\mu_i x}\right), \quad \mu_i \geq 0, \quad (3.7)$$

$$\phi_i^{k-}(x) = \exp\left(-\frac{1}{2} - \frac{1}{2}e^{2\mu_i x}\right), \quad \phi_i^{a-}(x) = -\exp\left(-\frac{1}{2} - \frac{1}{2}e^{2\mu_i x}\right), \quad \mu_i < 0, \quad (3.8)$$

so that $\phi_i^{a+}(0) = \phi_i^{k-}(0) = 1/e$, $\phi_i^{k+}(0) = \phi_i^{a-}(0) = -1/e$ and $\phi_i^{a+}(x) = \phi_i^{k-}(-x) = -\phi_i^{k+}(x) = -\phi_i^{a-}(-x)$. The asymptotic limits are therefore

$$\lim_{x \rightarrow -\infty} \phi_i^{a+}(x) = \lim_{x \rightarrow \infty} \phi_i^{k-}(x) = \frac{1}{\sqrt{e}}, \quad \lim_{x \rightarrow -\infty} \phi_i^{k+}(x) = \lim_{x \rightarrow \infty} \phi_i^{a-}(x) = -\frac{1}{\sqrt{e}}, \quad (3.9)$$

$$\lim_{x \rightarrow \infty} \phi_i^{a+}(x) = \lim_{x \rightarrow \infty} \phi_i^{k+}(x) = \lim_{x \rightarrow -\infty} \phi_i^{a-}(x) = \lim_{x \rightarrow -\infty} \phi_i^{k-}(x) = 0. \quad (3.10)$$

Hence, using the expression for the pre-potential (2.4) we obtain for all combinations the same real energy as function of μ_1, μ_2

$$E^{\phi_1^{pn}, \phi_2^{qm}}(\mu_1, \mu_2) = \frac{|\mu_1| + |\mu_2|}{2e}, \quad p, q = k, a; \quad n, m = \pm; \quad \mu_1, \mu_2 \in \mathbb{R}. \quad (3.11)$$

In the non-Hermitian scenario, when $\lambda \neq 0$, we solve the two sets of coupled BPS equations (3.3) and (3.4) numerically. Some sample computations are presented in figure 1.




Figure 1: Complex BPS kink and antikink solutions of the two pairs of coupled BPS equations (3.3) and (3.4) associated to the potential (3.2) with initial values $\phi_1^{k+}(0) = \phi_2^{k+}(0) = \phi_1^{a-}(0) = \phi_2^{a-}(0) = -1/e$ and $\phi_1^{a+}(0) = \phi_2^{a+}(0) = \phi_1^{k-}(0) = \phi_2^{k-}(0) = 1/e$ for $\mu_1 = 0.2, \mu_2 = 0.1, \lambda = 0.1$.

We observe that for increasing values of the coupling constants μ_i the real parts of ϕ_i approach $H(-x)/\sqrt{e}$ with $H(x)$ denoting the Heaviside step function. The imaginary parts keep oscillating with increased amplitudes and crucially vanish at $x \rightarrow \pm\infty$, which means that the energy is given by the expression in (3.11) for all values of λ . The analytical solution for $\lambda = 0$ are smooth kinks and antikinks who also approach a Heaviside step function for increasing values of μ_i .

We also observe from our numerical solutions in figure 1 that the solutions realize the \mathcal{CPT}_- -symmetry as

$$\phi_1^{k+}(x) = -[\phi_1^{k-}(-x)]^\dagger, \quad \phi_1^{a+}(x) = -[\phi_1^{a-}(-x)]^\dagger, \quad \phi_2^{k\pm}(x) = [\phi_2^{a\mp}(-x)]^\dagger. \quad (3.12)$$

Using now the properties of the kink and antikink solutions (3.12) we derive for the potential

$$\mathcal{V}_\lambda [\phi_1^{k+}(x), \phi_2^{k+}(x)] = \mathcal{V}_\lambda \left\{ -[\phi_1^{k-}(-x)]^\dagger, [\phi_2^{a-}(-x)]^\dagger \right\} = \mathcal{V}_\lambda^\dagger \left\{ [\phi_1^{k-}(-x)], [\phi_2^{a-}(-x)] \right\}, \quad (3.13)$$

and similarly for the others pairs of solutions. Changing the initial conditions we may also construct solutions that manifest the \mathcal{CPT}_+ -symmetry. The relation in (3.13) is precisely the second option in (2.8) that relates solutions of the self-dual system to solutions of the anti-self-dual system. As the energies in both systems must be the same it is guaranteed to be real.

Next we will identify which vacua are interpolated by which kind of BPS solution. It is easy to check that the real part of the potential has nine minima at

$$v^{\pm\pm} = (\pm e^{-1/2}, \pm e^{-1/2}), \quad v^{0\pm} = (0, \pm e^{-1/2}), \quad v^{\pm 0} = (\pm e^{-1/2}, 0), \quad v^{00} = (0, 0), \quad (3.14)$$

corresponding to the fixed points of the dynamical system (3.3) and (3.4) as solutions of $F_1^\pm(\phi_1, \phi_2) = F_2^\pm(\phi_1, \phi_2) = 0$. Next we compute the eigenvalues of the Jacobian matrix at these fixed points

$$J = \left(\begin{array}{cc} \partial_{\phi_1} F_1^\pm & \partial_{\phi_2} F_1^\pm \\ \partial_{\phi_1} F_2^\pm & \partial_{\phi_2} F_2^\pm \end{array} \right) \Big|_{v^{i,j}}, \quad i, j = 0, +, -, \quad (3.15)$$

in order to determine their stability. For the F^+ -system with $\mu_i > 0$ we find that $J(v^{\pm\pm})$ has two positive eigenvalues, $J(v^{00})$ has two negative eigenvalues and $J(v^{0\pm})$, $J(v^{\pm 0})$ have a positive and a negative eigenvalue. For $\phi_i \rightarrow 0$ we have to evaluate the values in an ε -neighbourhood. This means, see e.g. [33], that $v^{\pm\pm}$ are unstable fixed points, $v^{0\pm}$ and $v^{\pm 0}$ are saddle points and v^{00} is the only stable fixed point. For the F^- -system still with $\mu_i > 0$ all signs of the eigenvalues are reversed. Changing the sign of μ_i will also reverse the sign of one eigenvalue. Using the solutions from above as represented in figure 1, we have the following interpolations between the different vacua

$$v^{--} \xrightarrow{\phi_1^{k+} \phi_2^{k+}} v^{00}, \quad v^{00} \xrightarrow{\phi_1^{a-} \phi_2^{k-}} v^{-+}, \quad v^{00} \xrightarrow{\phi_1^{k-} \phi_2^{a-}} v^{+-}, \quad v^{++} \xrightarrow{\phi_1^{a+} \phi_2^{a+}} v^{00}. \quad (3.16)$$

This behaviour is also confirmed by the gradient flow for F^+ that is indicated in figure 2 superimposed onto the potential. We obtain similar relations for the F^- -system.




Figure 2: Real part of the potential $\mathcal{V}(\phi_1, \phi_2)$ in (3.2) as a function of $\text{Re} \phi_1$ and $\text{Re} \phi_2$ with the gradient flow of the real parts of F^+ superimposed in white. The kink-kink, kink-antikink, antikink-kink and antikink-antikink interpolate between the different types of stable and unstable vacua as specified in (3.16)

When passing from the \mathcal{V}_+ -theory to the \mathcal{V}_- -theory we pass through the special point $\mu_1 = \mu_2 = 0$. The energy (3.11) is defined for all values and does not become complex. To

investigate this point further the next model is designed in such a way that it appears to have an exceptional point, which, however, turns out to be not genuine.

4. A non-Hermitian coupled sine-Gordon model

Next we consider a modified version of a model whose real variant has been investigated recently in [15]. We generalize that model to one involving a complex non-Hermitian potential with a complex two-component scalar field and add an additional term designed in such a way that we apparently obtain an exceptional point [18]. We shall demonstrate that the system possesses complex solutions to its BPS equations with real energies in a certain region in the parameter space where the topological charge of the system is well-defined and real. There is also a region in which the energy is not well defined and not finite on the entire real x -axis.

Choosing the target space metric and the pre-potential as

$$\eta = \begin{pmatrix} 1 & -i\lambda \\ -i\lambda & 1 \end{pmatrix}, \quad \text{and} \quad U(\phi_1, \phi_2) = -(\cos \phi_1 + \mu \phi_1 + \cos \phi_2), \quad \lambda, \mu \in \mathbb{R}, \quad (4.1)$$

respectively, the potential resulting from the expression in (2.3) is derived as

$$\mathcal{V}(\phi_1, \phi_2) = \frac{1}{2(1+\lambda^2)} \left[(\sin \phi_1 - \mu)^2 + 2i\lambda (\sin \phi_1 - \mu) \sin \phi_2 + \sin^2 \phi_2 \right]. \quad (4.2)$$

We note that the singularity at $\lambda = 1$ present in the real version of this model discussed in [15] has been removed. The static versions of the BPS equations (2.6) obtained from (4.2) are the pairs of complex coupled first order equations

$$BPS_1^\pm : \quad \partial_x \phi_1 = \pm \frac{1}{1+\lambda^2} (\sin \phi_1 - \mu + i\lambda \sin \phi_2) =: G_1^\pm, \quad (4.3)$$

$$BPS_2^\pm : \quad \partial_x \phi_2 = \pm \frac{1}{1+\lambda^2} [i\lambda (\sin \phi_1 - \mu) + \sin \phi_2] =: G_2^\pm. \quad (4.4)$$

These equations are compatible under the modified \mathcal{CPT} -transformation

$$\mathcal{CPT} : \quad \phi_1(x) \rightarrow [\phi_1(-x)]^\dagger, \quad \phi_2(x) \rightarrow -[\phi_2(-x)]^\dagger, \quad \Leftrightarrow \quad BPS_i^\pm \rightarrow (BPS_i^\mp)^*. \quad (4.5)$$

Notice that we require again both signs to achieve consistency under the \mathcal{CPT} -conjugation. It is precisely this symmetry that is needed to derive the second relation in (2.8). Trying instead to realize the compatibility of BPS_i^+ or BPS_i^- with itself requires just a modified \mathcal{CT} -transformation $\phi_1(x) \rightarrow [\phi_1(x)]^\dagger$, $\phi_2(x) \rightarrow -[\phi_2(x)]^\dagger$, which as for the previous model is, however, not sufficient to be used in the argument in (2.7) that ensures the reality of the energy.

In the Hermitian limit, when $\lambda = 0$, the two pairs of BPS equations decouple and are easily solved by the kink and antikink solutions for the upper and lower sign, respectively,

$$\phi_1^{\pm(n)}(x) = 2 \arctan \left\{ \frac{1}{\mu} \left[1 + \sqrt{1-\mu^2} \tanh \left[\frac{1}{2} \sqrt{1-\mu^2} (\pm x + \kappa_1) \right] \right] \right\} + 2\pi n, \quad (4.6)$$

$$\phi_2^{\pm(n)}(x) = 2 \arctan (e^{\pm x + \kappa_2}) + 2\pi n, \quad (4.7)$$

where $n \in \mathbb{Z}$ and integration constants $\kappa_1, \kappa_2 \in \mathbb{R}$. From the asymptotic limits

$$\lim_{x \rightarrow \infty} \phi_1^{+(n)}(x) = \lim_{x \rightarrow -\infty} \phi_1^{-(n)}(x) = 2n\pi + \text{sign}(\mu)\pi - \arcsin(\mu), \quad (4.8)$$

$$\lim_{x \rightarrow -\infty} \phi_1^{+(n)}(x) = \lim_{x \rightarrow \infty} \phi_1^{-(n)}(x) = 2n\pi + \text{sign}(\mu) \arcsin(\mu), \quad (4.9)$$

$$\lim_{x \rightarrow \pm\infty} \phi_2^{+(n)}(x) = \lim_{x \rightarrow \mp\infty} \phi_2^{-(n)}(x) = 2n\pi + \frac{\pi \pm \pi}{2}, \quad (4.10)$$

for $|\mu| \leq 1$, we obtain from (2.4) for both signs the same expression for the energy as a function of μ

$$E^\pm(\mu) = 2 \left[1 + \sqrt{1 - \mu^2} - \mu \arctan \left(\frac{\sqrt{1 - \mu^2}}{\mu} \right) \right]. \quad (4.11)$$

For $|\mu| > 1$ the limits $\lim_{x \rightarrow \pm\infty} \phi_i(x)$ are not well defined as the solutions become periodic in this case. Limiting this case to a theory on a finite interval will, however, still give real energies. For instance, for an interval $[a, b]$ with $\kappa_1 = \kappa_2 = n = 0$ we compute the energy

$$E^\pm(\mu) = \pm \mu z(x) - \cos[z(x)] \pm \tanh x \Big|_{x=a}^b. \quad (4.12)$$

with $z(x) = 2 \arctan \left\{ [\mp 1 + \sqrt{\mu^2 - 1} \tan(x/2\sqrt{\mu^2 - 1})] / \mu \right\}$. This is real and well defined as long as one avoids $a, b = (2n\pi + \pi) / \sqrt{\mu^2 - 1}$, $n \in \mathbb{Z}$.

In the non-Hermitian scenario, when $\lambda \neq 0$, we solve the coupled equations (4.3) and (4.4) numerically, see figure 3 for some sample behaviours.




Figure 3: Complex BPS kink and antikink solutions of the pair of BPS equations (4.3) and (4.4) with initial values $\phi_1^{k+}(0) = \phi_2^{k+}(0) = \phi_2^{k-}(0) = \phi_1^{a-}(0) = \pi/2$ and $\phi_1^{a+}(0) = \phi_2^{a+}(0) = \phi_1^{k-}(0) = \phi_2^{a-}(0) = -\pi/2$ for $\lambda = 3$, $\mu = 0.5$.

We observe that the real parts are perturbed versions of the smooth kink and antikink solution of the Hermitian case, which exhibit more and more oscillations near the origin as λ increases. Asymptotically the solutions of the Hermitian and non-Hermitian cases tend to the same value. Crucially, we read off the \mathcal{CPT} -symmetry (4.5) for the solutions

$$\phi_1^{k\pm}(x) = [\phi_1^{a\mp}(-x)]^\dagger, \quad \phi_2^{k+}(x) = -[\phi_2^{k-}(-x)]^\dagger, \quad \phi_2^{a+}(x) = -[\phi_2^{a-}(-x)]^\dagger, \quad (4.13)$$

from which we derive for the potential

$$\mathcal{V}_\lambda \left[\phi_1^{k+}(x), \phi_2^{k+}(x) \right] = \mathcal{V}_\lambda \left\{ \left[\phi_1^{a-}(-x) \right]^\dagger, - \left[\phi_2^{k-}(-x) \right]^\dagger \right\} = \mathcal{V}_\lambda^\dagger \left\{ \phi_1^{a-}(-x), \phi_2^{k-}(-x) \right\}. \quad (4.14)$$

This is once more the second option in (2.8). Thus assuming the energies of kinks and antikinks in the + system are the same as the antikinks and kinks in the – system, respectively, this energy is guaranteed to be real.

Since the limits $x \rightarrow \pm\infty$ for these solutions are the same as for $\lambda = 0$, the expression for the energy $E(\mu)$ in (4.11) holds for all values of λ . Considering the expression in (4.11) it appears that $\mu = 1$ is an exceptional point of the system and that for $|\mu| > 1$ one might obtain complex conjugate pairs of eigenvalues. However, just as in the previous model, when the threshold is passed into that region the asymptotic limits of the kink solutions are no longer defined so that the expression for the energy becomes meaningless. Moreover, when defining the theory on a finite interval in space the energy is actually still real and does not occur in complex conjugate pairs. For an exceptional point to emerge we would also expect that the antilinear \mathcal{CPT} -symmetry (4.13) becomes broken when passing a genuine exceptional point. However, this symmetry is still preserved in the regime $|\mu| > 1$. Hence we conclude that $\mu = 1$ is not an exceptional point.

Next we identify the precise relation on which vacua are connected by which of the various BPS solutions. The infinite amount of vacua of the potential (4.2) are easily found to be

$$v_1^{(n,m)} = (\arcsin \mu + 2\pi n, m\pi), \quad \text{and} \quad v_2^{(n,m)} = (\pi - \arcsin \mu + 2n\pi, m\pi), \quad (4.15)$$

corresponding to the fixed points of the dynamical system (4.3) and (4.4), that are the solutions of $G_1^\pm(\phi_1, \phi_2) = G_2^\pm(\phi_1, \phi_2) = 0$. Computing once more the eigenvalues of the Jacobian matrix at these fixed points

$$J = \left(\begin{array}{cc} \partial_{\phi_1} G_1^\pm & \partial_{\phi_2} G_1^\pm \\ \partial_{\phi_1} G_2^\pm & \partial_{\phi_2} G_2^\pm \end{array} \right) \Big|_{v_j^{(n,m)}}, \quad (4.16)$$

with $j = 1, 2$, we find for the + system that $J(v_1^{(n,2m)})$ has two positive eigenvalues, $J(v_2^{(n,2m+1)})$ has two negative eigenvalues and $J(v_1^{(n,2m+1)})$, $J(v_2^{(n,2m)})$ have a positive and a negative eigenvalue. For the – system the signs are reversed. Thus the vacua $v_1^{(n,2m+1)}$, $v_2^{(n,2m)}$ are always saddle points, $v_1^{(n,2m)}$ are unstable/stable nodes (G^+/G^-) and $v_2^{(n,2m+1)}$ are stable/unstable nodes (G^-/G^+). Hence the kink and antikink solutions only interpolate between the vacua $v_1^{(n,2m)}$ and $v_2^{(n,2m+1)}$ as indicated for an example in figure 4 with the accompanying gradient flow.

The solutions depicted in figure 3 interpolate the vacua $v_i^{(n,m)}$ as

$$v_1^{(0,0)} \xrightarrow{\phi_1^{k+} \phi_2^{k+}} v_2^{(0,1)}, \quad v_1^{(0,0)} \xrightarrow{\phi_1^{a+} \phi_2^{a+}} v_2^{(-1,1)}, \quad v_1^{(0,0)} \xrightarrow{\phi_1^{a-} \phi_2^{k-}} v_2^{(0,-1)}, \quad v_1^{(0,0)} \xrightarrow{\phi_1^{k-} \phi_2^{a-}} v_2^{(-1,1)}, \quad (4.17)$$

hence confirming the consistency of the above. The other vacua $v_i^{(n,m)}$ for different choices of n and m are obtained by including the n -dependence into the solutions.




Figure 4: Real part of the potential $\mathcal{V}(\phi_1, \phi_2)$ as a function of $\text{Re } \phi_1$ and $\text{Re } \phi_2$ with the gradient flow of the real parts of G^+ superimposed in white. The kink solutions $\phi_1^{k+}(x)$, $\phi_2^{k+}(x)$ interpolate between the vacua $v_1^{(0,0)}$ and $v_2^{(0,1)}$ (red dots) as indicated by the red solid trajectory.

In both of our previous examples we have directly analyzed the complex non-Hermitian systems. In analogy to the treatment of many quantum systems, such an approach is especially meaningful under the assumption that there exists an equivalent Hermitian system with the same energy. In the next section we present such a system and thus further justify our approach.

5. Complex extended sine-Gordon model and its Hermitian partner

In this section we investigate a model with two complex fields consisting of two copies of sine-Gordon models of which one is complex \mathcal{PT} -symmetrically extended

$$\mathcal{V}(\phi_1, \phi_2) = \frac{m^2}{2\mu^2} \left[\sqrt{1 - \varepsilon^2} - \cos(\mu\phi_1) - i\varepsilon \sin(\mu\phi_1) \right] + \frac{m^2}{\mu^2} \sin^2\left(\frac{\mu}{2}\phi_2\right), \quad (5.1)$$

with constants $m, \mu \in \mathbb{R}$ and $|\varepsilon| \leq 1$. For simplicity we have not introduced an interaction term between ϕ_1 and ϕ_2 as the feature we are trying to illustrate can even be shown for a theory with one field only. We just keep a second field to maintain a similarity with the previously discussed systems and to allow for a direct comparison between the BPS solutions for the two fields. The constant term proportional to $\sqrt{1 - \varepsilon^2}$ is introduced for convenience. In order to find a Hermitian partner potential \mathfrak{v} to the non-Hermitian potential \mathcal{V} we employ now a Dyson map originally found in [22]

$$\tilde{\eta} = \exp \left[\frac{\text{arctanh } \varepsilon}{\mu} \int dx \pi_1(x, t) \right]. \quad (5.2)$$

Here the spacial momentum operator $\pi_1(x, t) := \partial_t \phi_1(x, t)$ satisfies the canonical equal time commutation relation $[\phi_1(x, t), \pi_1(y, t)] = i\delta(x - y)$. The inverse adjoint action of $\tilde{\eta}$ on \mathcal{V} then leads to

$$\mathfrak{v}(\phi_1, \phi_2) = \tilde{\eta}^{-1} \mathcal{V} \tilde{\eta} = \frac{m^2}{\mu^2} \left[\sqrt{1 - \varepsilon^2} \sin^2 \left(\frac{\mu}{2} \phi_1 \right) + \sin^2 \left(\frac{\mu}{2} \phi_2 \right) \right], \quad (5.3)$$

whereas the kinetic term remains unchanged as $\tilde{\eta}$ commutes with it. Even though we are here mainly interested in the properties of classical solutions, we briefly drew on the quantum field theory version of the model in order to carry out the similarity transformation. The effect of the adjoint action of $\tilde{\eta}$ on any smooth function of the fields (ϕ_1, ϕ_2) is $(\phi_1, \phi_2) \rightarrow (\phi_1 + i/\mu \operatorname{arctanh} \varepsilon, \phi_2)$.

We shall now demonstrate that the energies of the BPS solutions for the system involving the non-Hermitian potential \mathcal{V} and the Hermitian potential \mathfrak{v} are identical and real. Following the procedure of the previous sections we first note that the potential \mathcal{V} can be derived from the pre-potential

$$U(\phi_1, \phi_2) = -\frac{2^{3/2} m}{\mu^2} \left[(1 - \varepsilon^2)^{1/4} \cos \left(\frac{\mu}{2} \phi_1 - \frac{i}{2} \operatorname{arctanh} \varepsilon \right) + \cos \left(\frac{\mu}{2} \phi_2 \right) \right], \quad (5.4)$$

when taking the metric of the target space simply to be diagonal $\operatorname{diag} \eta = (1, 1)$. According to (2.6) the two pairs of coupled BPS equations are therefore

$$BPS_1^\pm : \quad \partial_x \phi_1 = \pm \frac{m\sqrt{2}(1 - \varepsilon^2)^{1/4}}{\mu} \sin \left(\frac{\mu \phi_1}{2} - \frac{i}{2} \operatorname{arctanh} \varepsilon \right), \quad (5.5)$$

$$BPS_2^\pm : \quad \partial_x \phi_2 = \pm \frac{m\sqrt{2}}{\mu} \sin \left(\frac{\mu}{2} \phi_2 \right). \quad (5.6)$$

Once again we can identify a pair of modified \mathcal{CPT} -transformations under which these equations are compatible

$$\mathcal{CPT}_\pm : \quad \phi_1(x) \rightarrow -[\phi_1(-x)]^\dagger, \quad \phi_2(x) \rightarrow \pm[\phi_2(-x)]^\dagger, \quad \Leftrightarrow \quad BPS_i^\pm \rightarrow (BPS_i^\mp)^*. \quad (5.7)$$

We solve the equations (5.5) and (5.6) by

$$\phi_1^{k/a+}(x) = -[\phi_1^{k/a-}(-x)]^* = \pm \frac{4}{\mu} \arctan \left[e^{mx(1-\varepsilon^2)^{1/4}/\sqrt{2} + \mu\kappa_1/2} \right] + \frac{i}{\mu} \operatorname{arctanh} \varepsilon, \quad (5.8)$$

$$\phi_2^{k/a+}(x) = -[\phi_2^{k/a-}(-x)]^* = \pm \frac{4}{\mu} \arctan \left[e^{mx/\sqrt{2} + \mu\kappa_2/2} \right], \quad (5.9)$$

with integration constants $\kappa_1, \kappa_2 \in \mathbb{C}$. The solution respect the \mathcal{CPT}_- -symmetry as indicated, which leads to the relation

$$\mathcal{V} \left[\phi_1^{k/a+}(x), \phi_2^{k/a+}(x) \right] = \mathcal{V}^* \left[\phi_1^{k/a-}(-x), \phi_2^{k/a-}(-x) \right], \quad (5.10)$$

for the potential that guarantees the reality of the energy when arguing along the same lines as above.

We may of course also compute the energies directly from the asymptotic limits of the solutions. For $|\varepsilon| \leq 1$ we find

$$\lim_{x \rightarrow \pm\infty} \phi_j^{k+}(x) = \lim_{x \rightarrow \mp\infty} \phi_j^{a-}(x) = \frac{\pi}{\mu} \pm \frac{\pi}{\mu} + \delta_{1j} \frac{i}{\mu} \operatorname{arctanh} \varepsilon, \quad (5.11)$$

$$\lim_{x \rightarrow \pm\infty} \phi_j^{a+}(x) = \lim_{x \rightarrow \mp\infty} \phi_j^{k-}(x) = -\frac{\pi}{\mu} \mp \frac{\pi}{\mu} + \delta_{1j} \frac{i}{\mu} \operatorname{arctanh} \varepsilon, \quad (5.12)$$

which by (2.4) gives the real energies

$$E^{\phi_1^{pn}, \phi_2^{qn}}(m, \mu, \varepsilon) = \frac{4\sqrt{2}m}{\mu^2} \left[1 + (1 - \varepsilon^2)^{1/4} \right], \quad p, q = k, a; \quad n = \pm; \quad m, \mu \in \mathbb{R}. \quad (5.13)$$

The special point $\varepsilon = 1$ is not an exceptional point as the BPS solutions for ϕ_1 and ϕ_2 have no definite asymptotic values. For $|\varepsilon| > 1$ the energies become complex, albeit not complex conjugate. The reason for the latter is that the \mathcal{CPT} -symmetry is not just broken for the solutions, but also at the level of the Hamiltonian. It is now easy to verify that the pre-potential $u(\phi_1, \phi_2)$ leading to the real potential $v(\phi_1, \phi_2)$ is simply obtained as $u = \tilde{\eta}^{-1} U \tilde{\eta}$. The solutions to the real BPS equations are then given by (5.8) and (5.9) with $(\phi_1, \phi_2) \rightarrow (\phi_1 - i/\mu \operatorname{arctanh} \varepsilon, \phi_2)$. The expression for the energy $E = \lim_{x \rightarrow \infty} u(\phi_1, \phi_2) - \lim_{x \rightarrow -\infty} u(\phi_1, \phi_2)$ is then the same as the one in (5.13).

6. Conclusions

By assuming the non-Hermitian field theories to possess self-dual and anti-self-dual fields we have derived their BPS soliton equations. We have solved these equations for their complex kink and antikink solutions for three different types of systems. We demonstrated that the solutions found exhibit different types of modified antilinear \mathcal{CPT} -symmetries relating the two versions of the BPS soliton equations. These symmetries were shown to lead to real energies on general grounds in certain regimes of the parameter space. For each of the systems we computed the topological energy that saturates the Bogomolny bound confirming the generic result. We observed that despite the fact that the BPS solutions are complex the corresponding energies are real. Technically this is due to the fact that contributions to the imaginary parts of the pre-potential are the same at spacial plus and minus infinity. Crucially we found that the \mathcal{CPT} -symmetries can not be utilized directly on the self-dual part of the BPS equation. However, taking both signs in (2.2) into account the symmetries can be identified.

For two of the systems we demonstrated explicitly how the kink/antikink solutions interpolate between certain types of vacua corresponding always to unstable fixed points at negative spacial infinity and stable fixed points at positive spacial infinity. For the complex extended sine-Gordon model we made the pseudo-Hermitian approach explicit and mapped the corresponding non-Hermitian Hamiltonian to a Hermitian partner Hamiltonian by means of a Dyson map. As the energies are preserved in this process and the Hermitian theory always possesses real energies, this establishes the reality of the energy computed from complex BPS solutions in the non-Hermitian theory.

There are evidently a number of interesting follow up problems and open question. Since in none of the models we treated the transition point in parameter space from real to complex or ill-defined energies led to a genuine exceptional point, it remains an open question whether such type of systems can be constructed. As our scheme is very general it should be applicable to all non-Hermitian field theories that admit the described self-dual and anti-self dual symmetries. It would be worth exploring whether there exist models or even general arguments that could ensure that modified antilinear CPT -symmetries may also be utilized to guarantee the reality of the energies by just using the self-dual sector of the theory. Hence it would be interesting to see the working of the above for more involved theories, possibly with a larger field content. It would also be interesting to further compare with an alternative approach to non-Hermitian field theories pursued in [27, 28, 29, 30, 31, 32] and investigate whether the BPS soliton solutions derived in that framework also posses real energies.

References

- [1] G. t Hooft, Magnetic monopoles in unified theories, Nucl. Phys. B **79**(CERN-TH-1876), 276–284 (1974).
- [2] A. P. Polyakov, Particle spectrum in the quantum field theory, JETP Letters **20**(6), 194–195 (1974).
- [3] E. B. Bogomolny, The stability of classical solutions, Sov. J. Nucl. Phys.(Engl. Transl.);(United States) **24**(4) (1976).
- [4] M. K. Prasad and C. M. Sommerfield, Exact classical solution for the't Hooft monopole and the Julia-Zee dyon, Phys. Rev. Lett. **35**(12), 760 (1975).
- [5] A. Fring and T. Taira, 't Hooft-Polyakov monopoles in non-Hermitian quantum field theory, Phys. Lett. B **807**, 135583 (2020).
- [6] A. Fring, A note on the integrability of non-Hermitian extensions of Calogero-Moser-Sutherland models, Mod. Phys. Lett. **21**, 691–699 (2006).
- [7] A. Fring, \mathcal{PT} -Symmetric deformations of the Korteweg-de Vries equation, J. Phys. A **40**, 4215–4224 (2007).
- [8] P. E. G. Assis and A. Fring, Integrable models from \mathcal{PT} -symmetric deformations, J. of Phys. A **42**, 105206 (2009).
- [9] A. Fring and M. Smith, Antilinear deformations of Coxeter groups, an application to Calogero models, J. Phys. A **43**, 325201 (2010).
- [10] A. Cavaglia, A. Fring, and B. Bagchi, \mathcal{PT} -symmetry breaking in complex nonlinear wave equations and their deformations, J. Phys. A **44**, 325201(42) (2011).
- [11] A. Fring, \mathcal{PT} -symmetric deformations of integrable models, Phil. Trans. Royal Soc. London A: Math., Phys. and Eng. Sci. **371**(1989), 20120046 (2013).
- [12] J. Cen and A. Fring, Complex solitons with real energies, J. Phys. A: Math. Theor. **49**(36), 365202 (2016).
- [13] J. Cen, F. Correa, and A. Fring, Time-delay and reality conditions for complex solitons, J. of Math. Phys. **58**(3), 032901 (2017).

- [14] C. Adam, L. A. Ferreira, E. Da Hora, A. Wereszczynski, and W. J. Zakrzewski, Some aspects of self-duality and generalised BPS theories, *JHEP* **2013**(8), 62 (2013).
- [15] L. A. Ferreira, P. Klimas, A. Wereszczyński, and W. J. Zakrzewski, Some comments on BPS systems, *J. Phys. A: Math. and Theor.* **52**(31), 315201 (2019).
- [16] C. Adam, K. Oles, J. M. Queiruga, T. Romanczukiewicz, and A. Wereszczynski, Solvable self-dual impurity models, *JHEP* **2019**(7), 150 (2019).
- [17] P. Klimas and W. J. Zakrzewski, Further comments on BPS systems, arXiv preprint arXiv:1908.02100 (2019).
- [18] W. D. Heiss, The physics of exceptional points, *J. of Phys. A: Math. and Theor.* **45**(44), 444016 (2012).
- [19] C. M. Bender, P. E. Dorey, C. Dunning, A. Fring, D. W. Hook, H. F. Jones, S. Kuzhel, G. Levai, and R. Tateo, *PT Symmetry: In Quantum and Classical Physics*, (World Scientific, Singapore) (2019).
- [20] A. Mostafazadeh, Pseudo-Hermitian Representation of Quantum Mechanics, *Int. J. Geom. Meth. Mod. Phys.* **7**, 1191–1306 (2010).
- [21] P. Kumar, A. Khare, and A. Saxena, A model field theory with $(\psi \ln \psi)^2$ potential: Kinks with super-exponential profiles, arXiv preprint arXiv:1908.04978 (2019).
- [22] C. M. Bender, H. F. Jones, and R. J. Rivers, Dual \mathcal{PT} -symmetric quantum field theories, *Phys. Lett.* **B625**, 333–340 (2005).
- [23] P. D. Mannheim, Goldstone bosons and the Englert-Brout-Higgs mechanism in non-Hermitian theories, *Phys. Rev. D* **99**(4), 045006 (2019).
- [24] A. Fring and T. Taira, Goldstone bosons in different PT-regimes of non-Hermitian scalar quantum field theories, *Nucl. Phys. B* **950**, 114834 (2020).
- [25] A. Fring and T. Taira, Pseudo-Hermitian approach to Goldstone’s theorem in non-Abelian non-Hermitian quantum field theories, *Phys. Rev. D* **101**(4), 045014 (2020).
- [26] A. Fring and T. Taira, Massive gauge particles versus Goldstone bosons in non-Hermitian non-Abelian gauge theory, arXiv preprint arXiv:2004.00723 (2020).
- [27] J. Alexandre, P. Millington, and D. Seynaeve, Symmetries and conservation laws in non-Hermitian field theories, *Phys. Rev. D* **96**(6), 065027 (2017).
- [28] J. Alexandre, P. Millington, and D. Seynaeve, Consistent description of field theories with non-Hermitian mass terms, in *J. Phys. Conf. Ser.*, volume 952, page 012012, 2018.
- [29] J. Alexandre, J. Ellis, P. Millington, and D. Seynaeve, Gauge invariance and the Englert-Brout-Higgs mechanism in non-Hermitian field theories, *Phys. Rev. D* **99**(7), 075024 (2019).
- [30] J. Alexandre, J. Ellis, P. Millington, and D. Seynaeve, Spontaneously breaking non-Abelian gauge symmetry in non-Hermitian field theories, *Phys. Rev. D* **101**(3), 035008 (2020).
- [31] J. Alexandre, J. Ellis, P. Millington, and D. Seynaeve, Spontaneously breaking non-Abelian gauge symmetry in non-Hermitian field theories, *Physical Review D* **101**(3), 035008 (2020).
- [32] J. Alexandre, J. Ellis, and P. Millington, Discrete spacetime symmetries and particle mixing in non-Hermitian scalar quantum field theories, arXiv preprint arXiv:2006.06656 (2020).
- [33] D. Arrowsmith and C. M. Place, *Dynamical systems: differential equations, maps, and chaotic behaviour*, volume 5, CRC Press, 1992.