



Comparison of the oxygen isotope signatures in speleothem records and iHadCM3 model simulations for the last millennium

Janica C. Bühler¹, Carla Roesch¹, Moritz Kirschner¹, Louise Sime², Max D. Holloway³, and Kira Rehfeld¹

¹Institute of Environmental Physics, Ruprecht-Karls-Universität Heidelberg, INF 229, 69120 Heidelberg, Germany

²British Arctic Survey, High Cross, Madingley Road, Cambridge, CB3 0ET, United Kingdom

³Scottish Association for Marine Science, Scottish Marine Institute, Oban, Argyll, PA37 1QA, United Kingdom

Correspondence: Janica Bühler (jbuehler@iup.uni-heidelberg.de), Kira Rehfeld (krehfeld@iup.uni-heidelberg.de)

Abstract.

Global changes in the climate, especially the warming trend in mean temperature, have received increasing public and scientific attention. Improving the understanding of changes in the mean and variability of climate variables as well as their interrelation is crucial for reliable climate change projections.

5 Comparisons between general circulation models and paleoclimate archives using indirect proxies for temperature and/or precipitation have been used to test and validate the capability of climate models to represent climate changes. The oxygen isotopic ratio $\delta^{18}\text{O}$ is routinely measured in speleothem samples at decadal or higher resolution and single specimens can cover full Glacial-Interglacial cycles. The calcium carbonate cave deposits are precisely dateable and provide well preserved (semi-) continuous, albeit multivariate climate signals in the lower and mid-latitudes, where the measured $\delta^{18}\text{O}$ in the mineral
10 does not directly represent temperature or precipitation. Therefore, speleothems represent suitable archives to assess simulated climate model abilities for the simulation of climate variability beyond the timescales covered by meteorological observations ($10^1 - 10^2\text{yr}$).

Here, we present three transient isotope enabled simulations from the Hadley Center Climate Model version 3 (iHadCM3) covering the last millennium (850-1850CE) and compare these to a large global dataset of speleothem $\delta^{18}\text{O}$ records from the
15 Speleothem Isotopes Synthesis and AnaLysis (SISAL) database version 2 (Comas-Bru et al., 2020). We evaluate systematically offsets in mean and variance of simulated $\delta^{18}\text{O}$ and test for the main climate drivers for individual records or regions.

The time-mean spatial offsets between the simulated $\delta^{18}\text{O}$ and the speleothem data are fairly small. However, using robust filters and spectral analysis, we show that the observed proxy-based variability of $\delta^{18}\text{O}$ is lower (higher) than simulated by iHadCM3 on decadal (centennial) timescales. Most of this difference can likely be attributed to the records' lower tempo-
20 ral resolution and averaging processes affecting the $\delta^{18}\text{O}$ signal. Using cross-correlation analyses at site-level and modeled gridbox level, we find evidence for highly variable but generally low signal-to-noise ratios in the proxy data. This points at a high influence of cave-internal processes and regional climate particularities and could suggest low regional representativity of individual sites. Long-range strong positive correlations dominate the speleothem correlation network but are much weaker in the simulation. One reason for this could lie in a lack of longterm internal climate variability in these model simulations, which
25 could be tested by repeating similar comparisons with other isotope-enabled climate models and paleoclimate databases.



1 Introduction

The impacts of a changing climate have been observed over the last century (IPCC, 2013) and indicate a strong sensitivity of natural and human systems to changes in climate. While the mean state of the climate is well observed, direction and magnitude of potential changes to its variability are still largely unclear (Franzke et al., 2020). However, changes in variability influence the occurrence of extreme temperature and precipitation events (Katz and Brown, 1992) and have major impacts on society, economy (Hänsel et al., 2020), and ecosystems (Vasseur et al., 2014).

Past climate changes provide a testbed to evaluate climate models and to better understand projected changes in the future (Schmidt et al., 2012; Braconnot et al., 2012). Instrumental records only cover a short period of time, since systematic observations of climate variables only began in 1750 CE (black line in Fig. 1a, Morice et al., 2012). Therefore, for model evaluation on longer than centennial time scales, we have to rely on evidence from paleoclimate archives, such as trees, ice cores, foraminifera from marine sediment cores, or speleothems. The $\delta^{18}\text{O}$ -proxy can be measured on these, and quite a few other paleoclimate archives with high precision (Schmidt et al., 2014). The climatic interpretation of $\delta^{18}\text{O}$ changes, however, are not always straightforward (Fairchild and Baker, 2012). Speleothem archives, which we rely on here, allow sampling of a wide range of climates in the low- to mid-latitudes and provide (semi-) continuous precisely dated time series of oxygen isotope ratios.

Few other transient model-data comparison studies focused on $\delta^{18}\text{O}$ (e.g., Wackerbarth, 2012; Dee et al., 2015; Parker et al., 2020). For example, Sjolte et al. (2018) compared Greenland ice cores to the ECHAM5/MPI-OM model over the last millennium and were able to differentiate between solar and volcanic forcing effects from their reconstructions. On orbital timescales (150,000 yr), Caley et al. (2014) compared speleothem records from South East Asia to a transient isotope-enabled simulation with the model of intermediate complexity iLOVECLIM. They found model-data similarity for the broad temporal trends, but differences at shorter timescales, highlighting the role of seasonality.

For our model-data comparison, we focus on the time period of the last millennium (850-1850AD Taylor et al. (2012)) for which a fairly high number of well-preserved datasets are available. This time period is characterized by stable, close-to-present-day, boundary conditions (fairly constant greenhouse gas concentrations and sea level) and climate variability due to natural, solar and volcanic, forcings (Schurer et al., 2014; PAGES2k-Consortium, 2019; Taylor et al., 2012; Neukom et al., 2019). It is also one of the key paleoclimate periods included in the joint experiments of the Paleoclimate Model Intercomparison Project Phase 3 and 4 (PMIP3/PMIP4, Jungclaus et al., 2010; Kageyama et al., 2018) and the overarching Coupled Model Intercomparison Project Phase 5 and 6 (CMIP5/CMIP6, Taylor et al., 2012; Eyring et al., 2016).

Modeled climate variability can be a consequence of either internal interactions and processes (internal variability) or of radiative forcings such as depicted in Fig. 1c-e (forced external variability), e.g. greenhouse gases, volcanic eruptions, or total solar irradiance. Previous studies have suggested that simulated temperature variability is systematically too low, especially on the regional scale (Laepplé and Huybers, 2014a). This has been attributed to too high diffusivity (Laepplé and Huybers, 2014b) or to missing processes and feedbacks (Rehfeld et al., 2016). Variability induced by external radiative forcing only accounts for a small fraction of the regional climate variance (Goosse et al., 2005; Laepplé and Huybers, 2014b). Discrepancies




Figure 1. Climate and main climatic drivers over the last millennium. (a) Global annual mean surface temperature (GMST) as modeled (iHadCM3, blue), reconstructed (PAGES2k, red) (PAGES2k-Consortium, 2019) and observed (HadCRUT4, black) (Morice et al., 2012). (b) Annual mean $\delta^{18}\text{O}$ in precipitation at Bunker Cave, Germany as modeled (iHadCM3, green) and measured calcite $\delta^{18}\text{O}$ (drip water equivalent) from SISAL entity 240 (dark blue) and 242 (light blue) at the Bunker cave site (Comas-Bru et al., 2019; Fohlmeister et al., 2012). Comparison plots for all entities are given in supplementary figures SF1-2. (c) Atmospheric CO_2 concentration, (d) volcanic forcing in units of aerosol optical depth (AOD) (Crowley and Unterman, 2013) and (e) total solar irradiance (TSI) as used in the model simulations (Steinhilber et al., 2009; Wang et al., 2005).

60 increase towards longer timescales (Laepple and Huybers, 2014a), and are substantial already at the multidecadal to centennial timescales that we target here.

The incorporation of an isotopic water cycle into isotope-enabled General Circulation Models (iGCM) provides additional means for understanding the hydrology of the climate system (Werner et al., 2016; Sturm et al., 2010; Tindall et al., 2009). The
 65 ratio of H_2^{18}O to H_2^{16}O is an indicator of evaporation temperature, precipitation amount, and altitude, as well as distance to source water (Dansgaard, 1964). It is given in the δ -notation as

$$\delta^{18}\text{O} = \left(\frac{\frac{^{18}\text{O}}{^{16}\text{O}}_{\text{sample}}}{\frac{^{18}\text{O}}{^{16}\text{O}}_{\text{standard}}} - 1 \right) \cdot 1000\text{‰},$$

where standard indicates the Vienna Standard Mean Ocean Water standard V-SMOW (Kendall and Caldwell, 1998).



On monthly to decadal time scales, the Global Network of Isotopes in Precipitation (GNIP) database (IAEA/WMO, 2020) provides measurements of $\delta^{18}\text{O}$ in collected precipitation water and these have been used in model-data comparisons for the present climate (Tindall et al., 2009; Werner et al., 2011; Comas-Bru et al., 2020). On decadal and longer timescales, paleoclimate archives such as speleothems are crucial. $\delta^{18}\text{O}$ variations in stalagmites, to first order, represent changes in $\delta^{18}\text{O}$ in the meteoric precipitation above the cave.

Speleothem cave deposits form in karst regions (Fairchild and Baker, 2012) under climatic conditions spanning from extremely cold (Lauritzen and Lundberg, 1999) and very arid (Neff et al., 2001) to extremely hot and humid conditions (Partin et al., 2007). As a terrestrial climate archive, they are able to store information on continental climate changes. They form as calcite or aragonite matrix from calcium dissolved in acidic drip water and do, therefore, archive the oxygen isotope from precipitation water in accumulated growth layers (Fairchild and Treble, 2009). $\delta^{18}\text{O}$ can be regarded as a proxy for surface temperature variations in higher latitudes, and precipitation amount in the tropics (Dansgaard, 1964), overlaid with distinct observable signatures of source water evaporation and transportation over longer distances (Bradley, 1999; Dansgaard, 1964). These signatures are also visible in speleothem records, including additionally a fractionation process involved in the calcite formation, which is primarily temperature-dependent (Urey, 1948; McCrea, 1950). The climatic interpretation of speleothem $\delta^{18}\text{O}$ variations in calcite or aragonite (hereafter $\delta^{18}\text{O}_{\text{speleo}}$) can be hampered by non-linear growth processes (Dreybrodt, 1980), and multiple cave-specific parameters such as vegetation cover (Haude, 1954; Wackerbarth et al., 2010), karst (Jean-Baptiste et al., 2019), and inner cave processes (Fairchild et al., 2006), which influence $\delta^{18}\text{O}_{\text{speleo}}$ as well as dating uncertainties (Breitenbach et al., 2012). For speleothems, in particular, positive correlations to ice core $\delta^{18}\text{O}$, which is considered a proxy for temperature, have been reported (McDermott et al., 2001) but also negative correlations to local annual mean temperatures at the cave site (e.g. (Lauritzen and Lundberg, 1999)). This highlights the complexity of the system and the potential regionality of the signal. In studies on drip water, $\delta^{18}\text{O}$ and annual mean temperature, regions with different dominant climate controls could be distinguished (Baker et al., 2019).

Here, we present three new last millennium isotope-enabled simulations from the iGCM version 3 of the Hadley Model (iHadCM3) and test how similar the $\delta^{18}\text{O}$ variations in iHadCM3 and speleothem records are (Sec. 4.1). A characterization of the datasets and relevant forcing can be found in Fig. 1. The robustness of the findings and methods are evaluated over the last millennium, for which a large number of high-resolution proxy datasets from the SISAL v.2. database (Comas-Bru et al., 2020) are available.

We explore these similarities on both spatial and temporal scales, to distinguish patterns of the mean state (Sec. 4.1), the variability (Sec. 4.2 and Sec. 4.3), and the spatial representativity of speleothem climate records (Sec. 4.4 and Sec. 4.5). We examine the simulation's capability to simulate and the records' capability to capture variability on different time scales to improve our understanding of processes and uncertainties of both.



100 2 Data

2.1 Model description and simulation overview

In this study, we use the coupled atmosphere-ocean isotope-enabled GCM iHadCM3, which has been widely used to simulate present and future climate (Sime et al., 2008; Tindall et al., 2009; IPCC, 2013), as well as for past climates such as the late Holocene and Last Glacial Maximum (Holloway et al., 2016), the last interglacial (Sime et al., 2009, 2013; Holloway et al., 105 2016, 2018) and the Eocene (Tindall et al., 2010).

The model consists of several components: the atmosphere model HadAM3 (Pope et al., 2000), the ocean model HadOM3 (Gordon et al., 2000), a sea ice model (Valdes et al., 2017) and a dynamic land surface and vegetation model (Cox, 2001). The atmospheric component is run at a horizontal resolution of $2.5^\circ \times 3.75^\circ$, 19 vertical levels and time steps of 30min. The oceanic output has a horizontal resolution of $1.25^\circ \times 1.25^\circ$, 20 vertical levels and time steps of 1h. For the isotope-enabled 110 version, water isotopes HD^{16}O and H_2^{18}O were added as two separate water species in the atmospheric model, and as tracers in the ocean model. Fixed isotope fractions are added to a fixed volume gridbox of the ocean and experience changes due to evaporation, precipitation, and runoff through a virtual isotope flux, altering the $\delta^{18}\text{O}$ ratio in the top level of the ocean accordingly (Tindall et al., 2009). The land surface and vegetation evolves dynamically and is based on TRIFFID (Cox, 2001) with timesteps of 5yr.

115 Compared to instrumental observations, the model represents sea surface temperature (SST), ice sheet, and ocean heat content well (Gordon et al., 2000). The freshwater hydrological cycle in the model shows only a slight overestimation in the local evaporation (Pardaens et al., 2003). The model simulates the major isotopic fractionation effects defined by Dansgaard (1964) (e.g. the latitude effect, the amount effect, and the continental effect) appropriately compared to GNIP data (Zhang et al., 2012). Additionally, a broad agreement in isotopic output with GNIP data in the general spatial distribution can be observed 120 and the general oxygen isotopic ratio features like latitude effect, amount effect, or the continental effect are captured well (Tindall et al., 2009). As such, iHadCM3 captures large scale features of climate and oxygen isotope ratios while remaining computationally efficient for the simulation of timescales such as the last millennium. The basic characteristics and boundary conditions of the last millennium simulations used in this analysis, which are identified with the *LM* prefix, are listed in Tab. 1.

125 2.2 The speleothem isotope dataset

The oxygen isotope ratio measured in speleothems is subject to many processes, starting from the source water which is influenced by the atmospheric circulation and climate. Therefore, the amount of precipitation, its composition, the annual mean temperature, and the variability of these events is in part imprinted in the archive. A comprehensive summary of the processes involving speleothem growth can be found in Fairchild and Baker (2012).

130 Vegetation above the cave has an impact on the source water through evapotranspiration, where the meteoric $\delta^{18}\text{O}$ is subject to additional fractionation processes and seasonal effects (Haude, 1954; Thornthwaite and Mather, 1957; Wackerbarth et al., 2010). Filter processes and transportation through the soil and upper karst influence the signal and may lead to varying transit



Table 1. Basic characterization of the *LM1*, *LM2* and *LM3* last millennium simulations.

Years	850-1850 CE 1100 - 100BP
Orography	fixed to 0BP
Orbital Parameter	fixed to 0BP
GHG	well mixed CO ₂ , CH ₄ , NO ₂ and other trace gases Schurer et al. (2014), Schmidt et al. (2012)
Vegetation	dynamic (TRIFFID (Cox, 2001))
Total Solar Irradiance	Steinhilber et al. (2009) Wang et al. (2005) Schurer et al. (2014)
Volcanic Forcing	Crowley and Unterman (2013)

times between several minutes and multiple years (Jean-Baptiste et al., 2019) at different drip sites within the same cave. The change in the partial CO₂ pressure conditions between the cave environment and the surface (Fairchild and Baker, 2012) in the lower epikarst makes the drip water degas and precipitate calcite in a fractionation process, which consequently forms a speleothem (Tremaine et al., 2011).

Varying environmental conditions within the cave can also be imprinted in the isotopic signal and may pronounce or attenuate the climate signal (Fairchild and Baker, 2012). Inside the speleothem, interactions with the cave environment or water inclusions within the mineral are still possible and, therefore, may further change the $\delta^{18}\text{O}_{\text{speleo}}$ archived in the speleothem.

The oxygen isotope composition of dripwater is influenced by all above-mentioned factors. Due to the multivariate processes impacting speleothem growth, the interpretation of the $\delta^{18}\text{O}_{\text{speleo}}$ signal is not straightforward, although systematic evaluation has identified patterns of similar climate influence based on modern observations (Baker et al., 2019). Proxy System Models (PSMs), where the input signal modification is modeled based on known processes in the karst may also help with the interpretation (Evans et al., 2013; Dee et al., 2015). PSMs of varying complexity have been proposed from the simple exponential decay filter, mimicking karst mixing (Dee et al., 2015) with the delay time as the single tunable parameter, to full-blown karst system models with numerous parameters describing soil water and gas equilibration or carbonate bedrock dissolution (Owen et al., 2018).




Figure 2. Site locations of the SISAL database on a global karst map (brown shadings Williams and Ford, 2006). The sites with entities that pass the prerequisites for our analysis are marked in colored triangles. These entities cover at least a period of 600yr within the last millennium and have a minimum of 30 (red), 20 (orange), or 10 (purple) $\delta^{18}\text{O}$ measurements and two dating points in this period. All other sites in the SISAL database v.2. are marked with a black dot. The nine clusters used in the analysis here are North America (c1, 12 entities), South America (c2, 13 entities), Europe with North Africa (c3, 21 entities), Southern Africa (c4, 2 entities - too few for systematic analysis), Middle East (c5, 6 entities), India and Central Asia (c6, 8 entities), East Asia (c7, 18 entities), South East Asia (c8, 4 entities), and New Zealand (c9, 3 entities).

The Speleothem Isotopes Synthesis and Analysis is an international working group, collecting speleothem datasets in a quality-controlled and cross-references database with rich metadata for samples and dating procedures (Atsawawanunt et al., 2018; Comas-Bru et al., 2020). The second version of the database SISAL v.2. includes measurements of stable ^{13}C and ^{18}O isotopes on speleothems of 691 individual entities from 294 globally distributed sites (Comas-Bru et al., 2020). In order to provide a comprehensive and reliable analysis, we only use data from entities which are not superseded (*entity_state = current*) and that cover at least a 600y period within the analysis period (850-1850CE). Furthermore, records considered must have at least two radiometric dates, or one radiometric date (in the analysis period) and be marked as actively forming at the time of collection, or be lamina counted. We only check for dates that are marked as *used*, indicating that they are known to have been used in the original chronology in the database. Samples without sample or depth information, are omitted. In the analysis, we filter the database adapted to the requirements of the different analyses, as depicted in Fig. 2. For the last millennium, we remain with 112 records from 92 different sites with at least ten isotopic measurements, that we used for the assessment of the mean $\delta^{18}\text{O}$ offset, and 85 records from 71 sites with at least 30 isotope samples for the correlation and network analyses.

For each U/Th-dated speleothem, SISAL v.2. provides the original age model (if available), and new possible age-models based on up to seven methods. On average, four new age-models are available. Methods include linear interpolation, linear regression, Bchron (Haslett and Parnell, 2008) as adapted by Roesch and Rehfeld (2019), Bacon (Blaauw and Christeny, 2011), Oxcal (Ramsey, 2009), copRa (Comas-Bru et al., 2020, modified R version after Breitenbach et al. (2012)), and StalAge (Scholz and Hoffmann, 2011). Details on the automated age modeling procedure are given in Roesch and Rehfeld (2019) and Comas-Bru et al. (2020). For each entity and ensemble method, one median best-fit estimate with confidence intervals, and



between 129 and 7737 age-models based on perturbations of the radiometric ages are available (Comas-Bru et al., 2020). These ensembles are available for 69 of the 87 entities that we used in the network analysis, resulting in a total of 464383 ensemble age-models in our analysis.

170 3 Methods

3.1 Speleothem analysis and drip water conversion

To increase the robustness of the results, we maximize the number of records by adaptive filtering of the database (Fig. 2). For calculations involving the time-averaged $\delta^{18}\text{O}$ -values, we only use speleothem data with at least 10 $\delta^{18}\text{O}_{\text{speleo}}$ measurements and two dating points within a 600yr period in 850-1850CE. For variance analyses, we demand at least 20 and for spectral as
175 well as correlation and network analyses 30 $\delta^{18}\text{O}_{\text{speleo}}$ measurements. For the investigation of spatial correlation patterns by network analysis, as explained in detail in Sec. 3.3, the set of speleothems is divided into nine regional clusters (Fig. 2). We primarily use the chronologies provided by the original authors, but test for the sensitivity to age-modeling choice by considering the age model ensembles (details below in Sec. 3.3.)

180 Within the last millennium, we remain with 17 aragonite and 89 calcite speleothems with 10 or more $\delta^{18}\text{O}$ sampled. We exclude six speleothems of mixed mineralogy. The $\delta^{18}\text{O}_{\text{speleo}}$ signal of calcite and aragonite speleothems is converted to its drip-water equivalent ($\delta^{18}\text{O}_{\text{dw.eq}}$) relative to the V-SMOW standard as in Comas-Bru et al. (2019). For calcite, we use the empirically-based fractionation formula of Tremaine et al. (2011)

$$\delta^{18}\text{O}_{\text{dw.eq}} = \delta^{18}\text{O}_{\text{calcite}} - \left(\left(\frac{16.1 \cdot 1000}{T} \right) - 24.6 \right), \quad (1)$$

185 where T is in K and $\delta^{18}\text{O}$ in units of ‰. For aragonite, we use the fractionation factor from Grossman and Ku (1986)

$$\delta^{18}\text{O}_{\text{dw.eq}} = \delta^{18}\text{O}_{\text{arag.}} - \left(\left(\frac{18.34 \cdot 1000}{T} \right) - 31.954 \right). \quad (2)$$

Here, temperature values T represent the local cave temperature in units of K. These are often not available. The annual mean temperature on the surface above the cave can, however, serve as a surrogate for local cave air temperatures (Fairchild and Baker, 2012). Both for aragonite and calcite drip water conversion, we use the simulated annual mean temperatures at the
190 cave location, down-sampled to the temporal resolution of the record. Note that, as a consequence, the conversion changes the time-averaged mean and the variance in our analysis. Finally, the V-PDB to V-SMOW conversion from Coplen et al. (1983) is used.

$$\delta^{18}\text{O}_{\text{SMOW}} = 1.03092 \cdot \delta^{18}\text{O}_{\text{PDB}} + 30.92. \quad (3)$$



Whenever we directly compare simulation output values with the speleothem records, e.g. when comparing means, variances, or spectra, we use $\delta^{18}\text{O}_{\text{dw.eq}}$, accounting for the different mineralogies. The conversion would, however, add an extra source of uncertainty in correlation analyses, as it implicitly builds on transient simulation data. Therefore, we denote the raw values of $\delta^{18}\text{O}$ measured directly in calcite or aragonite matrix by $\delta^{18}\text{O}_{\text{speleo}}$ and focus on those in the network and correlation analyses.

3.2 Statistical tests and time series processing

Speleothems form naturally, and therefore provide irregular time series with reconstructed and uncertain observation time series (Rehfeld and Kurths, 2014).

We account for this in our assessment as outlined below. Temperature, precipitation, and isotopic data are extracted from the simulation at cave locations by bi-linear interpolation. Annual mean values for temperature and precipitation are formed by averaging over all months from April onwards to March of the following year. Differences in mean are given in $\Delta\delta^{18}\text{O} = \delta^{18}\text{O} - \delta^{18}\text{O}_{\text{dw.eq}}$ (model-data difference) and variance ratios in the record's variance divided by the variance of the simulation at the cave location ($\text{Var}_{\text{Rec}}/\text{Var}_{\text{Sim}}$). If not explicitly stated otherwise, we always provide 90% confidence intervals by bootstrapping (Efron and Tibshirani, 1986) with 1000 repetitions. To reduce potential bias due to the irregular spatial distribution of cave sites, we use *area-weighting* in spatial mean estimates, where stated. This is done by calculating gridbox-means of all speleothems within a $3.75^\circ \times 2.5^\circ$ gridbox similar to the simulation, which is then area-weighted across latitudes, following Marcott et al. (2013).

While the simulation data is available at monthly basis, the proxy time series are irregular and at annual or lower resolution. Therefore, the simulation data at cave location is *down-sampled* to the record's reconstructed time axis by block averaging. The *power spectral density (PSD)* of a time series over a finite interval of time describes the distribution of power in frequency components of the time series. The integration over all spectral components yields the variance of the time series (Chatfield, 2003). For spectral analyses, the proxy records are interpolated to their mean resolution in a double interpolation and filtering procedure (following Laepple and Huybers, 2014a, b; Rehfeld et al., 2018; Dolman et al., 2020). Spectra of sufficient resolution can then be averaged to a mean spectrum over a certain frequency range (Kunz et al., 2020).

We test the impact of karst storage of drip water (Gelhar and Wilson, 1974; Dee et al., 2015) by applying an additional simplified aquifer recharge model style filter (hereafter *karst filter*). The impulse response of the Green's function depends solely on the transit time τ , as $g(t) = 1/\tau \cdot e^{-t/\tau}$, with $t > 0$. The Green's function is convolved with the simulated input $\delta^{18}\text{O}$ or $\delta^{18}\text{O}_{pw}$ signal to obtain the simulated karst-filtered signal in the cave.

The *correlation* of irregular time series is estimated by Person-correlation for irregular time series (Rehfeld et al., 2011; Rehfeld and Kurths, 2014). The *signal-to-noise ratio (SNR)* is estimated from the estimated cross-correlation \hat{r}_{ij} between two time series i and j by calculating $\text{SNR} = \hat{r}_{ij}/(1 - \hat{r}_{ij})$, as described by Fisher et al. (1985). If more than two estimates are available, e.g. at the gridbox level, the median between all possible combinations of cross-correlations between the time series is used. We account for age-model sensitivity by calculating cross-correlation estimates for all possible combinations of



available age-model ensembles (Comas-Bru et al., 2020). The resulting *best selection tuning* indicates the strongest absolute significant correlation that can be obtained from age-model selection a pair of records.

230 3.3 Spatial correlation via network analysis

Networks are practical representations for complex systems with interacting components and can be used to analyze dynamics in the climate system (Tsonis et al., 2006; Tupikina et al., 2014; Rehfeld et al., 2013). Here we use a network with n nodes, where n is the number of SISAL v.2. entities that fulfill the sampling criteria. The speleothem entities are joined in pairs by edges or links, where the $n \cdot (n - 1)$ links are formed if the cross-correlations $\hat{r}_{i,j}$ between two speleothem entities i and j , are
235 significantly different from zero with a p-value of $p_{i,j}$.

We split the network into sub-networks by hierarchical distance-based clustering of the node locations. The cluster that includes all East Asian caves is manually split into two clusters, one for East Asia (all caves above 20°N) and a cluster of South East Asia (all caves below 20°N). Links in the plots (Fig. 8) are visualized if they are stronger than a certain threshold $|r| > r_{5\%}$, where $r_{5\%}$ is minimum correlation strength of the 5% absolute strongest correlations ('fixed link density').

240 4 Results

4.1 Assessing model-data differences in time-averaged $\delta^{18}\text{O}$

We first compare the mean SISAL v.2.-record $\delta^{18}\text{O}_{\text{dw.eq}}$ and iHadCM3 $\delta^{18}\text{O}$ to assess potential model biases. Annual mean temperature and precipitation fields (Fig. 3a,b), and the mean modeled $\delta^{18}\text{O}$ together with the mean $\delta^{18}\text{O}_{\text{dw.eq}}$ in the SISAL records (Fig. 3c) is shown in Fig. 3. Shown and described are the fields and results for LMI, results for the other ensemble
245 members are generally very similar and given in supplementary table ST1. The major oxygen isotope ratio depletion features as described by Dansgaard (1964), can be distinguished. Modeled values show progressive depletion towards higher latitudes, the interior of continents, and towards regions with high precipitation amounts.

These offsets between modeled and measured $\delta^{18}\text{O}$ ($\Delta\delta^{18}\text{O} = \delta^{18}\text{O} - \delta^{18}\text{O}_{\text{dw.eq}}$) show a heterogeneous pattern (Fig. 3d). Generally, modeled values appear to be more depleted overall than the mean values of speleothem $\delta^{18}\text{O}_{\text{dw.eq}}$, except in the NH
250 extratropics. There are some localized clusters and individual sites with large positive and negative differences. One example is site 159 (eID = 531, Qunf cave in Oman) which is visible as a dark blue dot in Fig. 3d. The $\delta^{18}\text{O}_{\text{dw.eq}}$ record shows a mean enrichment in calcite ($\delta^{18}\text{O}_{\text{dw.eq}} = 0.49\text{‰}$), while the simulation shows much more depletion ($\delta^{18}\text{O} = -6.9\text{‰}$). This results in a fairly large model-data difference of $\Delta\delta^{18}\text{O} = -7.4\text{‰}$. The site stands out against the regional cluster, pointing towards a local site-specific factor influencing the offset. Site 277 (eID = 598, Huagapo cave in Peru) visible as a dark red dot in Fig. 3d,
255 shows a strong depletion in calcite ($\delta^{18}\text{O}_{\text{dw.eq}} = -13.7\text{‰}$) while the simulation is not as strongly depleted ($\delta^{18}\text{O} = -5.3\text{‰}$). This results in a large positive offset of $\Delta\delta^{18}\text{O} = 8.4\text{‰}$. The cave is located at an altitude of 3850m above sea level, whereas model altitude at the gridbox is close to sea-level. This should explain part of the offset.




Figure 3. Characterization of the mean state of the simulation (LM1): Shown are (a) mean annual surface temperature, (b) precipitation, and (c) $\delta^{18}\text{O}_{pw}$, including $\delta^{18}\text{O}_{dw.eq}$ at cave sites in drip water equivalents. Note the logarithmic color scale. Point-wise differences between the mean simulated $\delta^{18}\text{O}$ and proxy-based $\delta^{18}\text{O}_{dw.eq}$ (d) show anomalies. Spatially aggregated differences at the global and cluster level for simulations LM1-LM3 are given in supplementary table ST1.

At the regional scale, the largest cluster offset can be seen over China and East Asia (c7) $\Delta\delta^{18}\text{O} = +2.2\text{‰}$ ($-0.5, 4.9$, 90% confidence interval). However, the most consistent negative difference is visible over neighboring Indonesia (c8) $\Delta\delta^{18}\text{O} = -2.7\text{‰}$ ($-4.9, -0.6$). The smallest differences are found in Europe with $\Delta\delta^{18}\text{O} = +0.5\text{‰}$ ($-1.8, 2.7$) and India and Central Asia $\Delta\delta^{18}\text{O} = 0.7\text{‰}$ ($-4.4, 5.7$). Overall, the simulated $\delta^{18}\text{O}$ is smaller than the $\delta^{18}\text{O}_{dw.eq}$ measured in speleothems ($\Delta\delta^{18}\text{O} = -0.1\text{‰}$ ($-4.6, 4.4$), with no clear spatial pattern for the offsets distinguishable from the map. The gridbox-level area weighted global mean difference is -0.02‰ ($-0.22, 1.00$) for LM1.

We further explore the impact of site conditions on the model-data offset (Fig. 4). We find a decreasing $\delta^{18}\text{O}_{dw.eq}$ towards northern higher latitudes (Fig. 4a), and most notably, a dependency of $\delta^{18}\text{O}_{dw.eq}$ on the local mean annual temperature (Fig. 4b). We see more positive offsets in the northern hemisphere and mostly negative offsets in the southern hemispheres (Fig. 4c).

The offsets also show a strong influence of temperature (Fig. 4d) and elevation (Fig. 4i), which are both controlling factors during the isotopic fractionation process. The elevation difference between the simulation and the record spans from a 1332m higher elevation in the simulation (eID = 538 in Shenqi cave in China) and 3065m higher elevation in the records (eID = 598 in Huagapo cave in Peru, visible outlier in Fig. 4c,d). Here, the offsets increase with increasing absolute difference (Fig. 4j). The offset shows a weak correlation with precipitation (Fig. 4f), both in the annual mean and for the boreal winter/summer season




Figure 4. Systematic comparison of climate variables and cave parameters on $\delta^{18}\text{O}_{\text{dw.eq}}$ and the offset $\Delta\delta^{18}\text{O}$ to the simulation. Shown are the absolute values of $\delta^{18}\text{O}_{\text{dw.eq}}$ against: (a) site latitude, and (b) simulated local annual mean temperature, and the model-data difference against (c) latitude, (d) simulated mean annual temperature, (e) geology surrounding the cave ('?' means unknown geology), (f) mean annual (g) DJF and (h) JJA precipitation amount as well as (i) cave elevation, (j) the elevation difference between the model grid and actual cave, and (k) the overall cover thickness above cave. Symbols denote calcite (black circles) or aragonite (blue triangles) specimens. An unweighted linear regression (red line) is added for illustration, but without consideration of significance.

(see DJF and JJA precipitation in Fig. 4g,h). No relation can be seen with mineralogy, parent rock (Fig. 4e) or cover thickness
 275 (Fig. 4k).

4.2 Assessing model-data differences in the local variance of $\delta^{18}\text{O}$

To analyze how similar the variability of the isotopic signal is in speleothems and the iHadCM3 climate model, we compare the total variance in both over the last millennium. The global distribution of variance ratios between $\delta^{18}\text{O}_{\text{dw.eq}}$ and $\delta^{18}\text{O}$ (Fig. 5a) shows overall higher variability in the speleothem records than in the simulation, with local exceptions. This is also
 280 corroborated by the density plots of the ratio for both $\delta^{18}\text{O}$ and $\delta^{18}\text{O}_{\text{pw}}$ in Fig. 5b,c. Generally, the observed proxy variance is roughly two times higher than that of the down-sampled simulation $\delta^{18}\text{O}$ at the cave location (median of the histogram at 2.0 (1.4,2.8) in Fig. 5b,c). This is consistent with the predominance of red-shaded variance ratio visualizations in the




Figure 5. (a) Spatial visualization of the site-based dimensionless variance ratio V_{Rec}/V_{Sim} based on LM1. Aggregated density plots of the variance ratio of $\delta^{18}O$ (b) and precipitation-weighted $\delta^{18}O$ (c) for the raw records ('full', black lines) and where the simulation has been down-sampled to the record resolution ('down-sampled', red lines) illustrate the variance loss due to temporal averaging in the archive (uses LM1-3).

spatial view indicating $Var_{Rec}/Var_{Sim} > 1$ (Fig. 5a). However, there is a clear impact of averaging on the total variance, as down-sampling results in a variance ratio above unity. Overall, this shows a discrepancy between the variance observed in $\delta^{18}O_{dw.eq}$ and the simulated variance at the cave location over the total time period.

The highest variance ratio for down-sampled $\delta^{18}O_{pw}$ is found in Jiuxian cave in China (eID 330, with a variance ratio of 41.2), the lowest variance ratio in Dandak cave in India (eID 130, with a variance ratio of 0.2), while neighboring caves show very different variance ratios. As the modeled patterns are fairly smooth (Fig. 5a), this indicates a large heterogeneity of the speleothem data from the cave environment. We find no strong or significant relationship between variance or variance ratios to any tested climate or cave parameter (SF4 where we show a similar figure to Fig. 4 but for variance ratios).

4.3 Assessing $\delta^{18}O$ variability at interannual to centennial timescales

We extend the analysis of total variance (Fig. 5) to the time scale dependent variance (Fig. 6) to better explore variability on interannual, decadal, and centennial time scales as compared to the total variance over the last millennium. The spectra in Fig. 6d give an insight into the variability over different time scales and the representativity of records for reconstruction resolution.

On the left side (Fig. 6a-c), the time series of $\delta^{18}O_{dw.eq}$ of eID 240 (Bunker cave, Germany) is depicted (Fig. 6a), together with the simulated $\delta^{18}O$ at the cave site at different temporal resolutions (Fig. 6b,c), including karst-filtered $\delta^{18}O$. Comparing Fig. 6a-c visually, different levels of variance can already be distinguished e.g. between the filtered and unfiltered simulated data. The iHadCM3 $\delta^{18}O$ spectrum of the yearly resolved signal has similar variance over all frequencies, and shows a fairly constant PSD (Fig. 6d). Variance at decadal timescales (i.e. the PSD for higher frequencies) is just as high as the variance on centennial time scales (i.e the PSD for lower frequencies).




Figure 6. Variability on different time scales through comparison of measured $\delta^{18}\text{O}_{\text{dw.eq}}$ and simulated $\delta^{18}\text{O}$ time series as well as of their spectra. (a-c) Example time series of eID 240 in Bunker cave (Germany) (Fohlmeister et al., 2012). (a) The measured $\delta^{18}\text{O}_{\text{dw.eq}}$ in the speleothem, (b) the iHadCM3 simulated $\delta^{18}\text{O}_{pw}$ at the cave location with two filters (3 and 9 years) and (c) the simulated $\delta^{18}\text{O}_{pw}$ but down-sampled to the same temporal resolution as in (a) with 3 year filter. (d) Power spectral density (PSD) of mean spectra of simulated $\delta^{18}\text{O}_{pw}$ at the cave site in yearly resolution (blue), down-sampled to the caves resolution (red) and mean spectrum of the $\delta^{18}\text{O}_{\text{dw.eq}}$ of the records (black), including the karst-filters as shown in (a-c). The spectra are area-weighted and averaged over the three simulations (LM1, LM2 and LM3). The colors for the example eID in (a-c) correspond to the colors of the mean spectra over all entities in (d).

After down-sampling to the irregular resolution of the record, the simulated spectrum loses power in the higher frequency range. Comparing for example the time series in Fig. 6c to the spectra in Fig. 6d, the down-sampled spectrum indicates lower variability than the annual resolution spectrum on decadal timescales. On centennial timescales, both spectra display similar variability. Contrasting Fig. 6b to Fig. 6c, this loss in decadal time scale variability is also visible on the time series level.

The proxies' spectra have even fewer frequency components in the high frequency range, due to the lower temporal resolution. They do, however, show a higher PSD at lower frequencies. The records are, therefore, less variable on decadal time scales, and more variable than both the down-sampled and the full resolution simulated $\delta^{18}\text{O}_{pw}$ on centennial time scales.

An additional impact of karst processes and storage on the $\delta^{18}\text{O}$ variability could be expected. To test the impact of this, we apply simple karst filters (see Sec. 3) with increasing filter length and test whether they reduce the spectral mismatch. Filters of different length resulted in increasing spectral slopes with increasing transit times. A 3yr filter for the down-sampled $\delta^{18}\text{O}$ achieves equivalent variance trends as the record spectrum with less power on decadal timescales. It eventually flattens again for longer timescales, without exceeding the PSD of the unfiltered signal, such that it is less variable than the proxies on longer time scales. A full set of individual spectra (full simulation, down-sampled, record spectrum, and all filters) for all entities used in this analysis can be found in SF5 and SF6.




Figure 7. Correlation fields of simulated $\delta^{18}\text{O}$ and the related climate variables surface temperature (a) and precipitation (b) for simulation LM1 ($|c| > 0.2, p < 0.1$). Colored symbols give the correlation between simulated climate variables and the $\delta^{18}\text{O}_{\text{dw.eq}}$ of the speleothem records. Empty tiles mask insignificant correlations. Black dots show cave locations with insignificant correlations.

4.4 Climatic drivers of $\delta^{18}\text{O}$ variability

To distinguish main important climatic drivers for specific areas for $\delta^{18}\text{O}$ both in the simulation and in speleothems, we correlate simulated $\delta^{18}\text{O}$ with the simulated temperature (Fig. 7a) and the precipitation signal (Fig. 7b) on a gridbox level after temporal down-sampling. Grey (empty) tiles indicate insignificant correlation estimates. The correlation between $\delta^{18}\text{O}_{\text{speleo}}$ and the climate variable is shown.

We see strong correlations of simulated $\delta^{18}\text{O}$ to simulated temperature at high latitudes as well as over some landmasses (background in Fig. 7). The speleothem signals show positive as well as strong negative correlations. The absolute highest correlation is found for eID 124 in Leviathan cave in the USA ($c = -0.4$ ($-0.7, 0.1$)). In the simulation, this correlation is locally positive, which indicates that the simulated temperature is a positive $\delta^{18}\text{O}$ driver in the general area in the model. The correlation of the simulated climate and the record's $\delta^{18}\text{O}_{\text{speleo}}$ is, however, negative.

The correlation between the simulated precipitation and $\delta^{18}\text{O}$ is especially strong in the tropics. We find the highest absolute correlation for eID 523 in Gempa Bumi cave in Indonesia ($c = -0.5$ ($-0.7, -0.1$)). Here, the background also shows a negative correlation.

Comparing the two proposed climatic drivers of $\delta^{18}\text{O}$ variability, we observe that the correlations to temperature are higher in the higher latitudes, while correlations to the precipitation appear more important in the tropics. A fairly clear zonal structure of correlations between the climate and oxygen isotope ratio fields are visible in the model. However, only few records show a significant correlation ($p < 0.1$). We find 18, 15 and 22 significant correlations from 87 entities for temperature for the three LM ensemble runs respectively, and 14, 7, and 10, respectively, for precipitation. No clear climatic driver can, therefore, be extracted alone from record correlation results. Fewer records show significant correlation to both climate variables. The direct correlation of the time series of the simulated and proxy-based $\delta^{18}\text{O}$ results in only 19, 17, and 19 significant correlations from 87, i.e. at around 20% of the sites.




Figure 8. Network spanned by the 5% strongest absolute correlations of simulated iHadCM3 $\delta^{18}\text{O}$ at the SISAL cave sites (a, down-sampled). All model-based between-site-correlations are shown in the distance-binned boxplot (b). Network visualizations (c) and distance-binned boxplot (d) of the cross-correlations between SISAL site $\delta^{18}\text{O}_{\text{speleo}}$ for the original age models. The color values indicate the 5% strongest correlations in network and boxplot. The LOESS smoother (span = 0.2) in the boxplots indicate the correlation for the original chronology (red) as well as the absolute highest correlation through selection of age-models (blue).

4.5 Similarity measures and network analysis

Computing all statistical similarity between the $\delta^{18}\text{O}_{\text{speleo}}$ signals within a cave ('site-level-correlation') or across nearby caves (regional or gridbox-level correlation) yields a measure of representativity useful for model comparison and uncertainty
340 assessment. The networks in Fig. 8 are based on the simulated signal (down-sampled) $\delta^{18}\text{O}$ (Fig. 8a,b) and for $\delta^{18}\text{O}_{\text{speleo}}$ (Fig. 8c,d).

Network links are based on the 5% highest absolute correlations. The highest correlations are found at close proximity for the models (Fig. 8a,b), whereas links across a wide range of distance can be seen in the proxy data (Fig. 8c,d). High local correlations for the model data can be expected, as the simulated $\delta^{18}\text{O}$ within one cave will be the same, and only differs on a
345 temporal scale after the adjustment to the entity's temporal resolution (down-sampling). The mean absolute correlation for the 5% strongest significant links in Fig. 8a) is $c = 0.42$ (0.41, 0.43). Comparing the down-sampled distance-to-correlation plot (Fig. 8b) to that on annual resolution analysis, an additional scattering of correlation estimates at longer than 2000km distance is visible. Results for annual resolution can be found in SF7.

The network of $\delta^{18}\text{O}_{\text{speleo}}$ does not display large-scale spatial patterns and no observable relationship between correlation
350 and distance. The mean absolute correlation for the 5% strongest significant links shown in Fig. 8c) is $c = 0.52$ (0.52, 0.53). Computing the networks based on the ensemble age models, and selecting the age models that maximize the absolute correlation between sites, amplifies both positive and negative correlation estimates, but does not change the correlation-to-distance relationship. Comparing the results for simulated $\delta^{18}\text{O}$ (Fig. 8a,b), and $\delta^{18}\text{O}_{\text{speleo}}$ (Fig. 8c,d), we obtain low correlations at the




Figure 9. Cross-correlation on site, gridbox, clusters and global scale for speleothem records and the locally interpolated model output for $\delta^{18}\text{O}$. 12 (18) sites (gridboxes) contain more than one speleothem entity with a total of 27 (45). At each aggregation level the correlations between all entities is shown for $\delta^{18}\text{O}_{\text{speleo}}$ (white bars), and the down-sampled model output $\delta^{18}\text{O}$ at cave locations (blue bars). Different temporal scales (original resolution and 100yr-timescale (t.sc)) are compared as well as the age-model ensemble that gives the highest absolute correlation (dark green bars). Clusters are indicated with the number of speleothem entities in brackets, where c4, c5, c8, and c9 are not included because they contain too few entities. c6/ICA is the India and Central Asia cluster, c7/CEA is the China and Eastern Asia cluster.

local scale.

355

We can also investigate relationships using regional networks. For this, we look at correlation on different spatial levels and separate the network analysis from Fig. 8 by sites, gridboxes, and clusters. Cluster c4, c8, and c9 contain less than four entities and are excluded. We check for representativity on different time scales of record resolution (white, and dark blue) and a 100yr Gaussian smoothing filter (grey and light blue) and on different spatial scales using boxplots (Fig. 9).

360

At the site level, we find 27 entities within 12 sites that contain at least 2 entities. The median correlation of these 28 pairs is $c_{raw} = 0.25$ ($-0.17, 0.33$). On a 100yr timescale, this increases to $c_{100} = 0.42$ ($-0.46, 0.49$). The SNR gives a measure of the relative importance of non-climatic overprints on the proxy signal. We obtain a local SNR estimate of 0.5 ($0.4, 1.1$). On the gridbox level (45 entities in 18 gridboxes), we find a median correlation of $c_{raw} = 0.23$ ($0.2, 0.25$) (100yr timescale: $c_{100} = 0.34$ ($0.27, 0.47$)). As on spatial resolutions below gridbox level, correlations between simulated $\delta^{18}\text{O}$ are not meaningful, which is why the analysis in Fig. 9 shows only the correlations between record $\delta^{18}\text{O}_{\text{speleo}}$ and not those of the simulated $\delta^{18}\text{O}$ on a site and gridbox level.

365

For regional clusters, the correlation between proxies shows positive and negative median values. In the simulation, the median values are always positive. For clusters containing more than ten records detailed correlation maps including correlation matrices are depicted for Europe (SF8), China and Eastern Asia (SF9), South America (SF10), and North America (SF11) where very red maps and matrices can be found, indicating mostly positive correlations for the down-sampled simulation when

370



compared to more blue ones in the records, indicating that also negative correlations are present. On the global scale, the median correlation between all records is slightly positive ($c_{raw} = 0.1$ $(-0.09, 0.11)$), 100yr timescale: $c_{100} = 0.13$ $(-0.14, 0.17)$), whereas for the simulation this median is positive ($c_{raw} = 0.06$ $(0.06, 0.06)$) and strongly enhanced at centennial timescale to $c_{100} = 0.76$ $(0.73, 0.81)$.

375 By the selection of the age-model that maximizes the absolute correlation, we obtain a significant positive correlation at site level and a stronger significantly positive correlation at gridbox level. A detailed table with median correlations and SNR using the original chronologies as well as using the described age-model selection on different spatial levels is shown in ST2. Calculating correlations for different age-model ensembles was only done for the 69 entities, where both age-model ensembles were available (U/Th-dated entities) in Comas-Bru et al. (2020), and our strongest criteria were matched.

380 5 Discussion

5.1 $\delta^{18}\text{O}$ model-data comparison in mean and variance

In our study, we found the last millennium mean iHadCM3-simulated $\delta^{18}\text{O}$ to agree well with the mean state of the measured $\delta^{18}\text{O}_{\text{dw.eq}}$ (Fig. 3). The average unweighted offset of $\Delta\delta^{18}\text{O} = 0.1\text{‰}$ $(-4.6, 4.4)$ was small compared to the total $\delta^{18}\text{O}$ and the area weighted standard deviation of $\sigma^2 = 0.78\text{‰}^2$ $(0.77, 0.8)$ of the global simulated mean $\delta^{18}\text{O}_{pw}$. Measured $\delta^{18}\text{O}_{\text{dw.eq}}$
385 followed general isotopic signature patterns as described by Dansgaard (1964) but spatial pattern for the offsets were not distinguishable.

Baker et al. (2019) distinguished between temperature zones of climatic controls on $\delta^{18}\text{O}$ in offset analyses on drip water. They find a stronger influence of seasonality of precipitation in warmer climates, highlighting the importance of a karst-recharge model. Here, we also observed a strong temperature dependency reflected in the offset and $\delta^{18}\text{O}_{\text{dw.eq}}$ over the last
390 millennium, showing the influence of fractionation and other cave internal processes on the $\delta^{18}\text{O}$ in drip water (Fig. 4). The highest offsets on the Northern Hemisphere possibly indicate a stronger influence of the continental effect. Still, from the records alone and with no karst-recharge information, we were not able to distinguish specific climatic control regions. This requires a more thorough analysis including monitoring data.

We found no evidence that the variance ratio between record variance and simulated variance is related to the offset between
395 simulation and records (SF4 is similar to Fig. 4, with variance ratios instead of $\Delta\delta^{18}\text{O}$). Specifically, there is no correlation between site-level offset and site-level variance ratio (results not shown, $r = 0.1$ $(0.0, 0.3)$, $p = 0.2$). In general, the results based on the total variance over the last millennium are highly consistent, where variability in models is mostly due to volcanic forcing (Jungclauss et al., 2010). Differences in variance can, to some extent, be attributed to the sample resolution of the records, whereas down-sampling of simulated $\delta^{18}\text{O}$ decreases the variability on decadal time scales. The resolution to which
400 the simulation is temporally aggregated impacts whether the variance in the simulation appears to be larger or smaller than in the records. The variance over the last millennium in the records is, overall 2.0 $(1.4, 2.8)$ times as high as the simulated down-sampled variance in Fig. 5.



Furthermore, the simulated $\delta^{18}\text{O}$ time series at the cave sites show less variability on centennial timescales than the time series of the records. This is true even when comparing the same temporal resolutions (time scale dependent variance depicted
405 in SF12 for $\delta^{18}\text{O}_{\text{dw.eq}}$, yearly-resolution $\delta^{18}\text{O}$ and down-sampled $\delta^{18}\text{O}$). This is in agreement with the findings of Laepple and Huybers (2014b), who compared simulated and reconstructed temperature variability across different timescales and found that the model-data discrepancies increased with time scale, particularly on a regional level.

If we assume that paleoclimate archives record climate variability correctly, and that the proxy-climate relationships are not timescale-dependent or transient, discrepancies at the centennial timescale could in part be explained by the models' underestimation of variability, in particular on centennial time scales. However, we find little regional consistency and high
410 heterogeneity in the variance estimates from the speleothem records. These findings point to the strong influence of cave internal processes or the impact of seasonally filtered data captured by speleothems, which is in agreement with McDermott et al. (2001).

5.2 Influence of the karst-filter

By delaying the simulated down-sampled signal through a simplified karst-filter with a transit time of 3yr, we obtained match-
415 ing equivalent power spectra for the simulation and the records. Studies observing transit time in karst systems find increases in drip rate after an increase in precipitation e.g. after days (Riechelmann et al., 2011). More complex tritium measurements show actual transit times of e.g. years for the Bunker cave in Germany (Kluge et al., 2010) to decades in the Villars cave in France (Jean-Baptiste et al., 2019), depending on the karst hydrology. The karst filter effectively reduces the temporal resolution of
420 the record beyond the nominal median of 5.6yr (Fig. 6). Such low-pass filtering to model drip water transit times has been used (Wackerbarth et al., 2010; Dee et al., 2015; Lohmann et al., 2013) to produce similar time lags of 2 – 10yr, indicating that the best fit mean time lag for our karst filter of 3yr (down-sampled) is a realistic estimate for transit times.

We find low interannual to decadal variability in the $\delta^{18}\text{O}_{\text{speleo}}$ signals recorded by speleothems (Fig. 6). In part, this is likely due to the average resolution of the records, which lies close to these timescales. Furthermore, mixing processes in the
425 soil and karst could play an important role, where soil $\delta^{18}\text{O}$ is found to have much lower variability than precipitation $\delta^{18}\text{O}$ (Tang and Feng, 2001).

On decadal timescales (shorter than 50yr), the karst filter reduced the resolution-adjusted variance by 34% (20,43), on longer timescales (longer than 50yr) by 4% (3,4) of the non-filtered down-sampled variance. The total filtered and down-sampled variance over the last millennium decreased by 14% (9,27) of the unfiltered down-sampled variance. Still, this is
430 equivalent to only 29% (23,38) of the record variance, as the filter only decreases variance on annual to decadal time scales. On centennial time scales the filter has little to no effect, so the record's variance on these time scales is not strongly affected.

5.3 Representativity of $\delta^{18}\text{O}$ at different spatial levels

A clear picture of the relationship between the climatic drivers for the simulation was distinguishable. However, no systematic pattern and few significant correlations were found for the speleothem records (Fig. 7). Accounting for seasonal sensi-
435 tivity could enhance the number of simulation-to-record correlations of SF13, which shows the selected strongest seasonal



correlation. However, this does neither enhance the overall correlation (histogram of correlation distribution using annual down-sampled time series and seasonal down-sampled time series in SF14), nor the SNR (results not shown). Still, the strong influence of seasonality suggests a dependency of $\delta^{18}\text{O}_{\text{speleo}}$ on certain seasons rather than the annual mean. Supporting this, SF15 shows a correlation map with the strongest seasonal correlation of $\delta^{18}\text{O}_{\text{speleo}}$ to the simulated climate variables temperature, precipitation, and $\delta^{18}\text{O}$ in precipitation. Further dripwater monitoring studies that characterize the seasonality of individual caves could help the understanding of which climatic signal is captured by speleothems and enhance comparability between different caves.

We found low spatial representativity of individual speleothems for sites, gridboxes, and regions when compared to the simulation (Fig. 8 and Fig. 9). We obtained stronger correlations between entities by selection of the best-matching age-model ensemble for entities where these ensembles were available. This age-model ensemble 'tuning' increased the median of correlations on site and gridbox-level by roughly a factor of 2, while also increasing the SNR by a factor of 3 and 2.5, respectively. However, no improvement could be observed on the cluster and global level. A detailed table of correlations and SNRs using the original chronologies as well as using the age-model ensemble selection that gives the highest absolute correlation is showed in the supplement table ST2. Testing other 'tuning' options, such as the consideration of only the 50% of the records at closest proximity within a cluster, or the 50% with the smallest mean offset showed no improvement (boxplot similar to Fig. 9 for the other selection criteria in SF16). We also found no correlation between the total variance and the number of significant links in the network ($c = -0.02(-0.23, 0.19), p = 0.8$).

A strong between-site variability has been attributed to controls of regional atmospheric circulation according to Lachniet (2009). We also find a strong heterogeneity in the recorded variance of $\delta^{18}\text{O}$ at the gridbox and cluster levels. In part, this can be due to heterogeneous temporal resolution, but could also be influenced by non-climatic overprints on the $\delta^{18}\text{O}$ signal up to the centennial scale.

This could be investigated by comparing the $\delta^{18}\text{O}$ and the $\delta^{13}\text{C}$ signal recorded within the cave to vegetation, climate, and landscape evolution archives in the region. However, representativity tests across Western Europe noted coherent $\delta^{18}\text{O}_{\text{speleo}}$ trends on glacial-interglacial time scales, where trends are less clearly expressed during the Holocene (Lechleitner et al., 2018). Therefore, this study could be extended to longer time scales, when longer transient isotope-enabled simulations become available.

5.4 Limitations

Simulated isotope variability is primarily dictated by the model's climatology and the complexity of its dynamics and hydrological cycle. We use a three member initial-condition ensemble from a single iGCM in this study. Therefore, all results relate to these iHadCM3 last millennium ensemble runs and the chosen radiative forcings. In this respect, a more thorough comparison with more simulations is needed in order to estimate the capability of models to simulate variability and to find common biases. However, the establishment of isotope-enabled GCMs requires substantial work for the addition of isotopic tracers and their evaluation, and the computational costs increase. This still inhibits the simulation of large transient ensembles with iGCMs over centennial to millennial and orbital time scales. Nevertheless, the three-member ensemble we provide could



470 also be used to test offline data assimilation methods, as suggested by Dalaiden et al. (2020). This might also help to better identify the climate factors that govern the speleothem archiving of $\delta^{18}\text{O}$ and its variability.

An uncertainty factor in our study comes from the temperature-dependence of the calcite- and aragonite-to-drip-water conversion. We calculated the adjustment factors using the simulated annual mean temperature at the cave location, sampled to the speleothem's temporal resolution. We take this simulated temperature as a surrogate for the longterm-changes of the inside-
475 cave air temperature. Knowing the actual temperature history of the caves better could strongly reduce the uncertainty, as a bias of $\Delta 1^\circ\text{C}$ in the simulated temperature would account for a change in $\delta^{18}\text{O}_{\text{dw.eq}}$ of approximately $\Delta 2\text{‰}$. A bias of $\Delta 1\text{‰}$ in the $\delta^{18}\text{O}_{\text{dw.eq}}$ however, accounts for a temperature change of 0.1°C for the lowest simulated annual mean cave temperature (3.1°C in Norway), and a change of 13.1°C for the highest simulated annual mean cave temperature (32.5°C in the tropics).

Also, our study focussed solely on $\delta^{18}\text{O}$ as one particular proxy for climate and environmental changes and not other
480 geochemical proxies that can be measured on speleothem samples (Kaufmann, 2003; Schwarcz et al., 1976) or a combination of proxies, which have the potential of a more thorough interpretation of a climate signal. A multi-proxy approach, such as in Fohlmeister et al. (2017) or Baker et al. (2017) who also include $\delta^{13}\text{C}$ along with $\delta^{18}\text{O}$, could offer deeper insights. Many proxies for climate processes, such as $\delta^{13}\text{C}$ cannot (yet) be implemented in GCMs, as their representation of the biology, physics and ecology is not sufficiently complex. Therefore, in order to consider those in the evaluation, time series have to be
485 calibrated to climatic and environmental parameters that are explicitly modeled. This would introduce additional uncertainty, that could counteract the added value of considering multiple proxies in the first place.

We have considered a regional to global view on speleothem $\delta^{18}\text{O}$ signal. Therefore, influences and processes known for individual cave systems could not be considered. For example, Kluge et al. (2013) account for kinetic fractionation changes over time via clumped isotope measurements, and Jean-Baptiste et al. (2019) were able to extract transit times of dripwater in
490 Villar cave. Considering these and other local factors might give deeper insight into individual speleothem records, but it is difficult to scale quantitatively and systematically. Nevertheless, including monitoring datasets from different caves globally might give deeper insight into the filter and fractionation processes involved, and PSM studies informed by the monitoring and local expertise throughout the database could help in further comparison studies.

6 Conclusions

495 We presented an ensemble of iHadCM3 last millennium simulations and compared the oxygen isotope ratios, temperature, and precipitation variability to oxygen isotope ratio observations from a large speleothem dataset (a subset of SISAL v.2.). Overall, time-mean patterns of oxygen isotope ratio were fairly similar in both. Considering total variance as well as the variability on different time scales, we observed that the effects of resolution adjustment and a convolution karst filter were sufficient to bring simulated and observed $\delta^{18}\text{O}$ spectra into good agreement. Still, total variability in the speleothem records is much higher than
500 in the simulation. This supports previous studies that found that climate models currently do not capture appropriate variability on centennial time scales.



505 However, we find that the climatological and environmental interpretation of $\delta^{18}\text{O}_{\text{speleo}}$ is not straightforward. We found low signal-to-noise ratios, which imply a low spatial representativity of individual entities. Furthermore, while regional climatic signals were distinguished in the simulation, the main climatic drivers for $\delta^{18}\text{O}_{\text{speleo}}$ at the regional scale were difficult to isolate. It is difficult to establish the size of the spatial footprint of representativity, the seasonality, and the relevant climatological and environmental parameters for reconstructions. Here, expert knowledge on local cave processes, environmental history, and, in particular, the availability of monitoring data are crucial to aid the interpretation of the climate signal. Inner cave and karst processes, which influence the seasonality of the input signal above the cave and inside the cave, may need to be taken under consideration. However, monitoring data for evaluation and potentially calibration of reconstructions are currently only available for a few sites (e.g. Tremaine et al. (2011)). Furthermore, some parameters, such as transit times, are difficult to measure (Jean-Baptiste et al., 2019).

515 Proxy system models that account for the cave internal fractionation processes may give a deeper insight into how climate variability is captured in speleothem archives. To gain a deeper understanding of the underlying concepts that influence the capability of speleothems to capture and resolve climate variability and the capability of models to simulate them, further model-data comparison studies are required.

Code and data availability. Code to reproduce figures and analyses in this paper are provided at <https://github.com/paleovar/iHadCM3LastMill> (Bühler and Rehfeld, 2020, (accessed September 10, 2020)). Model data has been uploaded to Pangea. The SISAL v.2. database (Comas-Bru et al., 2020) can be downloaded at <https://researchdata.reading.ac.uk/256/> (Comas-Bru et al., 2020, (accessed September 10, 2020)). We use R for the data analysis (R Core Team, 2020). The main packages are tidyverse (Wickham et al., 2019), ncd4 (Pierce, 2019), ggplot2 (Wickham, 2016), and raster (Hijmans, 2019). We use the nest R-package (<https://github.com/krehfeld/nest> (Rehfeld et al., 2011; Rehfeld and Kurths, 2014)) and the PaleoSpec package (github.com/EarthSystemDiagnostics/PaleoSpec (Kunz et al., 2020)).

Author contributions. KR and JB conceived this study. KR conducted the model simulations, with the advice of MH and LS. JB, MK and KR prepared the model data, and JB, CR and KR prepared the speleothem data. JB and KR wrote the paper. CR, MK and MH contributed to revisions. All authors approved of the final version of the manuscript.

525 *Competing interests.* The authors declare that they have no conflict of interest.

Acknowledgements. We thank the SISAL (Speleothem Isotopes Synthesis and Analysis) working group of the Past Global Changes (PAGES) programme. We thank PAGES for their support of their activity. We acknowledge Laia Comas-Bru, Denis Scholz, Nils Weitzel, Martin Werner, Eric Wolff for advice and helpful discussions and Beatrice Ellerhoff, Elisa Ziegler and Moritz Adam for helpful comments on text and figures. This work was carried out using the ARCHER UK National Supercomputing Service (<http://www.archer.ac.uk>). Financial



530 support for this study was given by the Deutsche Forschungsgemeinschaft (DFG, project RE3994-1/1 and RE3994-2/1) as well as the German Federal Ministry of Education and Research (BMBF, grant no. 01LP19260).



References

- Atsawawanunt, K., Comas-Bru, L., and Mozhdehi, S. A.: The SISAL database: a global resource to document oxygen and carbon isotope records from speleothems, *Earth System Science Data Discussions*, 2018.
- 535 Baker, A., Hartmann, A., Duan, W., Hankin, S., Comas-Bru, L., Cuthbert, M. O., Treble, P. C., Banner, J., Genty, D., Baldini, L. M., Bartolomé, M., Moreno, A., Pérez-Mejías, C., and Werner, M.: Global analysis reveals climatic controls on the oxygen isotope composition of cave drip water, *Nature Communications*, 10, 2984, <https://doi.org/10.1038/s41467-019-11027-w>, 2019.
- Baker, J. L., Lachniet, M. S., Chervyatsova, O., Asmerom, Y., and Polyak, V. J.: Holocene warming in western continental Eurasia driven by glacial retreat and greenhouse forcing, *Nature Geoscience*, 10, 430–435, <https://doi.org/10.1038/ngeo2953>, 2017.
- 540 Blaauw, M. and Christeny, J. A.: Flexible paleoclimate age-depth models using an autoregressive gamma process, *Bayesian Analysis*, 6, 457–474, <https://doi.org/10.1214/11-BA618>, 2011.
- Braconnot, P., Harrison, S. P., Kageyama, M., Bartlein, P. J., Masson-Delmotte, V., Abe-Ouchi, A., Otto-Bliesner, B., and Zhao, Y.: Evaluation of climate models using palaeoclimatic data, *Nature Climate Change*, 2, 417–424, <https://doi.org/10.1038/nclimate1456>, 2012.
- Bradley, R. S.: *Paleoclimate: reconstructing climates of the Quaternary*, Elsevier, 1999.
- 545 Breitenbach, S. F., Rehfeld, K., Goswami, B., Baldini, J. U., Ridley, H. E., Kennett, D. J., Prufer, K. M., Aquino, V. V., Asmerom, Y., Polyak, V. J., Cheng, H., Kurths, J., and Marwan, N.: Constructing proxy records from age models (COPRA), *Climate of the Past*, 8, 1765–1779, <https://doi.org/10.5194/cp-8-1765-2012>, 2012.
- Bühler, J. C. and Rehfeld, K.: iHadCM3LastMill Github Repository, <https://github.com/paleovar/iHadCM3LastMill>, 2020, (accessed September 10, 2020).
- 550 Caley, T., Roche, D. M., and Renssen, H.: Orbital Asian summer monsoon dynamics revealed using an isotope-enabled global climate model, *Nature Communications*, 5, 6–11, <https://doi.org/10.1038/ncomms6371>, 2014.
- Chatfield, C.: *The Analysis of Time Series - An Introduction*, Chapman & Hall/CRC, sixth edit edn., 2003.
- Comas-Bru, L., Harrison, S. P., Werner, M., Rehfeld, K., Scroxtton, N., and Veiga-Pires, C.: Evaluating model outputs using integrated global speleothem records of climate change since the last glacial, *Climate of the Past Discussions*, pp. 1–53, <https://doi.org/10.5194/cp-2019-25>,
555 2019.
- Comas-Bru, L., Rehfeld, K., Roesch, C., Amirnezhad-Mozhdehi, S., Harrison, S. P., Atsawawanunt, K., Masood Ahmad, S., Ait Ibrahim, Y., Baker, A., Bosomworth, M., Breitenbach, S. F., Burstyn, Y., Columbu, A., and Deininger, M.: SISALv2: A comprehensive speleothem isotope database with multiple age-depth, *Earth System Science Data Discussions*, 225, 1–47, <https://doi.org/10.5194/essd-2020-39>, 2020.
- Comas-Bru, L., Atsawawanunt, K., Harrison, S., and working group Members, S.: SISAL database version 2.0, <https://doi.org/10.17864/1947.256>, <https://researchdata.reading.ac.uk/256/>, 2020, (accessed September 10, 2020).
- 560 Coplen, T. B., Kendall, C., and Hopple, J.: Comparison of stable isotope reference samples, *Nature*, 302, 28–30, 1983.
- Cox, P. M.: Description of the "TRIFFID" dynamic global vegetation model, Hadley Centre. Technical Note 24, p. 17, 2001.
- Crowley, T. J. and Unterman, M. B.: Technical details concerning development of a 1200 yr proxy index for global volcanism, *Earth System Science Data*, 5, 187–197, <https://doi.org/10.5194/essd-5-187-2013>, 2013.
- 565 Dalaiden, Q., Goosse, H., Klein, F., Lenaerts, J. T., Holloway, M., Sime, L., and Thomas, E. R.: How useful is snow accumulation in reconstructing surface air temperature in Antarctica? A study combining ice core records and climate models, *Cryosphere*, 14, 1187–1207, <https://doi.org/10.5194/tc-14-1187-2020>, 2020.
- Dansgaard, W.: Stable isotopes in precipitation, *Tellus*, 16, 436–468, <https://doi.org/10.3402/tellusa.v16i4.8993>, 1964.



- Dee, S., Emile-Geay, J., Evans, M. N., Allam, A., Steig, E. J., and Thompson, D. M.: PRYSM: An open-source framework for PRoxY
570 System Modeling, with applications to oxygen-isotope systems, *Journal of Advances in Modeling Earth Systems*, 7, 1220–1247,
<https://doi.org/10.1002/2015MS000447>, 2015.
- Dolman, A., Kunz, T., Groeneveld, J., and Laepple, T.: Estimating the timescale-dependent uncertainty of paleoclimate records—a spectral
approach. Part II: Application and interpretation, *Climate of The Past Discussions*, pp. 1–38, <https://doi.org/10.5194/cp-2019-153>, 2020.
- Dreybrodt, W.: Deposition of calcite from thin films of natural calcareous solutions and the growth of speleothems, *Chemical Geology*,
575 [https://doi.org/10.1016/0009-2541\(80\)90007-8](https://doi.org/10.1016/0009-2541(80)90007-8), 1980.
- Efron, B. and Tibshirani, R.: Bootstrap methods for standard errors, confidence intervals, and other measures of statistical accuracy, *Statistical
Science*, 1, 54–75, <https://doi.org/10.1214/ss/1177013815>, 1986.
- Evans, M. N., Tolwinski-ward, S. E., Thompson, D. M., and Anchukaitis, K. J.: Applications of proxy system modeling in high resolution
paleoclimatology, *Quaternary Science Reviews*, 76, 16–28, <https://doi.org/10.1016/j.quascirev.2013.05.024>, 2013.
- 580 Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model
Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geoscientific Model Development*, 9, 1937–1958,
<https://doi.org/10.5194/gmd-9-1937-2016>, 2016.
- Fairchild, I. J. and Baker, A.: *Speleothem science: from process to past environments*, John Wiles & Sons, 2012.
- Fairchild, I. J. and Treble, P. C.: Trace elements in speleothems as recorders of environmental change, *Quaternary Science Reviews*, 28,
585 449–468, <https://doi.org/10.1016/j.quascirev.2008.11.007>, 2009.
- Fairchild, I. J., Smith, C. L., Baker, A., Fuller, L., Spötl, C., Matthey, D., and McDermott, F.: Modification and preservation of environmental
signals in speleothems, *Earth-Science Reviews*, 75, 105–153, <https://doi.org/10.1016/j.earscirev.2005.08.003>, 2006.
- Fisher, D. A., Reeh, N., and Clausen, H.: Stratigraphic Noise in Time Series Derived from Ice Cores, *Annals of Glaciology*, 7, 76–83,
<https://doi.org/10.3189/s0260305500005942>, 1985.
- 590 Fohlmeister, J., Schröder-Ritzrau, a., Scholz, D., Spötl, C., Riechelmann, D. F. C., Mudelsee, M., Wackerbarth, a., Gerdes, a., Riechel-
mann, S., Immenhauser, a., Richter, D. K., and Mangini, a.: Bunker Cave stalagmites: an archive for central European Holocene climate
variability, *Climate of the Past*, 8, 1751–1764, <https://doi.org/10.5194/cp-8-1751-2012>, 2012.
- Fohlmeister, J., Plessen, B., Dudashvili, A. S., Tjallingii, R., Wolff, C., Gafurov, A., and Cheng, H.: Winter precipitation changes
during the Medieval Climate Anomaly and the Little Ice Age in arid Central Asia, *Quaternary Science Reviews*, 178, 24–36,
595 <https://doi.org/10.1016/j.quascirev.2017.10.026>, 2017.
- Franzke, C. L. E., Barbosa, S., Blender, R., Fredriksen, H., Laepple, T., Lambert, F., Nilsen, T., Rypdal, K., Rypdal, M., Scotto, M. G.,
Vannitsem, S., Watkins, N. W., Yang, L., and Yuan, N.: The Structure of Climate Variability Across Scales, *Reviews of Geophysics*, 58,
<https://doi.org/10.1029/2019rg000657>, 2020.
- Gelhar, L. W. and Wilson, J. L.: Ground-Water Quality Modeling a, *Groundwater*, 12, 399–408, 1974.
- 600 Goosse, H., Renssen, H., Timmermann, A., and Bradley, R. S.: Internal and forced climate variability during the last
millennium: A model-data comparison using ensemble simulations, *Quaternary Science Reviews*, 24, 1345–1360,
<https://doi.org/10.1016/j.quascirev.2004.12.009>, 2005.
- Gordon, C., Cooper, C., Senior, C. A., Banks, H. T., Gregory, J. M., Johns, T. C., Mitchell, J. F. B., and Wood, R. A.: The simulation of SST,
sea ice extents and ocean heat transports in a version of the Hadley Centre coupled model without flux adjustments, *Climate Dynamics*,
605 16, 147–168, <https://doi.org/https://doi.org/10.1007/s003820050010>, 2000.



- Grossman, E. L. and Ku, T. L.: Oxygen and carbon isotope fractionation in biogenic aragonite: Temperature effects, *Chemical Geology: Isotope Geoscience Section*, [https://doi.org/10.1016/0168-9622\(86\)90057-6](https://doi.org/10.1016/0168-9622(86)90057-6), 1986.
- Haslett, J. and Parnell, A.: A simple monotone process with application to radiocarbon-dated depth chronologies, *Journal of the Royal Statistical Society. Series C: Applied Statistics*, 57, 399–418, <https://doi.org/10.1111/j.1467-9876.2008.00623.x>, 2008.
- 610 Haude, W.: Zur praktischen Bestimmung aktuellen und potentiellen Evaporation und Evapotranspiration, *Mitteilungen des Deutschen Wetterdienstes*, Band 1, 1954.
- Hijmans, R. J.: raster: Geographic Data Analysis and Modeling, <https://CRAN.R-project.org/package=raster>, r package version 3.0-7, 2019.
- Holloway, M. D., Sime, L. C., Singarayer, J. S., Tindall, J. C., and Valdes, P. J.: Reconstructing paleosalinity from $\delta^{18}\text{O}$: Coupled model simulations of the Last Glacial Maximum, Last Interglacial and Late Holocene, *Quaternary Science Reviews*, 131, 350–364, 615 <https://doi.org/10.1016/j.quascirev.2015.07.007>, 2016.
- Holloway, M. D., Sime, L. C., Singarayer, J. S., Tindall, J. C., and Valdes, P. J.: Simulating the 128-ka Antarctic Climate Response to Northern Hemisphere Ice Sheet Melting Using the Isotope-Enabled HadCM3, *Geophysical Research Letters*, 45, 11,911–921,929, <https://doi.org/10.1029/2018GL079647>, 2018.
- Hänsel, M. C., Drupp, M. A., Johansson, D. J. A., Nesje, F., Azar, C., Freeman, M. C., Groom, B., and Sterner, T.: Climate economics support 620 for the UN climate targets, *Nature Climate Change*, pp. 1758–6798, <https://doi.org/10.1038/s41558-020-0833-x>, 2020.
- IAEA/WMO: Global Network of Isotopes in Precipitation. The GNIP Database, Accessible at: <http://www.iaea.org/water>, 2020.
- IPCC: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, <https://doi.org/10.1017/CBO9781107415324>, 2013.
- Jean-Baptiste, P., Genty, D., Fourré, E., and Régnier, E.: Tritium dating of dripwater from Villars Cave (SW-France), *Applied Geochemistry*, 625 107, 152–158, <https://doi.org/10.1016/j.apgeochem.2019.06.005>, 2019.
- Jungclauss, J. H., Lorenz, S. J., Timmreck, C., Reick, C. H., Brovkin, V., Six, K., Segschneider, J., Giorgetta, M. A., Crowley, T. J., Pongratz, J., Krivova, N. A., Vieira, L. E., Solanki, S. K., Klocke, D., Botzet, M., Esch, M., Gayler, V., Haak, H., Raddatz, T. J., Roeckner, E., Schnur, R., Widmann, H., Claussen, M., Stevens, B., and Marotzke, J.: Climate and carbon-cycle variability over the last millennium, *Climate of the Past*, 6, 723–737, <https://doi.org/10.5194/cp-6-723-2010>, 2010.
- 630 Kageyama, M., Braconnot, P., Harrison, S. P., Haywood, A. M., Jungclauss, J. H., Otto-Bliesner, B. L., Abe-Ouchi, A., Albani, S., Bartlein, P. J., Brierley, C., Crucifix, M., Dolan, A., Fernandez-Donado, L. L., Fischer, H., Hopcroft, P. O., Ivanovic, R. F., Lambert, F., Lunt, D. J., Mahowald, N. M., Richard Peltier, W., Phipps, S. J., Roche, D. M., Schmidt, G. A., Tarasov, L., Valdes, P. J., Zhang, Q., and Zhou, T.: The PMIP4 contribution to CMIP6 - Part 1: Overview and over-arching analysis plan, *Geoscientific Model Development*, 11, 1033–1057, <https://doi.org/10.5194/gmd-11-1033-2018>, 2018.
- 635 Katz, R. W. and Brown, B. G.: Extreme events in a changing climate: Variability is more important than averages, *Climatic Change*, 21, 289–302, <https://doi.org/10.1007/BF00139728>, 1992.
- Kaufmann, G.: Stalagmite growth and palaeo-climate: The numerical perspective, *Earth and Planetary Science Letters*, 214, 251–266, [https://doi.org/10.1016/S0012-821X\(03\)00369-8](https://doi.org/10.1016/S0012-821X(03)00369-8), 2003.
- Kendall, C. and Caldwell, E. A.: Chapter 2 - Fundamentals of Isotope Geochemistry, in: *Fundamentals of Isotope Geochemistry*, edited by 640 KENDALL, C. and McDONNELL, J. J. B. T. I. T. i. C. H., pp. 51–86, Elsevier, Amsterdam, <https://doi.org/https://doi.org/10.1016/B978-0-444-81546-0.50009-4>, 1998.
- Kluge, T., Riechelmann, D. F., Wieser, M., Spötl, C., Sültenfuß, J., Schröder-Ritzrau, A., Niggemann, S., and Aeschbach-Hertig, W.: Dating cave drip water by tritium, *Journal of Hydrology*, 394, 396–406, <https://doi.org/10.1016/j.jhydrol.2010.09.015>, 2010.



- 645 Kluge, T., Affek, H. P., Marx, T., Aeschbach-Hertig, W., Riechelmann, D. F. C., Scholz, D., Riechelmann, S., Immenhauser, A., Richter, D. K., Fohlmeister, J., Wackerbarth, A., Mangini, A., and Spötl, C.: Reconstruction of drip-water $\delta 18\text{O}$ based on calcite oxygen and clumped isotopes of speleothems from Bunker Cave (Germany), *Climate of the Past*, 9, 377–391, <https://doi.org/10.5194/cp-9-377-2013>, 2013.
- Kunz, T., Dolman, A. M., and Laepple, T.: Estimating the timescale-dependent uncertainty of paleoclimate records – a spectral approach. Part I: Theoretical concept, *Climate of the Past Discussions*, 2020, 1–38, <https://doi.org/10.5194/cp-2019-150>, 2020.
- 650 Lachniet, M. S.: Climatic and environmental controls on speleothem oxygen-isotope values, *Quaternary Science Reviews*, 28, 412–432, <https://doi.org/10.1016/j.quascirev.2008.10.021>, 2009.
- Laepple, T. and Huybers, P.: Ocean surface temperature variability: Large model–data differences at decadal and longer periods, *Proceedings of the National Academy of Sciences*, 111, 16 682–16 687, <https://doi.org/10.1073/pnas.1412077111>, 2014a.
- Laepple, T. and Huybers, P.: Global and regional variability in marine surface temperatures, *Geophysical Research Letters*, 41, 2528–2534, 655 <https://doi.org/10.1002/2014GL059345>, 2014b.
- Lauritzen, S.-E. and Lundberg, J.: Calibration of the speleothem delta function: an absolute temperature record for the Holocene in northern Norway, *The Holocene*, 9, 659–669, 1999.
- Lechleitner, F. A., Amirnezhad-Mozhdehi, S., Columbu, A., Comas-Bru, L., Labuhn, I., Pérez-Mejías, C., and Rehfeld, K.: The Potential of Speleothems from Western Europe as Recorders of Regional Climate: A Critical Assessment of the SISAL Database, *Quaternary*, 1, 30, 660 <https://doi.org/10.3390/quat1030030>, 2018.
- Lohmann, G., Wackerbarth, A., Langebroek, P. M., Werner, M., Fohlmeister, J., Scholz, D., and Mangini, A.: Simulated European stalagmite record and its relation to a quasi-decadal climate mode, *Climate of the Past*, 9, 89–98, <https://doi.org/10.5194/cp-9-89-2013>, 2013.
- Marcott, S. A., Shakun, J. D., Clark, P. U., and Mix, A. C.: A reconstruction of regional and global temperature for the past 11,300 years., *Science (New York, N.Y.)*, 339, 1198–201, <https://doi.org/10.1126/science.1228026>, 2013.
- 665 McCrea, J. M.: On the isotopic chemistry of carbonates and a paleotemperature scale, *The Journal of Chemical Physics*, 18, 849–857, <https://doi.org/10.1063/1.1747785>, 1950.
- McDermott, F., Matthey, D. P., and Hawkesworth, C.: Centennial-scale holocene climate variability revealed by a high-resolution speleothem $\delta 18\text{O}$ record from SW Ireland, *Science*, 294, 1328–1331, <https://doi.org/10.1126/science.1063678>, 2001.
- Morice, C. P., Kennedy, J. J., Rayner, N. A., and Jones, P. D.: Quantifying uncertainties in global and regional temperature change using an ensemble of observational estimates: The HadCRUT4 data set, *Journal of Geophysical Research Atmospheres*, 117, 1–22, 670 <https://doi.org/10.1029/2011JD017187>, 2012.
- Neff, U., Burns, S. J., Mangini, A., Mudelsee, M., Fleitmann, D., and Matter, A.: Strong coherence between solar variability and the monsoon in Oman between 9 and 6 kyr ago, *Nature*, 411, 290–293, <https://doi.org/10.1038/35077048>, 2001.
- Neukom, R., Steiger, N., Gómez-Navarro, J. J., Wang, J., and Werner, J. P.: No evidence for globally coherent warm and cold periods over 675 the preindustrial Common Era, *Nature*, 571, 550–554, <https://doi.org/10.1038/s41586-019-1401-2>, 2019.
- Owen, R., Day, C. C., and Henderson, G. M.: CaveCalc: a new model for speleothem chemistry & isotopes, *Computers and Geosciences*, 2018.
- PAGES2k-Consortium: Consistent multidecadal variability in global temperature reconstructions and simulations over the Common Era, *Nature Geoscience*, 12, 643–649, <https://doi.org/10.1038/s41561-019-0400-0>, 2019.
- 680 Pardaens, A. K., Banks, H. T., Gregory, J. M., and Rowntree, P. R.: Freshwater transports in HadCM3, *Climate Dynamics*, 21, 177–195, <https://doi.org/10.1007/s00382-003-0324-6>, 2003.



- Parker, S. E., Harrison, S. P., Comas-Bru, L., Kaushal, N., LeGrande, A. N., and Werner, M.: A data-model approach to interpreting speleothem oxygen isotope records from monsoon regions on orbital timescales, *Climate of the Past Discussions*, 2020, 1–30, <https://doi.org/10.5194/cp-2020-78>, 2020.
- 685 Partin, J. W., Cobb, K. M., Adkins, J. F., Clark, B., and Fernandez, D. P.: Millennial-scale trends in west Pacific warm pool hydrology since the Last Glacial Maximum, *Nature*, 449, 452–455, <https://doi.org/10.1038/nature06164>, 2007.
- Pierce, D.: ncd4: Interface to Unidata netCDF (Version 4 or Earlier) Format Data Files, <https://CRAN.R-project.org/package=ncdf4>, r package version 1.17, 2019.
- Pope, V. D., Gallani, M. L., Rowntree, P. R., and Stratton, R. A.: The impact of new physical parametrizations in the Hadley Centre climate
690 model: HadAM3, *Climate Dynamics*, 16, 123–146, <https://doi.org/10.1007/s003820050009>, 2000.
- R Core Team: R: A Language and Environment for Statistical Computing, R Foundation for Statistical Computing, Vienna, Austria, <https://www.R-project.org/>, 2020.
- Ramsey, C. B.: Bayesian analysis of radiocarbon dates, *Radiocarbon*, 51, 337–360, <https://doi.org/10.1017/s0033822200033865>, 2009.
- Rehfeld, K. and Kurths, J.: Similarity estimators for irregular and age-uncertain time series, *Climate of the Past*, pp. 107–122,
695 <https://doi.org/10.5194/cp-10-107-2014>, 2014.
- Rehfeld, K., Marwan, N., Heitzig, J., and Kurths, J.: Comparison of correlation analysis techniques for irregularly sampled time series, *Nonlinear Processes in Geophysics*, pp. 389–404, <https://doi.org/10.5194/npg-18-389-2011>, 2011.
- Rehfeld, K., Marwan, N., Breitenbach, S. F., and Kurths, J.: Late Holocene Asian summer monsoon dynamics from small but complex networks of paleoclimate data, *Climate Dynamics*, 41, 3–19, <https://doi.org/10.1007/s00382-012-1448-3>, 2013.
- 700 Rehfeld, K., Trachsel, M., Telford, R. J., and Laepple, T.: Assessing performance and seasonal bias of pollen-based climate reconstructions in a perfect model world, *Climate of the Past*, 12, 2255–2270, <https://doi.org/10.5194/cp-12-2255-2016>, 2016.
- Rehfeld, K., Münch, T., Ho, S. L., and Laepple, T.: Global patterns of declining temperature variability from the Last Glacial Maximum to the Holocene, *Nature*, 554, 356–359, <https://doi.org/10.1038/nature25454>, 2018.
- Riechelmann, D. F. C., Schröder-Ritzrau, A., Scholz, D., Fohlmeister, J., Spötl, C., Richter, D. K., and Mangini, A.: Monitoring Bunker Cave
705 (NW Germany): A prerequisite to interpret geochemical proxy data of speleothems from this site, *Journal of Hydrology*, 409, 682–695, <https://doi.org/10.1016/j.jhydrol.2011.08.068>, 2011.
- Roesch, C. and Rehfeld, K.: Automatising construction and evaluation of age-depth models for hundreds of speleothems, 9th International Workshop on Climate Informatics, 2019.
- Schmidt, G. A., Jungclaus, J. H., Ammann, C. M., Bard, E., Braconnot, P., Crowley, T. J., Delaygue, G., Joos, F., Krivova, N. A., Muscheler,
710 R., Otto-Bliesner, B. L., Pongratz, J., Shindell, D. T., Solanki, S. K., Steinhilber, F., and Vieira, L. E. A.: Climate forcing reconstructions for use in PMIP simulations of the Last Millennium (v1.1), *Geoscientific Model Development*, 5, 185–191, <https://doi.org/10.5194/gmd-5-185-2012>, 2012.
- Schmidt, G. A., Annan, J. D., Bartlein, P. J., Cook, B. I., Guilyardi, E., Hargreaves, J. C., Harrison, S. P., Kageyama, M., Legrande, A. N., Konecky, B., Lovejoy, S., Mann, M. E., Masson-Delmotte, V., Risi, C., Thompson, D., Timmermann, A., and Yiou, P.: Using palaeo-
715 climate comparisons to constrain future projections in CMIP5, *Climate of the Past*, 10, 221–250, <https://doi.org/10.5194/cp-10-221-2014>, 2014.
- Scholz, D. and Hoffmann, D. L.: StalAge - An algorithm designed for construction of speleothem age models, *Quaternary Geochronology*, 6, 369–382, <https://doi.org/10.1016/j.quageo.2011.02.002>, <http://dx.doi.org/10.1016/j.quageo.2011.02.002>, 2011.



- Schurer, A. P., Tett, S. F. B., and Hegerl, G. C.: Small influence of solar variability on climate over the past millennium, *Nature Geoscience*, 7, 104–108, <https://doi.org/10.1038/ngeo2040>, 2014.
- Schwarz, H. P., Harmon, R. S., Thompson, P., and Ford, D. C.: Stable isotope studies of fluid inclusions in speleothems and their paleoclimatic significance, *Geochimica et Cosmochimica Acta*, 40, 657–665, [https://doi.org/10.1016/0016-7037\(76\)90111-3](https://doi.org/10.1016/0016-7037(76)90111-3), 1976.
- Sime, L. C., Tindall, J. C., Wolff, E. W., Connolley, W. M., and Valdes, P. J.: Antarctic isotopic thermometer during a CO₂ forced warming event, *Journal of Geophysical Research Atmospheres*, 113, 1–16, <https://doi.org/10.1029/2008JD010395>, 2008.
- 725 Sime, L. C., Wolff, E. W., Oliver, K. I. C., and Tindall, J. C.: Evidence for warmer interglacials in East Antarctic ice cores, *Nature*, 462, 342–345, <https://doi.org/10.1038/nature08564>, 2009.
- Sime, L. C., Kohfeld, K. E., Le Quéré, C., Wolff, E. W., de Boer, A. M., Graham, R. M., and Bopp, L.: Southern Hemisphere westerly wind changes during the Last Glacial Maximum: Model-data comparison, *Quaternary Science Reviews*, <https://doi.org/10.1016/j.quascirev.2012.12.008>, 2013.
- 730 Sjolte, J., Sturm, C., Adolphi, F., Vinther, B. M., Werner, M., Lohmann, G., and Muscheler, R.: Solar and volcanic forcing of North Atlantic climate inferred from a process-based reconstruction, *Climate of the Past*, 14, 1179–1194, <https://doi.org/10.5194/cp-14-1179-2018>, 2018.
- Steinhilber, F., Beer, J., and Fröhlich, C.: Total solar irradiance during the Holocene, *Geophysical Research Letters*, 36, 1–5, <https://doi.org/10.1029/2009GL040142>, 2009.
- Sturm, C., Zhang, Q., and Noone, D.: An introduction to stable water isotopes in climate models: Benefits of forward proxy modelling for paleoclimatology, *Climate of the Past*, 6, 115–129, <https://doi.org/10.5194/cp-6-115-2010>, 2010.
- 735 Tang, K. and Feng, X.: The effect of soil hydrology on the oxygen and hydrogen isotopic compositions of plants' source water, *Earth and Planetary Science Letters*, 185, 355–367, [https://doi.org/10.1016/S0012-821X\(00\)00385-X](https://doi.org/10.1016/S0012-821X(00)00385-X), 2001.
- Taylor, K. E., Stouffer, R. J., and Meehl, G. A.: An overview of CMIP5 and the experiment design, *Bulletin of the American Meteorological Society*, 93, 485–498, <https://doi.org/10.1175/BAMS-D-11-00094.1>, 2012.
- 740 Thornthwaite, C. W. and Mather, J. R.: Instructions and tables for computing potential evapotranspiration and the water balance, Tech. rep., Centerton, 1957.
- Tindall, J., Flecker, R., Valdes, P., Schmidt, D. N., Markwick, P., and Harris, J.: Modelling the oxygen isotope distribution of ancient seawater using a coupled ocean-atmosphere GCM: Implications for reconstructing early Eocene climate, *Earth and Planetary Science Letters*, <https://doi.org/10.1016/j.epsl.2009.12.049>, 2010.
- 745 Tindall, J. C., Valdes, P. J., and Sime, L. C.: Stable water isotopes in HadCM3: Isotopic signature of El Niño-Southern Oscillation and the tropical amount effect, *Journal of Geophysical Research Atmospheres*, 114, 1–12, <https://doi.org/10.1029/2008JD010825>, 2009.
- Tremaine, D. M., Froelich, P. N., and Wang, Y.: Speleothem calcite formed in situ: Modern calibration of $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ paleoclimate proxies in a continuously-monitored natural cave system, *Geochimica et Cosmochimica Acta*, <https://doi.org/10.1016/j.gca.2011.06.005>, 2011.
- 750 Tsonis, A. A., Swanson, K. L., and Roebber, P. J.: What do networks have to do with climate?, *Bulletin of the American Meteorological Society*, 87, 585–595, <https://doi.org/10.1175/BAMS-87-5-585>, 2006.
- Tupikina, L., Rehfeld, K., Molkenhain, N., Stolbova, V., Marwan, N., and Kurths, J.: Characterizing the evolution of climate networks, *Nonlinear Processes in Geophysics*, 21, 705–711, <https://doi.org/10.5194/npg-21-705-2014>, 2014.
- Urey, H. C.: Oxygen Isotopes in Nature and in the Laboratory, *Science*, 108, 489–496, 1948.
- 755 Valdes, P. J., Armstrong, E., Badger, M. P. S., Bradshaw, C. D., Bragg, F., Crucifix, M., Davies-Barnard, T., Day, J., Farnsworth, A., Gordon, C., Hopcroft, P. O., Kennedy, A. T., Lord, N. S., Lunt, D. J., Marzocchi, A., Parry, L. M., Pope, V., Roberts, W. H. G., Stone, E. J.,



- Tourte, G. J. L., and Williams, J. H. T.: The BRIDGE HadCM3 family of climate models: HadCM3@Bristol v1.0, *Geoscientific Model Development*, 10, 3715–3743, <https://doi.org/10.5194/gmd-10-3715-2017>, 2017.
- 760 Vasseur, D. A., DeLong, J. P., Gilbert, B., Greig, H. S., Harley, C. D., McCann, K. S., Savage, V., Tunney, T. D., and O'Connor, M. I.: Increased temperature variation poses a greater risk to species than climate warming, *Proceedings of the Royal Society B: Biological Sciences*, 281, <https://doi.org/10.1098/rspb.2013.2612>, 2014.
- Wackerbarth, A.: Supplementary Material, pp. 90–92, 2012.
- Wackerbarth, A., Scholz, D., Fohlmeister, J., and Mangini, A.: Modelling the delta 18 O value of cave drip water and speleothem calcite, *Earth and Planetary Science Letters*, 299, 387–397, <https://doi.org/10.1016/j.epsl.2010.09.019>, 2010.
- 765 Wang, Y., Lean, J. L., and Sheeley, Jr., N. R.: Modeling the Sun's Magnetic Field and Irradiance since 1713, *The Astrophysical Journal*, 625, 522–538, <https://doi.org/10.1086/429689>, 2005.
- Werner, M., Langebroek, P. M., Carlsen, T., Herold, M., and Lohmann, G.: Stable water isotopes in the ECHAM5 general circulation model: Toward high-resolution isotope modeling on a global scale, *Journal of Geophysical Research Atmospheres*, 116, 1–14, <https://doi.org/10.1029/2011JD015681>, 2011.
- 770 Werner, M., Haese, B., Xu, X., Zhang, X., Butzin, M., and Lohmann, G.: Glacial–interglacial changes in H218O, HDO and deuterium excess – results from the fully coupled ECHAM5/MPI-OM Earth system model, *Geoscientific Model Development*, 9, 647–670, <https://doi.org/10.5194/gmd-9-647-2016>, 2016.
- Wickham, H.: *ggplot2: Elegant Graphics for Data Analysis*, Springer-Verlag New York, <https://ggplot2.tidyverse.org>, 2016.
- Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L. D., François, R., Golemund, G., Hayes, A., Henry, L., Hester, J., Kuhn, M., 775 Pedersen, T. L., Miller, E., Bache, S. M., Müller, K., Ooms, J., Robinson, D., Seidel, D. P., Spinu, V., Takahashi, K., Vaughan, D., Wilke, C., Woo, K., and Yutani, H.: Welcome to the tidyverse, *Journal of Open Source Software*, 4, 1686, <https://doi.org/10.21105/joss.01686>, 2019.
- Williams, P. W. and Ford, D. C.: Global distribution of carbonate rocks, *Zeitschrift für Geomorphologie Supplementband*, 147, 1, 2006.
- Zhang, X., Sun, Z., Guan, H., Zhang, X., Wu, H., and Huang, Y.: GCM simulations of stable isotopes in the water cycle in comparison with 780 GNIP observations over east asia, *Acta Meteorologica Sinica*, 26, 420–437, <https://doi.org/10.1007/s13351-012-0403-x>, 2012.