

EDGE-ORDERED RAMSEY NUMBERS

M. BALKO AND M. VIZER

ABSTRACT. We introduce and study a variant of Ramsey numbers for *edge-ordered graphs*, that is, graphs with linearly ordered sets of edges. The *edge-ordered Ramsey number* $\overline{R}_e(\mathfrak{G})$ of an edge-ordered graph \mathfrak{G} is the minimum positive integer N such that there exists an edge-ordered complete graph \mathfrak{K}_N on N vertices such that every 2-coloring of the edges of \mathfrak{K}_N contains a monochromatic copy of \mathfrak{G} as an edge-ordered subgraph of \mathfrak{K}_N .

We prove that the edge-ordered Ramsey number $\overline{R}_e(\mathfrak{G})$ is finite for every edge-ordered graph \mathfrak{G} and we obtain better estimates for special classes of edge-ordered graphs. In particular, we prove $\overline{R}_e(\mathfrak{G}) \leq 2^{O(n^3 \log n)}$ for every bipartite edge-ordered graph \mathfrak{G} on n vertices. We also introduce a natural class of edge-orderings, called *lexicographic edge-orderings*, for which we can prove much better upper bounds on the corresponding edge-ordered Ramsey numbers.

1. INTRODUCTION

An *edge-ordered graph* $\mathfrak{G} = (G, \prec)$ consists of a graph $G = (V, E)$ and a linear ordering \prec of the set of edges E . We sometimes use the term *edge-ordering of G* for the ordering \prec and also for \mathfrak{G} . An edge-ordered graph (G, \prec_1) is an *edge-ordered subgraph* of an edge-ordered graph (H, \prec_2) if G is a subgraph of H and \prec_1 is a suborder of \prec_2 . We say that (G, \prec_1) and (H, \prec_2) are *isomorphic* if there is a graph isomorphism between G and H that also preserves the edge-orderings \prec_1 and \prec_2 .

For a positive integer k , a *k-coloring* of the edges of a graph G is any function that assigns one of the k colors to each edge of G . The *edge-ordered Ramsey number* $\overline{R}_e(\mathfrak{G})$ of an edge-ordered graph \mathfrak{G} is the minimum positive integer N such that there exists an edge-ordering \mathfrak{K}_N of the complete graph K_N on N vertices such that every 2-coloring of the edges of \mathfrak{K}_N contains a monochromatic copy of \mathfrak{G} as an edge-ordered subgraph of \mathfrak{K}_N . More generally, for two edge-ordered graphs \mathfrak{G} and \mathfrak{H} , we use $\overline{R}_e(\mathfrak{G}, \mathfrak{H})$ to denote the minimum positive integer N such that there exists an edge-ordering \mathfrak{K}_N of K_N such that every 2-coloring of the edges of

Received June 6, 2019.

2010 *Mathematics Subject Classification*. Primary 05C55, 05D10.

The first author was supported by the grant no. 18-13685Y of the Czech Science Foundation (GAČR) and by the Center for Foundations of Modern Computer Science (Charles University project UNCE/SCI/004).

The second author was supported by the Hungarian National Research, Development and Innovation Office – NKFIH under the grant SNN 129364 and KH 130371.

\mathfrak{K}_N with colors red and blue contains a red copy of \mathfrak{G} or a blue copy of \mathfrak{H} as an edge-ordered subgraph of \mathfrak{K}_N .

To our knowledge, Ramsey numbers of edge-ordered graphs were not considered in the literature. On the other hand, Ramsey numbers of graphs with ordered vertex sets have been quite extensively studied recently; for example, see [2, 3, 6]. For questions concerning extremal problems about vertex-ordered graphs consult the recent surveys [14, 15]. A *vertex-ordered graph* $\mathcal{G} = (G, \prec)$ (or simply an *ordered graph*) is a graph G with a fixed linear ordering \prec of its vertices. We use the term *vertex-ordering* of G to denote the ordering \prec as well as the ordered graph \mathcal{G} . An ordered graph (G, \prec_1) is a *vertex-ordered subgraph* of an ordered graph (H, \prec_2) if G is a subgraph of H and \prec_1 is a suborder of \prec_2 . We say that (G, \prec_1) and (H, \prec_2) are *isomorphic* if there is a graph isomorphism between G and H that also preserves the vertex-orderings \prec_1 and \prec_2 . Unlike in the case of edge-ordered graphs, there is a unique vertex-ordering \mathcal{K}_N of K_N up to isomorphism. The *ordered Ramsey number* $\overline{R}(\mathcal{G})$ of an ordered graph \mathcal{G} is the minimum $N \in \mathbb{N}$ such that every 2-coloring of the edges of \mathcal{K}_N contains a monochromatic copy of \mathcal{G} as a vertex-ordered subgraph of \mathcal{K}_N .

For an n -vertex graph G , let $R(G)$ be the Ramsey number of G . It is easy to see that $R(G) \leq \overline{R}(\mathcal{G})$ and $R(G) \leq \overline{R}_e(\mathfrak{G})$ for each vertex-ordering \mathcal{G} of G and edge-ordering \mathfrak{G} of G . We also have $\overline{R}(G) \leq \overline{R}(\mathcal{K}_n) = R(K_n)$ and thus ordered Ramsey numbers are always finite. Proving that $\overline{R}_e(\mathfrak{G})$ is always finite seems to be more challenging; see Theorem 2.1.

The Turán numbers of edge-ordered graphs were recently introduced in [7]. The authors of [7] proved, for example, a variant of the Erdős–Stone–Simonovits Theorem for edge-ordered graphs, and also investigated the Turán numbers of small edge-ordered paths, star forests, and 4-cycles; see the last section of [15].

2. OUR RESULTS

We study the growth rate of edge-ordered Ramsey numbers with respect to the number of vertices for various classes of edge-ordered graphs. As our first result, we show that edge-ordered Ramsey numbers are always finite and thus well-defined.

Theorem 2.1. *For every edge-ordered graph \mathfrak{G} , the number $\overline{R}_e(\mathfrak{G})$ is finite.*

Theorem 2.1 also follows from a recent deep result of Hubička and Nešetřil [9, Theorem 4.33] about Ramsey numbers of general relational structures. In comparison, our proof of Theorem 2.1 is less general, but it is much simpler and produces better and more explicit bound on $\overline{R}_e(\mathfrak{G})$. It is a modification of the proof of Theorem 12.13 [12, Page 138], which is based on the Graham–Rothschild Theorem [8]. In fact, the proof of Theorem 2.1 yields a stronger induced-type statement where additionally the ordering of the vertex set is fixed. Theorem 2.1 can also be extended to k -colorings with $k > 2$.

Due to the use of the Graham–Rothschild Theorem, the bound on the edge-ordered Ramsey numbers obtained in the proof of Theorem 2.1 is still enormous. It follows from a result of Shelah [13, Theorem 2.2] that this bound on $\overline{R}_e(\mathfrak{G})$ is

primitive recursive, but it grows faster than, for example, a tower function of any fixed height. Thus we aim to prove more reasonable estimates on edge-ordered Ramsey numbers, at least for some classes of edge-ordered graphs.

As our second main result, we show that one can obtain a much better upper bound on edge-ordered Ramsey numbers of two edge-ordered graphs, provided that one of them is bipartite. For $d \in \mathbb{N}$, we say that a graph G is d -degenerate if every subgraph of G has a vertex of degree at most d .

Theorem 2.2. *Let \mathfrak{H} be a d -degenerate edge-ordered graph on n' vertices and let \mathfrak{G} be a bipartite edge-ordered graph with m edges and with both parts containing n vertices. If $d \leq n$ and $n' \leq t^{d+1}$ for $t = 3n^{10}m!$, then*

$$\overline{R}_e(\mathfrak{H}, \mathfrak{G}) \leq (n')^2 t^{d+1}.$$

In particular, if \mathfrak{G} is a bipartite edge-ordered graph on n vertices, then $\overline{R}_e(\mathfrak{G}) \leq 2^{O(n^3 \log n)}$. We believe that the bound can be improved. In fact, it is possible that $\overline{R}_e(\mathfrak{G})$ is at most exponential in the number of vertices of \mathfrak{G} for every edge-ordered graph \mathfrak{G} . We note that, for every graph G and its vertex-ordering \mathcal{G} , both the standard Ramsey number $R(G)$ and the ordered Ramsey number $\overline{R}(\mathcal{G})$ grow at most exponentially in the number of vertices of G .

In general, the difference between edge-ordered Ramsey numbers and ordered Ramsey numbers with the same underlying graph can be very large. Let M_n be a *matching* on n vertices, that is, a graph formed by a collection of $n/2$ disjoint edges. There are ordered matchings $\mathcal{M}_n = (M_n, <)$ with super-polynomial ordered Ramsey numbers $\overline{R}(\mathcal{M}_n)$ in n [2, 6]. In fact this is true for almost all ordered matchings on n vertices [6]. On the other hand, all edge-orderings of M_n are isomorphic as edge-ordered graphs and thus $\overline{R}_e(\mathfrak{M}_n) = R(M_n) \leq O(n)$ for every edge-ordering \mathfrak{M}_n of M_n .

We consider a special class of edge-orderings, which we call *lexicographic edge-orderings*, for which we can prove much better upper bounds on their edge-ordered Ramsey numbers and which seem to be quite natural.

An ordering \prec of edges of a graph $G = (V, E)$ is *lexicographic* if there is a one-to-one correspondence $f: V \rightarrow \{1, \dots, |V|\}$ such that any two edges $\{u, v\}$ and $\{w, t\}$ of G with $f(u) < f(v)$ and $f(w) < f(t)$ satisfy $\{u, v\} \prec \{w, t\}$ if either $f(u) < f(w)$ or if $(f(u) = f(w) \ \& \ f(v) < f(t))$. We say that such mapping f is *consistent* with \prec . Note that, for every vertex u , the edges $\{u, v\}$ with $f(u) < f(v)$ form an interval in \prec . Also observe that there is a unique (up to isomorphism) lexicographic edge-ordering $\mathfrak{K}_n^{\text{lex}}$ of K_n . Setting $\{u, v\} \prec' \{w, t\}$ if either $f(u) < f(w)$ or if $(f(u) = f(w) \ \& \ f(v) > f(t))$ we obtain the *max-lexicographic* edge-ordering \prec' of G .

For a linear ordering $<$ on some set X , we use $<^{-1}$ to denote the *inverse ordering* of $<$, that is, for all $x, y \in X$, we have $x <^{-1} y$ if and only if $y < x$.

The lexicographic and max-lexicographic edge-orderings are natural, as Nešetřil and Rödl [11] showed that these orderings are canonical in the following sense.

Theorem 2.3 ([11]). *For every $n \in \mathbb{N}$, there is a positive integer $T(n)$ such that every edge-ordered complete graph on $T(n)$ vertices contains a copy of K_n such that*

the edges of this copy induce one of the following four edge-orderings: lexicographic edge-ordering \prec , max-lexicographic edge-ordering \prec' , \prec^{-1} , or $(\prec')^{-1}$.

Theorem 2.3 is also an unpublished result of Leeb; see [10].

It is thus natural to consider the following variant of edge-ordered Ramsey numbers, which turns out to be more tractable than general edge-ordered Ramsey numbers. The *lexicographic edge-ordered Ramsey number* $\overline{R}_{\text{lex}}(\mathfrak{G})$ of a lexicographically edge-ordered graph \mathfrak{G} is the minimum N such that every 2-coloring of the edges of $\mathfrak{K}_N^{\text{lex}}$ contains a monochromatic copy of \mathfrak{G} as an edge-ordered subgraph of $\mathfrak{K}_N^{\text{lex}}$. Observe that $\overline{R}_e(\mathfrak{G}) \leq \overline{R}_{\text{lex}}(\mathfrak{G})$ for every lexicographically edge-ordered graph \mathfrak{G} .

For every lexicographically edge-ordered graph $\mathfrak{G} = (G, \prec)$, the lexicographic edge-ordered Ramsey number $\overline{R}_{\text{lex}}(\mathfrak{G})$ can be estimated from above with the ordered Ramsey number of some vertex-ordering of G . More specifically,

$$(1) \quad \overline{R}_{\text{lex}}(\mathfrak{G}) \leq \min_f \overline{R}(\mathcal{G}_f),$$

where the minimum is taken over all one-to-one correspondences $f: V \rightarrow \{1, \dots, |V|\}$ that are consistent with the lexicographic edge-ordering \mathfrak{G} and \mathcal{G}_f is the vertex-ordering of G determined by f . Since $\overline{R}(\mathcal{K}_n) = R(K_n)$, it follows from (1) and from the well-known bound $R(K_n) \leq 2^{2^n}$ that the numbers $\overline{R}_{\text{lex}}(\mathfrak{G})$ are always at most exponential in the number of vertices of G . In fact, we have $\overline{R}_{\text{lex}}(\mathfrak{K}_n^{\text{lex}}) = \overline{R}(\mathcal{K}_n) = R(K_n)$ for every n . The equality is achieved in (1), for example, for graphs with a unique vertex-ordering determined by the lexicographic edge-ordering. Such graphs include graphs where each edge is contained in a triangle. Additionally, combining (1) with a result of Conlon et al. [6, Theorem 3.6] gives the estimate

$$\overline{R}_{\text{lex}}(\mathfrak{G}) \leq 2^{O(d \log_2^2(2n/d))}$$

for every d -degenerate lexicographically edge-ordered graph \mathfrak{G} on n vertices. In particular, $\overline{R}_{\text{lex}}(\mathfrak{G})$ is at most quasi-polynomial in n if d is fixed.

We note that the bound (1) is not always tight. For example, $R(K_{1,n}) = \overline{R}_{\text{lex}}(\mathfrak{K}_{1,n})$ for every edge-ordering $\mathfrak{K}_{1,n}$ of $K_{1,n}$, as any two edge-ordered stars $K_{1,n}$ are isomorphic as edge-ordered graphs. However, the Ramsey number $R(K_{1,n})$ is known to be strictly smaller than $\overline{R}(\mathcal{K}_{1,n})$ for n even and for any vertex-ordering $\mathcal{K}_{1,n}$ of $K_{1,n}$; see [4] and [1, Observation 11 and Theorem 12].

Using the inequality (1) we obtain asymptotically tight estimate on the following lexicographic edge-ordered Ramsey numbers of paths. The *edge-monotone path* $\mathfrak{P}_n = (P_n, \prec)$ is the edge-ordered path on vertices v_1, \dots, v_n , where $\{v_1, v_2\} \prec \dots \prec \{v_{n-1}, v_n\}$.

Proposition 2.4. *For every integer $n > 2$, we have $\overline{R}_{\text{lex}}(\mathfrak{P}_n) \leq 2n - 3 + \sqrt{2n^2 - 8n + 11}$.*

The proof of Proposition 2.4 uses the fact that the one-to-one correspondence f consistent with the lexicographic edge-ordering of P_n is not determined uniquely. Indeed, we can choose the mapping f so that it determines the vertex-ordering \mathcal{P}_n

of P_n where edges are between consecutive pairs of vertices. Such vertex-ordering \mathcal{P}_n is called *monotone path*. However, it is known that $\overline{R}(\mathcal{P}_n) = (n-1)^2 + 1$ [5] and thus we cannot apply (1) to this ordering to obtain a linear bound on $\overline{R}_{\text{lex}}(\mathfrak{P}_n)$. Instead we choose a different mapping f that determines a vertex-ordering of P_n with linear ordered Ramsey number.

Finally, we show an upper bound on edge-ordered Ramsey numbers of two graphs, where one of them is bipartite and suitably lexicographically edge-ordered. This result uses a stronger assumption about \mathfrak{G} than Theorem 2.2, but gives much better estimate. For $m, n \in \mathbb{N}$, let $\mathfrak{K}_{m,n}^{\text{lex}}$ be the lexicographic edge-ordering of $K_{m,n}$ that induces a vertex-ordering, in which both parts of $K_{m,n}$ form an interval.

Theorem 2.5. *Let \mathfrak{H} be a d -degenerate edge-ordered graph on n' vertices and let \mathfrak{G} be an edge-ordered subgraph of $\mathfrak{K}_{n',n'}^{\text{lex}}$. Then*

$$\overline{R}_e(\mathfrak{H}, \mathfrak{G}) \leq (n')^2 n^{d+1}.$$

3. OPEN PROBLEMS

Many questions about edge-ordered Ramsey numbers remain open, for example proving a better upper bound on edge-ordered Ramsey numbers than the one obtained in the proof of Theorem 2.1. For general upper bounds, it suffices to focus on edge-ordered complete graphs. It is possible that edge-ordered Ramsey numbers of edge-ordered complete graphs do not grow significantly faster than the standard Ramsey numbers.

Problem 3.1. Is there a constant C such that, for every $n \in \mathbb{N}$ and every edge-ordered complete graph \mathfrak{K}_n on n vertices, we have $\overline{R}_e(\mathfrak{K}_n) \leq 2^{Cn}$?

It might also be interesting to consider sparser graphs and try to prove better upper bounds on their edge-ordered Ramsey numbers.

Another interesting open problem is to determine the growth rate of the function $T(n)$ from Theorem 2.3. The current upper bound on $T(n)$ is quite large as the proof of Nešetřil and Rödl [11] uses Ramsey's theorem for quadruples and $6! = 720$ colors.

REFERENCES

1. Balko M., Cibulka J., Král K. and Kynčl J., *Ramsey numbers of ordered graphs*, [arxiv:1310.7208](https://arxiv.org/abs/1310.7208).
2. Balko M., Cibulka J., Král K. and Kynčl J., *Ramsey numbers of ordered graphs*, *Electron. Notes Discrete Math.* **49** (2015), 419–424.
3. Balko M., Jelínek V. and Valtr P., *On ordered Ramsey numbers of bounded-degree graphs*, *J. Combin. Theory Ser. B* **134** (2019), 179–202.
4. Burr S. A. and Roberts J. A., *On Ramsey numbers for stars*, *Util. Math.* **4** (1973), 217–220.
5. Choudum S. A. and Ponnusamy B., *Ordered Ramsey numbers*, *Discrete Math.* **247** (2002), 79–92.
6. Conlon D., Fox J., Lee C. and Sudakov B., *Ordered Ramsey numbers*, *J. Combin. Theory Ser. B* **122** (2017), 353–383.
7. Gerbner D., Methuku A., Nagy, T. D. Pálvölgyi, D. Tardos, G. and Vizer M., *Edge ordered Turán problems*, manuscript, 2019.

8. Graham R. L. and Rothschild B. L., *Ramsey's theorem for n -parameter sets*, Trans. Amer. Math. Soc. **159** (1971), 257–292.
9. Hubička J. and Nešetřil J., *All those Ramsey classes (Ramsey classes with closures and forbidden homomorphisms)*, [arxiv:1606.07979](https://arxiv.org/abs/1606.07979).
10. Nešetřil J., Prömel H. J., Rödl V. and Voigt B., *Canonizing ordering theorems for Hales-Jewett structures*, J. Combin. Theory Ser. A **40** (1985), 394–408.
11. Nešetřil J. and Rödl V., *Statistics of orderings*, Abh. Math. Semin. Univ. Hambg. **87** (2017), 421–433.
12. Prömel H. J., *Ramsey Theory for Discrete Structures*, Springer, Cham, 2013.
13. Shelah S., *Primitive recursive bounds for van der Waerden numbers*, J. Amer. Math. Soc. **1** (1988), 683–697.
14. Tardos G., *Extremal theory of ordered graphs*, in: Proceedings of the International Congress of Mathematics, vol. 3, 2018.
15. Tardos G., *The extremal theory of vertex ordered and edge ordered graphs*, Surveys in Combinatorics (2019), to appear.

M. Balko, Department of Applied Mathematics, Faculty of Mathematics and Physics, Charles University, Czech Republic,
e-mail: balko@kam.mff.cuni.cz

M. Vizer, Alfréd Rényi Institute of Mathematics, Hungarian Academy of Sciences, Budapest, Hungary,
e-mail: vizermate@gmail.com