

ENDOVASCULAR TREATMENT OF ISCHEMIC STROKE

Treat the right patient, at the right time, in the right place



Esmee Venema

ENDOVASCULAR TREATMENT OF ISCHEMIC STROKE

Treat the right patient, at the right time, in the right place

Esmee Venema

ISBN: 978-94-6380-858-3

Cover design & lay-out: Wendy Schoneveld || www.wenziD.nl

Printed by: ProefschriftMaken || www.proefschriftmaken.nl

The research described in this thesis was performed at the Department of Public Health and the Department of Neurology, Erasmus MC University Medical Center, Rotterdam, the Netherlands. The research was partly funded by the Erasmus MC program for Cost-Effectiveness Research.

Financial support by the Erasmus University Rotterdam, the Department of Public Health, and the Dutch Heart Foundation for the publication of this thesis is gratefully acknowledged.

© Esmee Venema, 2020

All rights reserved. No part of this thesis may be reproduced, stored or transmitted in any form or by any means without prior permission of the author, or the copyright-owning journals for previously published chapters.

Endovascular Treatment of Ischemic Stroke

Treat the right patient, at the right time, in the right place

Endovasculaire behandeling van het herseninfarct

Behandel de juiste patiënt, op het juiste moment en de juiste plaats

Proefschrift

ter verkrijging van de graad van doctor aan de
Erasmus Universiteit Rotterdam
op gezag van de rector magnificus

Prof.dr. R.C.M.E. Engels

en volgens besluit van het College voor Promoties.

De openbare verdediging zal plaatsvinden op
woensdag 30 september 2020 om 13.30 uur

door

Esmee Venema

geboren te Rotterdam

Promotiecommissie

Promotoren: Prof.dr. E.W. Steyerberg
Prof.dr. D.W.J. Dippel

Overige leden: Prof.dr.ir. H. Boersma
Prof.dr. W.H. van Zwam
Prof.dr. F. Lecky

Copromotoren: Dr. H.F. Lingsma
Dr. B. Roozenbeek

Contents

Chapter 1	General introduction	9
-----------	----------------------	---

Part I Treat the right patient

Chapter 2	Towards personalized endovascular treatment of patients with ischemic stroke: a study protocol for development and validation of a clinical decision aid. <i>BMJ Open, 2017</i>	25
Chapter 3	Selection of patients for endovascular treatment of ischemic stroke: development and validation of a clinical decision tool in two randomized trials. <i>BMJ, 2017</i>	37
Chapter 4	Improving selection of patients for endovascular treatment of ischemic stroke: validation and updating of MR PREDICTS with data from 4,398 patients. <i>Submitted</i>	57
Chapter 5	Multivariable outcome prediction after endovascular treatment for ischemic stroke (MR PREDICTS@24H): a post-procedural tool to predict functional outcome at 3 months. <i>In preparation</i>	79

Part II Treat at the right time

Chapter 6	Workflow and factors associated with delay in the delivery of endovascular treatment for ischemic stroke in the MR CLEAN trial. <i>J Neurointerv Surg, 2018</i>	101
Chapter 7	Effect of inter-hospital transfer on endovascular treatment for ischemic stroke. <i>Stroke, 2019</i>	117

Chapter 7.1	Response by Venema et al to letter regarding article, “Effect of inter-hospital transfer on endovascular treatment for ischemic stroke”. <i>Stroke, 2019</i>	133
-------------	---	-----

Chapter 8	Effect of workflow improvements in endovascular stroke treatment: a systematic review and meta-analysis. <i>Stroke, 2019</i>	137
-----------	---	-----

Part III Treat in the right place

Chapter 9	Personalized prehospital triage in ischemic stroke: a decision-analytic model. <i>Stroke, 2019</i>	171
-----------	---	-----

Chapter 10	Prehospital triage strategies for the transportation of suspected stroke patients in the United States. <i>Accepted for publication</i>	191
------------	--	-----

Chapter 11	Prehospital triage of patients with suspected stroke symptoms (PRESTO): protocol of a prospective observational study. <i>BMJ Open, 2019</i>	209
------------	---	-----

Chapter 12	General discussion	221
------------	--------------------	-----

Appendices

Summary	247
Samenvatting	253
Acknowledgments	259
Dankwoord	265
List of publications	271
PhD portfolio	277
About the author	281



Chapter 1

General introduction

One in six people will ever in their life suffer a stroke and over 35,000 stroke patients are annually admitted to a hospital in the Netherlands.^{1,2} The effect of stroke is devastating: it is a significant cause of long-term disability and the second leading cause of death worldwide.^{3,4}

Stroke is caused by a sudden interruption in the blood supply to the brain due to a thrombus that occludes an artery (ischemic stroke) or by a ruptured artery that leads to a bleeding in the brain tissue (hemorrhagic stroke). The vast majority of strokes are ischemic (87%), with atherosclerosis and cardio-embolism as main underlying causes.³ In ischemic stroke, the impaired blood flow results in a lack of oxygen and glucose in the underlying brain tissue. Neurons will be damaged and, due to their high energy demand, die quickly. The loss of neurological function depends on the size and location of the brain tissue that is affected, and may include paralysis of one side of the body, sensory disturbances, impaired vision, and speech and language difficulties.

Acute treatment of ischemic stroke

During the acute phase of ischemic stroke, on average 1.9 million neurons are destroyed each minute that the artery is blocked.⁵ Since neurological function loss can be reversible if the blood flow is restored in time, treatment has to be started as soon as possible. Intravenous treatment with alteplase (IVT) to dissolve the blood clot is standard of care for ischemic stroke patients presenting within 4.5 hours after stroke onset.⁶⁻⁸ However, IVT is less effective in the subgroup of patients with a large vessel occlusion (LVO), a thrombus in one of the proximal intracranial arteries in the anterior circulation, which account for approximately 24% to 46% of all ischemic strokes.⁹⁻¹¹ These patients are often severely affected and have a poor prognosis despite treatment with IVT.^{12,13}

Endovascular treatment

A more effective treatment option for patients with LVO is endovascular treatment (EVT), which consists of mechanical clot removal (thrombectomy, Figure 1.1), and originally of delivery of a thrombolytic agent at the site of the occlusion. From 1998, multiple trials demonstrated that this treatment is effective in reopening the occluded vessel and restoring the blood flow, but these studies were not able to show an effect on functional outcome of patients.¹⁴⁻¹⁹ The breakthrough came in 2015, when The Multicenter Randomized Clinical Trial of Endovascular Treatment for Acute Ischemic Stroke in the Netherlands (MR CLEAN) was the first to prove the safety and clinical effectiveness of EVT in patients presenting within 6 hours after onset of stroke.²⁰ Four other randomized controlled trials (RCTs) were stopped early after publication of the MR CLEAN results and showed similar effectiveness.²¹⁻²⁴ EVT soon became standard of care in developed countries.²⁵




Figure 1.1 Illustration of mechanical thrombectomy, reprinted with permission of MayfieldClinic.com. All rights reserved. No reuse permitted.

Treatment benefit

Endovascular treatment aims to improve functional outcome of ischemic stroke patients. Therefore, the primary outcome in most EVT trials was the modified Rankin Scale (mRS), an ordinal scale that measures the degree of disability during daily life activities.²⁶ This scale ranges from 0 (no symptoms) to 5 (severe disability), with an extra category of 6 to account for death (Table 1.1).

Table 1.1 Modified Rankin Scale.

Grade	Description
0	No remaining symptoms
1	No significant disability despite symptoms; able to perform all usual activities
2	Slight disability; unable to perform all previous activities, but independent
3	Moderate disability; requiring some help, but able to walk without assistance
4	Moderately severe disability; unable to walk without assistance and unable to attend to own bodily needs without assistance
5	Severe disability; bedridden, requiring constant nursing care and attention
6	Dead

In a patient-level pooled meta-analysis of five RCTs, the chance of achieving a good functional outcome, defined as an mRS score of 0–2, increased with EVT from 26.5% to 46%. The analysis revealed an astonishingly low number needed to treat of 2.6 to reduce disability by at least one level on the mRS.²⁷ However, this is the average treatment effect for the overall group of patients included in the trials, while the effect will likely vary between individual patients. Some patients will benefit more from EVT than other patients due to heterogeneity of baseline risk and relative treatment effect.^{28–31} The baseline risk of good outcome without treatment, or in other words, the natural history of disease, can be affected by prognostic factors such as age or severity of symptoms. Changes in baseline risk will affect the absolute benefit of a certain treatment, ie, the difference between outcome with and without treatment (Figure 1.2A). The relative treatment effect can be modified by predictive factors, for example when the treatment has a larger effect if started earlier after onset of symptoms (Figure 1.2B). Relative effects appear to be more stable across populations with different baseline risks, and are therefore useful when comparing two treatments or when combining the results of different trials in a meta-analysis. However, the absolute treatment benefit is what matters for a patient and is therefore more relevant for clinical decision-making.³² As an extreme example, a relative risk of 5 might increase the probability of the outcome with only 0.04% if that outcome is very rare (baseline risk = 0.01%).

Subgroup analysis are often performed to compare the relative treatment effect between different subgroups of patients within a trial population.^{33–35} However, these analyses are mostly underpowered, assess only one variable at a time, without taking into account a patient's full baseline risk, and are prone to false-negative and false-positive results.^{36,37} When multiple patient characteristics are evaluated simultaneously, more clinically relevant heterogeneity in treatment effect between individual patients will be found.^{28,29,38}

Treatment delay and workflow

Early initiation of EVT is associated with better clinical outcomes as the treatment effect strongly declines over time. Every hour of delay between symptom onset and start of EVT results in a 3–5% decreased probability of achieving functional independence.^{39,40} It is estimated that every 20 minutes decrease in time to treatment may lead to an average benefit equivalent to 3 months of disability-free life.⁴¹ Reducing delay will also increase the number of stroke patients that can be treated within the recommended 6 hour time-window. Although recent trials have shown that EVT can also be beneficial 6–24 hours after onset of symptoms, this applies to a selected group of patients with sufficient viable brain tissue on additional imaging only.^{42,43} Efficient workflow processes that decrease the time from onset to treatment are therefore important to increase the number of patients eligible for EVT and improve the overall outcome of treated patients.




Figure 1.2 Example to illustrate the hypothetical effect of (A) decreasing baseline risk with older age, assuming the same relative treatment effect (relative risk (RR) = 3) for all age categories; and (B) smaller relative treatment effect with increasing time to treatment, assuming a constant baseline risk.

Previously, several quality improvement initiatives resulted in significantly shorter door-to-needle times and a higher percentage of patients treated with IVT.⁴⁴⁻⁴⁶ These studies showed that in-hospital workflow processes can be streamlined using pre-notification of hospitals by the emergency medical services, rapid stroke team activation, readily available imaging facilities, and frequent feedback to the stroke team on time performance measures. Efficient workflow processes for EVT will also require interdisciplinary teamwork and communication between the emergency department and the neuro-interventional team. In the prehospital setting, potential EVT candidates should be recognized by the emergency medical services and transported to a hospital without any further delay. Potential workflow improvements include direct transportation to an intervention center with facilities for EVT instead of transportation to the closest hospital, and the use of air transportation or mobile stroke units. However, it is unknown how these interventions affect the delivery of EVT.

Prehospital triage

In current clinical practice, most suspected stroke patients are transported to the nearest primary stroke center for rapid IVT and further evaluation. When eligible for EVT, patients have to be transferred to an intervention center. Due to the importance of early treatment, it has been suggested that patients with ischemic stroke due to LVO would benefit from direct transportation to a center capable of performing EVT.⁴⁷ LVO can only be assessed with computed tomography (CT) imaging in the hospital, but several prehospital stroke scales have been developed to estimate the likelihood of LVO in patients presenting with stroke symptoms in the ambulance, based on the neurological examination and severity of symptoms.^{48,49} Although potentially beneficial for LVO patients, bypassing the nearest primary stroke center might be harmful for the majority of stroke patients because of the time-depending effect of IVT.^{6,50} Prehospital triage of suspected stroke patients therefore requires a trade-off between the harm of delaying IVT versus the potential benefit of rapid EVT.

Methods used in this thesis

Prediction modeling

A clinical prediction model estimates the probability of an individual to have a certain disease (diagnosis) or develop a certain clinical outcome (prognosis) based on the combination of a number of characteristics.⁵¹ Such models enable researchers or clinicians to make predictions for individual patients based on the effect of multiple factors combined. It can be used to inform individuals about their expected outcome and to select the right patients for a certain treatment or study.⁵² In contrast to etiological studies, a prediction model is not used to suggest a causal relationship between predictors and outcome. Also, it does not provide relative risk estimates such as an odds ratio or risk ratio, but it provides the absolute probability of a certain disease or outcome for an individual patient.

The development of a prediction model consists of several important steps, including careful predictor selection and model specification.⁵²⁻⁵⁴ The validity of the model should be evaluated at least in the derivation cohort (internal validation), and preferably also with external validation to assess generalizability of the model in other populations or settings.⁵⁵⁻⁵⁷ Performance measures often used in validation studies are discrimination and calibration. Discrimination assesses whether models are able to distinguish between patient with low risk and high risk of the outcome, while calibration describes the agreement between observed and predicted values.⁵³

Decision analyses

Decision analyses are designed to compare strategies in situations with decisional uncertainty. It provides a framework to combine all available evidence and uncertainties, to balance the harms and benefits of each alternative, and to make informed decisions.⁵⁸⁻⁶¹ A decision tree is a visual representation of all these alternatives and the consequence of

each choice. The expected outcome per strategy is calculated by multiplying the outcome values with the probability that the outcome will occur. According to the basic principles of decision analyses, the strategy with the highest outcome value would be preferred. Uncertainty concerning estimated model parameters and assumptions can be explored using sensitivity analyses in which parameters are varied over a range of estimates to assess its effect on the decision.⁶²

Table 1.2 Overview of the data sources used in this thesis.

Study	Design	Location of participating centers	Time frame of patient inclusion	Number of patients used for the analyses in this thesis
MR CLEAN	Phase III, multicenter RCT with open-label treatment and blinded outcome evaluation	The Netherlands	December 2010 – March 2014	500 (Chapters 3 and 6)
MR CLEAN Registry	Nationwide, prospective, observational study	The Netherlands	March 2014 – November 2017	3156 (Chapter 4) 3260 (Chapter 5) 1526 (Chapter 7)
IMS III	Phase III, multicenter RCT with open-label treatment and blinded outcome evaluation	The United States, Canada, Australia, and several countries in Europe	August 2006 – April 2012	260 (Chapter 3)
HERMES	Individual patient data from seven RCTs	The United States, Canada, Australia, New Zealand, Korea, and several countries in Europe	2010 – 2015	1242 (Chapter 4) 781 (Chapter 5)

Data sources

The analyses in this thesis were performed using clinical data from multiple sources (Table 1.2).

- The **MR CLEAN trial** randomized patients with ischemic stroke due to a proximal LVO between EVT (within 6 hours after stroke onset) plus usual care, versus usual care alone.^{20,63}
- The **MR CLEAN Registry** included all consecutive patients treated in the Netherlands after the final MR CLEAN inclusion, to monitor the implementation, outcome and safety of EVT in routine clinical practice.⁶⁴
- The **IMS III trial** (Interventional Management of Stroke) randomized ischemic stroke patients to EVT after IVT versus IVT alone. Patient inclusion was not restricted to patients with a proven LVO on non-invasive vessel imaging and the trial was stopped early because of futility.^{17,65}

- The **HERMES collaboration** (Highly Effective Reperfusion Using Multiple Endovascular Devices) combined data from seven international randomized controlled trials (RCTs), including MR CLEAN.^{20-24,66,67} Patient enrollment was performed according to the specific inclusion and exclusion criteria of each trial.

Aim and outline of this thesis

The overall aim of this thesis is to increase the benefit of endovascular treatment for ischemic stroke by optimizing prediction of outcome and treatment effect, reducing treatment delay, and improving prehospital triage strategies.

This translates into the following research questions:

1. Which are the right patients to treat?
 - a. Can we reliably and accurately predict outcome and treatment benefit of EVT for individual patients?
2. How can we treat patients at the right time?
 - a. What are the main causes of prehospital and in-hospital delay of EVT?
 - b. How do workflow improvements effect treatment delay and outcome?
3. How to direct patients to the right place?
 - a. Which factors should influence the decision to transport individual patients directly to an intervention center?
 - b. What is the optimal prehospital triage strategy for suspected stroke patients?

The first part of this thesis covers the development and validation of prediction models for outcome and treatment benefit of EVT (“treat the right patient”). **Chapters 2 and 3** describe the development and first external validation of a clinical decision tool to predict outcome with and without EVT (MR PREDICTS). In **Chapter 4**, this model is externally validated and updated with data from the HERMES collaboration and the MR CLEAN Registry. **Chapter 5** contains the development and validation of MR PREDICTS@24H, a post-procedural tool to predict functional outcome at 3 months more accurate with clinical data available within 24 hours after EVT.

The second part of this thesis is focused on rapid initiation of EVT (“treat at the right time”). **Chapter 6** aims to identify treatment delay in the MR CLEAN trial and factors associated with such delay. In **Chapter 7**, the effect of inter-hospital transfer on time to treatment and functional outcome is assessed by comparing patients transferred from a primary stroke center with patients directly admitted to an intervention center in the MR CLEAN Registry. **Chapter 8** reports the results of a systematic review and meta-analysis on the effectiveness of workflow improvements on time to treatment and outcome.

The third part of this thesis evaluates prehospital triage strategies for suspected stroke patients to reduce treatment delay and further improve effectiveness of EVT (“treat in the right place”). **Chapter 9** describes a decision-analytic model to determine the optimal prehospital transportation strategy for individual patients and to assess the factors that should influence this decision. In **Chapter 10**, this model is applied to the United States to evaluate the effect of several policies on outcomes of the ischemic stroke population. **Chapter 11** contains the study protocol of PRESTO, a multicenter observational cohort study to prospectively validate prehospital stroke scales for the prediction of LVO in the prehospital setting.

The main results of this thesis are summarized and discussed in **Chapter 12**, providing recommendations for future research and current clinical practice.

References

1. Seshadri S, Beiser A, Kelly-Hayes M, et al. The lifetime risk of stroke: estimates from the Framingham Study. *Stroke* 2006; **37**(2): 345-50.
2. de Boer AR, van Dis I, Visseren FLJ, Vaartjes I, Bots ML. Ziekte en sterfte aan hart- en vaatziekten. In: de Boer AR, van Dis I, Visseren FLJ, Vaartjes I, Bots ML Hart- en vaatziekten in Nederland 2019, cijfers over incidentie, prevalentie, ziekte en sterfte. Den Haag: Hartstichting; 2019.
3. Benjamin EJ, Virani SS, Callaway CW, et al. Heart Disease and Stroke Statistics-2018 Update: A Report From the American Heart Association. *Circulation* 2018; **137**(12): e67-e492.
4. Feigin VL, Norrving B, Mensah GA. Global Burden of Stroke. *Circ Res* 2017; **120**(3): 439-48.
5. Saver JL. Time is brain--quantified. *Stroke* 2006; **37**(1): 263-6.
6. Emberson J, Lees KR, Lyden P, et al. Effect of treatment delay, age, and stroke severity on the effects of intravenous thrombolysis with alteplase for acute ischaemic stroke: a meta-analysis of individual patient data from randomised trials. *Lancet* 2014; **384**(9958): 1929-35.
7. Wardlaw JM, Murray V, Berge E, del Zoppo GJ. Thrombolysis for acute ischaemic stroke. *Cochrane Database Syst Rev* 2014; (7): CD000213.
8. Adams HP, Jr., Brott TG, Furlan AJ, et al. Guidelines for Thrombolytic Therapy for Acute Stroke: a Supplement to the Guidelines for the Management of Patients with Acute Ischemic Stroke. A statement for healthcare professionals from a Special Writing Group of the Stroke Council, American Heart Association. *Stroke* 1996; **27**(9): 1711-8.
9. Christou I, Burgin WS, Alexandrov AV, Grotta JC. Arterial status after intravenous TPA therapy for ischaemic stroke. A need for further interventions. *Int Angiol* 2001; **20**(3): 208-13.
10. Bhatia R, Hill MD, Shobha N, et al. Low rates of acute recanalization with intravenous recombinant tissue plasminogen activator in ischemic stroke: real-world experience and a call for action. *Stroke* 2010; **41**(10): 2254-8.
11. Rennert RC, Wali AR, Steinberg JA, et al. Epidemiology, Natural History, and Clinical Presentation of Large Vessel Ischemic Stroke. *Neurosurgery* 2019; **85**(suppl_1): S4-S8.
12. Lima FO, Furie KL, Silva GS, et al. Prognosis of untreated strokes due to anterior circulation proximal intracranial arterial occlusions detected by use of computed tomography angiography. *JAMA Neurol* 2014; **71**(2): 151-7.
13. Heldner MR, Zubler C, Mattle HP, et al. National Institutes of Health stroke scale score and vessel occlusion in 2152 patients with acute ischemic stroke. *Stroke* 2013; **44**(4): 1153-7.
14. del Zoppo GJ, Higashida RT, Furlan AJ, Pessin MS, Rowley HA, Gent M. PROACT: a phase II randomized trial of recombinant pro-urokinase by direct arterial delivery in acute middle cerebral artery stroke. PROACT Investigators. Prolyse in Acute Cerebral Thromboembolism. *Stroke* 1998; **29**(1): 4-11.
15. Furlan A, Higashida R, Wechsler L, et al. Intra-arterial prourokinase for acute ischemic stroke. The PROACT II study: a randomized controlled trial. Prolyse in Acute Cerebral Thromboembolism. *JAMA* 1999; **282**(21): 2003-11.
16. Ogawa A, Mori E, Minematsu K, et al. Randomized trial of intraarterial infusion of urokinase within 6 hours of middle cerebral artery stroke: the middle cerebral artery embolism local fibrinolytic intervention trial (MELT) Japan. *Stroke* 2007; **38**(10): 2633-9.
17. Broderick JP, Palesch YY, Demchuk AM, et al. Endovascular therapy after intravenous t-PA versus t-PA alone for stroke. *N Engl J Med* 2013; **368**(10): 893-903.
18. Ciccone A, Valvassori L, Nichelatti M, et al. Endovascular treatment for acute ischemic stroke. *N Engl J Med* 2013; **368**(10): 904-13.
19. Kidwell CS, Jahan R, Gornbein J, et al. A trial of imaging selection and endovascular treatment for ischemic stroke. *N Engl J Med* 2013; **368**(10): 914-23.
20. Berkhemer OA, Fransen PS, Beumer D, et al. A randomized trial of intraarterial treatment for acute ischemic stroke. *N Engl J Med* 2015; **372**(1): 11-20.
21. Campbell BC, Mitchell PJ, Kleinig TJ, et al. Endovascular therapy for ischemic stroke with

- perfusion-imaging selection. *N Engl J Med* 2015; **372**(11): 1009-18.
22. Goyal M, Demchuk AM, Menon BK, et al. Randomized assessment of rapid endovascular treatment of ischemic stroke. *N Engl J Med* 2015; **372**(11): 1019-30.
 23. Jovin TG, Chamorro A, Cobo E, et al. Thrombectomy within 8 hours after symptom onset in ischemic stroke. *N Engl J Med* 2015; **372**(24): 2296-306.
 24. Saver JL, Goyal M, Bonafe A, et al. Stent-retriever thrombectomy after intravenous t-PA vs. t-PA alone in stroke. *N Engl J Med* 2015; **372**(24): 2285-95.
 25. Powers WJ, Derdeyn CP, Biller J, et al. 2015 American Heart Association/American Stroke Association Focused Update of the 2013 Guidelines for the Early Management of Patients With Acute Ischemic Stroke Regarding Endovascular Treatment: A Guideline for Healthcare Professionals From the American Heart Association/American Stroke Association. *Stroke* 2015; **46**(10): 3020-35.
 26. van Swieten JC, Koudstaal PJ, Visser MC, Schouten HJ, van Gijn J. Interobserver agreement for the assessment of handicap in stroke patients. *Stroke* 1988; **19**(5): 604-7.
 27. Goyal M, Menon BK, van Zwam WH, et al. Endovascular thrombectomy after large-vessel ischaemic stroke: a meta-analysis of individual patient data from five randomised trials. *Lancet* 2016; **387**(10029): 1723-31.
 28. Kent DM, Steyerberg E, van Klaveren D. Personalized evidence based medicine: predictive approaches to heterogeneous treatment effects. *BMJ* 2018; **363**: k4245.
 29. Kent DM, Paulus JK, van Klaveren D, et al. The Predictive Approaches to Treatment effect Heterogeneity (PATH) Statement. *Ann Intern Med* 2019.
 30. Rothwell PM. Treating individuals: from randomised trials to personalised medicine. London: Elsevier; 2007.
 31. Kravitz RL, Duan N, Braslow J. Evidence-based medicine, heterogeneity of treatment effects, and the trouble with averages. *Milbank Q* 2004; **82**(4): 661-87.
 32. Sormani MP, Bruzzi P. Estimating a treatment effect: Choosing between relative and absolute measures. *Mult Scler* 2017; **23**(2): 197-200.
 33. Yusuf S, Wittes J, Probstfield J, Tyroler HA. Analysis and interpretation of treatment effects in subgroups of patients in randomized clinical trials. *JAMA* 1991; **266**(1): 93-8.
 34. Assmann SF, Pocock SJ, Enos LE, Kasten LE. Subgroup analysis and other (mis)uses of baseline data in clinical trials. *Lancet* 2000; **355**(9209): 1064-9.
 35. Rothwell PM. Treating individuals 2. Subgroup analysis in randomised controlled trials: importance, indications, and interpretation. *Lancet* 2005; **365**(9454): 176-86.
 36. Brookes ST, Whitley E, Peters TJ, Mulheran PA, Egger M, Davey Smith G. Subgroup analyses in randomised controlled trials: quantifying the risks of false-positives and false-negatives. *Health Technol Assess* 2001; **5**(33): 1-56.
 37. Brookes ST, Whitley E, Egger M, Smith GD, Mulheran PA, Peters TJ. Subgroup analyses in randomised trials: risks of subgroup-specific analyses; power and sample size for the interaction test. *J Clin Epidemiol* 2004; **57**(3): 229-36.
 38. Hayward RA, Kent DM, Vijan S, Hofer TP. Multivariable risk prediction can greatly enhance the statistical power of clinical trial subgroup analysis. *BMC Med Res Methodol* 2006; **6**: 18.
 39. Mulder M, Jansen IGH, Goldhoorn RB, et al. Time to Endovascular Treatment and Outcome in Acute Ischemic Stroke: MR CLEAN Registry Results. *Circulation* 2018; **138**(3): 232-40.
 40. Saver JL, Goyal M, van der Lugt A, et al. Time to Treatment With Endovascular Thrombectomy and Outcomes From Ischemic Stroke: A Meta-analysis. *JAMA* 2016; **316**(12): 1279-88.
 41. Meretoja A, Keshkaran M, Tatlisumak T, Donnan GA, Churilov L. Endovascular therapy for ischemic stroke: Save a minute-save a week. *Neurology* 2017; **88**(22): 2123-7.
 42. Albers GW, Marks MP, Kemp S, et al. Thrombectomy for Stroke at 6 to 16 Hours with Selection by Perfusion Imaging. *N Engl J Med* 2018; **378**(8): 708-18.
 43. Nogueira RG, Jadhav AP, Haussen DC, et al. Thrombectomy 6 to 24 Hours after Stroke with a Mismatch between Deficit and Infarct. *N Engl J Med* 2018; **378**(1): 11-21.
 44. Meretoja A, Strbian D, Mustanoja S, Tatlisumak T, Lindberg PJ, Kaste M. Reducing in-hospital

- delay to 20 minutes in stroke thrombolysis. *Neurology* 2012; **79**(4): 306-13.
45. Fonarow GC, Zhao X, Smith EE, et al. Door-to-needle times for tissue plasminogen activator administration and clinical outcomes in acute ischemic stroke before and after a quality improvement initiative. *JAMA* 2014; **311**(16): 1632-40.
46. Dirks M, Niessen LW, van Wijngaarden JD, et al. Promoting thrombolysis in acute ischemic stroke. *Stroke* 2011; **42**(5): 1325-30.
47. Chartrain AG, Shoimah H, Jauch EC, Mocco J. A review of acute ischemic stroke triage protocol evidence: a context for discussion. *J Neurointerv Surg* 2018; **10**(11): 1047-52.
48. Smith EE, Kent DM, Bulsara KR, et al. Accuracy of Prediction Instruments for Diagnosing Large Vessel Occlusion in Individuals With Suspected Stroke: A Systematic Review for the 2018 Guidelines for the Early Management of Patients With Acute Ischemic Stroke. *Stroke* 2018; **49**(3): e111-e22.
49. Krebs W, Sharkey-Toppen TP, Cheek F, et al. Prehospital Stroke Assessment for Large Vessel Occlusions: A Systematic Review. *Prehosp Emerg Care* 2018; **22**(2): 180-8.
50. Hacke W, Donnan G, Fieschi C, et al. Association of outcome with early stroke treatment: pooled analysis of ATLANTIS, ECASS, and NINDS rt-PA stroke trials. *Lancet* 2004; **363**(9411): 768-74.
51. Moons KG, Royston P, Vergouwe Y, Grobbee DE, Altman DG. Prognosis and prognostic research: what, why, and how? *BMJ* 2009; **338**: b375.
52. Steyerberg EW. *Clinical Prediction Models: A Practical Approach to Development, Validation and Updating* (2nd ed.). New York: Springer; 2019.
53. Steyerberg EW, Vergouwe Y. Towards better clinical prediction models: seven steps for development and an ABCD for validation. *Eur Heart J* 2014; **35**(29): 1925-31.
54. Royston P, Moons KG, Altman DG, Vergouwe Y. Prognosis and prognostic research: Developing a prognostic model. *BMJ* 2009; **338**: b604.
55. Justice AC, Covinsky KE, Berlin JA. Assessing the generalizability of prognostic information. *Ann Intern Med* 1999; **130**(6): 515-24.
56. Collins GS, de Groot JA, Dutton S, et al. External validation of multivariable prediction models: a systematic review of methodological conduct and reporting. *BMC Med Res Methodol* 2014; **14**: 40.
57. Altman DG, Vergouwe Y, Royston P, Moons KG. Prognosis and prognostic research: validating a prognostic model. *BMJ* 2009; **338**: b605.
58. Hunink MGM, Weinstein MC, Wittenberg E, et al. *Decision Making in Health and Medicine; Integrating Evidence and Values* (2nd ed.). Cambridge, UK: Cambridge University Press; 2001.
59. Sox H, Higgins MC, Owens DK. *Medical Decision Making* (2nd ed.). New Jersey: Wiley-Blackwell; 2013.
60. Kassirer JP. The principles of clinical decision making: an introduction to decision analysis. *Yale J Biol Med* 1976; **49**(2): 149-64.
61. Schwartz WB, Gorry GA, Kassirer JP, Essig A. Decision analysis and clinical judgment. *Am J Med* 1973; **55**(3): 459-72.
62. Bilcke J, Beutels P, Brisson M, Jit M. Accounting for methodological, structural, and parameter uncertainty in decision-analytic models: a practical guide. *Med Decis Making* 2011; **31**(4): 675-92.
63. Fransen PS, Beumer D, Berkhemer OA, et al. MR CLEAN, a multicenter randomized clinical trial of endovascular treatment for acute ischemic stroke in the Netherlands: study protocol for a randomized controlled trial. *Trials* 2014; **15**: 343.
64. Jansen IGH, Mulder M, Goldhoorn RB, investigators MCR. Endovascular treatment for acute ischaemic stroke in routine clinical practice: prospective, observational cohort study (MR CLEAN Registry). *BMJ* 2018; **360**: k949.
65. Khatri P, Hill MD, Palesch YY, et al. Methodology of the Interventional Management of Stroke III Trial. *Int J Stroke* 2008; **3**(2): 130-7.
66. Bracad S, Ducrocq X, Mas JL, et al. Mechanical thrombectomy after intravenous alteplase versus alteplase alone after stroke (THRACE): a randomised controlled trial. *Lancet Neurol* 2016; **15**(11): 1138-47.
67. Muir KW, Ford GA, Messow CM, et al. Endovascular therapy for acute ischaemic stroke: the Pragmatic Ischaemic Stroke Thrombectomy Evaluation (PISTE) randomised, controlled trial. *J Neurol Neurosurg Psychiatry* 2017; **88**(1): 38-44.



Part I

Treat the right patient



Chapter 2

Towards personalized endovascular treatment of patients with ischemic stroke: a study protocol for development and validation of a clinical decision aid

BMJ Open, 2017

Esmee Venema*, Maxim J.H.L. Mulder*, Bob Roozenbeek, Joseph P. Broderick, Sharon D. Yeatts, Pooja Khatri, Olvert A. Berkhemer, Yvo B.W.E.M. Roos, Charles B.L.M. Majoie, Robert J. van Oostenbrugge, Wim H. van Zwam, Aad van der Lugt, Ewout W. Steyerberg, Diederik W.J. Dippel, and Hester F. Lingsma

*equal contribution

Abstract

Introduction

Endovascular treatment (EVT) proved to be overall beneficial in patients with ischemic stroke due to a proximal occlusion in the anterior circulation. However, heterogeneity in treatment benefit may be relevant for personalized clinical decision making. Our aim is to improve selection of patients for EVT by predicting individual treatment benefit or harm.

Methods and analysis

We will use data collected in the Multicenter Randomized Clinical Trial of Endovascular Treatment for Acute Ischemic Stroke in the Netherlands (MR CLEAN) trial to analyze the effect of baseline characteristics on outcome and treatment effect. A multivariable proportional odds model with interaction terms will be developed to predict outcome for each individual patient, both with and without EVT. Model performance will be expressed as discrimination and calibration, after bootstrap resampling and shrinkage of regression coefficients to correct for optimism. External validation will be conducted on data of patients in the Interventional Management of Stroke III trial (IMS III). Primary outcome will be the modified Rankin Scale (mRS) at 90 days after stroke.

Ethics and dissemination

The proposed study will provide an internationally applicable clinical decision aid for EVT. Findings will be disseminated widely through peer-reviewed publications, conference presentations and in an online web-application tool. Formal ethical approval was not required as primary data were already collected.

Introduction

In 2015, five consecutive randomized controlled trials (RCTs) showed that endovascular treatment (EVT) within 6 hours after stroke onset, improves functional outcome of patients with a proximal occlusion in the anterior circulation.¹⁻⁶ This was a major breakthrough in the field, and EVT is now implemented in updated guidelines on ischemic stroke management.⁷

Ideally, EVT will be targeted at patients who are expected to have optimal benefit: personalized treatment. In this study protocol we present seven steps for development and validation of a clinical decision aid to predict which individual patients with ischemic stroke will benefit most from EVT.^{8,9}

Strengths and limitations of this study

- *Multiple characteristics will be evaluated simultaneously to show clinically relevant heterogeneity in treatment benefit between patients.*
- *Multivariable prediction modelling substantially increases statistical power compared to other approaches and is more robust, especially in small datasets.*
- *We will use a relatively small cohort for the development of a prediction model.*
- *Using a proportional odds model requires the assumption that the odds ratio are the same for each cut-off of the modified Rankin Scale.*

Methods and analysis

Step 1: Problem definition and data inspection

Problem definition

RCTs provide estimates of treatment effects for average patients. However, it is important to take potential heterogeneity of treatment effects into account. Clinically relevant differences in the absolute effect of a treatment can be caused by 1) differences in the relative treatment effect (predictive effects) and 2) differences in baseline risk on the outcome of interest (prognostic effects).^{10,11} For example, in the Multicenter Randomized Clinical Trial of Endovascular Treatment for Acute Ischemic Stroke in the Netherlands (MR CLEAN) trial, there is no predictive effect of age; the relative treatment effect is constant across age subgroups.¹ This is demonstrated by a non-significant test for interaction between age and treatment (Figure 2.1A). However, variation in baseline risk on favorable outcome according to age results in a larger absolute treatment benefit in younger patients (Figure 2.1B).




Figure 2.1. Relative risk (A) and absolute risk difference (B) for functional independence (mRS 0-2) in MR CLEAN sort by age. MR CLEAN, Multicenter Randomized Clinical Trial of Endovascular Treatment for Acute Ischemic Stroke in the Netherlands; mRS, modified Rankin Scale

Conventional subgroup analyses are focused mainly on predictive effects and assess the effect of only one variable at a time. If predictive and prognostic effects of multiple characteristics are evaluated simultaneously in multivariable prediction modelling, it is likely that larger heterogeneity in treatment benefit between individual patients will be found. Our aim is to improve selection of patients for EVT by predicting treatment benefit or harm for individual stroke patients.

Development data

We will use data of the MR CLEAN trial (n=500), which was a phase 3, multicenter clinical trial with randomized treatment group assignment, open-label treatment, and blinded end-point evaluation. EVT plus usual care (which could include intravenous administration of alteplase) was compared with usual care alone. EVT consisted of arterial catheterization with a microcatheter to the level of occlusion and delivery of a thrombolytic agent, mechanical thrombectomy, or both.¹

Severity of stroke was assessed at baseline with the National Institutes of Health Stroke Scale (NIHSS; range 0–42). Baseline Computed tomography (CT) was evaluated with the Alberta Stroke Program Early Computed Tomography Score (ASPECTS; range 0–10). Baseline imaging (CT angiography) was used to determine the location of occlusion and to grade the quality of collateral flow to the ischemic area with a 4-point scale. Detailed information about the MR CLEAN trial can be found in the study protocol and the publication of the main results.^{1,12}

Endpoints of interest

Primary outcome will be the modified Rankin Scale (mRS), a 7-point scale ranging from 0 (no symptoms) to 6 (death) at 90 days after stroke.¹³ We will provide estimates of treatment benefit as the absolute increase in probability on functional independence (defined as mRS 0–2) and survival (defined as mRS 0–5).

Step 2: Coding of variables

As variables, we will use patient characteristics that are expected to predict outcome, or that are expected to interact with treatment, based on expert opinion and the recent literature (Table 2.1). Non-linearity of continuous variables will be tested by comparing the 2 log likelihood of models with linear and restricted cubic spline (RCS) functions.¹⁴

Timing of treatment is an essential predictor of outcome. Since time to randomization was not a reliable indicator for time to treatment in the MR CLEAN trial and will not be applicable in clinical practice, we will use time from stroke onset to groin puncture. Since time to groin puncture is not observable in the control group, we will explore imputation approaches based on the correlation with time to randomization. All other baseline variable values are more than 98% complete in the MR CLEAN data, so we choose simple imputation by the mean for continuous variables and simple imputation by the mode for categorical variables.

Step 3 and 4: Model specification and estimation

We will test the effect of variables on functional outcome and treatment effect with proportional odds regression modelling. All variables from Table 2.1 will be tested for effect on outcome and interaction with treatment effect. Prognostic variables (main effects) and predictive variables (interaction effects) with a p-value of 0.15 in univariable and multivariable analyses will be included in our final model. A p-value of 0.15 was chosen to make the predictor selection less data driven and prevent overfitting.^{14,15} We will perform shrinkage of all regression coefficients with ridge regression to prevent overfitting of the model.¹⁴ Predicted probabilities for each of the mRS categories, with and without treatment, will be derived from the ordinal model. All statistical analyses will be performed within the computing environment R version 3.2.2 (The R Foundation).

Table 2.1. Patient characteristics that are expected to predict outcome (prognostic), or that are expected to interact with treatment (predictive).

	% of data complete in MR CLEAN	Prognostic	Predictive
Clinical			
Age ^{6,24}	100%	X	
Baseline NIHSS ^{25,26}	100%	X	
History of diabetes mellitus ²⁷	100%	X	
History of previous stroke ²⁸	100%	X	
History of atrial fibrillation ²⁹	100%	X	
Pre-stroke mRS score ²⁸	100%	X	
Systolic blood pressure ³⁰	100%	X	
IV treatment with alteplase ³¹⁻³³	100%	X	
Time from stroke onset to groin puncture ^{34,35}	100%*	X	X
Radiological			
ASPECTS ^{6,36}	99.2%	X	
Location of intracranial occlusion on non-invasive vessel imaging ^{37,38}	99.8%	X	
Collateral score on CTA ^{38,39}	98.4%	X	X

*Of patients undergoing endovascular treatment.

ASPECTS, Alberta Stroke Program Early CT score; CTA, computed tomography angiography; IV, intravenous; MR CLEAN, Multicenter Randomized Clinical Trial of Endovascular Treatment for Acute Ischemic Stroke in the Netherlands; mRS, modified Rankin Scale; NIHSS, National Institutes of Health Stroke Scale.

Step 5: Model performance

Model performance will be expressed in discrimination and calibration. Discrimination will be quantified with the c-statistic. The c-statistic is similar to the area under the curve (AUC) for binary outcomes and estimates the probability that out of two randomly chosen patients, the patient with the higher predicted probability of a good outcome will indeed have a better outcome. Calibration refers to the agreement between predicted and observed risks and will be assessed graphically with calibration plots, and expressed as calibration slope and intercept. The calibration slope describes the relative overall effect of the variables in the validation sample, and is ideally equal to 1.

The intercept indicates whether predictions are systematically too high or too low, and should ideally be zero.¹⁶ We will calculate a general c-statistic to express the performance of our ordinal model and additional calibration plots with specific c-statistics for the predictions of favorable functional outcome (mRS 0-2) and survival (mRS 0-5).

Step 6: Model validity

The c-statistic will be internally validated with a bootstrap procedure (500 samples with replacement) to estimate the degree of optimism in parameter estimates.⁸ After penalization of the regression coefficients we will externally validate the model on data of patients in the Interventional Management of Stroke III trial (IMS III) with an occlusion in the anterior circulation on non-invasive vessel imaging.¹⁷ Coefficients of the final model will be fitted on the combined development and validation datasets.

After validation, we will assess whether the model can be used to discriminate between patients with low and high expected benefit by making individual predictions of outcome for all patients included in the development and validation data.

Step 7: Model presentation

The final model will be online available to be used in clinical practice, both for mobile devices and as a web-application. It will provide predictions of all mRS categories for each individual patient, both with and without EVT.

Ethics and dissemination

Findings will be disseminated widely through peer-reviewed publications, conference presentations and in an online web-application tool. Formal ethical approval was not required for this study as primary data were already collected.

Discussion

Compared to the current subgroup analyzes on the effect of EVT, our modelling approach has multiple advantages. First, it accounts for the fact that patients have multiple characteristics that simultaneously affect the likelihood of treatment benefit.¹⁸ Thus, our model will show more clinically relevant heterogeneity in treatment benefit between patients. Second, a multivariable prediction model substantially increases statistical power to identify heterogeneity in treatment effects compared to other approaches.¹⁹ These include neural network and decision trees. We use regression modelling since it is considered more robust, especially in relatively small datasets.^{20,21}

There are some differences between patients included in MR CLEAN and IMS III that may influence the external validity of our model. IMS III had different inclusion criteria, used older devices and used older treatment paradigms than MR CLEAN. In order to overcome these limitations, we will use only those patients in IMS III with an occlusion in the intracranial anterior circulation on noninvasive vessel imaging. We will compare the baseline

characteristics of the derivation and validation cohort and describe relevant differences that might lead to an under- or overestimation of the model performance. Interestingly, a substantial treatment effect in the IMS III patients with proven intracranial large vessel occlusion has been reported.²²

Furthermore, even though the MR CLEAN trial has included most patients of the recent RCTs, the cohort remains relatively small for the development of a prediction model, especially for the selection of both main effect and interaction effects. We will reduce regression coefficients to prevent overfitting and we will perform external validation. In the future, we will further validate and update our model in the pooled individual patient data of the Highly Effective Reperfusion evaluated in Multiple Endovascular Stroke Trials (HERMES) collaboration, harboring data of all patients from recent randomized trials regarding EVT (over 1700 patients in total). Moreover, we aim to investigate the validity of our model predicting outcome after treatment in clinical practice. Our model will therefore be tested by applying it to recently treated patients in all Dutch neurovascular centers participating in the MR CLEAN Registry.

We will use a proportional odds model to analyze the full mRS score as outcome. Formally this model requires the assumption that the odds ratio are the same for each cut-off of the mRS. However, previous studies have shown that even if the proportionality assumption is violated, proportional odds analysis is still more efficient than dichotomization.²³ In addition, all recent RCTs on the effect of EVT used the full mRS and analyzed their results with proportional odds regression.

Conclusion

The proposed study will provide an internationally applicable clinical decision aid for the selection of patients for EVT. We consider this study an important next step towards personalized treatment of ischemic stroke patients.

References

- Berkhemer OA, Fransen PS, Beumer D, et al. A randomized trial of intraarterial treatment for acute ischemic stroke. *N Engl J Med* 2015; **372**(1): 11-20.
- Campbell BC, Mitchell PJ, Kleinig TJ, et al. Endovascular therapy for ischemic stroke with perfusion-imaging selection. *N Engl J Med* 2015; **372**(11): 1009-18.
- Campbell BC, Donnan GA, Lees KR, et al. Endovascular stent thrombectomy: the new standard of care for large vessel ischaemic stroke. *Lancet Neurol* 2015; **14**(8): 846-54.
- Jovin TG, Chamorro A, Cobo E, et al. Thrombectomy within 8 hours after symptom onset in ischemic stroke. *N Engl J Med* 2015; **372**(24): 2296-306.
- Saver JL, Goyal M, Bonafe A, et al. Stent-retriever thrombectomy after intravenous t-PA vs. t-PA alone in stroke. *N Engl J Med* 2015; **372**(24): 2285-95.
- Goyal M, Menon BK, van Zwam WH, et al. Endovascular thrombectomy after large-vessel ischaemic stroke: a meta-analysis of individual patient data from five randomised trials. *Lancet* 2016; **387**(10029): 1723-31.
- Powers WJ, Derdeyn CP, Biller J, et al. 2015 American Heart Association/American Stroke Association Focused Update of the 2013 Guidelines for the Early Management of Patients With Acute Ischemic Stroke Regarding Endovascular Treatment: A Guideline for Healthcare Professionals From the American Heart Association/American Stroke Association. *Stroke* 2015; **46**(10): 3020-35.
- Steyerberg EW. Clinical prediction models: a practical approach to development, validation, and updating. *New York: Springer*, 2009.
- Steyerberg EW, Vergouwe Y. Towards better clinical prediction models: seven steps for development and an ABCD for validation. *Eur Heart J* 2014; **35**(29): 1925-31.
- Rothwell PM. *Treating Individuals: from randomised trials to personalised medicine*. London: Elsevier, 2007.
- van Klaveren D, Vergouwe Y, Farooq V, Serruys PW, Steyerberg EW. Estimates of absolute treatment benefit for individual patients required careful modeling of statistical interactions. *J Clin Epidemiol* 2015; **68**(11): 1366-74.
- Fransen PS, Beumer D, Berkhemer OA, et al. MR CLEAN, a multicenter randomized clinical trial of endovascular treatment for acute ischemic stroke in the Netherlands: study protocol for a randomized controlled trial. *Trials* 2014; **15**: 343.
- van Swieten JC, Koudstaal PJ, Visser MC, Schouten HJ, van Gijn J. Interobserver agreement for the assessment of handicap in stroke patients. *Stroke* 1988; **19**(5): 604-7.
- Harrell FE. Regression modeling strategies: with applications to linear models, logistic regression, and survival analysis. *New York: Springer-Verlag* 2001.
- Steyerberg EW, Eijkemans MJ, Harrell FE, Jr., Habbema JD. Prognostic modelling with logistic regression analysis: a comparison of selection and estimation methods in small data sets. *Stat Med* 2000; **19**(8): 1059-79.
- Steyerberg EW, Vickers AJ, Cook NR, et al. Assessing the performance of prediction models: a framework for traditional and novel measures. *Epidemiology* 2010; **21**(1): 128-38.
- Broderick JP, Palesch YY, Demchuk AM, et al. Endovascular therapy after intravenous t-PA versus t-PA alone for stroke. *N Engl J Med* 2013; **368**(10): 893-903.
- Kent DM, Rothwell PM, Ioannidis JP, Altman DG, Hayward RA. Assessing and reporting heterogeneity in treatment effects in clinical trials: a proposal. *Trials* 2010; **11**: 85.
- Hayward RA, Kent DM, Vijan S, Hofer TP. Multivariable risk prediction can greatly enhance the statistical power of clinical trial subgroup analysis. *BMC Med Res Methodol* 2006; **6**: 18.
- van der Ploeg T, Austin PC, Steyerberg EW. Modern modelling techniques are data hungry: a simulation study for predicting dichotomous endpoints. *BMC Med Res Methodol* 2014; **14**: 137.
- van der Ploeg T, Nieboer D, Steyerberg EW. Modern modeling techniques had limited

- external validity in predicting mortality from traumatic brain injury. *J Clin Epidemiol* 2016.
22. Demchuk AM, Goyal M, Yeatts SD, et al. Recanalization and clinical outcome of occlusion sites at baseline CT angiography in the Interventional Management of Stroke III trial. *Radiology* 2014; **273**(1): 202-10.
 23. McHugh GS, Butcher I, Steyerberg EW, et al. A simulation study evaluating approaches to the analysis of ordinal outcome data in randomized controlled trials in traumatic brain injury: results from the IMPACT Project. *Clin Trials* 2010; **7**(1): 44-57.
 24. Wardlaw JM, Murray V, Berge E, et al. Recombinant tissue plasminogen activator for acute ischaemic stroke: an updated systematic review and meta-analysis. *Lancet* 2012; **379**(9834): 2364-72.
 25. Broderick JP, Berkhemer OA, Palesch YY, et al. Endovascular Therapy Is Effective and Safe for Patients With Severe Ischemic Stroke: Pooled Analysis of Interventional Management of Stroke III and Multicenter Randomized Clinical Trial of Endovascular Therapy for Acute Ischemic Stroke in the Netherlands Data. *Stroke* 2015; **46**(12): 3416-22.
 26. Frankel MR, Morgenstern LB, Kwiatkowski T, et al. Predicting prognosis after stroke - A placebo group analysis from the National Institute of Neurological Disorders and Stroke rt-PA Stroke Trial. *Neurology* 2000; **55**(7): 952-9.
 27. Luitse MJ, Biessels GJ, Rutten GE, Kappelle LJ. Diabetes, hyperglycaemia, and acute ischaemic stroke. *Lancet Neurol* 2012; **11**(3): 261-71.
 28. Karlinski M, Kobayashi A, Czlonkowska A, et al. Intravenous Thrombolysis for Stroke Recurring Within 3 Months From the Previous Event. *Stroke* 2015; **46**(11): 3184-9.
 29. Sanak D, Herzig R, Kral M, et al. Is atrial fibrillation associated with poor outcome after thrombolysis? *J Neurol* 2010; **257**(6): 999-1003.
 30. Jauch EC, Saver JL, Adams HP, Jr., et al. Guidelines for the early management of patients with acute ischemic stroke: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke* 2013; **44**(3): 870-947.
 31. Tissue plasminogen activator for acute ischemic stroke. The National Institute of Neurological Disorders and Stroke rt-PA Stroke Study Group. *N Engl J Med* 1995; **333**(24): 1581-7.
 32. Hacke W, Kaste M, Bluhmki E, et al. Thrombolysis with alteplase 3 to 4.5 hours after acute ischemic stroke. *N Engl J Med* 2008; **359**(13): 1317-29.
 33. Mulder MJ, Berkhemer OA, Franssen PS, et al. Treatment in patients who are not eligible for intravenous alteplase: MR CLEAN subgroup analysis. *Int J Stroke* 2016.
 34. Emberson J, Lees KR, Lyden P, et al. Effect of treatment delay, age, and stroke severity on the effects of intravenous thrombolysis with alteplase for acute ischaemic stroke: a meta-analysis of individual patient data from randomised trials. *Lancet* 2014; **384**(9958): 1929-35.
 35. Franssen PS, Berkhemer OA, Lingsma HF, et al. Time to Reperfusion and Treatment Effect for Acute Ischemic Stroke: A Randomized Clinical Trial. *JAMA Neurol* 2015: 1-7.
 36. Barber PA, Demchuk AM, Zhang J, Buchan AM. Validity and reliability of a quantitative computed tomography score in predicting outcome of hyperacute stroke before thrombolytic therapy. ASPECTS Study Group. Alberta Stroke Programme Early CT Score. *Lancet* 2000; **355**(9216): 1670-4.
 37. Puetz V, Dzialowski I, Hill MD, et al. Intracranial thrombus extent predicts clinical outcome, final infarct size and hemorrhagic transformation in ischemic stroke: the clot burden score. *International Journal of Stroke* 2008; **3**(4): 230-6.
 38. Tan IYL, Demchuk AM, Hopyan J, et al. CT Angiography Clot Burden Score and Collateral Score: Correlation with Clinical and Radiologic Outcomes in Acute Middle Cerebral Artery Infarct. *Am J Neuroradiol* 2009; **30**(3): 525-31.
 39. Berkhemer OA, Jansen IG, Beumer D, et al. Collateral Status on Baseline Computed Tomographic Angiography and Intra-Arterial Treatment Effect in Patients With Proximal Anterior Circulation Stroke. *Stroke* 2016; **47**(3): 768-76.





Chapter 3

Selection of patients for endovascular treatment of ischemic stroke: development and validation of a clinical decision tool in two randomized trials

BMJ, 2017

Esmee Venema*, Maxim J.H.L. Mulder*, Bob Roozenbeek, Joseph P. Broderick, Sharon D. Yeatts, Pooja Khatri, Olvert A. Berkhemer, Bart J. Emmer, Yvo B.W.E.M. Roos, Charles B.L.M. Majoie, Robert J. van Oostenbrugge, Wim H. van Zwam, Aad van der Lugt, Ewout W. Steyerberg, Diederik W.J. Dippel, and Hester F. Lingsma

*equal contribution

Abstract

Objective

To improve the selection of patients with acute ischemic stroke for endovascular treatment (EVT) using a clinical decision tool to predict individual treatment benefit.

Design

Multivariable regression modeling with data from two randomized controlled clinical trials.

Setting

Sixteen hospitals in the Netherlands (derivation cohort) and 58 hospitals in the United States, Canada, Australia and Europe (validation cohort).

Participants

500 patients from the Multicenter Randomized Clinical Trial of Endovascular Treatment for Acute Ischemic Stroke in the Netherlands (MR CLEAN) trial (derivation cohort) and 260 patients with proven intracranial occlusion from the Interventional Management of Stroke III (IMS III) trial (validation cohort).

Main outcome measures

The primary outcome was the modified Rankin Scale (mRS) at 90 days after stroke. We constructed an ordinal logistic regression model to predict outcome and treatment benefit, defined as the difference between the predicted probability of good functional outcome (mRS 0–2) with and without EVT.

Results

Eleven baseline clinical and radiological characteristics were included in the model. The externally validated c-statistic was 0.68 (95% confidence interval (CI) 0.64 to 0.73) for the ordinal model and 0.73 (95% CI 0.67 to 0.79) for the prediction of good functional outcome, indicating moderate discriminative ability. The mean predicted treatment benefit varied between patients in the combined derivation and validation cohort from -2.3% to 24.3%. There was benefit of EVT predicted for some individual patients from groups in which no treatment effect was found in previous subgroup analyses, such as those with no or poor collaterals.

Conclusion

The proposed clinical decision tool combines multiple baseline clinical and radiological characteristics and shows large variations in treatment benefit between patients. The tool is clinically useful as it aids in distinguishing between individual patients who may experience benefit from EVT and those who will not.

Trial registration

clinicaltrials.gov NCT00359424 (IMS III) and isrctn.com ISRCTN10888758 (MR CLEAN).

Box 3.1 Research in context.**What is already known on this topic**

- *Endovascular treatment improves functional outcome in patients with acute ischemic stroke caused by a proximal occlusion.*
- *There is large variation in the selection of candidates for endovascular treatment in current practice because of the uncertainty of treatment benefit in specific subgroups.*

What this study adds

- *A newly developed clinical decision tool combines multiple baseline clinical and radiological characteristics and shows large variations in treatment benefit between patients.*
- *Selection of individual patients for endovascular treatment should therefore not be based on single patient characteristics.*
- *This model is the first step towards individualized selection of patients for endovascular treatment of ischemic stroke and may be used as a tool for assisting clinical decision making.*

Box 3.2 Descriptions of ASPECTS and collateral score.**ASPECTS**

A quantitative grading system to assess early ischemic changes on a non-contrast CT scan. Scores ranges from 0 to 10, with 10 points for a normal CT scan and 1 point subtracted for every defined region with evidence of early ischemic changes.¹⁶

Collateral score

A 4 point scale to grade the collateral flow of the occluded territory on vessel imaging, with 0 representing absent collateral flow, 1 representing poor collateral flow (<50% filling), 2 representing moderate collateral flow (between 50% and 100% filling), and 3 representing good collateral flow (100% filling).¹⁷

Background

Stroke is the second most common cause of mortality world-wide and of disability in high-income countries.¹ In Western countries, 80% of strokes are ischemic.² Ischemic strokes caused by a proximal occlusion in the intracranial cerebral arteries result in poor outcome.^{3,4} Endovascular treatment (EVT) improves functional outcome in patients with acute ischemic stroke caused by such a proximal occlusion,⁵⁻¹¹ with a number needed to treat of 5 (odds ratio 2.35, 95% confidence interval (CI) 1.85 to 2.98).¹² However, this is an average treatment effect and it is likely that treatment benefit will vary between individual patients.^{13,14} In current practice there is debate on the selection of candidates for EVT because of uncertainty of treatment benefit in specific subgroups and patients not included in the trials.^{12,15}

Clinicians combine multiple clinical features in their clinical decision making on how to treat an individual patient. For example, consider a 70-year old man who is admitted 40 minutes after onset of symptoms, with a severe left hemisphere ischemic stroke and a National Institutes of Health Stroke Scale (NIHSS) score of 22, an Alberta Stroke Program Early Computed Tomography Score (ASPECTS) of 7 and a M1 occlusion but no collaterals on computed tomography angiography (CTA). A previous subgroup analysis using data of the MR CLEAN trial suggested no treatment effect for patients with no or poor collaterals.¹⁵ But if this man can be treated very early after onset of stroke, will he benefit from EVT? Or consider a diabetic woman with high systolic blood pressure, aged 80, who arrived in a primary stroke center too late for treatment with intravenous (IV) alteplase, with a NIHSS score of 22, ASPECTS of 9, and a carotid-T occlusion with good collaterals on CTA. Should she be transferred to an intervention center 40 miles away if EVT just within the 6-hour time window is possible?

We developed and validated a clinical decision tool to provide individualized predictions of the effect of EVT based on multiple characteristics. Such a tool may be helpful to support clinical judgement when making complicated treatment decisions.

Methods

In short, we developed a multivariable prediction model in patients included in the MR CLEAN trial (Multicenter Randomized Clinical Trial of Endovascular Treatment for Acute Ischemic Stroke in the Netherlands, n=500) and validated this model in a subgroup of patients with an occlusion on CTA in the IMS III trial (Interventional Management of Stroke III, n=260). The primary outcome was the modified Rankin Scale (mRS) at 90 days after stroke. We constructed an ordinal logistic regression model to predict outcome and treatment benefit. This benefit was defined as the difference between the predicted probability of good functional outcome (mRS 0-2) with and without EVT. Variables were selected using univariable and multivariable selection steps (P<0.15).

Derivation cohort

We used data from all 500 patients of MR CLEAN (derivation cohort) for the development of our model.⁵ MR CLEAN was a phase III multicenter clinical trial with randomized treatment-group assignment, open-label treatment and blinded outcome evaluation. EVT plus usual care was compared with usual care alone (control group). Usual care could include treatment with IV alteplase if eligible. Enrolled patients were 18 years or older (no upper age limit), had a score of 2 or higher on the NIHSS (range 0 to 42), an occlusion of the proximal intracranial carotid artery, middle cerebral artery (M1 or M2), or anterior cerebral artery (A1 or A2), established with CTA. The start of EVT had to be possible within six hours after stroke onset. The imaging committee evaluated the findings on baseline non-contrast computed tomography (NCCT) for the ASPECTS and non-invasive baseline vessel imaging (CTA, magnetic resonance angiography, or digital subtraction angiography) for the location of the occlusion and collateral score.

More detailed information about MR CLEAN can be found in the study protocol and the publication of the main results.^{5,18}

Model development

Patient characteristics obtained before treatment that are expected to predict outcome or to interact with treatment, based on expert opinion or recent literature, were specified in our statistical analysis plan.¹⁹ We used ordinal logistic regression modeling, which assumes proportional odds, to test the effect of age, baseline NIHSS score, systolic blood pressure, treatment with IV alteplase, history of ischemic stroke, atrial fibrillation, diabetes mellitus, prestroke mRS, ASPECTS, location of occlusion, collateral score and time to treatment, as well as the corresponding interactions with treatment. Primary outcome was the mRS score, a 7-point scale ranging from 0 (no symptoms) to 6 (death), at 90 days after stroke.²⁰ For additional analyses, we derived the probabilities for good functional outcome (mRS 0–2) from the ordinal model. Treatment benefit was defined as the difference between the predicted probability of good functional outcome with and without EVT.

In our final multivariable model we selected the main effects or interaction terms with a P value of <0.15 in univariable and multivariable analyses. Location of occlusion was analyzed categorically and ASPECTS and collateral score were analyzed continuously. Continuous variables were not dichotomized. Non-linearity of continuous variables was tested with restricted cubic spline functions.²¹ In the final model we used restricted cubic spline functions for age and systolic blood pressure. As a measure for time to treatment we used the time from stroke onset to groin puncture. Since groin puncture was not performed in control subjects, time to groin puncture was not observable in the control arm. Single imputation based on regression using age, NIHSS score, inter-hospital transfer, hospital of first presentation and time to randomization, was used to assign time to expected groin puncture ($R^2=0.89$). Since all other variables were more than 98% complete within the derivation

cohort, we used simple imputation by the mean for continuous variables and simple imputation by the mode for categorical variables.

Internal validation with bootstrapping was used to estimate the degree of optimism in the final model. Regression coefficients were reduced with penalized regression to correct for this optimism.^{21,22} Coefficients of non-linear terms and interaction terms were reduced with a larger penalty than the main effects.²²

External validation

External validation of our model was performed in the Interventional Management of Stroke III (IMS III) trial.²³ The IMS III trial (n=656) was a phase III multicenter clinical trial with randomized treatment group assignment, open label treatment, and blinded outcome evaluation. The trial tested the approach of IV alteplase followed by EVT, as compared with standard IV alteplase. Further details on the methods used in the trial have been reported extensively.^{23,24}

We included patients with proven occlusion in the anterior circulation on non-invasive vessel imaging, and an available mRS score at 90 days in the validation cohort (n=260). Missing collateral scores because of insufficient CTA imaging (n=68) were replaced by single imputation with regression using age, history of diabetes mellitus and presence of internal carotid T occlusion. Single imputation for time to groin puncture (n=102; primarily control patients) was performed using age, NIHSS, time to randomization and inter-hospital transfer. All other variables were more than 98% complete and missing values of these variables were imputed with the mean (continuous variables) or mode (categorical variables).

Model performance in the validation cohort was expressed by discrimination and calibration. Discrimination was quantified with the concordance or c-statistic, which varies between 0.5 for a non-informative model and 1 for a perfectly discriminating model.²⁵ We calculated the general c-statistic of our ordinal model and an additional c-statistic for the predictions of good functional outcome (mRS 0–2).

Calibration refers to the level of agreement between predicted risks and observed outcome; this was assessed graphically with a validation plot for the prediction of good functional outcome (mRS 0–2) expressed as calibration slope and intercept. The calibration slope describes the effect of the predictors in the validation sample versus the derivation sample, and is ideally equal to 1. The intercept indicates whether predictions are systematically too high or too low, and should ideally be zero.²⁶

After external validation, the regression coefficients were fitted on a dataset combining all patients in the derivation and validation cohort. To assess if our model could be used to select individual patients for EVT, we estimated the individual predictions for all 760 patients

included in this combined dataset. We created a scatterplot with the predicted probabilities of good functional outcome (mRS 0–2) for these patients without EVT on the x-axis and the predicted probabilities with EVT on the y-axis. We made additional plots for the predictions of patients with no or poor collaterals and patients with low ASPECTS scores, since pre-specified subgroup analyses showed that these groups had no or very limited benefit of treatment.

All statistical analyses were performed with R statistical software (version 3.2.2) and the rms library (version 4.4-0). The web-application was developed with the R Shiny package (shiny version 0.13.0).

Patient involvement

No patients were involved in setting the research question or the outcome measures, nor were they involved in the design and implementation of the study. No patients were asked to advise on interpretation or writing up of results. There are plans to disseminate the results of the research to the relevant patient community.

Results

Table 3.1 shows that baseline patient characteristics and important characteristics of workflow and outcome were similar between the derivation cohort (n=500) and validation cohort (n=260). The validation cohort was somewhat more homogeneous, by not including patients with baseline disability (premorbid mRS ≥ 3) or patients not treated with IV alteplase.

Most variables were predictors of outcome (Table 3.2). The strongest predictors in multivariable analysis were age ($P < 0.001$), baseline NIHSS ($P < 0.001$), systolic blood pressure ($P < 0.001$), history of ischemic stroke ($P = 0.03$), diabetes mellitus ($P = 0.02$), prestroke mRS ($P = 0.003$), ASPECTS ($P = 0.001$), location of occlusion ($P = 0.03$) and collateral score ($P < 0.001$). Interactions with relative treatment effect were found in univariable analysis for history of ischemic stroke, atrial fibrillation, collateral score and time to groin puncture (all $P \leq 0.10$, Figure 3.1). In the multivariable model, the effects of EVT were similar as in univariable analysis, with larger effects in patients without previous ischemic stroke ($P = 0.07$), in patients with better collateral scores ($P = 0.07$), and in patients with shorter times to groin puncture ($P = 0.13$). Atrial fibrillation was not significant in multivariable analysis as either a main effect ($P = 0.67$) or interaction effect ($P = 0.27$), and was therefore excluded from the model.

The final multivariable model included age, baseline NIHSS score, systolic blood pressure, IV treatment with alteplase, history of ischemic stroke, diabetes mellitus, pre-stroke mRS, ASPECTS, location of occlusion, collateral score and time from onset to groin puncture. We added terms representing the interaction between treatment and each of previous stroke,

Table 3.1. Overview of the derivation and validation cohort.

	Derivation cohort (n=500)	Validation cohort (n=260)
Age, years – mean (SD)	65 (14)	67 (12)
Male sex	292 (58%)	135 (52%)
Baseline NIHSS – median (IQR)	18 (14–22)	17 (14–21)
Systolic blood pressure, mmHg – mean (SD)	145 (25)	149 (26)
Treatment with IV alteplase	445 (89%)	260 (100%)
Allocation to EVT	233 (47%)	174 (67%)
Medical history		
Ischemic stroke	54 (11%)	28 (11%)
Atrial fibrillation	135 (27%)	89 (35%)
Diabetes mellitus	68 (14%)	49 (19%)
Pre-stroke mRS		
0	404 (81%)	231 (89%)
1	50 (10%)	22 (8%)
2	25 (5%)	7 (3%)
≥3	21 (4%)	0
Imaging		
ASPECTS on NCCT – median (IQR)	9 (8–10)	8 (6–9)
Location of occlusion on non-invasive vessel imaging		
ICA-(T)	138 (28%)	66 (25%)
M1	319 (64%)	144 (55%)
M2	39 (8%)	50 (19%)
A1	3 (1%)	0
Workflow		
Onset to randomization, min – median (IQR)	200 (150–261)	143 (120–170)
Onset to groin puncture, min – median (IQR)	260 (210–311)	205 (168–235)
Onset to reperfusion, min – median (IQR)	340 (274–395)	275 (238–319)
Outcome		
Recanalization (mTICI 2B/3)	116 (59%)	69 (45%)
mRS at 90 days		
0	7 (1%)	27 (10%)
1	36 (7%)	46 (18%)
2	84 (17%)	39 (15%)
3	87 (17%)	36 (14%)
4	133 (27%)	44 (17%)
5	45 (9%)	18 (7%)
6 (mortality)	108 (22%)	50 (19%)

Values are numbers (percentages) unless stated otherwise.

ASPECTS, Alberta Stroke Program Early CT score; EVT, endovascular treatment; ICA-(T), internal carotid artery (with terminal segment); IQR, interquartile range; IV, intravenous; mRS, modified Rankin Scale; mTICI, modified thrombolysis in cerebral infarction scale; NCCT, non-contrast computed tomography; NIHSS, National Institutes of Health Stroke Scale.

collateral score and time to groin puncture. The internally validated c-statistic for ordinal outcome was 0.74 without interaction terms and this increased to 0.75 by adding interaction with treatment. The c-statistic for good functional outcome was 0.79.

External validation

Similar effects were found for most variables in the validation cohort except for systolic blood pressure, diabetes mellitus and the interaction between history of ischemic stroke and treatment effect. The externally validated c-statistic was 0.69 (95% CI 0.64 to 0.73) for the ordinal model and 0.73 (0.67 to 0.79) for the prediction of good functional outcome (Figure 3.2).

Table 3.2. Main effects in derivation cohort (n=500).

	Univariable model		Multivariable model	
	Common odds ratio (95% CI)	p-value	Common odds ratio (95% CI)	p-value
Intra-arterial treatment	1.66 (1.21 to 2.28)	0.002	1.86 (1.34 to 2.59)	<0.001
Age (per year)		<0.001		<0.001
<65 years	0.97 (0.95 to 0.99)		1.00 (0.97 to 1.02)	
>=65 years	0.92 (0.89 to 0.94)		0.92 (0.89 to 0.95)	
Baseline NIHSS (per point)	0.91 (0.88 to 0.94)	<0.001	0.93 (0.90 to 0.96)	<0.001
Systolic blood pressure (per 10 mmHg)		<0.001		<0.001
<130 mmHg	1.12 (0.88 to 1.41)		1.24 (0.97 to 1.59)	
>=130 mmHg	0.76 (0.70 to 0.83)		0.77 (0.70 to 0.85)	
Treatment with IV alteplase	1.85 (1.12 to 3.08)	0.02	1.62 (0.94 to 2.79)	0.08
History of ischemic stroke	0.48 (0.29 to 0.80)	0.005	0.53 (0.31 to 0.92)	0.03
Atrial fibrillation	0.52 (0.36 to 0.73)	<0.001	0.92 (0.62 to 1.36)	0.67
Diabetes mellitus	0.37 (0.23 to 0.59)	<0.001	0.54 (0.33 to 0.90)	0.02
Prestroke mRS	0.63 (0.52 to 0.77)	<0.001	0.72 (0.58 to 0.90)	0.003
ASPECTS (per point)	1.16 (1.07 to 1.26)	<0.001	1.16 (1.06 to 1.28)	0.001
Level of occlusion on non-invasive imaging		0.02		0.03
ICA-(T)	1.0 (reference)		1.0 (reference)	
M1	1.53 (1.08 to 2.17)		1.43 (0.98 to 2.07)	
M2	2.11 (1.15 to 3.88)		2.35 (1.20 to 4.60)	
Collateral score	1.95 (1.62 to 2.36)	<0.001	1.61 (1.31 to 1.96)	<0.001
Time from onset stroke to groin puncture (per 30 minutes)	0.94 (0.88 to 1.00)	0.07	0.93 (0.86 to 1.00)	0.04

Presented common odds ratios reflect the effect on the reversed modified Rankin Scale (odds ratio >1 corresponds with better functional outcome).

ASPECTS, Alberta Stroke Program Early CT score; CI, confidence interval; CTA, computed tomography angiography; EVT, endovascular treatment; ICA-(T), internal carotid artery (with terminal segment); IQR, interquartile range; IV, intravenous; mRS, modified Rankin Scale; NCCT, non-contrast computed tomography; NIHSS, National Institutes of Health Stroke Scale.




Figure 3.1. Univariable interaction effects in the derivation cohort (n=500). Interaction with treatment is expressed as the log odds for good functional outcome (modified Rankin Scale 0–2) with and without endovascular treatment on the y-axis. Variables on the x-axis are expressed continuously (time to groin puncture) or categorically (previous stroke, atrial fibrillation and collateral score).

The expected benefit of EVT varied largely between patients in the combined derivation and validation cohort (Figure 3.3A). Mean predicted absolute treatment benefit was an 11.8% higher probability of mRS 0–2 compared to the probability without EVT, and varied from -2.3% to 24.3% between individual patients in the combined derivation and validation cohort. The individual predictions for patients with no or poor collaterals (score 0–1) or low ASPECTS (score 0–5) illustrate the substantial variation in outcome and treatment benefit in these groups (Figures 3.3B and 3.3C). For some patients, who have multiple characteristics that negatively affect treatment benefit, the model predicts no benefit or even harm.

We calculated the predicted probabilities of good functional outcome with and without EVT for the two patients described in the introduction (Figure 3.4). The first patient is expected to benefit from EVT despite absent collaterals and moderate ASPECTS. The probability of achieving good functional outcome increases by 11 percentage points, from 16% without EVT to 27% with EVT. The predictions for the second patient illustrate that a good collateral score does not guarantee a large treatment benefit. The 80-year old patient has a low probability of achieving good functional outcome (3% without EVT and 5% with EVT), with some shift on the total mRS scale.




Figure 3.2. Calibration plot for predicted good functional outcome, defined as modified Rankin Scale (mRS) 0–2, in the validation cohort (n=260). The calibration slope reflects the strength of the predictors. The calibration intercept reflects the calibration-in-the-large, indicating whether predicted probabilities are systematically too low or too high. The overall observed proportion of patients with mRS 0–2 in the validation cohort was higher as to be expected using our model. The linear bar chart shows the distribution of patients with (=1) or without (=0) an observed outcome of mRS 0–2. Discrimination between low and high likelihood of good functional outcome was moderate (c-statistic 0.73, 95% CI 0.67 to 0.79).

We implemented our model in a web-application that provides predictions of outcome for individual patients based on baseline clinical and radiological characteristics for use in clinical practice. It shows bar charts with the expected distribution of mRS categories with and without EVT, the predicted probabilities of good functional outcome and the predicted absolute treatment benefit (Figure 3.4). This web-application was made accessible online at www.mrpredicts.com.




Figure 3.3. (A) Predicted probabilities of good functional outcome (modified Rankin Scale (mRS) score 0–2) for all individual patients in combined derivation and validation cohort (n=760). Each dot represents one individual patient, with the probability of good functional outcome (mRS score 0–2) without endovascular treatment (EVT) expressed on x-axis, and probability for good functional outcome with EVT on y-axis. Above the diagonal line the predicted probability of good functional outcome with EVT is higher than that without EVT. The farther above this line, the larger the predicted effect of treatment. (B) Patients highlighted with no or poor collaterals (score 0–1). (C) Patients highlighted with low Alberta Stroke Program Early CT Score (ASPECTS, score 0–5).




Figure 3.4. A stylized representation of the clinical decision tool. Baseline characteristics and predicted probabilities of good functional outcome (modified Rankin Scale (mRS) score 0–2) for two examples (see introduction). ASPECT, Alberta Stroke Program Early Computed Tomography Score; EVT, endovascular treatment; NIHSS, National Institutes of Health Stroke Scale; ICA, internal carotid artery; CTA, computed tomography angiography.

3

Discussion

We developed and externally validated a clinical decision tool to predict the benefit of EVT for individual patients with ischemic stroke, based on multiple patient characteristics. The predicted treatment benefit varied substantially between individual patients with different risk profiles.

Strengths and weaknesses in relation to other studies

Two risk scores have been described previously for the prediction of functional outcome after EVT.^{27,28} These scores are of limited value because they were developed on older cohorts of patients who were treated before the introduction of stent-retrievers and contain only a small number of clinical variables. Furthermore, they do not provide individual predictions and most of the variables and outcome measures in these studies had been dichotomized, which is considered to be statistically inefficient and biologically implausible.²⁹ Our model combines eleven baseline clinical and radiological characteristics simultaneously to provide individualized predictions of the effect of EVT. In contrast, conventional subgroup analyses focus mainly on predictive effects and assess the effect of only one variable at a time. Previous subgroup analyses of EVT trials have tested whether there are differences in effect of EVT based on for example: time to treatment,³⁰⁻³⁴ stroke severity,^{12,35} and collateral score.¹⁵ Analyzing one variable at a time may provide mechanistic insights to inform future studies and shape clinical considerations. However, they are of limited value in individual patient care, because treatment benefit is influenced by multiple individual factors simultaneously.^{13,14} Furthermore, even with similar relative treatment effect, individual patients may have different absolute treatment effects due to different baseline risks. More targeted individual treatment decisions can be obtained by using a more complex multivariable modeling approach to identify individual patients with large or small expected treatment benefit.¹³

We found modest interaction with treatment for history of ischemic stroke, collateral score and time from stroke onset to groin puncture. For collateral score and time to groin puncture, interaction with effect of EVT was already shown in previous subgroup analyses.^{15,30} Both variables are clinically likely to cause interaction with EVT. However, previous stroke has not been studied for interaction with treatment before, and was an unexpected finding in our study. It may be a chance finding, since it was not reproduced in the validation cohort and we have no clinical explanation. When the regression coefficients were fitted on data of the combined derivation and validation cohort and the coefficients of interaction terms were reduced to prevent overfitting, the interaction effect for previous stroke in the final model was small. Further validation should reveal whether the relative effect of EVT is modified by experience of a previous stroke.

Our study has several limitations. The discriminative ability of the model in the external validation was modest. It should be emphasized however, that the c-statistic for the ordinal

outcome is a conservative measure. It assesses discrimination between exact categories of the mRS, instead of discrimination between 2 groups with different outcome (for example mRS 0–2 versus mRS 3–6). Externally validated c-statistics of all cut-offs were better than the ordinal c-statistic (eg, 0.73 for good functional outcome and 0.75 for mortality). Nevertheless, the relatively small sample size and inclusion of interaction terms in the model may have resulted in some optimism and overfitting, despite shrinkage of the regression coefficients. The calibration was also suboptimal; despite the fact that most patients were treated with first generation thrombectomy devices, patients in the IMS III trial had a better outcome than predicted by our model. This could be explained by the patient selection in IMS III (eg, pre-morbid mRS 0–2, age <82 years, treatment with IV alteplase),³⁶ which resulted in a better prognosis overall. Patients in the IMS III control group had better outcomes than patients in the control group in MR CLEAN (mRS 0–2 = 39% (IMS III with occlusion on CTA) versus 19% (MR CLEAN)), leading to inadequate calibration of our model.^{5,36}

Implications for clinicians

Despite its limitations, the currently developed model is the first to predict the effect of EVT for individual patients upon arrival at the emergency department. When compared to other models used in neurovascular practice, HAS-BLED (c-statistic = 0.65) and CHA₂DS₂-VASc (c-statistic = 0.61), it performs accurately.^{37,38} The predictions made by our decision tool often agree with clinical intuition, which should not be surprising. However, estimates derived from large datasets are preferable to the subjective opinion of a physician, whose experience, no matter how vast, can never match the information contained in large datasets.³⁹

Currently, some centers withhold EVT in specific subgroups of patients (eg, low ASPECTS, no collaterals, age >80 years, or M2 occlusion). Indeed, our model predicts no benefit of EVT for some individual patients, especially when a patient has more than one characteristic that negatively affects the effect of EVT. The decision not to treat may be particularly relevant in patients who have to be transferred to an intervention center. The model may help to identify patients without expected benefit of EVT and topple the balance in favor of no treatment. But, perhaps more importantly, our study shows that treatment should not be withheld based on a single characteristic. Some patients belonging to one of the subgroups that are considered as having no benefit of EVT, such as poor collaterals or low ASPECTS, may still benefit from EVT substantially if other characteristics are favorable. This emphasizes the importance of making personalized treatment decisions, instead of using average treatment effects, and shows the need for combining multiple clinical and radiological baseline characteristics instead of withholding treatment based on a single characteristic.⁴⁰

This is the first model for EVT decision making. The predictions of our model should be considered as a starting point for clinical decision making, and not as a final recommendation. Our model was developed using the MR CLEAN database, consisting of an unselected population with few selection criteria. Therefore, our model is likely applicable in centers

that use few clinical and radiological selection criteria. Future analyses within larger studies may refine the current recommendations and improve the validity of the model.

Conclusion

The proposed clinical decision tool combines multiple baseline clinical and radiological characteristics and shows large variations in treatment benefit between patients. The model is clinically useful as it aids in distinguishing between individual patients who may experience benefit from endovascular treatment for acute ischemic stroke and those who will not.

References

1. Lopez AD, Mathers CD, Ezzati M, Jamison DT, Murray CJL. Global and regional burden of disease and risk factors, 2001: systematic analysis of population health data. *Lancet* 2006; **367**(9524): 1747-57.
2. van der Worp HB, van Gijn J. Clinical practice. Acute ischemic stroke. *N Engl J Med* 2007; **357**(6): 572-9.
3. Smith WS, Lev MH, English JD, et al. Significance of large vessel intracranial occlusion causing acute ischemic stroke and TIA. *Stroke* 2009; **40**(12): 3834-40.
4. Lima FO, Furie KL, Silva GS, et al. Prognosis of untreated strokes due to anterior circulation proximal intracranial arterial occlusions detected by use of computed tomography angiography. *JAMA Neurol* 2014; **71**(2): 151-7.
5. Berkhemer OA, Fransen PS, Beumer D, et al. A randomized trial of intraarterial treatment for acute ischemic stroke. *N Engl J Med* 2015; **372**(1): 11-20.
6. Campbell BC, Mitchell PJ, Kleinig TJ, et al. Endovascular therapy for ischemic stroke with perfusion-imaging selection. *N Engl J Med* 2015; **372**(11): 1009-18.
7. Goyal M, Demchuk AM, Menon BK, et al. Randomized assessment of rapid endovascular treatment of ischemic stroke. *N Engl J Med* 2015; **372**(11): 1019-30.
8. Jovin TG, Chamorro A, Cobo E, et al. Thrombectomy within 8 hours after symptom onset in ischemic stroke. *N Engl J Med* 2015; **372**(24): 2296-306.
9. Saver JL, Goyal M, Bonafe A, et al. Stent-retriever thrombectomy after intravenous t-PA vs. t-PA alone in stroke. *N Engl J Med* 2015; **372**(24): 2285-95.
10. Bracard S, Ducrocq X, Mas JL, et al. Mechanical thrombectomy after intravenous alteplase versus alteplase alone after stroke (THRACE): a randomised controlled trial. *Lancet Neurol* 2016; **15**(11): 1138-47.
11. Muir KW, Ford GA, Messow CM, et al. Endovascular therapy for acute ischaemic stroke: the Pragmatic Ischaemic Stroke Thrombectomy Evaluation (PISTE) randomised, controlled trial. *J Neurol Neurosurg Psychiatry* 2017; **88**(1): 38-44.
12. Goyal M, Menon BK, van Zwam WH, et al. Endovascular thrombectomy after large-vessel ischaemic stroke: a meta-analysis of individual patient data from five randomised trials. *Lancet* 2016; **387**(10029): 1723-31.
13. Kent DM, Rothwell PM, Ioannidis JP, Altman DG, Hayward RA. Assessing and reporting heterogeneity in treatment effects in clinical trials: a proposal. *Trials* 2010; **11**: 85.
14. Rothwell PM. Treating individuals 2. Subgroup analysis in randomised controlled trials: importance, indications, and interpretation. *Lancet* 2005; **365**(9454): 176-86.
15. Berkhemer OA, Jansen IG, Beumer D, et al. Collateral Status on Baseline Computed Tomographic Angiography and Intra-Arterial Treatment Effect in Patients With Proximal Anterior Circulation Stroke. *Stroke* 2016; **47**(3): 768-76.
16. Barber PA, Demchuk AM, Zhang J, Buchan AM. Validity and reliability of a quantitative computed tomography score in predicting outcome of hyperacute stroke before thrombolytic therapy. ASPECTS Study Group. Alberta Stroke Programme Early CT Score. *Lancet* 2000; **355**(9216): 1670-4.
17. Tan IYL, Demchuk AM, Hopyan J, et al. CT Angiography Clot Burden Score and Collateral Score: Correlation with Clinical and Radiologic Outcomes in Acute Middle Cerebral Artery Infarct. *Am J Neuroradiol* 2009; **30**(3): 525-31.
18. Fransen PS, Beumer D, Berkhemer OA, et al. MR CLEAN, a multicenter randomized clinical trial of endovascular treatment for acute ischemic stroke in the Netherlands: study protocol for a randomized controlled trial. *Trials* 2014; **15**: 343.
19. Mulder MJ, Venema E, Roozenbeek B, et al. Towards personalised intra-arterial treatment of patients with acute ischaemic stroke: a study protocol for development and validation of a clinical decision aid. *BMJ Open* 2017; **7**(3): e013699.
20. van Swieten JC, Koudstaal PJ, Visser MC, Schouten HJ, van Gijn J. Interobserver

- agreement for the assessment of handicap in stroke patients. *Stroke* 1988; **19**(5): 604-7.
21. Harrell FE. Regression modeling strategies: with applications to linear models, logistic regression, and survival analysis. *New York: Springer-Verlag* 2001.
 22. Steyerberg EW. Clinical prediction models: a practical approach to development, validation, and updating. *New York: Springer*, 2009.
 23. Broderick JP, Palesch YY, Demchuk AM, et al. Endovascular therapy after intravenous t-PA versus t-PA alone for stroke. *N Engl J Med* 2013; **368**(10): 893-903.
 24. Khatri P, Hill MD, Palesch YY, et al. Methodology of the Interventional Management of Stroke III Trial. *Int J Stroke* 2008; **3**(2): 130-7.
 25. Steyerberg EW, Vickers AJ, Cook NR, et al. Assessing the performance of prediction models: a framework for traditional and novel measures. *Epidemiology* 2010; **21**(1): 128-38.
 26. Steyerberg EW, Vergouwe Y. Towards better clinical prediction models: seven steps for development and an ABCD for validation. *Eur Heart J* 2014; **35**(29): 1925-31.
 27. Halleivi H, Barreto AD, Liebeskind DS, et al. Identifying patients at high risk for poor outcome after intra-arterial therapy for acute ischemic stroke. *Stroke* 2009; **40**(5): 1780-5.
 28. Flint AC, Cullen SP, Faigeles BS, Rao VA. Predicting long-term outcome after endovascular stroke treatment: the totaled health risks in vascular events score. *AJNR Am J Neuroradiol* 2010; **31**(7): 1192-6.
 29. Scott SC, Goldberg MS, Mayo NE. Statistical assessment of ordinal outcomes in comparative studies. *J Clin Epidemiol* 1997; **50**(1): 45-55.
 30. Franssen PS, Berkhemer OA, Lingsma HF, et al. Time to Reperfusion and Treatment Effect for Acute Ischemic Stroke: A Randomized Clinical Trial. *JAMA Neurol* 2015: 1-7.
 31. Goyal M, Jadhav AP, Bonafe A, et al. Analysis of Workflow and Time to Treatment and the Effects on Outcome in Endovascular Treatment of Acute Ischemic Stroke: Results from the SWIFT PRIME Randomized Controlled Trial. *Radiology* 2016; **279**(3): 888-97.
 32. Menon BK, Sajobi TT, Zhang Y, et al. Analysis of Workflow and Time to Treatment on Thrombectomy Outcome in the Endovascular Treatment for Small Core and Proximal Occlusion Ischemic Stroke (ESCAPE) Randomized, Controlled Trial. *Circulation* 2016; **133**(23): 2279-86.
 33. Ribo M, Molina CA, Cobo E, et al. Association Between Time to Reperfusion and Outcome Is Primarily Driven by the Time From Imaging to Reperfusion. *Stroke* 2016; **47**(4): 999-1004.
 34. Saver JL, Goyal M, van der Lugt A, et al. Time to Treatment With Endovascular Thrombectomy and Outcomes From Ischemic Stroke: A Meta-analysis. *JAMA* 2016; **316**(12): 1279-88.
 35. Broderick JP, Berkhemer OA, Palesch YY, et al. Endovascular Therapy Is Effective and Safe for Patients With Severe Ischemic Stroke: Pooled Analysis of Interventional Management of Stroke III and Multicenter Randomized Clinical Trial of Endovascular Therapy for Acute Ischemic Stroke in the Netherlands Data. *Stroke* 2015; **46**(12): 3416-22.
 36. Demchuk AM, Goyal M, Yeatts SD, et al. Recanalization and clinical outcome of occlusion sites at baseline CT angiography in the Interventional Management of Stroke III trial. *Radiology* 2014; **273**(1): 202-10.
 37. Lip GY, Frison L, Halperin JL, Lane DA. Comparative validation of a novel risk score for predicting bleeding risk in anticoagulated patients with atrial fibrillation: the HAS-BLED (Hypertension, Abnormal Renal/Liver Function, Stroke, Bleeding History or Predisposition, Labile INR, Elderly, Drugs/Alcohol Concomitantly) score. *J Am Coll Cardiol* 2011; **57**(2): 173-80.
 38. Lip GY, Nieuwlaat R, Pisters R, Lane DA, Crijns HJ. Refining clinical risk stratification for predicting stroke and thromboembolism in atrial fibrillation using a novel risk factor-based approach: the euro heart survey on atrial fibrillation. *Chest* 2010; **137**(2): 263-72.
 39. Lingsma HF, Roozenbeek B, Steyerberg EW, Murray GD, Maas AI. Early prognosis in traumatic brain injury: from prophecies to predictions. *Lancet Neurol* 2010; **9**(5): 543-54.
 40. Kent DM, Nelson J, Dahabreh IJ, Rothwell PM, Altman DG, Hayward RA. Risk and treatment effect heterogeneity: re-analysis of individual participant data from 32 large clinical trials. *Int J Epidemiol* 2016; **45**(6): 2075-88.





Chapter 4

Improving selection of patients for endovascular treatment of ischemic stroke: validation and updating of MR PREDICTS with data from 4,398 patients

Submitted

Esmee Venema, Bob Roozenbeek, Maxim J.H.L. Mulder, Scott Brown, Charles B.L.M. Majoie, Ewout W. Steyerberg, Andrew M. Demchuk, Keith W. Muir, Antoni Dávalos, Peter J. Mitchell, Serge Bracard, Olvert A. Berkhemer, Geert J. Lycklama à Nijeholt, Robert J. van Oostenbrugge, Yvo B.W.E.M. Roos, Wim H. van Zwam, Aad van der Lugt, Michael D. Hill, Philip White, Bruce C.V. Campbell, Francis Guillemin, Jeffrey L. Saver, Tudor G. Jovin, Mayank Goyal, Diederik W.J. Dippel, and Hester F. Lingsma, on behalf of the HERMES collaborators and MR CLEAN Registry Investigators



Chapter 5

Multivariable outcome prediction after endovascular treatment for ischemic stroke (MR PREDICTS@24H): a post-procedural tool to predict functional outcome at 3 months

In preparation

Vicky Chalos, Esmee Venema, Maxim J.H.L. Mulder, Bob Roozenbeek, Ewout W. Steyerberg, Marieke J.H. Wermer, Geert J. Lycklama à Nijeholt, H. Bart van der Worp, Mayank Goyal, Bruce C.V. Campbell, Keith W. Muir, Francis Guillemin, Serge Bracard, Philip White, Antoni Dávalos, Tudor G. Jovin, Michael D. Hill, Peter J. Mitchell, Andrew M. Demchuk, Jeffrey L. Saver, Aad van der Lugt, Scott Brown, Hester F. Lingsma, and Diederik W.J. Dippel, on behalf of the HERMES collaborators and MR CLEAN Registry investigators

Part II

Treat at the right time



Chapter 6

Workflow and factors associated with delay in the delivery of endovascular treatment for ischemic stroke in the MR CLEAN trial

J Neurointerv Surg, 2018

Esmee Venema*, Nikki Boodt*, Olvert A. Berkhemer, Pleunie P.M. Rood, Wim H. van Zwam, Robert J. van Oostenbrugge, Aad van der Lugt, Yvo B.W.E.M. Roos, Charles B.L.M. Majoie, Hester F. Lingsma, and Diederik W.J. Dippel, on behalf of the MR CLEAN investigators

*equal contribution

Abstract

Objective

The effect of endovascular treatment (EVT) for acute ischemic stroke is highly time-dependent. We investigated delay of EVT and factors associated with such delay.

Methods

MR CLEAN was a randomized trial of EVT plus usual care versus usual care alone (n=500). With multivariable linear regression, we analyzed the effect of intravenous treatment, general anesthesia, off-hours and inter-hospital transfer on time to admission at the emergency department (ED) of the intervention center and time to treatment. Furthermore, we assessed compliance with a target of 75 minutes for time from ED to treatment, and calculated the potential absolute increase of patients with a good outcome (modified Rankin Scale score ≤ 2), if this target had been achieved in all treated patients.

Results

Inter-hospital transfer prolonged time to ED with 140 minutes (95% confidence interval 129 to 150), but reduced time from ED to treatment with 77 minutes (64 to 91). Time from ED to treatment was increased with 19 minutes by general anesthesia (5 to 33) and total time was increased with 23 minutes during off-hours (6 to 40). The in-hospital target was achieved in 11.5% (22/191) of patients. Full compliance with the target time of 75 minutes from ED to treatment would have increased the proportion of patients with good outcome with 7.6% (6.7% to 8.5%).

Conclusion

Inter-hospital transfer is an important cause of delay in the delivery of EVT and every effort should be made to avoid transfers and reduce transfer-related delay. Furthermore, in-hospital workflow should be optimized to improve functional outcome after EVT.

Introduction

Endovascular treatment (EVT) with the use of stent retrievers has been proven safe and effective for patients with acute ischemic stroke due to a proximal intracranial arterial occlusion in the anterior circulation.¹ The effect of treatment is highly time-dependent and increased time from stroke onset to reperfusion is associated with a decreased likelihood of good functional outcome.²⁻⁷ The Multicenter Randomized Clinical Trial of Endovascular Treatment for Acute Ischemic Stroke in the Netherlands (MR CLEAN) showed a decrease in the effect of EVT on good outcome (risk difference between intervention and control) of 6.4% per hour delay in time to expected reperfusion.³ It seems likely that reducing treatment delay will further increase the effect of EVT.

The timeline from stroke onset to reperfusion consists of several steps, and each step is prone to factors causing delay. An important potential cause of delay in the prehospital stage is referral to a community hospital without facilities for EVT (a primary stroke center), which necessitates inter-hospital transportation of patients to an intervention center. Other factors that may increase time to reperfusion are intravenous treatment with alteplase (IVT), the use of general anesthesia during the endovascular procedure, and inefficiency of in-hospital workflow, specifically outside working hours.⁸⁻¹⁰ In-hospital workflow consists of all protocols and procedures in the emergency department (ED) of the primary stroke center, the ED of the intervention center, and the interventional suite.

In addition to the previously defined benchmark time of 60 minutes for IVT, various lenient or strict target times for EVT are now being suggested to improve in-hospital workflow.¹¹⁻¹³ The Highly Effective Reperfusion Evaluated in Multiple Endovascular Stroke Trials (HERMES) collaborators reported the 25th percentile of ED admission to groin puncture, which amounted to 75 minutes, and could very well serve as a benchmark time.⁷

Our aim was to investigate prehospital and in-hospital delay in the delivery of EVT in the MR CLEAN trial and to assess the impact of inter-hospital transfer and other factors on such delay.

Methods

Study design

MR CLEAN was a multicenter randomized clinical trial in the Netherlands comparing EVT plus usual care (intervention group) with usual care alone (control group).^{14,15} MR CLEAN is registered with the ISRCTN registry (ISRCTN10888758).

Study setting and population

Patients of 18 years or older with a clinical diagnosis of acute stroke, a deficit on the National Institute of Health Severity of Stroke (NIHSS) scale of 2 points or more, and a proximal occlusion in the anterior circulation on vessel imaging were included. Detailed inclusion and exclusion criteria can be found in the study protocol.¹⁴ Usual care could include IVT if indicated. The endovascular procedure had to be initiated within 6 hours after onset of stroke symptoms and could include intra-arterial delivery of a thrombolytic agent at the level of occlusion, mechanical thrombectomy, or both. In total 500 patients were enrolled from December 2010 to March 2014. Of these patients, 233 were allocated to the intervention arm and 267 were allocated to the control arm. EVT consisted of stent thrombectomy in the majority of cases (195/233, 84%).^{14,15}

Study protocol

Inclusion in the study could occur in three settings: (1) patients arrived directly at an intervention center after onset of stroke symptoms; (2) patients were presented at a primary stroke center first before being transferred to an intervention center for study participation; or (3) patients suffering a stroke while already hospitalized. In the hospital of first presentation, a non-contrast CT (NCCT) was performed to exclude intracerebral hemorrhage. This was followed by standard medical treatment which could include IVT. Vessel imaging with computed tomographic angiography (CTA), magnetic resonance angiography (MRA), or digital-subtraction angiography (DSA), had to be performed before inclusion in the study in the intervention center, to assess the presence of an intracranial arterial occlusion.

Time points of various events were prospectively collected in the trial, including time of stroke onset, start of IVT, admission to the ED of the intervention center and time of randomization. In patients receiving EVT, timing of treatment was documented by the local interventionist. Time of imaging (ie, NCCT and CTA, MRA or DSA) was prospectively recorded in the intervention centers, but had to be collected retrospectively if performed in a primary stroke center.

Definitions and outcome measures

Stroke onset was defined as the time point when stroke symptoms were first noticed by the patient or an observer. In case the time of first symptoms was unknown, stroke onset was defined as the moment the patient was last seen well. We defined off-hours as weekend days or working days between 17:00 hours and 08:00 hours. Groin puncture was defined as the moment of placing a sheet in the groin, indicating the start of the endovascular procedure. Reperfusion was measured with the modified Thrombolysis in Cerebral Infarction (mTICI) score, which can range from grade 0 for no reperfusion, to grade 3 for full reperfusion.¹⁶ We defined time of reperfusion as the first moment a mTICI score of 2b or 3 was reached. When reperfusion was not achieved, we used the timepoint when the last angiography run was done, which indicated the end of the endovascular procedure.

Time intervals of primary interest were: (1) time from stroke onset to admission at the ED of the intervention center as a measure for prehospital delay; (2) time from admission at the ED of the intervention center to groin puncture as a measure for in-hospital delay; and (3) time from stroke onset to groin puncture. Additionally, we compared the following time intervals between transfer patients and direct patients: (4) time from stroke onset to IVT; (5) time from admission at ED to vessel imaging; (6) time from vessel imaging to randomization; and (7) time from groin puncture to reperfusion. If all vessel imaging was performed in the primary stroke center, the patient was excluded from analyses (5) and (6). Time of transportation from the primary stroke center to the intervention center was estimated for all transfer patients with the TomTom MyDrive application (version 4.1.4.3089, available at mydrive.tomtom.com).

Statistical analysis

For descriptive purposes, time intervals were reported as medians with interquartile range (IQR). To assess transfer-related treatment delay, we compared baseline characteristics and median time intervals of patients transferred from a primary stroke center versus patients directly arrived at the intervention center.

Using multivariable linear regression analyses we evaluated the effect of patient and workflow-related characteristics on time to treatment. Factors assessed were administration of IVT, use of general anesthesia during the endovascular procedure, admission during off-hours and inter-hospital transfer. We included the following potential confounders in these analyses: age, gender, baseline NIHSS score, history of ischemic stroke and prestroke modified Rankin Scale (mRS). Missing baseline characteristics were imputed with the mean (continue variables) or mode (categorical variables). Missing time variables were not imputed.

To assess the impact of in-hospital delay, we compared the time from admission at ED to groin puncture with the target time of 75 minutes. For each individual patient we estimated the excess delay by subtracting target time from the actual time. The mean excess delay of all treated patients was multiplied by an increase in treatment benefit (defined as the risk difference for good outcome (mRS score 0–2)) of 6.4% per hour, based on previous results of the MR CLEAN trial.³ This provides an estimate of the potential increase in treatment benefit if the target time would have been achieved in all treated patients.

All statistical analyses were performed with R software (version 3.2.1, R Foundation).

Results

Of the 500 patients included in MR CLEAN, 281 (56.2%) arrived directly at an intervention center and 219 (43.8%) were transferred from a primary stroke center. Transfer patients were significantly younger than those arriving directly at an intervention center (62.2 versus 67.0 years) and were admitted more frequently during off-hours (63.4 versus 51.6%). Other baseline characteristics are shown in Table 6.1.

Table 6.1. Baseline characteristics of patients in the MR CLEAN trial, according to mode of arrival at the emergency department of the intervention center.

	Direct (n=281)	Transfer (n=219)	p-value
Age, years – mean (SD)	67.0 (13.6)	62.2 (13.6)	<0.01
Male sex – n (%)	156 (55.5)	136 (62.1)	0.16
Medical history – n (%)			
Atrial fibrillation	77 (27.4)	58 (26.5)	0.90
Hypertension	135 (48.0)	92 (42.0)	0.21
Hyperlipidemia	71 (25.3)	58 (26.5)	0.84
Diabetes mellitus	41 (14.6)	27 (12.3)	0.55
Previous ischemic stroke	33 (11.7)	21 (9.6)	0.53
Prestroke mRS – n (%)			0.09
0	220 (78.3)	184 (84.0)	
1	27 (9.6)	23 (10.5)	
2	19 (6.8)	6 (2.7)	
>2	15 (5.3)	6 (2.7)	
NIHSS score – median (IQR)	17 (14–22)	18 (15–22)	0.13
Location of occlusion – n (%)*			0.24
Internal carotid artery	72 (25.7)	66 (30.1)	
M1 middle cerebral artery segment	180 (64.3)	139 (63.5)	
M2 middle cerebral artery segment	27 (9.6)	12 (5.5)	
A1 or A2 anterior cerebral artery segment	1 (0.4)	2 (0.9)	
IVT – n (%)	249 (88.6)	196 (89.5)	0.87
Admission during off hours – n (%)†	141 (51.6)	128 (63.4)	0.01
Allocation to EVT – n (%)	132 (47.0)	101 (46.1)	0.92
Use of general anesthesia – n (%)‡	39 (32.5)	41 (42.3)	0.18

*One missing value. †25 missing values. ‡Only endovascular treated patients.

EVT, endovascular treatment; IVT, intravenous treatment; mRS, modified Rankin Scale; NIHSS, National Institutes of Health Stroke Scale.

Table 6.2. Time intervals in the MR CLEAN trial according to mode of arrival at the emergency department of the intervention center, expressed as median (IQR).

All patients	Direct (n=281)	Transfer (n=219)
Stroke onset to IVT (n=444)	85 (65–110)	85 (66–110)
Stroke onset to admission to ED (n=471)	52 (34–75)	200 (166–245)
Admission to ED to vessel imaging (n=439)	39 (25–67)	24 (17–32)
Vessel imaging to randomization (n=460)	53 (34–80)	21 (15–33)
Endovascular treated patients	Direct (n=132)	Transfer (n=101)
Stroke onset to groin puncture (n=205)	233 (202–287)	315 (260–346)
Admission to ED to groin puncture (n=192)	170 (142–205)	96 (75–120)
Groin puncture to reperfusion (n=194)	56 (42–86)	61 (37–90)

ED, emergency department; IVT, intravenous treatment.

Median time from stroke onset to groin puncture was 233 minutes (IQR 202–287) in direct arrival patients, and 315 minutes (260–346) in transfer patients (Table 6.2). The median time of inter-hospital transportation was 30 minutes (23–41), but total time from onset to admission at the ED of the intervention center was 148 minutes longer in transfer patients than in direct arrival patients. In-hospital workflow was improved in transfer patients, with a decrease in median time from arrival to vessel imaging of 15 minutes and a decrease in time from vessel imaging to randomization of 32 minutes. Duration of the endovascular procedure from groin puncture to reperfusion was comparable (56 minutes in direct arrival patients versus 61 minutes in transfer patients). Overall, we observed an association between increasing time from onset of stroke to admission at ED and decreasing time from admission to groin puncture (Supplementary Figure 6.1). However, the average prehospital delay caused by inter-hospital transfer was only partially compensated by improved in-hospital workflow (Figure 6.1).

In multivariable analysis, the total time to treatment (groin puncture) was increased by 23 minutes during off-hours (95% confidence interval (CI) 6 to 40) and by 65 minutes due to inter-hospital transfer (95% CI 48 to 82) (Figure 6.2). Transfer prolonged time to admission to the ED by 140 minutes (129 to 150), but was associated with a shorter time to groin puncture (77 minutes, 95% CI 64 to 91). Time from ED to groin puncture was increased by an additional 19 minutes by the use of general anesthesia during the endovascular procedure (95% CI 5 to 33).




Figure 6.1. Mean time intervals in MR CLEAN (n=500) according to mode of arrival at the emergency department of the intervention center. CTA, computed tomography angiography; ED, emergency department; IVT, intravenous treatment.




Figure 6.2. Effect of clinical factors on time intervals in MR CLEAN (n=500). ED, emergency department (of the intervention center).

In patients receiving EVT, time from admission to the ED to groin puncture was on average 145 ± 61 minutes. The defined target time of 75 minutes was reached in only 11.5% (22/191) of the treated patients. However, if this target time had been achieved in all patients, mean time from onset to groin puncture would have been 71 minutes (95% CI 63 to 80) shorter. Because the risk difference of good functional outcome decreases with 6.4% per hour, the absolute treatment benefit (proportion of patients with mRS 0–2) in MR CLEAN could have been 7.6% (6.7% to 8.5%) higher.

Discussion

This study shows that inter-hospital transfer was the most important factor associated with treatment delay in MR CLEAN. Time from onset to treatment increases with more than one hour in patients transferred from a primary stroke center, despite improved in-hospital workflow times. Furthermore, only a few patients received EVT within the proposed target time of 75 minutes for time from admission to the ED to groin puncture, and enhancing this compliance by improving in-hospital workflow might further increase functional outcome after EVT.

Overall, the time intervals in MR CLEAN are longer than in several other trials on EVT, such as The Endovascular Treatment for Small Core and Proximal Occlusion Ischemic Stroke (ESCAPE), EXtending the time for Thrombolysis in Emergency Neurological Deficits with Intra-Arterial therapy (EXTEND-IA) and Solitaire With the Intention for Thrombectomy as Primary Endovascular Treatment (SWIFT PRIME).¹⁷⁻¹⁹ For example, median time to reperfusion was 340 minutes in MR CLEAN versus 241 minutes in ESCAPE. This can be explained by the strong focus on improving workflow and reducing time delays in ESCAPE, while MR CLEAN did not require the participating investigators to be trained in delivery of fast care and logistics. Furthermore, the study design of these trials allowed verbal or deferred informed consent, while in MR CLEAN a written informed consent was required before randomization.^{15,18} Trials with a shorter median time to groin puncture had a significant larger treatment effect (Supplementary Figure 6.2).

Total time to treatment was increased most by inter-hospital transfer, confirming the results of previous studies.^{4,5,9,10} Transfer patients arrived 148 minutes later at the ED, while the median transportation time in our study was only 30 minutes. This suggests that that most delay was caused during work-up in the primary hospital. However, increased time to arrival at the ED was clearly associated with a decrease in the following time to groin puncture. This paradoxical effect was likely caused by the fact that this group consisted mainly of transfer patients who had already undergone part of the workup, the intervention center was pre-notified and clinical information was available before arrival at the ED, which made it possible for the staff to mobilize the neuro-interventional team and for the angiosuite to

be prepared in advance. This finding suggests that improvement of in-hospital workflow might lead to a shorter time from ED to treatment for patients who are directly admitted to an intervention center.

Our study has a number of limitations. Our analysis was restricted to data that was collected during the trial. We had no information available on the time of arrival in the primary stroke center and the time of departure to the intervention center to estimate door-in to door-out times, while this might have provided more insight into the various elements of delay. Also, the reported workflow times from this randomized study might not be representative for daily clinical practice, since the process of obtaining written informed consent and randomization can increase treatment delay substantially. Furthermore, the vessel imaging in MR CLEAN was often performed or repeated in the intervention center, while in daily routine transfer patients can be directly transferred to the interventional suite without any delay of additional imaging. Since the distances between hospitals in the Netherlands are very short and the median transportation time in this study was only 30 minutes, our results cannot be easily extrapolated to other countries. Transportation times in larger countries are usually longer, and the effect of inter-hospital transfer on treatment delay might therefore be much stronger in those countries. Last, inter-hospital transfer has not only a disadvantageous effect on time to treatment, but also on the number of patients that might be eligible for treatment.²⁰ We had no data on the patients in primary stroke centers who were not included in the trial because of the need for a transfer and the associated delay. Therefore, we were not able to quantify the impact of transfer delay on the likelihood of receiving EVT.

Overall, it remains clear that inter-hospital transfer causes significant pre-hospital delays in EVT, and that adequate measures have to be taken to shorten transfer times and, if possible, to reduce the number of transferred patients. This could be achieved by transporting ischemic stroke patients directly to an intervention center using prehospital triaging scales to select patients with a high risk of a large vessel occlusion.²¹⁻²³ These scales have to be implemented in regional care systems and protocols for prehospital triage to minimize transfer delays depending on the characteristics of the area.²⁴⁻²⁶ In regions with very large distances, the additional use of air ambulance services for the transportation of EVT candidates might be beneficial. Another possibility would be to bypass the primary stroke center by using a mobile stroke unit; an ambulance with a CT scanner and a specialized team on board, allowing in-ambulance initiation and administration of IVT and direct transportation to the intervention center.²⁷⁻²⁹

The other important step in reducing treatment delay in EVT is improving in-hospital workflow. The reduced time from ED to groin puncture in transfer patients suggests that it should be possible to shorten the time delay in the ED. Previous studies have already shown that optimizing workflow and protocols can decrease time to treatment.^{30,31} Pre-notification

of the interventional team as soon as a patient with a suspected large vessel occlusion is announced, direct transport from door to scanner, a parallel nature of care processes and optimizing the processes around IVT administration and initiation of EVT are all potential factors to reduce ED delay.^{10,32,33} Providing reasonable time targets might also improve this process, as was shown with the introduction of door-to-balloon target times for percutaneous coronary intervention.³⁴


In summary, inter-hospital transfer is an important cause of delay in the delivery of EVT and every effort should be made to avoid transfer and reduce transfer-related delay. Furthermore, in-hospital workflow can and should be optimized to improve functional outcome after EVT.

References


1. Goyal M, Menon BK, van Zwam WH, et al. Endovascular thrombectomy after large-vessel ischaemic stroke: a meta-analysis of individual patient data from five randomised trials. *Lancet* 2016; **387**(10029): 1723-31.
2. Khatri P, Yeatts SD, Mazighi M, et al. Time to angiographic reperfusion and clinical outcome after acute ischaemic stroke: an analysis of data from the Interventional Management of Stroke (IMS III) phase 3 trial. *Lancet Neurol* 2014; **13**(6): 567-74.
3. Fransen PS, Berkhemer OA, Lingsma HF, et al. Time to Reperfusion and Treatment Effect for Acute Ischemic Stroke: A Randomized Clinical Trial. *JAMA Neurol* 2015: 1-7.
4. Goyal M, Jadhav AP, Bonafe A, et al. Analysis of Workflow and Time to Treatment and the Effects on Outcome in Endovascular Treatment of Acute Ischemic Stroke: Results from the SWIFT PRIME Randomized Controlled Trial. *Radiology* 2016; **279**(3): 888-97.
5. Menon BK, Sajobi TT, Zhang Y, et al. Analysis of Workflow and Time to Treatment on Thrombectomy Outcome in the Endovascular Treatment for Small Core and Proximal Occlusion Ischemic Stroke (ESCAPE) Randomized, Controlled Trial. *Circulation* 2016; **133**(23): 2279-86.
6. Ribo M, Molina CA, Cobo E, et al. Association Between Time to Reperfusion and Outcome Is Primarily Driven by the Time From Imaging to Reperfusion. *Stroke* 2016; **47**(4): 999-1004.
7. Saver JL, Goyal M, van der Lugt A, et al. Time to Treatment With Endovascular Thrombectomy and Outcomes From Ischemic Stroke: A Meta-analysis. *JAMA* 2016; **316**(12): 1279-88.
8. Asif KS, Lazzaro MA, Zaidat O. Identifying delays to mechanical thrombectomy for acute stroke: onset to door and door to clot times. *J Neurointerv Surg* 2014; **6**(7): 505-10.
9. Goyal M, Almekhlafi MA, Fan LQ, et al. Evaluation of Interval Times From Onset to Reperfusion in Patients Undergoing Endovascular Therapy in the Interventional Management of Stroke III Trial. *Circulation* 2014; **130**(3): 265-U120.
10. Menon BK, Almekhlafi MA, Pereira VM, et al. Optimal Workflow and Process-Based Performance Measures for Endovascular Therapy in Acute Ischemic Stroke Analysis of the Solitaire FR Thrombectomy for Acute Revascularization Study. *Stroke* 2014; **45**(7): 2024-9.
11. Casaubon LK, Boulanger JM, Blacquiere D, et al. Canadian Stroke Best Practice Recommendations: Hyperacute Stroke Care Guidelines, Update 2015. *Int J Stroke* 2015; **10**(6): 924-40.
12. Sacks D, Black CM, Cognard C, et al. Multisociety Consensus Quality Improvement Guidelines for Intraarterial Catheter-directed Treatment of Acute Ischemic Stroke, from the American Society of Neuroradiology, Canadian Interventional Radiology Association, Cardiovascular and Interventional Radiological Society of Europe, Society for Cardiovascular Angiography and Interventions, Society of Interventional Radiology, Society of NeuroInterventional Surgery, European Society of Minimally Invasive Neurological Therapy, and Society of Vascular and Interventional Neurology. *AJNR Am J Neuroradiol* 2013; **34**(4): E0.
13. McTaggart RA, Ansari SA, Goyal M, et al. Initial hospital management of patients with emergent large vessel occlusion (ELVO): report of the standards and guidelines committee of the Society of NeuroInterventional Surgery. *J Neurointerv Surg* 2015.
14. Fransen PS, Beumer D, Berkhemer OA, et al. MR CLEAN, a multicenter randomized clinical trial of endovascular treatment for acute ischemic stroke in the Netherlands: study protocol for a randomized controlled trial. *Trials* 2014; **15**: 343.
15. Berkhemer OA, Fransen PSS, Beumer D, et al. A Randomized Trial of Intraarterial Treatment for Acute Ischemic Stroke. *New Engl J Med* 2015; **372**(1): 11-20.
16. Zaidat OO, Yoo AJ, Khatri P, et al. Recommendations on angiographic revascularization grading standards for acute ischemic stroke: a consensus statement. *Stroke* 2013; **44**(9): 2650-63.
17. Campbell BCV, Mitchell PJ, Kleinig TJ, et al. Endovascular Therapy for Ischemic Stroke with Perfusion-Imaging Selection. *New Engl J Med* 2015; **372**(11): 1009-18.

18. Goyal M, Demchuk AM, Menon BK, et al. Randomized Assessment of Rapid Endovascular Treatment of Ischemic Stroke. *New Engl J Med* 2015; **372**(11): 1019-30.
19. Saver JL, Goyal M, Bonafe A, et al. Stent-Retriever Thrombectomy after Intravenous t-PA vs. t-PA Alone in Stroke. *New Engl J Med* 2015; **372**(24): 2285-95.
20. Prabhakaran S, Ward E, John S, et al. Transfer delay is a major factor limiting the use of intra-arterial treatment in acute ischemic stroke. *Stroke* 2011; **42**(6): 1626-30.
21. Perez de la Ossa N, Carrera D, Gorchs M, et al. Design and validation of a prehospital stroke scale to predict large arterial occlusion: the rapid arterial occlusion evaluation scale. *Stroke* 2014; **45**(1): 87-91.
22. Lima FO, Silva GS, Furie KL, et al. Field Assessment Stroke Triage for Emergency Destination: A Simple and Accurate Prehospital Scale to Detect Large Vessel Occlusion Strokes. *Stroke* 2016; **47**(8): 1997-2002.
23. Hastrup S, Damgaard D, Johnsen SP, Andersen G. Prehospital Acute Stroke Severity Scale to Predict Large Artery Occlusion: Design and Comparison With Other Scales. *Stroke* 2016; **47**(7): 1772-6.
24. Michel P. Prehospital Scales for Large Vessel Occlusion: Closing in on a Moving Target. *Stroke* 2017; **48**(2): 247-9.
25. Holodinsky JK, Williamson TS, Kamal N, Mayank D, Hill MD, Goyal M. Drip and Ship Versus Direct to Comprehensive Stroke Center: Conditional Probability Modeling. *Stroke* 2017; **48**(1): 233-8.
26. Milne MS, Holodinsky JK, Hill MD, et al. Drip 'n Ship Versus Mothership for Endovascular Treatment: Modeling the Best Transportation Options for Optimal Outcomes. *Stroke* 2017; **48**(3): 791-4.
27. Ebinger M, Winter B, Wendt M, et al. Effect of the use of ambulance-based thrombolysis on time to thrombolysis in acute ischemic stroke: a randomized clinical trial. *JAMA* 2014; **311**(16): 1622-31.
28. Rajan SS, Baraniuk S, Parker S, Wu TC, Bowry R, Grotta JC. Implementing a mobile stroke unit program in the United States: why, how, and how much? *JAMA Neurol* 2015; **72**(2): 229-34.
29. Walter S, Kostopoulos P, Haass A, et al. Diagnosis and treatment of patients with stroke in a mobile stroke unit versus in hospital: a randomised controlled trial. *Lancet Neurol* 2012; **11**(5): 397-404.
30. Aghaebrahim A, Streib C, Rangaraju S, et al. Streamlining door to recanalization processes in endovascular stroke therapy. *J Neurointerv Surg* 2017; **9**(4): 340-5.
31. Schregel K, Behme D, Tsogkas I, et al. Effects of Workflow Optimization in Endovascularly Treated Stroke Patients - A Pre-Post Effectiveness Study. *PLoS One* 2016; **11**(12): e0169192.
32. Zerna C, Assis Z, d'Esterre CD, Menon BK, Goyal M. Imaging, Intervention, and Workflow in Acute Ischemic Stroke: The Calgary Approach. *AJNR Am J Neuroradiol* 2016; **37**(6): 978-84.
33. Goyal M, Jadhav AP, Wilson AT, Nogueira RG, Menon BK. Shifting bottlenecks in acute stroke treatment. *J Neurointerv Surg* 2016; **8**(11): 1099-100.
34. Task Force on the management of S. T. segment elevation acute myocardial infarction of the European Society of Cardiology, Steg PG, James SK, et al. ESC Guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation. *Eur Heart J* 2012; **33**(20): 2569-619.

Supplementary material



Supplementary Figure 6.1. The relationship between time from stroke onset to admission to the emergency department (ED) of the intervention center and the following time to groin puncture. Each dot represents one patient, the blue line is the fitted regression line with 95% confidence bands.



Supplementary Figure 6.2. The relationship between the median time from stroke onset to groin puncture and the treatment effect on functional outcome (adjusted common odds ratio) of different trials. Each dot represents one trial, the dashed line is the fitted regression line with 95% confidence bands.



Chapter 7

Effect of inter-hospital transfer on endovascular treatment for ischemic stroke

Stroke, 2019

Esmee Venema*, Adrien E. Groot*, Hester F. Lingsma, Wouter Hinsenveld, Kilian M. Treurniet, Vicky Chalos, Sanne M. Zinkstok, Maxim J.H.L. Mulder, Inger R. de Ridder, Henk A. Marquering, Wouter J. Schonewille, Marieke J.H. Wermer, Charles B.L.M. Majoie, Yvo B.W.E.M. Roos, Diederik W.J. Dippel, Jonathan M. Coutinho*, and Bob Roozenbeek*, for the MR CLEAN Registry Investigators

*equal contribution

Abstract

Background and purpose

To assess the effect of inter-hospital transfer on time to treatment and functional outcome after endovascular treatment (EVT) for ischemic stroke, we compared patients transferred from a primary stroke center to patients directly admitted to an intervention center in a large nationwide registry.

Methods

The MR CLEAN (Multicenter Randomized Clinical Trial of Endovascular Treatment for Acute Ischemic Stroke in the Netherlands) Registry is an ongoing, prospective, observational study in all centers that perform EVT in the Netherlands. We included adult patients with an acute anterior circulation stroke who received EVT between March 2014 to June 2016. Primary outcome was time from arrival at the first hospital to arterial groin puncture. Secondary outcomes included the 90-day modified Rankin Scale (mRS) and functional independence (mRS 0–2).

Results

In total, 821/1526 patients (54%) were transferred from a primary stroke center. Transferred patients less often had pre-stroke disability (227/800 (28%) versus 255/699 (36%), $P=0.02$) and more often received intravenous treatment with alteplase (659/819 (81%) versus 511/704 (73%), $P<0.01$). Time from first presentation to groin puncture was longer for transferred patients (164 versus 104 minutes, $P<0.01$, adjusted delay 57 minutes (95% CI 51 to 62)). Transferred patients had worse functional outcome (adjusted common odds ratio 0.75 (95% CI 0.62 to 0.90)) and less often achieved functional independence (244/720 (34%) versus 289/681 (42%), absolute risk difference -8.5% (95% CI -8.7 to -8.3)).

Conclusions

Inter-hospital transfer of patients with acute ischemic stroke is associated with delay of EVT and worse outcomes in routine clinical practice, even in a country where between-center distances are short. Direct transportation of patients potentially eligible for endovascular treatment to an intervention center may improve functional outcome.

Introduction

Endovascular treatment (EVT) has been proven effective in patients with ischemic stroke due to an intracranial large vessel occlusion (LVO) of the anterior circulation.¹ The effect of EVT is highly time-dependent and a reduction in onset-to-treatment time increases the chance of good functional outcome for the patient.^{2,3}

Studies focusing on workflow characteristics identified inter-hospital transfer as one of the main causes of treatment delay in the recent randomized controlled trials.^{4,5} Patients with ischemic stroke are often first presented at the nearest primary stroke center to provide rapid treatment with intravenous thrombolytics (IVT). If the diagnostic work-up indicates eligibility for EVT, the patient is then transferred to an intervention center with facilities for EVT ('drip-and-ship'). Analysis of data from a US registry suggested that this drip-and-ship method increases time to treatment and decreases the odds of good functional outcome in daily clinical practice.⁶ However, the effect of inter-hospital transfer might be different in a more densely populated region with short between-center distances and a well-organized acute stroke care system.

We investigated the frequency of inter-hospital transfer and the characteristics of transferred patients in a large nationwide cohort in the Netherlands: the MR CLEAN (Multicenter Randomized Clinical trial of Endovascular treatment for Acute ischemic stroke in the Netherlands) Registry. We aimed to assess the effect of inter-hospital transfer on time to treatment and functional outcome in routine clinical practice.

Material and methods

Study design

Detailed methods of the MR CLEAN Registry have been reported previously.⁷ The MR CLEAN Registry is an ongoing, prospective, observational study in all centers that perform EVT in the Netherlands. Registration started in March 2014, directly after the final inclusion in MR CLEAN. All eighteen intervention centers in the Netherlands, of which sixteen centers participated in the MR CLEAN trial, registered their patients. All data were centrally collected and checked for completeness and consistency. The imaging assessment committee assessed imaging and the adverse event committee scored the safety parameters.

The central medical ethics committee of the Erasmus MC University Medical Center, Rotterdam, The Netherlands, evaluated the study protocol and granted permission to carry out the study as a registry. Data will not be made available to other researchers as no patient approval has been obtained for sharing coded data. However, syntax and output files of statistical analyses are available from the corresponding author on reasonable request.

Study population

All consecutive patients with ischemic stroke in the anterior and posterior circulation who received arterial groin puncture in the angiosuite have been registered. For the current analysis, we used data of patients treated between March 2014-June 2016. We included patients adhering to the following criteria: age \geq 18 years; treatment in a center that participated in the MR CLEAN trial; and presence of an intracranial proximal arterial occlusion in the anterior circulation (intracranial carotid artery (ICA/ICA-T), middle cerebral artery (M1/M2) or anterior cerebral artery (A1/A2)), demonstrated by computed tomography (CT) angiography. EVT could consist of arterial catheterization followed by mechanical thrombectomy and/or thrombus aspiration, with or without delivery of a thrombolytic agent. Patients arrived either directly at the intervention center or were transferred from one of the primary stroke centers (Figure 7.1). Emergency medical services in the Netherlands identify potential stroke patients using the Face, Arm, Speech Test (FAST) and transport patients with an onset time of less than 6 hours to the closest stroke center with IVT or EVT facilities.

Definitions and outcome measures

Time of stroke onset was defined as the moment of witnessed onset of stroke symptoms or, when exact onset was unknown, the moment that the patient was last seen well. Admission outside office hours was defined as time of arrival at the intervention center on working days between 17:00 hours and 8:00 hours, during weekend days or national festive days. All transfer patients received a non-contrast CT in the primary stroke center and most transfer patients received a CT angiography before transfer to an intervention center. Imaging was not routinely repeated upon arrival at the intervention center, but only in case of a clinical indication (e.g. deterioration or substantial improvement). Imaging characteristics were assessed based on the first scan made (for transferred patients in the primary stroke center, for direct patients in the intervention center). Transfer-related travel time between each primary stroke center and the receiving intervention center was estimated using the TomTom MyDrive application (version 4.2.1.3495, available on mydrive.tomtom.com). A ratio of 0.85 was subsequently applied on each calculated time to approach the transfer-related travel time that an ambulance would actually need. This ratio is based upon measured travel times of ambulances in a previous study in the Netherlands.⁸ The door-in-door-out time in the primary stroke center was then estimated by subtracting the transfer-related travel time from the reported time from door of the primary stroke center to door of the intervention center.

Primary outcome was the time from first presentation to start of treatment, defined as the interval between arrival at the first hospital to arterial groin puncture. Secondary outcomes included: time from arrival at intervention center to groin puncture; time from stroke onset to groin puncture; the modified Rankin Scale (mRS) at 90 days after stroke; functional independence, defined as mRS score 0–2; mortality at 90 days; successful reperfusion at




Figure 7.1. Map of the Netherlands including primary stroke centers, intervention centers and population density.

the end of EVT, defined as an extended Thrombolysis in Cerebral Infarction (eTICI) score $\geq 2B$,⁹ and occurrence of symptomatic intracerebral hemorrhage. Intracerebral hemorrhage was defined as symptomatic if patients died or deteriorated neurologically (an increase of ≥ 4 points on the National Institutes of Health Stroke Scale (NIHSS) score compared with the NIHSS before worsening), and the hemorrhage was related to the clinical deterioration (according to the Heidelberg Bleeding Classification).¹⁰

Statistical analysis

We compared patients transferred from a primary stroke center with patients who were directly admitted to an intervention center. Baseline characteristics were compared using Chi-square test for categorical variables, independent samples T-test for normally distributed

continuous variables, and Mann-Whitney U test for non-normal distributed continuous variables. Time intervals were expressed as medians with interquartile range (IQR). For regression analyses, missing data were imputed using multiple imputations by chained equations (MICE) based on relevant covariates and outcome.

Linear regression analyses were used to assess the effect of transfer on time intervals. Prespecified adjustments were made for age, baseline NIHSS score, prestroke mRS score, treatment with IVT, hypertension (baseline systolic blood pressure >185 mmHg and/or diastolic blood pressure >110 mmHg), location of occlusion, Alberta Stroke Program Early Computed Tomography Score (ASPECTS) on non-contrast CT, collateral score on CT angiography, the use of general anesthesia, and admission outside office hours. The effect of transfer on functional outcome was assessed using an ordinal logistic regression analysis for the shift in mRS score at 90 days. Binary logistic regression analyses were used for functional independence, mortality, successful reperfusion, and occurrence of symptomatic intracerebral hemorrhage. These analyses were adjusted for the following prespecified variables: age, baseline NIHSS score, prestroke mRS score, history of ischemic stroke or transient ischemic attack, treatment with IVT, location of occlusion, ASPECTS, collateral score, and time from onset to first hospital.

All analyses were performed using R software (Version 3.4.4, R Foundation) with the rms package (version 5.1-2).

Results

In total, 1627 consecutive patients treated with EVT for ischemic stroke were included in the MR CLEAN Registry between March 2014 and June 2016. We excluded 101 patients for the following reasons: age below 18 years (n=2); EVT performed in a non MR CLEAN center (n=20), and occlusion of the posterior circulation (n=79). The remaining 1526 patients were included in the analyses (Figure 7.2). Of these patients, 821 (53.8%) were transferred from a primary stroke center and 705 (46.2%) were presented directly to an intervention center. The percentage of transferred patients ranged between 3.4% and 77% per intervention center.

Baseline characteristics

Demographics and baseline stroke severity were similar in both groups: median age was 71 years for transferred patients versus 70 years for directly admitted patients, 53% of all patients were men, and median NIHSS at baseline was 16 for both groups (Table 7.1). Transferred patients were more often treated with IVT (80.5% versus 72.6%, $P<0.01$) and had less often pre-stroke disability (mRS \geq 1, 28.4% versus 36.5%, $P=0.02$). Imaging characteristics were less favorable for transferred patients, with lower ASPECTS (median 8 versus 9, $P=0.01$) and more often absent or poor collaterals (collateral score 0–1, 44.1% vs.




Figure 7.2. Flowchart of patient selection. EVT, endovascular treatment.

35.2%, $P < 0.01$). Transferred patients less often had an occlusion of segment 2 of the middle cerebral artery (10.6% versus 14.4%, $P = 0.01$). Imaging was repeated in 49/821 (6%) and 39/821 transfer patients (4.8%) for non-contrast CT and CT angiography, respectively.

Treatment times

The median time from arrival at the first hospital to groin puncture was longer for transferred patients than for patients who were directly admitted to an intervention center (164 versus 104 minutes, $P < 0.01$, Table 7.2). The adjusted difference was 57 minutes (95% confidence interval (CI) 51 to 62 minutes), in favor of patients who were admitted directly (Table 7.3). After subtraction of the mean estimated transfer-related travel time (23 minutes), this time difference was 39 minutes. Transferred patients arrived slightly earlier after stroke onset at the first hospital (50 versus 55 minutes, $P < 0.01$), but time from onset to arrival at the intervention center was much longer (174 versus 55 minutes, $P < 0.01$). Median door-to-needle times for patients that received IVT were slightly longer for transferred patients (26 versus 24 minutes, $P = 0.02$). Although the interval from arrival at the intervention center to start of treatment was shorter for transferred patients (47 versus 104 minutes, $P < 0.01$, adjusted difference -58 minutes (95% CI -62 to -54)), the total time from onset to groin puncture was still substantially longer in these patients (230 versus 170 minutes, $P < 0.01$, adjusted difference 40 minutes (95% CI 31 to 48)). The median duration of the endovascular procedure was similar in both groups (63 versus 62 minutes, $P = 0.79$).

Clinical outcome

There was a significant shift towards worse functional outcome in transferred patients (adjusted common odds ratio (OR) 0.75, 95% CI 0.62 to 0.90; Figure 7.3). Transferred

Table 7.1. Baseline characteristics.

	Direct presentation n = 705	Transferred patients n = 821	P-value
Age, median (IQR)	70 (59 – 79)	71 (60 – 80)	0.45
Male sex, n (%)	371/705 (52.6)	438/821 (53.3)	0.78
Baseline NIHSS*, median (IQR)	16 (11 – 20)	16 (12 – 20)	0.19
Systolic blood pressure†, mean±SD	149±25	150±25	0.41
Treatment with IVT, n (%)	511/704 (72.6)	659/819 (80.5)	<0.01
Presentation outside office hours, n (%)	441/705 (62.6)	541/821 (65.9)	0.17
General anesthesia, n (%)	180/662 (27.2)	205/759 (27.0)	0.94
Medical history			
Ischemic stroke, n (%)	118/703 (16.8)	135/814 (16.6)	0.92
Myocardial infarction, n (%)	113/697 (16.2)	120/798 (15.0)	0.53
Diabetes mellitus, n (%)	126/700 (18.0)	136/817 (16.6)	0.49
Atrial fibrillation, n (%)	147/697 (21.1)	188/807 (23.3)	0.31
Prestroke mRS, n (%)			0.02
0	444/699 (63.5)	573/800 (71.6)	
1	97/699 (13.9)	98/800 (12.3)	
2	66/699 (9.4)	49/800 (6.1)	
≥3	92/699 (13.2)	80/800 (10.0)	
Imaging characteristics			
ASPECTS on first NCCT‡, median (IQR)	9 (7 – 10)	8 (7 – 10)	0.01
Collateral score on first CTA, n (%)			<0.01
Grade 0	42/674 (6.2)	56/743 (7.5)	
Grade 1	195/674 (28.9)	272/743 (36.6)	
Grade 2	259/674 (38.4)	288/743 (38.8)	
Grade 3	178/674 (26.4)	127/743 (17.1)	
Location of occlusion, n (%)			0.01
ICA	39/693 (5.6)	46/765 (6.0)	
ICA-T	142/693 (20.5)	180/765 (23.5)	
Proximal M1	161/693 (23.2)	210/765 (27.5)	
Distal M1	231/693 (33.3)	240/765 (31.4)	
M2	100/693 (14.4)	81/765 (10.6)	
Other§	20/693 (2.9)	8/765 (1.0)	

Number of missing values: *30, †43, ‡67.

§Occlusion in segment 1 or 2 of the anterior cerebral artery (A1: n=3; A2: n=3), segment 3 of the middle cerebral artery (M3: n=9), or no occlusion visible (n=13) on CTA after adjudication by the imaging assessment committee. ASPECTS, Alberta Stroke Program Early CT Score; CTA, computed tomography angiography; M1, segment 1 of the middle cerebral artery; M2, segment 2 of the middle cerebral artery; mRS, modified Rankin Scale; NCCT, non-contrast computed tomography; NIHSS, National Institutes of Health Stroke Scale; ICA, internal carotid artery; ICA-T, internal carotid artery tandem; IVT, intravenous thrombolytics.

patients less often achieved functional independence when compared to directly admitted patients (mRS 0-2, 33.9% versus 42.4%, absolute risk difference -8.5% (95% CI -8.7 to -8.3)), also after adjustment (adjusted OR 0.69 (95% CI 0.54 to 0.89); Table 7.4). Mortality rates were not significantly different between the two groups (adjusted OR 1.27 (95% CI 0.97 to 1.66)). There were no significant differences in successful reperfusion rate (adjusted OR 1.13 (95% CI 0.91 to 1.39)) and occurrence of symptomatic intracerebral hemorrhage (adjusted OR 0.87 (95% CI 0.56 to 1.36)).

Table 7.2. Treatment times, presented as medians with interquartile range.

	Direct presentation n = 705	Transferred patients n = 821	P-value	Missing values n (%)
Onset to door first hospital	55 (37 - 105)	50 (35 - 80)	<0.01	197/1526 (12.9)
Onset to door intervention center	55 (37 - 105)	174 (139 - 220)	<0.01	30/1526 (2.0)
Onset to IVT	82 (63 - 125)	79 (60 - 115)	<0.01	218/1173 (18.6)
Door to needle	24 (18 - 34)	26 (20 - 36)	0.02	215/1173 (18.3)
Door to door*	NA	112 (91 - 140)	N/A	196/822 (23.5)
Primary stroke center to intervention center (estimated)	NA	21 (15 - 31)	NA	3/821 (0.4)
Door in door out† (estimated)	NA	88 (68 - 117)	NA	197/821 (24.0)
Door first hospital to groin puncture	104 (80 - 135)	164 (135 - 198)	<0.01	197/1526 (12.9)
Door intervention center to groin puncture	104 (80 - 135)	47 (31 - 70)	<0.01	30/1526 (2.0)
Onset to groin puncture	170 (135 - 246)	230 (190 - 277)	<0.01	0
Duration of procedure	62 (40 - 90)	63 (40 - 87)	0.79	163/1526 (10.7)
Onset to reperfusion	238 (185 - 314)	288 (244 - 343)	<0.01	91/1526 (6.0)

*Door primary stroke center to door intervention center. †Door primary stroke center to door intervention center, minus the estimated travel time from primary stroke center to intervention center. IVT, intravenous thrombolytics; NA, not applicable.

Table 7.3. Outcome measures: time intervals (in minutes).

	Direct presentation n = 705	Transferred patients n = 821	Unadjusted beta (95% CI)	Adjusted beta* (95% CI)
Door first hospital to groin puncture	104 (80 – 135)	164 (135 – 198)	55 (50 to 61)	57 (51 to 62)
Door intervention center to groin puncture	104 (80 – 135)	47 (31 – 70)	-59 (-63 to -55)	-58 (-62 to -54)
Onset to groin puncture	170 (135 – 246)	230 (190 – 277)	36 (27 to 44)	40 (31 to 48)

*Adjusted for age, baseline National Institute of Health Stroke Severity score, prestroke modified Rankin Scale, treatment with intravenous thrombolytics, hypertension (baseline systolic blood pressure >185 mmHg and/or diastolic blood pressure >110 mmHg), location of occlusion, Alberta Stroke Program Early Computed Tomography Score, collateral score, use of general anesthesia and admission outside office hours.

Table 7.4. Outcome measures: clinical outcome.

	Direct presentation n = 705	Transferred patients n = 821	Unadjusted OR (95% CI)	Adjusted OR* (95% CI)
Functional independence at 90 days (mRS 0-2)	289/681 (42.4)	244/720 (33.9)	0.71 (0.58 to 0.88)	0.69 (0.54 to 0.89)
Mortality at 90 days	180/681 (26.4)	227/720 (31.5)	1.24 (0.99 to 1.55)	1.27 (0.97 to 1.66)
Successful reperfusion (eTICI \geq 2B)	395/698 (56.6)	473/806 (58.7)	1.10 (0.90 to 1.35)	1.13 (0.91 to 1.39)
Symptomatic ICH	45/705 (6.4)	44/821 (5.4)	0.83 (0.54 to 1.27)	0.87 (0.56 to 1.36)

*Adjusted for age, baseline National Institute of Health Stroke Severity score, prestroke modified Rankin Scale, history of ischemic stroke, treatment with intravenous thrombolytics, location of occlusion, Alberta Stroke Program Early Computed Tomography Score, collateral score and time from onset to first hospital. eTICI, extended Thrombolysis In Cerebral Infarction scale; ICH, intracerebral hemorrhage; mRS, modified Rankin Scale.

Discussion

In this large, nationwide cohort study, more than half of the patients treated with EVT were transferred from a primary stroke center. These patients had significant longer time intervals between first presentation and start of treatment, between symptom onset and start of treatment, and a lower chance of achieving good functional outcome. Even when we accounted for the travel time required to transfer a patient to an intervention center, start of treatment was still delayed by half an hour in transferred patients compared to patients who were directly admitted to an intervention center.




Figure 7.3. Functional outcome measured with the modified Rankin Scale score at 90 days, for patients presented directly in intervention center versus transferred from a primary stroke center (125 missing values). Transferred patients had worse functional outcomes than directly presented patients (adjusted common odds ratio 0.75 (95% CI 0.62 to 0.90)).

From earlier studies, we know that treatment delay translates to a decreased chance of achieving a favorable outcome.² However, because of selection bias and time delay associated with informed consent procedures, data from randomized trials may not always reflect daily practice. The STRATIS (Systematic Evaluation of Patients Treated With Neurothrombectomy Devices for Acute Ischemic Stroke) Registry included 984 patients from a large region in the United States of America after the implementation of EVT as standard care.⁶ Similar to our study, the authors found that transferred patients had longer treatment times and worse functional outcomes when compared to direct admitted patients. However, their time analyses included only patients with successful reperfusion, which is not a fair representation of routine clinical practice. Also, distances between hospitals in rural areas of the USA are much larger than in our region, and time delays in primary hospitals – for instance because of initiation of IVT – are generally longer in the USA than in the Netherlands.^{8,11,12} Results from an earlier small study in a region with short distances, including only one primary stroke center and one intervention center, showed that treatment times were longer for transferred patients, and the difference between the two groups for functional outcome was similar to our study.¹³

The Netherlands is a small, densely populated country, with 82 hospitals that provide 24/7 acute stroke care. Approximately 99.8% of the Dutch population has the ability to reach an emergency department within 45 minutes.¹⁴ Our country has short between-center distances. This is emphasized by the fact that the mean transfer time was only 23 minutes, indicating that these results probably reflect the acute stroke care logistics in other highly populated regions. Although the observed in-hospital workflow times in this study were almost an hour shorter than in the MR CLEAN trial in the Netherlands,¹⁵ there is still room

for improvement. It has been suggested that increased experience, reflected in higher volumes of patients treated with EVT, leads to shorter treatment times and, subsequently, improved functional outcomes.¹⁶ Workflow times in our cohort do gradually decrease over time from 2014 until 2016. A decline in the median door-to-groin time (with the door of the first hospital as start) was observed for both transfer patients (187 to 149 minutes) and direct patients (123 to 97 minutes). However, the times achieved in our cohort are still longer when compared to other studies reporting workflow times. Additional interventions to optimize patient transfer management (eg, holding the initial ambulance primary stroke center until the decision of EVT eligibility has been made, streamlining transfer protocols, improving cloud-based image sharing, and transporting transfer patients directly to the angio-suite) might be useful to further reduce transfer-related delay in the future.¹⁷⁻²⁰

Pre-hospital transportation of stroke patients suspected of LVO directly to an intervention center decreases time to EVT, but the potential harm of delaying IVT should be taken into account as well. We know, however, that the chance of achieving recanalization with IVT prior to thrombectomy in patients with LVO is low.²¹ Therefore, we should focus on finding an algorithm to predict LVO early, so that emergency medical services can present these patients directly to the intervention center.^{22,23} Implementing such a triage protocol has already shown to be feasible and seems to improve treatment times.²⁴ One ongoing randomized trial in Catalonia, Spain, is evaluating the effect of the mothership vs. the drip-and-ship strategy on functional outcome, among patients with a high likelihood of having a LVO identified with the Rapid Arterial Occlusion Evaluation (RACE) scale (Direct Transfer to an Endovascular Center Compared to Transfer to the Closest Stroke Center in Acute Stroke Patients With Suspected Large Vessel Occlusion (RACECAT), URL: www.clinicaltrials.gov. Unique identifier: NCT02795962). While this trial will provide class I evidence on whether a mothership model improves functional outcome, it will be challenging to translate its results to regions with other geographical and demographical conditions. Modeling studies showed that differences in transportation times and treatment times affect the optimal transportation strategy and that triage protocols should therefore be based on regional characteristics and individual likelihood of LVO.²⁵⁻²⁸

There were a number of baseline imbalances between the two groups. First, transferred patients more often received IVT. One of the reasons that could explain this observation is that patients who passed the time window for IVT are probably more often directly transported to an intervention center. Second, transferred patients more often had proximal occlusions (ie, ICA, ICA-T, and proximal M1) when compared to directly admitted patients. It could be that physicians hesitate to transport a patient with a more distal occlusion towards the intervention center, as earlier studies showed that these occlusions generally respond well to IVT.²⁹ Another observation that might be explained by a selection mechanism in the intervention center is that transferred patients less often had prestroke disability. This suggests that patients with favorable characteristics are more likely to be transferred

to an intervention center. However, imaging characteristics, assessed at the hospital of first presentation, were slightly less favorable for transferred patients. After adjustment for these baseline imbalances, we still found an 8.5% decrease in the chance of achieving functional independence after inter-hospital transfer.

Strengths of our study include that the analyses were performed with data from one of the largest cohorts of consecutive patients treated with EVT, with individual patient data from all stroke intervention centers in the Netherlands. It strongly reflects daily clinical practice, including a relatively large number of transferred patients. A number of limitations also warrant comment. Despite the fact that the MR CLEAN Registry is a nationwide study with consecutive inclusion of patients, our study is not completely free from a risk of selection bias. Patients with an LVO who did not receive EVT, for example due to clinical improvement on arrival, were not included in the study. Some patients who were initially admitted to a primary stroke center may not have been transferred to an intervention center because the time window would have been passed by the moment they would arrive in the intervention center or because there was uncertainty about treatment eligibility in case of an M2 occlusion. These patients could have been treated if they had been transported directly to the intervention center. Since the MR CLEAN Registry only contains data of patients who actually received EVT, we cannot estimate the size and impact of this patient population. The negative effect of a drip-and-ship strategy might therefore be larger in real life. Another limitation of our study is that we had no recorded door in door out times available to distinguish between in-hospital delay in the primary stroke center and delay due to the actual travel time between the primary stroke center and intervention center. We estimated the average inter-hospital travel times using the TomTom MyDrive application and then calculated the estimated door in door out times. Also, we used multiple imputation for time from first presentation to groin puncture, since time of arrival in the first hospital was missing for 13% of the patients. The difference between transferred patients and directly admitted patients in time from first presentation to groin puncture was comparable with the difference in time from onset to groin puncture, a variable that was available for all patients. We therefore believe that it has little impact on the validity of our results. Last, even though we adjusted for differences in prognostic factors, residual confounding might still influence our results.

Conclusions

In conclusion, these results from a large national cohort show that inter-hospital transfer of ischemic stroke patients eligible for EVT is associated with longer treatment times and worse functional outcome in clinical practice, even in a country where between-center distances are short. Direct transportation of EVT candidates to an intervention center might reduce treatment delay and thereby may improve functional outcome.

References

1. Goyal M, Menon BK, van Zwam WH, et al. Endovascular thrombectomy after large-vessel ischaemic stroke: a meta-analysis of individual patient data from five randomised trials. *Lancet* 2016; **387**(10029): 1723-31.
2. Saver JL, Goyal M, van der Lugt A, et al. Time to Treatment With Endovascular Thrombectomy and Outcomes From Ischemic Stroke: A Meta-analysis. *JAMA* 2016; **316**(12): 1279-88.
3. Mulder M, Jansen IGH, Goldhoorn RB, et al. Time to Endovascular Treatment and Outcome in Acute Ischemic Stroke: MR CLEAN Registry Results. *Circulation* 2018; **138**(3): 232-40.
4. Menon BK, Sajobi TT, Zhang Y, et al. Analysis of Workflow and Time to Treatment on Thrombectomy Outcome in the Endovascular Treatment for Small Core and Proximal Occlusion Ischemic Stroke (ESCAPE) Randomized, Controlled Trial. *Circulation* 2016; **133**(23): 2279-86.
5. Dijkland SA, Voormolen DC, Venema E, et al. Utility-Weighted Modified Rankin Scale as Primary Outcome in Stroke Trials: A Simulation Study. *Stroke* 2018; **49**(4): 965-71.
6. Froehler MT, Saver JL, Zaidat OO, et al. Interhospital Transfer Before Thrombectomy Is Associated With Delayed Treatment and Worse Outcome in the STRATIS Registry (Systematic Evaluation of Patients Treated With Neurothrombectomy Devices for Acute Ischemic Stroke). *Circulation* 2017; **136**(24): 2311-21.
7. Jansen IGH, Mulder M, Goldhoorn RB, investigators MCR. Endovascular treatment for acute ischaemic stroke in routine clinical practice: prospective, observational cohort study (MR CLEAN Registry). *BMJ* 2018; **360**: k949.
8. Groot AE, van Schaik IN, Visser MC, et al. Association between i.v. thrombolysis volume and door-to-needle times in acute ischemic stroke. *J Neurol* 2016; **263**(4): 807-13.
9. Goyal M, Fargen KM, Turk AS, et al. 2C or not 2C: defining an improved revascularization grading scale and the need for standardization of angiography outcomes in stroke trials. *J Neurointerv Surg* 2014; **6**(2): 83-6.
10. von Kummer R, Broderick JP, Campbell BC, et al. The Heidelberg Bleeding Classification: Classification of Bleeding Events After Ischemic Stroke and Reperfusion Therapy. *Stroke* 2015; **46**(10): 2981-6.
11. Bray BD, Campbell J, Cloud GC, et al. Bigger, faster? Associations between hospital thrombolysis volume and speed of thrombolysis administration in acute ischemic stroke. *Stroke* 2013; **44**(11): 3129-35.
12. Fonarow GC, Smith EE, Saver JL, et al. Timeliness of tissue-type plasminogen activator therapy in acute ischemic stroke: patient characteristics, hospital factors, and outcomes associated with door-to-needle times within 60 minutes. *Circulation* 2011; **123**(7): 750-8.
13. Gerschenfeld G, Muresan IP, Blanc R, et al. Two Paradigms for Endovascular Thrombectomy After Intravenous Thrombolysis for Acute Ischemic Stroke. *JAMA Neurol* 2017; **74**(5): 549-56.
14. Kommer R, de Bruin-Kooistra M, Deuning C. Aanbod En Bereikbaarheid van de Spoedeisende Ziekenhuiszorg in Nederland 2017; Analyse Gevoelige Ziekenhuizen. 2017.
15. Franssen PS, Berkhemer OA, Lingsma HF, et al. Time to Reperfusion and Treatment Effect for Acute Ischemic Stroke: A Randomized Clinical Trial. *JAMA Neurol* 2016; **73**(2): 190-6.
16. Gupta R, Horev A, Nguyen T, et al. Higher volume endovascular stroke centers have faster times to treatment, higher reperfusion rates and higher rates of good clinical outcomes. *J Neurointerv Surg* 2013; **5**(4): 294-7.
17. Ng FC, Low E, Andrew E, et al. Deconstruction of Interhospital Transfer Workflow in Large Vessel Occlusion: Real-World Data in the Thrombectomy Era. *Stroke* 2017; **48**(7): 1976-9.
18. McTaggart RA, Yaghi S, Cutting SM, et al. Association of a Primary Stroke Center Protocol for Suspected Stroke by Large-Vessel Occlusion With Efficiency of Care and Patient Outcomes. *JAMA Neurol* 2017; **74**(7): 793-800.
19. Jadhav AP, Kenmuir CL, Aghaebrahim A, et al. Interfacility Transfer Directly to the Neuroangiography Suite in Acute Ischemic Stroke Patients Undergoing Thrombectomy. *Stroke* 2017; **48**(7): 1884-9.

20. Kansagra AP, Wallace AN, Curfman DR, et al. Streamlined triage and transfer protocols improve door-to-puncture time for endovascular thrombectomy in acute ischemic stroke. *Clin Neurol Neurosurg* 2018; **166**: 71-5.
21. Bhatia R, Hill MD, Shobha N, et al. Low rates of acute recanalization with intravenous recombinant tissue plasminogen activator in ischemic stroke: real-world experience and a call for action. *Stroke* 2010; **41**(10): 2254-8.
22. Heldner MR, Hsieh K, Broeg-Morvay A, et al. Clinical prediction of large vessel occlusion in anterior circulation stroke: mission impossible? *J Neurol* 2016; **263**(8): 1633-40.
23. Smith WS, Lev MH, English JD, et al. Significance of large vessel intracranial occlusion causing acute ischemic stroke and TIA. *Stroke* 2009; **40**(12): 3834-40.
24. Zaidi SF, Shawver J, Espinosa Morales A, et al. Stroke care: initial data from a county-based bypass protocol for patients with acute stroke. *J Neurointerv Surg* 2017; **9**(7): 631-5.
25. Holodinsky JK, Williamson TS, Demchuk AM, et al. Modeling Stroke Patient Transport for All Patients With Suspected Large-Vessel Occlusion. *JAMA Neurol* 2018; **75**(12): 1477-86.
26. Schlemm E, Ebinger M, Nolte CH, Endres M, Schlemm L. Optimal Transport Destination for Ischemic Stroke Patients With Unknown Vessel Status: Use of Prehospital Triage Scores. *Stroke* 2017; **48**(8): 2184-91.
27. Benoit JL, Khatri P, Adeoye OM, et al. Prehospital Triage of Acute Ischemic Stroke Patients to an Intravenous tPA-Ready versus Endovascular-Ready Hospital: A Decision Analysis. *Prehosp Emerg Care* 2018; **22**(6): 722-33.
28. Parikh NS, Chatterjee A, Diaz I, et al. Modeling the Impact of Interhospital Transfer Network Design on Stroke Outcomes in a Large City. *Stroke* 2018; **49**(2): 370-6.
29. Seners P, Turc G, Maier B, Mas JL, Oppenheim C, Baron JC. Incidence and Predictors of Early Recanalization After Intravenous Thrombolysis: A Systematic Review and Meta-Analysis. *Stroke* 2016; **47**(9): 2409-12.



Chapter 7.1

Response by Venema et al to letter regarding article,
"Effect of inter-hospital transfer on endovascular treatment
for ischemic stroke"

Stroke, 2019

In Response:

We read with interest the letter by Drs Seners and Baron regarding our recent publication about the effect of inter-hospital transfer on time to treatment and functional outcome of stroke patients undergoing endovascular treatment (EVT) in the MR CLEAN Registry.¹

We agree that our results apply only to those patients with a large vessel occlusion (LVO) who effectively received groin puncture in an intervention center. As we stated in our discussion, the MR CLEAN (Multicenter Randomized Clinical Trial of Endovascular Treatment for Acute Ischemic Stroke) Registry does not contain data of patients with an LVO who did not undergo EVT. Due to this limitation, it is impossible to fully estimate the effect of first presentation in a primary stroke center versus direct transportation to an intervention center for the total LVO stroke population. A subset of the patients presenting in a primary stroke center with an LVO might not receive EVT due to clinical improvement on arrival at the intervention center. Assuming that the thrombus has resolved due to prior IV treatment with alteplase, these patients are indeed likely to have a more favorable outcome, and excluding them will lead to an underestimation of good functional outcome in the total transfer population. Direct transportation to an intervention center could have a negative effect on functional outcome of these patients because it delays IV treatment. On the other hand, there might also be patients in the transfer group that did not receive EVT because the remaining time window did not allow for transportation to the intervention center or because there was uncertainty about treatment eligibility (eg, in case of an M2 occlusion). These patients could potentially have been treated with EVT if they had been transported directly to an intervention center. Excluding these patients will, therefore, lead to an overestimation of good functional outcome in the transfer population. In other words, due to the limitations inherent to our study population, our results could both overestimate and underestimate the true difference in clinical outcome of patients between the drip-and-ship and mothership models.

Drs Seners and Baron refer to a study of Gerschenfeld et al,² in which all patients that were transferred for EVT - including those who eventually did not receive EVT - were included. However, patients who received only IV alteplase and were not transferred to an intervention center, were not included in the analysis. The reasons for withholding EVT in these 144 patients in the drip-and-ship paradigm and 122 patients in the mothership paradigm are not clearly described. Also, due to the small sample size, this study was likely underpowered to detect a difference in clinical outcome in the first place.

In conclusion, our results suggest that, if treated with EVT, patients directly transported to an intervention center have a shorter time to treatment and better functional outcome compared to patients transferred from a primary stroke center. Caution is warranted when generalizing these results to the total LVO stroke population and, similar to our colleagues, we look forward to the results of ongoing randomized trials that directly compare the drip-

and-ship versus mothership approach. Still, even the results of these trials cannot be directly translated to regions with other geographical and organizational characteristics, and modeling studies will therefore be required.³

References

1. Venema E, Groot AE, Lingsma HF, Hinsenveld W, Treurniet KM, Chalos V, et al. Effect of interhospital transfer on endovascular treatment for acute ischemic stroke. *Stroke* 2019; **50**(4): 923-30.
2. Gerschenfeld G, Muresan IP, Blanc R, Obadia M, Abrivard M, Piotin M, et al. Two paradigms for endovascular thrombectomy after intravenous thrombolysis for acute ischemic stroke. *JAMA Neurol* 2017; **74**(5): 549-56.
3. Venema E, Lingsma HF, Chalos V, Mulder MJHL, Lahr MMH, van der Lugt A, et al. Personalized Prehospital Triage in Acute Ischemic Stroke. *Stroke* 2019; **50**(2): 313-20.

This letter was written in response to:

Seners P, Baron JC. Letter by Seners and Baron Regarding Article, "Effect of Interhospital Transfer on Endovascular Treatment for Acute Ischemic Stroke". *Stroke* 2019; **50**(9) :e259.



Chapter 8

Effect of workflow improvements in endovascular stroke treatment: a systematic review and meta-analysis

Stroke, 2019

Abstract

Background and purpose

Rapid initiation of endovascular stroke treatment is associated with better clinical outcome. The effect of specific improvements is not well known. We performed a systematic review and meta-analysis on the effectiveness of specific workflow improvements on time to treatment and outcome.

Methods

A random effects meta-analysis was used to evaluate the difference in mean time to treatment between intervention group and control group. Secondary outcomes included good functional outcome at 90 days (modified Rankin Scale score 0–2).

Results

Fifty-one studies (3 randomized controlled trials, 13 pre-post intervention studies, and 35 observational studies) with in total 8,467 patients were included. Most frequently reported workflow intervention types concerned anesthetic management (n=26), in-hospital patient transfer management (n=14), and prehospital management (n=11). Patients in the intervention group had shorter time to treatment intervals (weighted mean difference 26 minutes; 95% CI 19 to 33; $P<0.001$) compared with controls. Subgroup meta-analysis of intervention types also showed a shorter time to treatment in the intervention group: a mean difference of 12 minutes (95% CI 6 to 17; $P<0.001$) for anesthetic management, 37 minutes (95% CI 22 to 52; $P<0.001$) for prehospital management, 41 minutes (95% CI 27 to 54; $P<0.001$) for in-hospital patient transfer management, 47 minutes (95% CI 28 to 67; $P<0.001$) for teamwork, and 64 minutes (95% CI 24 to 104; $P=0.002$) for feedback. The mean difference in time to treatment of studies with multiple interventions implemented simultaneously was 50 minutes (95% CI 31 to 69; $P<0.001$) in favor of the intervention group. Patients in the intervention group had increased likelihood of favorable outcome (risk ratio 1.39; 95% CI 1.15 to 1.66; $P<0.001$).

Conclusions

Interventions in the workflow of endovascular stroke treatment lead to a significant reduction in time to treatment and results in an increased likelihood of favorable outcome. Acute stroke care should be reorganized by making use of the examples of workflow interventions described in this review to ensure the best medical care for stroke patients.

Introduction

Multiple trials have shown the benefit of endovascular recanalization therapy in selected stroke patients.¹⁻³ Earlier treatment is associated with better functional outcome.⁴ The time from symptom onset to treatment is influenced by prehospital and in-hospital processes. Healthcare systems are being reorganized to offer stroke patients rapid and effective medical care. Stroke services had already changed their workflow since treatment with intravenous alteplase for selected stroke patients was proven effective.⁵ Implementation of new strategies to improve the workflow process for treatment with intravenous alteplase has led to a significant reduction of in-hospital delay.⁶

Providing an optimal diagnostic process and rapid endovascular stroke treatment requires close collaboration of the emergency medical service, emergency department team, stroke team, neurointerventional team, and anesthesia team. Diagnostic imaging and endovascular treatment facilities should be available in very little time. Several strategies to reduce the time to endovascular stroke treatment have been proposed.⁷⁻⁹ However, the effect of individual and combined strategies on reducing time to treatment is unclear. We performed a systematic review and meta-analysis on the effectiveness of specific workflow improvement interventions for rapid delivery of endovascular stroke treatment.

Methods

This systematic review and meta-analysis was performed according to the Preferred Reporting Items for Systematic Review and Meta-Analysis (PRISMA) guidelines.¹⁰ All data and supporting materials are available within the article and its Supplementary material.

Search strategy

Medline, EMBASE, Cochrane Central, and Web of Science were searched for studies that evaluated the effect of one or more workflow interventions on time to endovascular stroke treatment, from database inception to November 14th, 2017. Google Scholar and Google were searched at November 14th, 2017, and the first 200 hits were included. We developed a broad search strategy consisting of a combination of the 2 main topics of this study: endovascular stroke treatment and workflow intervention. The complete search strategy is available in the Supplementary material. We restricted our search to studies published in English and excluded conference abstracts.

Eligibility criteria

Studies were included if ≥ 1 (prehospital or in-hospital) interventions in the workflow of endovascular stroke treatment were assessed and effect on time to treatment intervals was reported. Endovascular stroke treatment was defined as mechanical thrombectomy or intra-arterial fibrinolysis in an acute stroke patient with an intracranial large vessel occlusion.

Interventions only aimed at the duration of the endovascular treatment itself, for example type of mechanical thrombectomy device used, were excluded. Interventions intended only to increase the accuracy of patient selection, for example the introduction of a new imaging protocol, were also excluded. Studies were included in the systematic review when time to endovascular treatment was reported, from symptom onset to start treatment, or any time window between symptom onset and start treatment. Randomized and non-randomized controlled trials and pre-post intervention studies were included. Observational studies, or post-hoc analyses of observational data in trials were only included when a control group was reported. Reviews, editorials, and guidelines were excluded. Two authors (Drs Janssen and Venema) independently assessed the eligibility of all retrieved studies. Title and abstracts were first screened to identify potentially eligible articles, and then full texts were read to confirm inclusion. Reference lists of identified eligible papers and review papers were scanned for additional relevant studies.

Risk of bias assessment

The risk of bias of each included study was assessed against the following key criteria: random sequence generation; allocation concealment; blinding of participants, personnel and outcomes; incomplete outcome data; and selective outcome reporting; in accordance with the methods recommended by the Cochrane Library.¹¹ The following judgements were used: low risk, high risk, or unclear risk of bias (either lack of information or uncertainty on the potential for bias). Summary of risk of bias per key criterion was provided for all included articles separately.

Data extraction and outcome variables

Data was extracted from published reports by 2 authors (Drs Janssen and Venema). Workflow interventions were described and divided into six predefined categories: (A) anesthetic management, (B) prehospital management, (C) in-hospital patient transfer management, (D) teamwork, (E) feedback, and (F) other workflow interventions. Other collected data on study characteristics included study design, study period, stroke type (anterior or posterior circulation stroke, or both), and sample size.

The primary outcome measure in this study was the difference in time to treatment between the intervention group and control group. Other study outcomes were good functional outcome, defined as modified Rankin Scale (mRS) score 0–2 at 90 days after endovascular treatment, symptomatic intracranial hemorrhage, and mortality.

Statistical analysis

Mean time to treatment with standard deviation (SD) for the intervention group and control group was retrieved from each included study. When mean values with SD were not available in the publication nor obtained from the authors of the original publication, we used reported median time to treatment with interquartile range (IQR) to estimate the sample mean and SD using the method described by Wan et al.¹² The absolute difference of mean

time to treatment with 95% confidence intervals (CI) was calculated for each study using a two-sample t-test.

Studies were included in the meta-analysis when mean time to treatment with SD or median time to treatment with IQR was available for both groups. Weighted difference in mean time to treatment with 95% CI was calculated using a random-effects inverse variance model, with the estimate of heterogeneity being taken from the Mantel-Haenszel model. Subgroup analysis of the difference in mean time to treatment was performed for the predefined workflow intervention categories A to E and for studies implementing multiple interventions simultaneously.

Data on binary outcomes (good functional outcome, symptomatic intracranial hemorrhage, and mortality) was pooled using random-effect meta-analysis and expressed as risk ratios. Publication bias was assessed by constructing a funnel plot. All statistical analyses were conducted with Stata, version 15 (Statacorp LLC, College Station, TX).

Results

Our literature search identified 4,127 potentially relevant unique articles; 211 articles were retained for full-text review (Figure 8.1). A total of 51 studies met the inclusion criteria and were included in the qualitative synthesis.^{2,13-62} We contacted authors from 31 of 51 studies with requests for additional data necessary for our meta-analysis. These additional data were provided for 17 of 31 studies. The sample mean difference in time to treatment with SD could be estimated from published data from 8 of 31 studies. After exclusion of the remaining 6 studies because of lack of sufficient data, a total of 45 studies was included in the meta-analysis on effect of workflow interventions on the time to treatment.

Fifty-one studies with 8,467 patients (4,037 intervention group and 4,430 control group) reported the effect of 25 different workflow interventions on the time to endovascular treatment (Table 8.1 and 8.2). Two studies reported the effect on time to treatment of two interventions separately.^{50,55} Most frequently reported workflow intervention types concerned anesthetic management (n=26), in-hospital patient transfer management (n=14), and prehospital management (n=11). Ten studies reported the effect on time to treatment of multiple interventions implemented simultaneously. Time to treatment was shorter in the intervention group in 48 of 53 interventions (91%) reported in the 51 included studies. Included studies differed in study design, with 3 studies randomizing patients for the workflow intervention of interest in our study, 13 pre-post intervention studies, and the remaining 35 studies reporting observational data mostly from hospital stroke registries or randomized controlled trials investigating the effect of endovascular stroke treatment versus conservative treatment. Data collection was performed retrospectively in 34 studies, and 16 studies collected data from ≥ 1 center. Risk of bias assessment is reported in Supplementary Table 8.1.




Figure 8.1. Flowchart of included and excluded articles, following the Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA) guidelines.

Random effects meta-analysis of 45 studies (with 47 interventions), including 7,482 patients (3,480 intervention group and 4,002 control group) showed a difference in mean time to treatment of 26 minutes (95% CI 19 to 32; $P < 0.001$) in favor of the intervention group (Figure 8.2). I^2 value was 85.4%, and Chi-squared value was 314.87 (degrees of freedom 46; $P < 0.001$), indicating considerable heterogeneity between studies.

Table 8.1. Studies included in systematic review.

Author	Country	Study Design	Study Period*	Anterior, posterior circulation stroke or both	Intervention group (n)	Control group (n)	Type of intervention
Abou-Chebl et al ¹³	USA	Retrospective cohort study; multicenter	2005–2009	Anterior	552	428	A
Abou-Chebl et al ¹⁴	USA	Post hoc analysis retrospective NASA Registry; multicenter	2012–2013	Both	68	159	A
Abou-Chebl et al ¹⁵	Canada, Europe, USA	Post hoc analysis IMS III trial; multicenter	2006–2012	Both	269	147	A
Aghaebrahim et al ¹⁶	USA	Prospective pre-post study; single center	2012–2013/ 2013–2014	Both	108	178	B1, C1–3, D1–2, E1, F1
Alotaibi et al ¹⁷	Canada	Retrospective pre-post study; single center	2011–2014/ 2014–2016	Both	28	17	E2
Van den Berg et al ¹⁸	The Netherlands	Retrospective cohort study; multicenter	2002–2010	Anterior	278	70	A
Berkhemer et al ¹⁹	The Netherlands	Post-hoc analysis MR CLEAN trial; multicenter	2010–2014	Anterior	137	79	A
Bracard et al ²	France	Post-hoc analysis THRACE trial; multicenter	2010–2014	Both	74	69	A
Cerejo et al ²⁰	USA	Retrospective cohort study; single center	2014	Anterior	5	5	B2
Davis et al ²¹	Canada	Retrospective cohort study; single center	2003–2009	Both	37	39	A
Eesa et al ²²	Canada	Retrospective cohort study; single center	2005–2009	Both	71	30	A
Frei et al ²³	USA	Retrospective pre-post study; single center	2012–2013/ 2013–2015	Both	267	113	B1, D2–4, F1–3
Goyal et al ²⁴	Canada, Europe, USA	Post-hoc analysis IMS III trial; multicenter	2006–2012	Both	17	64	B3
Goyal et al ²⁵	Europe, USA	Post-hoc analysis SWIFT PRIME trial; multicenter	2012–2014	Anterior	61	35	A
Hassan et al ²⁶	USA	Retrospective cohort study; multicenter	2006–2010	Both	83	53	A
Henden et al ²⁷	Sweden	Randomized controlled trial; single center	2013–2016	Anterior	45	45	A
Herrmann et al ²⁸	Germany	Pre-post study; retrospective data pre-intervention, prospective data post-intervention; single center	2006–2009/ 2009–2010	Both	23	48	F4

Table 8.1. Continued

Author	Country	Study Design	Study Period*	Anterior, posterior circulation stroke or both	Intervention group (n)	Control group (n)	Type of intervention
Jadhav et al ²⁹	USA	Retrospective cohort study; single center	2013–2016	Both	111	150	C2
Jagani et al ³⁰	USA	Retrospective cohort study; single center	2008–2015	Both	61	38	A
Janssen et al ³¹	Germany	Retrospective cohort study; single center	2012–2014	Anterior	31	53	A
Jeon et al ³²	Korea	Retrospective pre-post study; single center	2014–2016/ 2016	Not specified	19	93	B1, C3, D2, E1, F1–2
John et al ³³	USA	Retrospective cohort study; single center	2008–2012	Anterior	99	91	A
Jumaa et al ³⁴	USA	Retrospective cohort study; single center	2006–2009	Anterior	73	53	A
Just et al ³⁵	Canada	Retrospective cohort study; single center	2000–2013	Both	67	42	A
Kamper et al ³⁶	Germany	Retrospective pre-post study; single center	2002–2006/ 2007–2010	Posterior	20	18	F5
Koge et al ³⁷	Japan	Retrospective pre-post study; single center	2008–2014/ 2014–2016	Not specified	23	19	D3–4, E1
Komatsubara et al ³⁸	Japan	Pre-post study; retrospective or prospective data collection not specified; single center	2012–2014/ 2014–2015	Both	14	14	E1, F1, F6
Li et al ³⁹	USA	Retrospective cohort study; single center	2006–2012	Both	74	35	A
Liang et al ⁴⁰	USA	Retrospective cohort study; single center	2015–2016	Not specified	22	17	B4
Mascitelli et al ⁴¹	USA	Retrospective pre-post study; single center	2014/ 2014–2015	Both	29	27	B1, E1–2, F1
McTaggart et al ⁴²	USA	Retrospective cohort study; multicenter	2015–2016	Anterior	22	48	B4–5, C2, D2
Mehta et al ⁴³	USA	Pre-post study; retrospective data pre-intervention, prospective data post-intervention; single center	2007–2011/ 2011–2013	Anterior	51	93	C3, D2–4
Menon et al ⁴⁴	Canada, Ireland, South Korea, UK, USA	Prespecified secondary analysis ESCAPE trial; multicenter	2013–2014	Anterior	136	15	A

Author	Country	Study Design	Study Period*	Anterior, posterior circulation stroke or both	Intervention group (n)	Control group (n)	Type of intervention
Miley et al ⁴⁵	USA	Retrospective cohort study; multicenter	2005–2008	Both	52	39	A
Mundiyanapurath et al ⁴⁶	Germany	Prospective cohort study; single center	2013–2014	Both	15	29	A
Nichols et al ⁴⁷	USA	Post-hoc analysis IMS II trial; multicenter	2003–2006	Anterior	40	17	A
Pedragosa et al ⁴⁸	Spain	Prospective cohort study; multicenter	2008–2010	Not specified	25	20	B6
Pfaff et al ⁴⁹	Germany	Prospective cohort study with historical controls; single center	2014	Both	3	16	C4
Pfaff et al ⁵⁰	Germany	Prospective cohort study with historical controls; single center	2014–2016	Anterior	22	28	A
Pfaff et al ⁵⁰	Germany	Prospective cohort study with historical controls; single center	2014–2016	Anterior	28	28	C4
Psychogios et al ⁵¹	Germany	Retrospective cohort study; single center	2016	Not specified	30	44	C4
Qureshi et al ⁵²	USA	Retrospective cohort study; multicenter	2007–2012	Not specified	66	117	C3
Ragoschke et al ⁵³	Germany	Pre-post study; retrospective data pre-intervention, prospective data post-intervention; single center	2006–2010/ 2010–2014	Both	174	81	C5
Rai et al ⁵⁴	USA	Prospective pre-post study; single center	2011–2014/ 2015	Both	30	64	B1, D2–4, F2
Ribo et al ⁵⁵	Spain	Retrospective cohort study; single center	2015–2016	Not specified	74	87	C1
Ribo et al ⁵⁵	Spain	Retrospective cohort study; single center	2015–2016	Not specified	40	87	C2
Schonenberger et al ⁵⁶	Germany	Randomized controlled trial; single center	2014–2016	Anterior	77	73	A
Schregel et al ⁵⁷	Germany	Retrospective pre-post study; single center	2008–2014/ 2014–2015	Both	90	278	C3, D2, E1
Simonsen et al ⁵⁸	Denmark	Randomized controlled trial, single center	2015–2017	Anterior	63	65	A
Singer et al ⁵⁹	Austria, Germany	Post-hoc analysis ENDOSTROKE registry; both retrospective and prospective data collection; multicenter	2011–2012	Both	36	691	A

Table 8.1. Continued

Author	Country	Study Design	Study Period*	Anterior, posterior circulation stroke or both	Intervention group (n)	Control group (n)	Type of intervention
Slezak et al ⁶⁰	Switzerland	Prospective cohort study; single center	2010–2015	Anterior	135	266	A
Sugg et al ⁶¹	USA	Retrospective cohort study; single center	2007–2009	Both	57	9	A
Tsujimoto et al ⁶²	Japan	Retrospective cohort study; single center	2011–2013	Both	6	16	B7

ENDOSTROKE, Endovascular Stroke Treatment; ESCAPE, Endovascular Treatment for Small Core and Anterior circulation Proximal Occlusion with Emphasis on Minimizing CT to Recanalization Times; IMS, Interventional Management of Stroke; MR CLEAN, Multicenter Randomized Clinical Trial of Endovascular Treatment for Acute Ischemic Stroke in The Netherlands; NASA, North American Solitaire Stent-Retriever Acute Stroke; SWIFT PRIME, Solitaire with the Intention for Thrombectomy as Primary Endovascular Treatment for Acute Ischemic Stroke; THRACE, Thrombectomie des Artères Cerebrales.

*Study period for pre/post intervention group.

The mean time to treatment was shorter in the intervention group compared to controls in the predefined workflow intervention categories (Table 6.3). The weighted difference in mean time to treatment was 12 minutes (95% CI 6 to 17; $P < 0.001$) for anesthetic management, 37 minutes (95% CI 22 to 52; $P < 0.001$) for prehospital management, 41 minutes (95% CI 27 to 54; $P < 0.001$) for in-hospital patient transfer management, 47 minutes (95% CI 28 to 67; $P < 0.001$) for teamwork, and 64 minutes (95% CI 24 to 104; $P = 0.002$) for feedback. The weighted difference in mean time to treatment of studies with multiple interventions implemented simultaneously was 50 minutes (95% CI 31 to 69; $P < 0.001$) in favor of the intervention group. Forest plots of the difference in mean time to treatment for each type of workflow intervention are showed in Supplementary Figure 8.1. The description of used time intervals in the studies, mean (SD) estimates for each study group, and a subgroup analysis per time interval is provided in Supplementary Table 8.2.

Twenty studies reported the occurrence of favorable outcome, defined as score 0–2 on the modified Rankin Scale at 90 days (Supplementary Table 8.3). Meta-analysis showed that patients in the intervention group had a higher likelihood of favorable outcome (absolute risk difference 12.2%; risk ratio (RR) 1.39; 95% CI 1.15 to 1.66; $P < 0.001$) in comparison with controls. Data from 21 studies reporting the prevalence of symptomatic intracranial hemorrhage showed no difference between patients in the intervention groups and controls (RR 0.88; 95% CI 0.71 to 1.09; $P = 0.239$). Mortality was assessed in 25 studies. Twelve studies reported in-hospital mortality, 2 studies reported mortality at 30 days, and 11 studies reported mortality at 3 months. Patients in the intervention groups had a lower risk of overall mortality (absolute risk difference 7.4%; RR 0.74; 95% CI 0.63 to 0.87; $P < 0.001$) compared with controls.

Table 8.2. Categories of workflow interventions in endovascular stroke treatment.

Anesthetic Management
A = Non-general anesthesia versus general anesthesia
Prehospital Management*
B1 = Pre-notification ED team, CT technologist, and stroke team by EMS
B2 = Mobile Stroke Treatment Unit with CT scanner, point of care laboratory testing, vascular neurologist available via telemedicine
B3 = 'Ship and drip' for transfer patients versus 'drip and ship'
B4 = CT-angiography at primary stroke center versus at intervention center
B5 = Cloud based image sharing between primary stroke center and intervention center
B6 = Use of telemedicine assessment by a stroke neurologist at primary stroke center
B7 = Air transfer versus ground transfer
In-hospital Patient Transfer
C1 = Transporting patients directly to CT scanner by EMS
C2 = Transporting (transfer) patients directly to angiosuite by EMS
C3 = 'No turn around' approach (not returning to ED after imaging for decision-making)
C4 = Single room used for CT, angiography, and EVT
C5 = Single room for patient evaluation, CT, angiography, and EVT
Teamwork
D1 = Early communication between ED team and stroke team regarding plan of care
D2 = Early activation neurointerventional team
D3 = Parallel processing from ED/hospital ward to CT: clinical assessment, laboratory tests, imaging, patient/family education by the teams in a parallel workflow
D4 = Parallel processing from CT to angiosuite: neurointerventional team meets patient at CT, teams evaluate CT/CTA and make treatment decision while angiosuite is set up, patient/family education
Feedback
E1 = Education and feedback all teams
E2 = Smartphone application/digital system for real-time window from stroke onset to puncture for all teams, visualizing performance metrics
Other
F1 = Limiting non-essential interventions (e.g. ECG, chest X-ray, additional venous access, bladder catheter placement)
F2 = Standard angiography set for all of the devices needed for EVT
F3 = No groin shaving
F4 = Standard operating procedure for intubation at the intensive care unit prior to EVT
F5 = Standard operating procedure for EVT
F6 = Not waiting for effect intravenous alteplase versus waiting for 1 hour

CT, computed tomography; CTA, computed tomography angiography; ECG, electrocardiogram; ED, emergency department; EMS, emergency medical service; EVT, endovascular treatment.

*Prehospital management includes all interventions performed before the patient arrives at the intervention center.

We found no evidence of a potential publication bias in the funnel plot that was constructed after exclusion of two studies with a very large absolute difference in time to treatment between intervention group and controls (Figure 8.3).^{34,36}




Figure 8.2. Forest plot of weighted difference in mean time to treatment for workflow interventions in endovascular stroke treatment, using random effect meta-analysis.




Figure 8.3. Funnel plot to detect potential publication bias in 43 studies of workflow interventions improvements in endovascular stroke treatment.

Table 8.3. Random-effects meta-analysis of difference in mean time to treatment for categories of workflow interventions in endovascular stroke treatment.

	Number of studies	Number of patients (intervention/control group)	Weighted mean difference, minutes (95% CI)
All interventions	47	3,480/4,002	26 (19 to 32) P<0.001
Anesthetic management	23	2,283/2,445	12 (6 to 17) P<0.001
Prehospital management	10	442/463	37 (22 to 52) P<0.001
In-hospital patient transfer management	13	730/1150	41 (27 to 54) P<0.001
Teamwork	7	502/708	47 (228 to 67) P<0.001
Feedback	4	161/417	64 (24 to 104) P=0.002
Multiple interventions simultaneously	8	531/735	50 (31 to 69) P<0.001

Discussion

Our systematic review and meta-analysis showed that interventions in the workflow of endovascular treatment for acute ischemic stroke led to a significant reduction in time to treatment. This applied to all categories of studied interventions, which were interventions aimed at using local anesthesia or conscious sedation, optimizing prehospital management, reducing in-hospital patient transfer, improving teamwork, and supplying feedback on achieved time intervals to the team. These workflow interventions led to higher likelihood of favorable functional outcome after 3 months.

The favorable effect of workflow interventions on the time to treatment are consistent with previous studies including acute stroke patients treated with intravenous alteplase. Implementation of a national quality improvement initiative organized by the American Heart Association/American Stroke Association, including more than 70,000 patients, resulted in significant shorter door-to-needle time and significant higher percentage of patients treated with intravenous alteplase within 60 minutes.⁶³ Workflow improvement strategies were promoting pre-notification of hospitals by EMS, rapid activation of the entire stroke team, rapid acquisition of brain imaging, and provision of feedback to the stroke team on performance. A single center study showed that the introduction of multiple concurrent strategies aimed at reducing in-hospital delay in treatment of acute stroke patients with intravenous alteplase led to a remarkable time reduction and final median door-to-needle time of 20 minutes.⁶

Our results are also consistent with studies on workflow improvement for reperfusion treatment of patients with myocardial infarction with ST-segment elevation. A study on time-saving strategies in the workflow for patients with acute myocardial infarction, including 365 hospitals, showed that rapid activation and availability of the entire team and use of real-time data feedback by the staff in the ED and angiography suite, reduced mean door-to-balloon time with 8 to 19 minutes.⁶⁴

The workflow interventions in this review can easily be implemented in any intervention center. A time-saving effect of >1 hour could be achieved by providing feedback on time-intervals to the entire team. Implementation of regular feedback in the four included studies in this meta-analysis was executed by supplying time intervals and outcome to the entire team daily using an online bulletin or email, reviewing each patient during weekly or monthly meetings, or comparing actually achieved times to target times every 3 months.^{32,37,41,57} Evaluation of time intervals can simply be added to existing regularly meetings at intervention hospitals. Optimizing in-hospital teamwork by using parallel processing instead of sequential processing in the workflow, and by early activation of all team members, requires multidisciplinary protocols or standard operating procedures. The time-investment to draft and implement such protocols seems well worthwhile, since our meta-analysis showed a

mean time reduction of 47 minutes.^{22,32,37,42,43,54,57} Effects of multiple interventions cannot be simply added, but implementing multiple interventions at the same time still led to a very large time reduction of 50 minutes and is probably preferred above implementing 1 intervention at a time.

Anesthetic management in endovascular stroke treatment is a much discussed topic, since it possibly influences both time to treatment intervals as cerebral perfusion and thereby indirect functional outcome. A meta-analysis including 4,716 patients undergoing endovascular stroke treatment showed a difference in time to treatment of 14 minutes in favor of patients receiving local anesthesia or conscious sedation compared with general anesthesia, and a higher odds of good functional outcome.⁶⁵ Which studies were used for comparing time to treatment by type of anesthesia management and the way missing data was handled was not disclosed. Our meta-analysis included additional studies on anesthetic management and showed a comparable difference in time to treatment of 12 minutes in favor of patients receiving local anesthesia or conscious sedation. Both meta-analyses included many observational studies with possible selection bias. Only 3 randomized controlled trials, randomizing patients for local anesthesia or conscious sedation versus general anesthesia, were included in our meta-analysis, showing a non-significant difference in treatment intervals in two studies,^{27,58} and a significant difference in time to treatment in 1 study of 10 minutes in favor of conscious sedation (95% CI 2 to 18).⁵⁶ We did not find studies comparing conscious sedation with local anesthesia. Regarding anesthetic management in endovascular stroke treatment and its effect on time to treatment, results of included randomized and non-randomized studies in our analysis varied between a significant positive effect or a significant negative effect of local anesthesia or conscious sedation, and a non-significant difference compared with general anesthesia. By combining these results in a meta-analysis, we showed a potential positive effect of non-general anesthesia on workflow.

The favorable effect of reducing time to treatment on functional outcome as described in previous studies is confirmed by our study.^{4,66} Analysis of 5 endovascular stroke treatment trials showed a 4% absolute risk difference for a good functional outcome per hour of delay between symptom onset and reperfusion.⁴ Our meta-analysis showed a difference in time to treatment effect of 26 minutes, with a total absolute risk difference of good functional outcome of 12%, which is higher compared to the ~2% absolute risk difference per half hour as seen in the meta-analysis of 5 endovascular stroke treatment trials. However, selection bias could have occurred in the non-randomized studies included in our meta-analysis and differences in baseline characteristics might have influenced our results. The effect of time to treatment on functional outcome might be stronger in clinical practice compared to a selected patient population from randomized controlled trials.⁶⁷ Furthermore, some workflow improvements, such as anesthetic management, have an effect on functional outcome which is not completely explained by the difference in time to treatment.¹

A meta-analysis of 5 large endovascular stroke trials showed no effect of time to treatment on rates of mortality and symptomatic intracranial hemorrhage.⁴ Our study showed no difference in rate of symptomatic intracranial hemorrhage, but a significantly lower mortality among patients in the intervention group. However, possible selection bias in the non-randomized studies included in our meta-analysis, could have influenced the effect of time to treatment on mortality.

This study has several limitations. To perform the meta-analysis, we estimated the mean time to treatment for 8 studies using the median time to treatment, IQR, and sample size. Since the meta-analysis is aimed at the difference in time to treatment between groups, rather than the actual time intervals per group, we assume that using the estimation of the mean time to treatment has no significant effect on the primary outcome. Considerable heterogeneity between included studies was observed. Therefore, we used a random-effects inverse variance model for our meta-analysis and categorized the interventions to perform separate analyses for each intervention type. Forty-eight of 51 included studies used a non-randomizing study design, with a high risk of selection bias. Furthermore, most data was collected retrospectively in a single center, without blinding of personnel and participants, possibly leading to performance bias. Multiple pre-post intervention studies were included in our meta-analysis, in which learning effect over time can also effect time to treatment. Therefore, generalizability is difficult to assess for the individual studies. However, since we included multiple studies on the same subject, these results can give us valuable insight on the possible effects in general practice, which is very promising. One of the purposes of a systematic review is to identify gaps in our knowledge and point out clinical areas that would benefit from more research. The 7 subcategories of pre-hospital intervention with only a limited number of studies, suggest that more work can be done in this area. Intervention studies and modelling of pre-hospital workflow may provide more insights and effective pre-hospital management strategies may have a relatively large effect on outcome.

In conclusion, interventions in the workflow of endovascular stroke treatment lead to a significant reduction in time to treatment. Reduction of any delay in time to treatment, by workflow interventions aimed at any interval between symptom onset and treatment, leads to a higher chance of good functional outcome for each individual patient. Acute stroke care should be reorganized by making use of the examples of workflow interventions described in this review to ensure the best medical care for patients with acute ischemic stroke.

Acknowledgments

We are grateful to Wichor Bramer, information specialist of the Medical Library of the Erasmus MC University Medical Center, Rotterdam, the Netherlands, for his help with the systematic search of the literature. We would also like to show our gratitude to the authors of the included articles in this review who contributed by sharing additional data from their studies with us to use in the meta-analysis:

D.P. Archer¹, G.L. Baird², D. Bar-Or³, L.A. van den Berg⁴, S. Brown⁵, M.J. Davis¹, K. Fassbender⁶, J.T. Fifi⁷, D. Frei⁸, M. Goyal⁹, S-B. Jeon¹⁰, M. Lesmeister⁶, J.R. Mascitelli⁷, C. McGraw³, R.A. McTaggart^{2,11}, B.K. Menon⁹, S. Mundiyanapurath¹², M. Möhlenbruch¹³, J. Pfaff¹³, M. Psychogios¹⁴, and S.D. Yeatts¹⁵

¹Department of Anesthesia, University of Calgary, Calgary, Canada

²Departments of Diagnostic Imaging, Brown University, Providence, USA

³Trauma Research LLC, Swedish Medical Center, Englewood, USA

⁴Department of Neurology, Academic Medical Center, Amsterdam, The Netherlands

⁵Altair Biostatistics, USA

⁶Department of Neurology, University of Saarland, Homburg, Germany

⁷Department of Neurosurgery, Icahn School of Medicine at Mount Sinai, New York, USA

⁸Radiology Imaging Associates PC, Englewood, USA

⁹Department of Radiology and Clinical Neurosciences, University of Calgary, Calgary, Canada

¹⁰Department of Neurology, Asan Medical Center, University of Ulsan College of Medicine, Seoul, Korea

¹¹Departments of Neurology and Neurosurgery, Brown University, Providence, USA

¹²Department of Neurology, University Hospital Heidelberg, Heidelberg, Germany

¹³Department of Neuroradiology, University Hospital Heidelberg, Heidelberg, Germany

¹⁴Department of Neuroradiology, University Medical Center Göttingen, Göttingen, Germany

¹⁵Department of Public Health Sciences, Medical University of South Carolina, Charleston

References

1. Goyal M, Menon BK, van Zwam WH, et al. Endovascular thrombectomy after large-vessel ischaemic stroke: a meta-analysis of individual patient data from five randomised trials. *Lancet* 2016; **387**(10029): 1723-31.
2. Bracard S, Ducrocq X, Mas JL, et al. Mechanical thrombectomy after intravenous alteplase versus alteplase alone after stroke (THRACE): a randomised controlled trial. *Lancet Neurol* 2016; **15**(11): 1138-47.
3. Muir KW, Ford GA, Messow CM, et al. Endovascular therapy for acute ischaemic stroke: the Pragmatic Ischaemic Stroke Thrombectomy Evaluation (PISTE) randomised, controlled trial. *J Neurol Neurosurg Psychiatry* 2017; **88**(1): 38-44.
4. Saver JL, Goyal M, van der Lugt A, et al. Time to Treatment With Endovascular Thrombectomy and Outcomes From Ischemic Stroke: A Meta-analysis. *JAMA* 2016; **316**(12): 1279-88.
5. National Institute of Neurological D, Stroke rt PASSG. Tissue plasminogen activator for acute ischemic stroke. *N Engl J Med* 1995; **333**(24): 1581-7.
6. Meretoja A, Strbian D, Mustanoja S, Tatlisumak T, Lindberg PJ, Kaste M. Reducing in-hospital delay to 20 minutes in stroke thrombolysis. *Neurology* 2012; **79**(4): 306-13.
7. Asif KS, Lazzaro MA, Zaidat O. Identifying delays to mechanical thrombectomy for acute stroke: onset to door and door to clot times. *J Neurointerv Surg* 2014; **6**(7): 505-10.
8. Assis ZA, Menon BK, Goyal M. Imaging department organization in a stroke center and workflow processes in acute stroke. *Eur J Radiol* 2017; **96**: 120-4.
9. Goyal M, Menon BK, Hill MD, Demchuk A. Consistently achieving computed tomography to endovascular recanalization <90 minutes: solutions and innovations. *Stroke* 2014; **45**(12): e252-6.
10. Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009; **6**(7): e1000097.
11. Higgins JP, Altman DG, Gotzsche PC, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ* 2011; **343**: d5928.
12. Wan X, Wang W, Liu J, Tong T. Estimating the sample mean and standard deviation from the sample size, median, range and/or interquartile range. *BMC Med Res Methodol* 2014; **14**: 135.
13. Abou-Chebl A, Lin R, Shazam Hussain M, et al. Conscious sedation versus general anesthesia during endovascular therapy for acute anterior circulation stroke: Preliminary results from a retrospective, multicenter study. *Stroke* 2010; **41**(6): 1175-9.
14. Abou-Chebl A, Zaidat OO, Castonguay AC, et al. North American SOLITAIRE stent-retriever acute stroke registry: Choice of anesthesia and outcomes. *Stroke* 2014; **45**(5): 1396-401.
15. Abou-Chebl A, Yeatts SD, Yan B, et al. Impact of General Anesthesia on Safety and Outcomes in the Endovascular Arm of Interventional Management of Stroke (IMS) III Trial. *Stroke* 2015; **46**(8): 2142-8.
16. Aghaebrahim A, Streib C, Rangaraju S, et al. Streamlining door to recanalization processes in endovascular stroke therapy. *J Neurointerv Surg* 2017; **9**(4): 340-5.
17. Alotaibi NM, Sarzetto F, Guha D, et al. Impact of Smartphone Applications on Timing of Endovascular Therapy for Ischemic Stroke: A Preliminary Study. *World Neurosurg* 2017; **107**: 678-83.
18. Berg LA, Koelman DLH, Berkhemer OA. Type of anesthesia and differences in clinical outcome after intra-arterial treatment for ischemic stroke. *Stroke* 2015.
19. Berkhemer OA, van den Berg LA, Fransen PS, et al. The effect of anesthetic management during intra-arterial therapy for acute stroke in MR CLEAN. *Neurology* 2016; **87**(7): 656-64.
20. Cerejo R, John S, Buletko AB, et al. A Mobile Stroke Treatment Unit for Field Triage of Patients for Intraarterial Revascularization Therapy. *J Neuroimaging* 2015; **25**(6): 940-5.
21. Davis MJ, Menon BK, Baghirzada LB, et al. Anesthetic management and outcome in patients during endovascular therapy for acute stroke. *Anesthesiology* 2012; **116**(2): 396-405.

22. Eesa M, Menon BK, Hill MD, Demchuk A, Goyal M. Achieving faster recanalization times by IA thrombolysis in acute ischemic stroke: where should we direct our efforts? *Interv Neuroradiol* 2011; **17**(2): 228-34.
23. Frei D, McGraw C, McCarthy K, et al. A standardized neurointerventional thrombectomy protocol leads to faster recanalization times. *J Neurointerv Surg* 2017; **9**(11): 1035-41.
24. Goyal M, Almekhlafi MA, Fan L, et al. Evaluation of interval times from onset to reperfusion in patients undergoing endovascular therapy in the interventional management of stroke III trial. *Circulation* 2014; **130**(3): 265-72.
25. Goyal M, Jadhav AP, Bonafe A, et al. Analysis of Workflow and Time to Treatment and the Effects on Outcome in Endovascular Treatment of Acute Ischemic Stroke: Results from the SWIFT PRIME Randomized Controlled Trial. *Radiology* 2016; **279**(3): 888-97.
26. Hassan AE, Chaudhry SA, Zacharatos H, et al. Increased rate of aspiration pneumonia and poor discharge outcome among acute ischemic stroke patients following intubation for endovascular treatment. *Neurocrit Care* 2012; **16**(2): 246-50.
27. Henden PL, Rentzos A, Karlsson JE, et al. General Anesthesia Versus Conscious Sedation for Endovascular Treatment of Acute Ischemic Stroke The AnStroke Trial (Anesthesia During Stroke). *Stroke* 2017; **48**(6): 1601-+.
28. Herrmann O, Hug A, Bosel J, et al. Fast-Track Intubation for Accelerated Interventional Stroke Treatment. *Neurocritical Care* 2012; **17**(3): 354-60.
29. Jadhav AP, Kenmuir CL, Aghaebrahim A, et al. Interfacility Transfer Directly to the Neuroangiography Suite in Acute Ischemic Stroke Patients Undergoing Thrombectomy. *Stroke* 2017; **48**(7): 1884-9.
30. Jagani M, Brinjikji W, Rabinstein AA, Pasternak JJ, Kallmes DF. Hemodynamics during anesthesia for intra-arterial therapy of acute ischemic stroke. *J Neurointerv Surg* 2016; **8**(9): 883-8.
31. Janssen H, Buchholz G, Killer M, Ertl L, Bruckmann H, Lutz J. General Anesthesia Versus Conscious Sedation in Acute Stroke Treatment: The Importance of Head Immobilization. *Cardiovasc Intervent Radiol* 2016; **39**(9): 1239-44.
32. Jeon SB, Ryoo SM, Lee DH, et al. Multidisciplinary Approach to Decrease In-Hospital Delay for Stroke Thrombolysis. *Journal of Stroke* 2017; **19**(2): 196-204.
33. John S, Thebo U, Gomes J, et al. Intra-arterial therapy for acute ischemic stroke under general anesthesia versus monitored anesthesia care. *Cerebrovasc Dis* 2014; **38**(4): 262-7.
34. Jumaa MA, Zhang F, Ruiz-Ares G, et al. Comparison of safety and clinical and radiographic outcomes in endovascular acute stroke therapy for proximal middle cerebral artery occlusion with intubation and general anesthesia versus the nonintubated state. *Stroke* 2010; **41**(6): 1180-4.
35. Just C, Rizek P, Tryphonopoulos P, Pelz D, Arango M. Outcomes of General Anesthesia and Conscious Sedation in Endovascular Treatment for Stroke. *Can J Neurol Sci* 2016; **43**(5): 655-8.
36. Kamper L, Meyn H, Rybacki K, et al. Workflow optimization in vertebrobasilar occlusion. *Cardiovasc Intervent Radiol* 2012; **35**(3): 491-7.
37. Koge J, Matsumoto S, Nakahara I, et al. Improving treatment times for patients with in-hospital stroke using a standardized protocol. *J Neurol Sci* 2017; **381**: 68-73.
38. Komatsubara K, Dembo T, Sato E, et al. Does reducing the duration from symptom onset to recanalization improve the results of intracranial mechanical thrombectomy in the elderly? *Neurol Med -Chir* 2017; **57**(3): 107-14.
39. Li FH, Deshaies EM, Singla A, et al. Impact of Anesthesia on Mortality During Endovascular Clot Removal for Acute Ischemic Stroke. *Journal of Neurosurgical Anesthesiology* 2014; **26**(4): 286-90.
40. Liang JW, Stein L, Wilson N, Fifi JT, Tuhim S, Dhamoon MS. Timing of vessel imaging for suspected large vessel occlusions does not affect groin puncture time in transfer patients with stroke. 2017.
41. Mascitelli JR, Wilson N, Shoirah H, et al. The impact of evidence: Evolving therapy for acute ischemic stroke in a large healthcare system. *J Neurointerv Surg* 2016; **8**(11): 1129-35.
42. McTaggart RA, Yaghi S, Cutting SM, et al. Association of a primary stroke center protocol for suspected stroke by large-vessel occlusion with efficiency of care and patient outcomes. *JAMA Neurol* 2017; **74**(7): 793-800.
43. Mehta BP, Leslie-Mazwi TM, Chandra RV, et al. Reducing door-to-puncture times for

- intra-arterial stroke therapy: A pilot quality improvement project. *J Am Heart Assoc* 2014; **3**(6).
44. Menon BK, Sajobi TT, Zhang Y, et al. Analysis of Workflow and Time to Treatment on Thrombectomy Outcome in the Endovascular Treatment for Small Core and Proximal Occlusion Ischemic Stroke (ESCAPE) Randomized, Controlled Trial. *Circulation* 2016; **133**(23): 2279-86.
 45. Miley JT, Memon MZ, Hussein HM, et al. A Multicenter Analysis of "Time to Microcatheter" for Endovascular Therapy in Acute Ischemic Stroke. *J Neuroimaging* 2011; **21**(2): 159-64.
 46. Mundiyanapurath S, Schonenberger S, Rosales ML, et al. Circulatory and Respiratory Parameters during Acute Endovascular Stroke Therapy in Conscious Sedation or General Anesthesia. *J Stroke Cerebrovasc Dis* 2015; **24**(6): 1244-9.
 47. Nichols C, Carrozzella J, Yeatts S, Tomsick T, Broderick J, Khatri P. Is periprocedural sedation during acute stroke therapy associated with poorer functional outcomes? *J Neurointerv Surg* 2010; **2**(1): 67-70.
 48. Pedragosa A, Alvarez-Sabín J, Rubiera M, et al. Impact of telemedicine on acute management of stroke patients undergoing endovascular procedures. *Cerebrovasc Dis* 2012; **34**(5-6): 436-42.
 49. Pfaff J, Herweh C, Pham M, et al. Mechanical thrombectomy using a combined CT/C-arm X-ray system. *J Neurointerv Surg* 2016; **8**(6): 621-5.
 50. Pfaff J, Schönenberger S, Herweh C, et al. Influence of a combined CT/C-arm system on periprocedural workflow and procedure times in mechanical thrombectomy. *Eur Radiol* 2017; **27**(9): 3966-72.
 51. Psychogios MN, Behme D, Schregel K, et al. One-Stop Management of Acute Stroke Patients: Minimizing Door-to-Reperfusion Times. *Stroke* 2017; **48**(11): 3152-5.
 52. Qureshi AI, Egila H, Adil MM, et al. "No turn back approach" to reduce treatment time for endovascular treatment of acute ischemic stroke. *J Stroke Cerebrovasc Dis* 2014; **23**(5): e317-e23.
 53. Ragoschke-Schumm A, Yilmaz U, Kostopoulos P, et al. Stroke Room': Diagnosis and Treatment at a Single Location for Rapid Intraarterial Stroke Treatment. *Cerebrovasc Dis* 2015; **40**(5-6): 251-7.
 54. Rai AT, Smith MS, Boo S, Tarabishy AR, Hobbs GR, Carpenter JS. The pit-crew model for improving door-to-needle times in endovascular stroke therapy: A Six-Sigma project. *J Neurointerv Surg* 2016; **8**(5): 447-52.
 55. Ribo M, Boned S, Rubiera M, et al. Direct transfer to angi suite to reduce door-to-puncture time in thrombectomy for acute stroke. *J Neurointerv Surg*. 2018; **10**(3): 221-4.
 56. Schonenberger S, Uhlmann L, Hacke W, et al. Effect of Conscious Sedation vs General Anesthesia on Early Neurological Improvement Among Patients With Ischemic Stroke Undergoing Endovascular Thrombectomy: A Randomized Clinical Trial. *Jama* 2016; **316**(19): 1986-96.
 57. Schregel K, Behme D, Tsogkas I, et al. Effects of Workflow Optimization in Endovascularly Treated Stroke Patients - A Pre-Post Effectiveness Study. *PLoS ONE* 2016; **11**(12).
 58. Simonsen CZ, Yoo AJ, Sorensen LH, et al. Effect of General Anesthesia and Conscious Sedation During Endovascular Therapy on Infarct Growth and Clinical Outcomes in Acute Ischemic Stroke: A Randomized Clinical Trial. *JAMA Neurol* 2018.
 59. Singer OC, Haring HP, Trenkler J, et al. Periprocedural aspects in mechanical recanalization for acute stroke: data from the ENDOSTROKE registry. *Neuroradiology* 2013; **55**(9): 1143-51.
 60. Slezak A, Kurmann R, Oppliger L, et al. Impact of anesthesia on the outcome of acute ischemic stroke after endovascular treatment with the Solitaire stent retriever. *Am J Neuroradiol* 2017; **38**(7): 1362-7.
 61. Sugg RM, Jackson AS, Holloway W, Martin CO, Akhtar N, Rymer M. Is mechanical embolectomy performed in nonanesthetized patients effective? *Am J Neuroradiol* 2010; **31**(8): 1533-5.
 62. Tsujimoto M, Yoshimura S, Enomoto Y, et al. Preliminary Experience with Air Transfer of Patients for Rescue Endovascular Therapy after Failure of Intravenous Tissue Plasminogen Activator. *Neurologia Medico-Chirurgica* 2015; **55**(3): 248-52.
 63. Fonarow GC, Zhao X, Smith EE, et al. Door-to-needle times for tissue plasminogen activator administration and clinical outcomes in acute ischemic stroke before and after a quality improvement initiative. *JAMA* 2014; **311**(16): 1632-40.

64. Bradley EH, Herrin J, Wang Y, et al. Strategies for reducing the door-to-balloon time in acute myocardial infarction. *N Engl J Med* 2006; **355**(22): 2308-20.
65. Brinjikji W, Pasternak J, Murad MH, et al. Anesthesia-Related Outcomes for Endovascular Stroke Revascularization: A Systematic Review and Meta-Analysis. *Stroke* 2017; **48**(10): 2784-91.
66. Khatri P, Yeatts SD, Mazighi M, et al. Time to angiographic reperfusion and clinical outcome after acute ischaemic stroke: an analysis of data from the Interventional Management of Stroke (IMS III) phase 3 trial. *Lancet Neurol* 2014; **13**(6): 567-74.
67. Mulder M, Jansen IGH, Goldhoorn RB, et al. Time to Endovascular Treatment and Outcome in Acute Ischemic Stroke: MR CLEAN Registry Results. *Circulation* 2018; **138**(3): 232-40.

Supplementary material

Literature search strategy


Embase

('cerebrovascular accident'/exp OR 'brain ischemia'/exp OR 'occlusive cerebrovascular disease'/de OR 'middle cerebral artery occlusion'/de OR 'basilar artery obstruction'/de OR 'stroke patient'/de OR 'brain infarction'/exp OR ((cerebrovascular NEAR/3 accident*) OR cva OR stroke* OR ((brain OR cerebr* OR 'basilar arter*') NEAR/3 (ischemi* OR ischaemi* OR occlu* OR infarct*)):ab,ti) AND ('thrombectomy'/exp OR 'endovascular surgery'/de OR 'intraarterial drug administration'/de OR (thrombectom* OR endovascular* OR intravascu* OR endo-vascular* OR intra-vascul* OR intraarterial* OR intra-arterial*):ab,ti) AND ('workflow'/de OR 'therapy delay'/de OR 'time to treatment'/de OR 'time factor'/de OR (workflow* OR work-flow* OR pathway* OR ((therap* OR treat* OR puncture* OR reperfusion* OR admission OR care) NEAR/6 (delay* OR time-to OR door-to OR picture-to OR arrival* OR onset*)) OR ((Streamlin* OR speed OR rapid OR optimi*) NEAR/3 (treat* OR therap* OR recanal* OR reperfus* OR revascular*)) OR ((Intrahospital* OR Intra-hospital* OR prehospital* OR pre-hospital*) NEAR/3 (time OR period* OR delay*)) OR (time NEAR/3 (factor* OR lapse)) OR timing):ab,ti) NOT ([Conference Abstract]/lim OR [Letter]/lim OR [Note]/lim OR [Editorial]/lim) AND [english]/lim


Medline Ovid

(exp stroke/ OR exp Brain Ischemia/ OR (Cerebrovascular Disorders/ AND Arterial Occlusive Diseases/) OR ((cerebrovascular ADJ3 accident*) OR cva OR stroke* OR ((brain OR cerebr* OR basilar arter*) ADJ3 (ischemi* OR ischaemi* OR occlu* OR infarct*)):ab,ti.) AND (Thrombectomy/ OR Endovascular Procedures/ OR (thrombectom* OR endovascular* OR intravascu* OR endo-vascular* OR intra-vascul* OR intraarterial* OR intra-arterial*):ab,ti.) AND (workflow/ OR Time-to-Treatment/ OR time factors/ OR (workflow* OR work-flow* OR pathway* OR ((therap* OR treat* OR puncture* OR reperfusion* OR admission OR care) ADJ6 (delay* OR time-to OR door-to OR picture-to OR arrival* OR onset*)) OR ((Streamlin* OR speed OR rapid OR optimi*) ADJ3 (treat* OR therap* OR recanal* OR reperfus* OR revascular*)) OR ((Intrahospital* OR Intra-hospital* OR prehospital* OR pre-hospital*) ADJ3 (time OR period* OR delay*)) OR (time ADJ3 (factor* OR lapse)) OR timing):ab,ti.) NOT (letter OR news OR comment OR editorial OR congresses OR abstracts).pt. AND english.la.

A. Anesthetic management




B. Prehospital management




Supplementary Figure 8.1. Forest plots of weighted difference in mean time to treatment for specific intervention types, using random effect meta-analysis.


C. In-hospital patient transfer management




D. Teamwork



E. Feedback



F. Multiple interventions implemented simultaneously



Supplementary Table 8.1. Risk of bias for included studies on effect of workflow improvements on time to treatment.

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias) (time to treatment)	Incomplete outcome data (attrition bias) (time to treatment)	Selective reporting (reporting bias)
Abou-Chebl ¹³	-	-	?	+	+	?
Abou-Chebl ¹⁴	-	-	-	+	+	?
Abou-Chebl ¹⁵	-	-	?	+	+	?
Aghaebrahim ¹⁶	-	-	+	+	+	+
Alotaibi ¹⁷	-	-	?	-	+	-
van den Berg ¹⁸	-	-	-	+	+	?
Berkhemer ¹⁹	-	-	-	+	+	?
Bracard ²	-	-	?	?	+	?
Cerejo ²⁰	-	-	+	+	+	?
Davis ²¹	-	-	+	+	?	?
Eesa ²²	-	-	+	+	+	-
Frei ²³	-	-	-	?	+	+
Goyal ²⁴	-	-	+	+	+	?
Goyal ²⁵	-	-	-	+	+	?
Hassan ²⁶	-	-	?	+	+	?
Henden ²⁷	+	+	+	+	+	+
Herrmann ²⁸	-	-	-	-	-	+
Jadhav ²⁹	-	-	+	+	+	?
Jagani ³⁰	-	-	+	+	?	?
Janssen ³¹	-	-	+	+	+	?
Jeon ³²	-	-	+	+	+	+
John ³³	-	-	+	+	+	?
Jumaa ³⁴	-	-	+	+	+	?
Just ³⁵	-	-	+	+	+	?
Kamper ³⁶	-	-	-	+	+	+
Koge ³⁷	-	-	+	+	+	+
Komatsubara ³⁸	-	-	+	?	+	+
Li ³⁹	-	-	-	+	+	?
Liang ⁴⁰	-	-	?	+	+	?
Mascitelli ⁴¹	-	-	+	+	+	+
McTaggart ⁴²	-	-	+	+	+	?
Mehta ⁴³	-	-	-	-	+	+

Supplementary Table 8.1. Contined

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias) (time to treatment)	Incomplete outcome data (attrition bias) (time to treatment)	Selective reporting (reporting bias)
Menon ⁴⁴	-	-	?	+	+	?
Miley ⁴⁵	-	-	?	+	+	?
Mundiyanapurath ⁴⁶	-	-	-	+	+	?
Nichols ⁴⁷	-	-	?	+	+	?
Pedragosa ⁴⁸	-	-	+	+	+	?
Pfaff ⁴⁹	-	-	-	?	+	?
Pfaff ⁵⁰	-	-	-	+	+	?
Psychogios ⁵¹	-	-	?	+	+	?
Qureshi ⁵²	-	-	?	+	+	?
Ragoschke ⁵³	-	-	-	-	+	+
Rai ⁵⁴	-	-	-	+	+	+
Ribo ⁵⁵	-	-	?	?	+	?
Schonenberger ⁵⁶	+	+	+	+	+	+
Schregel ⁵⁷	-	-	+	+	+	+
Simonsen ⁵⁸	+	+	+	+	+	+
Singer ⁵⁹	-	-	-	+	+	?
Slezak ⁶⁰	-	-	+	?	+	?
Sugg ⁶¹	-	-	+	+	+	?
Tsujimoto ⁶²	-	-	-	?	+	?
Total						
Low Risk	6%	6%	45%	80%	94%	29%
Unclear risk	0%	0%	24%	14%	4%	67%
High Risk	94%	94%	31%	6%	2%	4%

Low risk is '+' in green, high risk is '-' in red, unclear risk is '?' in yellow.

Supplementary Table 8.2. Mean time to treatment for each study group and pooled weighted mean time to treatment, for specific time intervals, using random effects meta-analysis .**A.** Onset to start endovascular treatment

Author	Time to treatment in minutes, mean (SD)	
	Intervention group	Control group
Abou-Chebl et al ¹³	296 (172)	306 (133)
van den Berg et al ¹⁸	231 (99)	269 (147)
Bracard et al ²	254 (57)	244 (60)
Davis et al ²¹	264 (93)	285 (115)
Goyal et al ²⁴	229 (49)	225 (42)
Henden et al ²⁷	190 (88)	199 (110)
Jagani et al ³⁰	256 (119)	282 (126)
Jumaa et al ³⁴	654 (804)	418 (291)
Li et al ³⁹	276 (120)	300 (138)
McTaggart et al ⁴²	112 (31)	168 (56)
Nichols et al ⁴⁷	232 (55)	237 (58)
Simonsen et al ⁵⁸	186 (72)	202 (71)
Slezak et al ⁶⁰	277 (126)	299 (157)
Tsujimoto et al ⁶²	233 (12)	273 (15)

Number of patients (intervention/control group): 1,466/1,243

Pooled weighted mean difference, minutes (95% CI): 18 (4 to 32), P=0.01

B. Door to start endovascular treatment

Author	Time to treatment in minutes, mean (SD)	
	Intervention group	Control group
Abou-Chebl et al ¹⁴	141 (91)	142 (91)
Berkhemer et al ¹⁹	134 (60)	162 (69)
Cerejo et al ²⁰	52 (24)	71 (39)
Frei et al ²³	61 (42)	76 (36)
Goyal et al ²⁵	97 (42)	103 (48)
Hassan et al ²⁶	287 (348)	310 (335)
Herrmann et al ²⁸	108 (58)	144 (42)
Jadhav et al ²⁹	22 (21)	81 (73)
Jeon et al ³²	993 (44)	202(179)
Kamper et al ³⁶	248 (165)	581 (823)
Liang et al ⁴⁰	43 (18)	58 (13)
Mascitelli et al ⁴¹	104 (53)	181 (110)
Mehta et al ⁴³	106 (28)	142 (44)

Supplementary Table 8.2. Continued

B.

Author	Time to treatment in minutes, mean (SD)	
	Intervention group	Control group
Pedragosa et al ⁴⁸	47 (31)	69 (45)
Psychogios et al ⁵¹	21 (6)	61 (35)
Qureshi et al ⁵²	158 (68)	180 (79)
Ragoschke et al ⁵³	106 (30)	121 (46)
Rai et al ⁵⁴	92 (37)	151 (51)
Ribo et al ⁵⁵ -1	60 (29)	90 (53)
Ribo et al ⁵⁵ -2	17 (8)	90 (53)
Schonenberger et al ⁵⁶	66 (20)	76 (29)
Schregel et al ⁵⁷	73 (42)	142 (93)
Singer et al ⁵⁹	92 (53)	86 (32)
Sugg et al ⁶¹	141 (52)	161 (66)

Number of patients (intervention/control group): 1,595/2,441

Pooled weighted mean difference, minutes (95% CI): 33 (23 to 43), P<0.001

C. Imaging to start endovascular treatment

Author	Time to treatment in minutes, mean (SD)	
	Intervention group	Control group
Janssen et al ³¹	59 (22)	79 (18)
Koge et al ³⁷	56 (24)	70 (30)
Menon et al ⁴⁴	56 (28)	72 (34)
Miley et al ⁴⁵	173 (57)	174 (66)
Pfaff et al ⁴⁹	39 (8)	61 (17)
Pfaff et al ⁵⁰ -1	42 (18)	45 (12)
Pfaff et al ⁵⁰ -2	45 (12)	69 (20)

Number of patients (intervention/control group): 305/198

Pooled weighted mean difference, minutes (95% CI): 15 (7 to 23), P<0.001

D. Onset to recanalization

Author	Time to treatment in minutes, mean (SD)	
	Intervention group	Control group
John et al ³³	436 (189)	510 (538)
Mundiyanapurath et al ⁴⁶	246 (90)	274 (99)

Number of patients (intervention/control group): 114/120

Supplementary Table 8.3. Efficacy and safety outcomes for workflow interventions aimed at reducing time to endovascular stroke treatment.

	Intervention group	Control group	Absolute risk difference (%)	Risk ratio (95% CI)
mRS 0–2 at 90 days	42.9% (946/2,206)	30.7% (690/2,246)	12.2	1.39 (1.15 to 1.66) P<0.001
Symptomatic intracranial hemorrhage	7.7% (183/2,368)	8.2% (154/1,883)	-0.5	0.88 (0.71 to 1.09) P=0.239
Mortality (all)	17.7% (411/2,325)	25.1% (535/2,133)	-7.4	0.74 (0.63 to 0.87) P<0.001
In-hospital mortality	15.0% (197/1,316)	24.2% (190/785)	-9.2	0.71 (0.54 to 0.94) P=0.017
Mortality at 30 days	14.4% (28/194)	19.8% (23/116)	-5.4	0.62 (0.31 to 1.24) P=0.175
Mortality at 3 months	22.8% (186/815)	26.1% (322/1,232)	-3.3	0.79 (0.64 to 0.96) P=0.019



Part III

Treat in the right place



Chapter 9

Personalized prehospital triage in acute ischemic stroke: a decision-analytic model

Stroke, 2019

Esmee Venema, Hester F. Lingsma, Vicky Chalos, Maxim J.H.L. Mulder,
Maarten M.H. Lahr, Aad van der Lugt, Adriaan C.G.M. van Es, Ewout W. Steyerberg,
M.G. Myriam Hunink, Diederik W.J. Dippel, and Bob Roozenbeek

Abstract

Background and purpose

Direct transportation to a center with facilities for endovascular treatment might be beneficial for patients with acute ischemic stroke, but it can also cause harm by delay of intravenous treatment. Our aim was to determine the optimal prehospital transportation strategy for individual patients and to assess which factors influence this decision.

Methods

We constructed a decision tree model to compare outcome of ischemic stroke patients after transportation to a primary stroke center versus a more distant intervention center. The optimal strategy was estimated based on individual patient characteristics, geographical location and workflow times. In the base case scenario, the primary stroke center was located at 20 minutes and the intervention center at 45 minutes. Additional sensitivity analyses included an urban scenario (10 versus 20 minutes) and a rural scenario (30 versus 90 minutes).

Results

Direct transportation to the intervention center led to better outcomes in the base case scenario when the likelihood of large vessel occlusion (LVO) was >33%. With a high likelihood of LVO (66%, comparable with a Rapid Arterial Occlusion Evaluation score of 5 or above), the benefit of direct transportation to the intervention center was 0.10 quality-adjusted life years (=36 days in full health). In the urban scenario, direct transportation to an intervention center was beneficial when the risk of LVO was 24% or higher. In the rural scenario, this threshold was 49%. Other factors influencing the decision included door-to-needle times, door-to-groin times and the door-in-door-out time.

Conclusions

The preferred prehospital transportation strategy for suspected stroke patients depends mainly on the likelihood of LVO, driving times, and in-hospital workflow times. We constructed a robust model that combines these characteristics and can be used to personalize prehospital triage, especially in more remote areas.

Introduction

Inter-hospital transfer for endovascular treatment (EVT) of patients with acute ischemic stroke due to an intracranial large vessel occlusion (LVO) is one of the major causes of treatment delay.¹⁻⁵ Delay of EVT is associated with poor functional outcome.^{6,7} Early identification of patients eligible for EVT followed by direct transportation to an endovascular-capable center might reduce transfer-related delay and thereby improve outcome. For this purpose, several prehospital stroke scales have been developed to identify patients who are at high risk of having an intracranial LVO based on their clinical symptoms. Currently, there is no evidence on superiority of one of these scales and more prospective validation in the prehospital setting is needed to reliably assess their accuracy.^{8,9}

Other factors that might be of importance for the prehospital triage of suspected stroke patients include the prognosis of an individual patient and the expected benefit of EVT. Especially since the large majority of ischemic stroke patients is not eligible for EVT and only benefits from rapid treatment with intravenous thrombolytics (IVT), the harm of delaying IVT should be taken into account as well. The time-dependent effect of both treatments requires a trade-off between reducing delay of IVT by transportation to the nearest primary stroke center, and avoiding transfer-related delay of EVT by direct transportation to an intervention center.

Our aim was to determine the optimal prehospital transportation strategy for individual patients with suspected ischemic stroke and to assess which factors influence this decision.

Methods

Decision model

We constructed a decision tree model to compare outcome of patients with ischemic stroke after two different prehospital strategies: transportation to the nearest primary stroke center versus direct transportation to a more distant intervention center (Figure 9.1). The outcome of each strategy was estimated based on individual patient characteristics, geographical location and treatment times. We combined a short-run model including 3 months outcome data from randomized controlled trials and a long-run Markov model that simulated 40 annual cycles. The benefit of direct transportation to the intervention center was defined as the average amount of quality-adjusted life years (QALYs) gained by this strategy. We considered a difference of more than 0.02 QALYs (=1 week in full health) to be clinically relevant.

Our study did not need approval by an ethics committee since we did not use individual patient data. Analytic methods and study materials that support the findings of this study are available from the corresponding author upon reasonable request.




Figure 9.1. Schematic overview of the model structure. The model starts with the initial decision of transportation to the nearest primary stroke center or to the nearest intervention center. The short-run model calculates the probability of every possible pathway and the associated distribution of the modified Rankin Scale (mRS) score after 3 months. It takes into account driving times, in-hospital workflow characteristics, and time-dependent treatment effects. In each annual cycle of the following Markov model, patients can remain in the same health state or die. These probabilities are based on the age and sex dependent annual mortality rates, adjusted for previously reported death hazard rate ratios of stroke patients. The decision node is represented with a square. The circles represent chance nodes, the circles marked with an M represent Markov models and the triangles represent terminal nodes. EVT, endovascular treatment; IVT, treatment with intravenous thrombolytics; LVO, large vessel occlusion.

Model parameters

Individual input parameters consisted of age, sex, time since onset of stroke symptoms, likelihood of LVO, driving time from scene to the primary stroke center, driving time from scene to the intervention center, and driving time between the primary stroke center and the intervention center. The base case used in our analyses was a 68-year old man with suspected stroke symptoms since one hour and an average risk of 30% to have an LVO causing the ischemic stroke. The nearest primary stroke center in the base case scenario was located at a 20 minutes drive by ambulance, while the intervention center was located at 45 minutes. Driving time between the centers was 35 minutes. Total time to treatment was calculated based on the driving times and in-hospital workflow characteristics (Tables 9.1 through 9.3). These parameters were varied in several sensitivity analyses.

Other model parameters were estimated based on previous literature and expert opinion of two neurovascular specialists (Drs Dippel and Roozenbeek, Supplementary Table 9.1). The β coefficients of treatment effect and time-dependent decline of treatment effect were

Table 9.1. Driving times used in the various analyses, in minutes.

	Base case scenario	Urban scenario	Rural scenario
From scene to primary stroke center	20	10	30
From scene to intervention center	45	20	90
From primary stroke center to intervention center	35	15	75

Table 9.2. In-hospital workflow characteristics used in the various analyses, in minutes.

	Base case analysis	Sensitivity analysis
Door-to-needle time (primary stroke center)	30	60
Door-to-needle time (intervention center)	30	30
Door-in-door-out time (primary stroke center)	60	90
Door-to-groin time (directly admitted)	80	80
Door-to-groin time (transferred)	50	50

Table 9.3. Calculations of treatment times in the model, in minutes.

Time to IVT (primary stroke center)	Time since onset of symptoms + driving time from scene to primary stroke center + door-to-needle time in primary stroke center
Time to IVT (intervention center)	Time since onset of symptoms + driving time from scene to intervention center + door-to-needle time in intervention center
Time to EVT (directly admitted)	Time since onset of symptoms + driving time from scene to intervention center + door-to-groin time for directly admitted patients
Time to EVT (transferred)	Time since onset of symptoms + driving time from scene to primary stroke center + door-in-door-out time + driving time from primary stroke center to intervention center + door-to-groin time for transferred patients

IVT, treatment with intravenous thrombolytics; EVT, endovascular treatment.

estimated with an ordinal logistic regression model using previously reported outcome distributions of treated patients and control patients in different time intervals.^{6,10} The effect of the uncertainty around these estimates was assessed with a probabilistic sensitivity analysis.

Likelihood of LVO

We modeled the entire range from 0% to 100% likelihood of having an LVO as cause of the ischemic stroke, and we calculated the threshold at which direct transportation to the intervention center would be beneficial. The average prevalence of LVO ranges mostly between 20% and 40% in different populations of ischemic stroke patients,¹¹ but prehospital stroke scales can be used in individual patients to distinguish between lower and higher

risk. To illustrate this, we calculated the likelihood of LVO in case of a positive test (positive predictive value) and in case of a negative test (1 minus the negative predictive value) for the Los Angeles Motor Scale (LAMS), the Rapid Arterial Occlusion Evaluation (RACE), and the 3-Item Stroke Scale (3I-SS), using the pooled sensitivity and specificity rates reported in a previous meta-analysis (Table 9.4).⁸ In the sensitivity analyses, we used the RACE scale as an example of low risk (14%) and high risk (66%).

Table 9.4. Likelihood of large vessel occlusion based on several prehospital stroke scales.

Stroke scale	Sensitivity*	Specificity*	Prior probability = 20%		Prior probability = 30%		Prior probability = 40%	
			PPV	1-NPV	PPV	1-NPV	PPV	1-NPV
LAMS ≥ 4	38%	87%	42%	15%	56%	23%	66%	32%
RACE ≥ 5	67%	85%	53%	9%	66%	14%	75%	21%
3I-SS ≥ 4	19%	97%	61%	17%	73%	26%	81%	36%

The different prior probabilities illustrate the prevalence of large vessel occlusion in different populations of ischemic stroke patients.

*Pooled sensitivity and specificity rates as reported previously.⁸

3I-SS, 3-Item Stroke Scale; LAMS, Los Angeles Motor Scale; NPV, negative predictive value; PPV, positive predictive value; and RACE, Rapid Arterial Occlusion Evaluation.

Outcome measures

Primary outcome of the short-run model was the distribution of modified Rankin Scale (mRS) scores after 3 months. The mRS is a 7-point scale to assess functional outcome and disability, and ranges from 0 (no symptoms) to 6 (death). The outcome distributions of untreated ischemic stroke patients with and without an LVO were based on the control groups of previously reported randomized clinical trials (Supplementary Figure 9.1).^{10,12} These distributions were shifted for different outcomes based on the estimated treatment effect. To do so, we first transformed the probabilities of each of the baseline mRS scores to the log-odds scale, using the formula $\log\text{-odds}(p) = \log(p/(1-p))$. We then adjusted these log-odds for the effect of IVT and EVT by adding the β coefficient (=log odds ratio) for the treatment effect at time 0 minus the decline in treatment effect over time (Tables 9.5 and 9.6). The adjusted log-odds were then converted back to probabilities using the inverse logit function: $p(\log\text{-odds}) = 1/(1+\exp(-\log\text{-odds}))$. This resulted in the adjusted mRS distributions for treated patients.

All mRS scores were considered to be separate health states with an associated utility score to calculate the average amount of QALYs per strategy. This is a commonly used measure to assess both the length and the quality of life.¹³ Utility scores represent the quality of life, and range from 1 (perfect quality of life) to 0 (death), or lower for states considered worse

than death. We used previously reported utility scores for ischemic stroke patients: 0.95 for mRS 0; 0.93 for mRS 1; 0.83 for mRS 2; 0.62 for mRS 3; 0.42 for mRS 4; and 0.11 for mRS 5.¹⁴ The average life expectancy was calculated for every mRS category with a long-run Markov model that simulated 40 annual cycles. Age and sex dependent annual mortality rates were adjusted for previously reported death hazard rate ratios of stroke patients (Supplementary Table 9.1).¹⁵ Life years were discounted with 3% per year. We calculated the total number of QALYs by multiplying the utility scores of each mRS category with the corresponding life expectancy in years.

Model assumptions

The model was used for scenarios in which the nearest hospital is not an intervention center, since there has to be decisional uncertainty about the optimal transportation strategy. Outcome was modeled for patients with ischemic stroke only, disregarding other diagnoses. A substantial proportion of all patients suspected of ischemic stroke have an intracerebral hemorrhage, a transient ischemic attack or a stroke-mimic, but we assumed the outcome of these patients to be unrelated to the transportation strategy. Because our analyses concerned the trade-off between IVT and EVT, we did not consider patients presenting >4.5 hours after onset of symptoms.

We assumed that IVT could be given within 4.5 hours after onset of symptoms and EVT within 6 hours after onset. We modeled the decrease in treatment effect of IVT and EVT consistently over time and similar for all patients. Since large pooled analyses showed no significant interaction between age or gender and treatment effect, we did not include age or gender specific treatment effects in our model.^{12,16} The relative treatment effect of IVT was not influenced by stroke severity, which implies a smaller absolute treatment effect for patients with a more severe stroke. The importance of these model assumptions was assessed in several sensitivity analyses.

Sensitivity analyses

In addition to the base case scenario, we also assessed examples of an urban scenario (10 minutes from scene to primary stroke center; 20 minutes from scene to intervention center; and 15 minutes inter-center driving time) and a more rural scenario (30, 90, and 75 minutes respectively). We used a Tornado-analysis to explore the relative importance of the model parameters, by varying each parameter at a time while the others were held constant. Additional sensitivity analyses were performed with: increased workflow times in the primary stroke center (Table 9.2); a female patient; a patient with contra-indications for IVT; an absent effect of IVT for patients with LVO; and utility weights as defined in a study of Chaisinanunkul et al.¹⁷

We performed a probabilistic sensitivity analysis using second-order Monte-Carlo simulations to assess decisional uncertainty around the model parameters. This involves running the

model 10,000 times and calculating the optimal transportation strategy every single time. In each simulation, the estimates of the model parameters were randomly sampled from the prespecified distribution of each parameter (β , (log)normal or triangular) (Supplementary Table 9.1). We reported the percentage of simulations in which transportation to the primary stroke center or direct transportation to the intervention center was preferred, and the median benefit of direct transportation to an intervention center with a 95% credible interval.

We constructed an online tool in which input factors and regional workflow times can be adjusted for an individual patient in a specific scenario. We used R statistical software (version 3.4.4) with the dampack package (version 0.0.0.9) and the R Shiny package (version 1.0.5).

Results

Base case analysis

Direct transportation to the intervention center was preferred for the 68-year old man in the base case scenario, with a primary stroke center located at 20 minutes and an intervention center at 45 minutes, when the likelihood of LVO was 34% or above (Figure 9.2). When the risk of having an LVO was 66%, comparable with a RACE score of 5 or more, the benefit of direct transportation to the intervention center was 0.10 QALYs (=36 days). When the risk of having an LVO was 14%, comparable with a RACE score below 5, transportation to the primary stroke center was preferred with a difference of 0.03 QALYs (=11 days).

Sensitivity analyses

In the urban scenario (10 minutes to the primary stroke center, 20 minutes to the intervention center), the threshold for direct transportation to the intervention center was lower than in the base case scenario (24%, Figure 9.2). In the rural scenario (30 minutes to the primary stroke center and 90 minutes to the intervention center), direct transportation to the intervention center was only preferred for patients with an LVO likelihood of 49% or above.

Other factors that strongly affected the decision threshold in the Tornado-analysis were the in-hospital workflow characteristics (Supplementary Figure 9.3). While remaining the other parameters of the base case constant, transportation to the primary stroke center was beneficial when the door-to-groin time for directly transported patients was above 102 minutes, the door-to-groin time for transferred patients was below 28 minutes, or the door-in-door-out time in the primary stroke center was <38 minutes. Age did not influence the preferred strategy, although it was important for the general prognosis of individual patients.

In the analysis with increased workflow times in the primary stroke center, transportation to the intervention center was more favorable (Supplementary Figure 9.4). As might be




Figure 9.2. The optimal transportation strategy based on the likelihood of large vessel occlusion. Primary stroke center: the nearest nonendovascular-capable stroke center; intervention center: the nearest endovascular-capable stroke center. A, represents the base case scenario (primary stroke center at 20 minutes and intervention center at 45 minutes); B, the urban scenario (10 and 20 minutes, respectively); and C, the rural scenario (30 and 90 minutes).

expected, we found that transportation to the primary stroke center was never preferred for a patient with contra-indications for IVT. In all other exploratory sensitivity analyses, the decision threshold remained relatively unchanged.

Probabilistic sensitivity analysis

The uncertainty around the estimates of the model parameters caused little decisional uncertainty in the probabilistic sensitivity analysis (Supplementary Table 9.2). When the risk of having an LVO in the base case scenario was high, transportation to the intervention center was associated with a clinically relevant benefit in 94% of the simulations with a median difference of 0.09 QALYs (=33 days in full health, 95% credible interval: 0.00 to 0.20). When the risk of having an LVO was low, direct transportation to the intervention center was preferred in only 6% of the simulations, with a median difference of -0.03 QALYs (95% credible interval: -0.09 to 0.03).

Online tool

The effect of simultaneous changes in individual patient characteristics and regional workflow characteristics on the optimal prehospital transportation strategy for individual patients can be further explored by using the online tool at mrpredicts.shinyapps.io/triage (Figure 9.3).

Personalized prehospital triage in acute ischemic stroke

A decision-analytic model




Figure 9.3. Screenshot of the online tool. This interactive tool can be used at mrpredicts.shinyapps.io/triage. It shows the effect of changes in individual patient characteristics and regional workflow times on the optimal transportation strategy based on the likelihood of large vessel occlusion.

Table 9.5. Calculations of treatment effect in the model.

Probability of receiving IVT	If treatment is possible within 4.5 hours after onset of symptoms: 55%, otherwise: 0%
Probability of early reperfusion	If LVO present and treated with IVT: 11%, otherwise: 0%
Probability of receiving EVT	If treatment is possible within 6 hours after onset of symptoms: 85%, otherwise 0%
Treatment effect IVT (β coefficient)*	0.56 (=odds ratio 1.75) at time 0, minus 0.0019 x time to IVT in minutes
Treatment effect EVT (β coefficient)*	1.35 (=odds ratio 3.85) at time 0, minus 0.0026 x time to EVT in minutes

*The time-dependent decrease in treatment effect is illustrated in Supplementary Figure 9.2. IVT, treatment with intravenous thrombolytics; EVT, endovascular treatment.

Table 9.6. Calculations of treatment effect in the model.

	Primary stroke center	Intervention center
Outcome with large vessel occlusion (log-odds)*		
Early reperfusion after IVT	1.35 + (-0.0026 x time to IVT (primary stroke center)) + log-odds from baseline mRS distribution with LVO	1.35 + (-0.0026 x time to IVT (intervention center)) + log-odds from baseline mRS distribution with LVO
EVT (with or without prior IVT)	1.35 + (-0.0026 x time to EVT (transferred)) + log-odds from baseline mRS distribution with LVO	1.35 + (-0.0026 x time to EVT (directly admitted)) + log-odds from baseline mRS distribution with LVO
No EVT	Log-odds from baseline mRS distribution with LVO	Log-odds from baseline mRS distribution with LVO
Outcome without large vessel occlusion (log-odds)*		
IVT	0.56 + (-0.0019 x time to IVT (primary stroke center)) + log-odds from baseline mRS distribution without LVO	0.56 + (-0.0019 x time to IVT (intervention center)) + log-odds from baseline mRS distribution without LVO
No IVT	Log-odds from baseline mRS distribution without LVO	Log-odds from baseline mRS distribution without LVO

*The baseline mRS distribution with LVO and without LVO is illustrated in Supplementary Figure 9.1. The corresponding probability of a certain outcome was calculated as: $1/(1+\exp(-\log\text{-odds}))$. EVT, endovascular treatment; IVT, treatment with intravenous thrombolytics; LVO, large vessel occlusion; and mRS, modified Rankin Scale.

Discussion

Our decision model shows that direct transportation to an intervention center can be beneficial for patients with a high risk of having an LVO, but will likely lead to worse outcomes when the risk is low, especially in scenarios with longer driving times. Combining individual likelihood of LVO and estimated driving times on a case-by-case basis could improve prehospital triage decisions, decrease treatment delay and thereby improve functional outcome of individual patients. In-hospital workflow characteristics such as the door-in-door-out and door-to-groin times have a large effect on the optimal strategy as well. Our online tool allows individual and regional input to inform prehospital triage strategies in different settings.

No single time threshold can be given to optimize triage decisions without considering the probability of being eligible for EVT, as was described by other models.¹⁸⁻²¹ As a measure for the likelihood of LVO, these models used previously reported cutoffs of one or multiple prehospital stroke scales. However, prospective prehospital validation is required to assess and compare the accuracy of these scales when used by emergency medical services in a broad population of suspected stroke patients.⁸ The full dependence of these models on insufficiently validated prehospital stroke scales limits the validity of their results. Therefore, we did not depend our model on a specific prehospital stroke scale, but assessed the entire range from 0% to 100% likelihood of LVO. Several prehospital stroke scales may be used to estimate this likelihood, preferably after more extensive validation.

Our study also has some limitations. Decision-analytic modeling requires the use of estimated model parameters and multiple assumptions. In extensive sensitivity analyses, we found no substantial decisional uncertainty. Nevertheless, the assumptions we made about the treatment effect of IVT and EVT might have influenced our results. We used reported outcomes and effect sizes of IVT based on studies that did not assess LVO status, since randomized trials on the effect of IVT were performed before the introduction of CT angiography as standard of care for acute stroke patients. To calculate a common odds ratio for the decay in effect of IVT, we used mRS distributions at different time points provided by a pooled analysis from 2004.¹⁰ A more recent study with a larger sample size only reported odds ratios specific for the cutoff of mRS 0-1,¹⁶ which are not valid to use in shift-analyses of total mRS score. Although the effect of IVT tends to diminish with more proximally located occlusions,^{22,23} we included a consistent relative treatment effect of IVT in the range of the National Institute of Health Stroke Scale score of 5 to 22.¹⁶ Since the baseline outcome distribution is less favorable for patients with an LVO, the same relative treatment effect will give a smaller absolute effect in LVO stroke. The precise benefit of IVT prior to EVT is still uncertain, since the available studies are flawed by confounding by indication.^{24,25} The results of ongoing randomized controlled trials comparing EVT and IVT with prior IVT have to provide more insight on this matter.

Outcome of stroke mimics, such as migraine, epilepsy and conversion disorder, were considered to be independent of time, since there are no time-dependent treatment options that would require transportation to the nearest hospital. Patients with an intracerebral hemorrhage might benefit from transportation to a specialized center, but there is currently insufficient evidence to include the effect of rapid treatment in our model. Recent studies showed that EVT can be effective in a subgroup of stroke patients presenting between 6 and 24 hours after onset.^{26,27} Other factors might be important to optimize triage of these patients and our results therefore only apply to patients presenting within 6 hours. Furthermore, we did not include costs in our model. Because outcome was measured based on functional outcome, we were only able to model harm due to delayed treatment. Inconvenience of unnecessary transportations for patients and their relatives, as well as inefficient resource utilization was not integrated in the model. It is likely that unnecessary transportation to an intervention center will cause crowding at the emergency department, which might have financial consequences and a negative effect on the in-hospital workflow. It is therefore important to further explore the cost-effectiveness of different prehospital triage strategies.

We constructed a decision model that predicts the optimal strategy based on individual patient characteristics, and that can easily be adjusted for differences in geographical location or in-hospital workflow characteristics. Our online tool forms the basis of personalized prehospital triage of suspected ischemic stroke patients. Clinicians and researchers can plug in the specific characteristics of their own region into this tool to guide local prehospital triage policies. Although a fixed threshold the likelihood of LVO might be suitable for triage in urbanized regions with small distances between centers, our model showed that a higher threshold should likely be considered in more remote areas. After more extensive validation of the prehospital stroke scales, the tool can be used to combine the individual's likelihood of LVO with estimated driving and local workflow times to improve prehospital triage decisions on a case-by-case basis. Future integration with a GPS-controlled navigation application will further facilitate personalized triage based on real-time information about driving and local workflow times.

Conclusions

The preferred prehospital transportation strategy for suspected stroke patients depends mainly on the likelihood of LVO, driving times and in-hospital workflow times. We constructed a robust model that combines these characteristics and can be used to personalize prehospital triage, especially in more remote areas.

References

1. Goyal M, Jadhav AP, Bonafe A, et al. Analysis of Workflow and Time to Treatment and the Effects on Outcome in Endovascular Treatment of Acute Ischemic Stroke: Results from the SWIFT PRIME Randomized Controlled Trial. *Radiology* 2016; **279**(3): 888-97.
2. Menon BK, Sajobi TT, Zhang Y, et al. Analysis of Workflow and Time to Treatment on Thrombectomy Outcome in the Endovascular Treatment for Small Core and Proximal Occlusion Ischemic Stroke (ESCAPE) Randomized, Controlled Trial. *Circulation* 2016; **133**(23): 2279-86.
3. Menon BK, Almekhlafi MA, Pereira VM, et al. Optimal workflow and process-based performance measures for endovascular therapy in acute ischemic stroke: analysis of the Solitaire FR thrombectomy for acute revascularization study. *Stroke* 2014; **45**(7): 2024-9.
4. Venema E, Boodt N, Berkhemer OA, et al. Workflow and factors associated with delay in the delivery of intra-arterial treatment for acute ischemic stroke in the MR CLEAN trial. *J Neurointerv Surg* 2017; **10**: 424-8.
5. Froehler MT, Saver JL, Zaidat OO, et al. Interhospital Transfer Before Thrombectomy Is Associated With Delayed Treatment and Worse Outcome in the STRATIS Registry (Systematic Evaluation of Patients Treated With Neurothrombectomy Devices for Acute Ischemic Stroke). *Circulation* 2017; **136**(24): 2311-21.
6. Saver JL, Goyal M, van der Lugt A, et al. Time to Treatment With Endovascular Thrombectomy and Outcomes From Ischemic Stroke: A Meta-analysis. *JAMA* 2016; **316**(12): 1279-88.
7. Mulder M, Jansen IGH, Goldhoorn RB, et al. Time to Endovascular Treatment and Outcome in Acute Ischemic Stroke: MR CLEAN Registry Results. *Circulation* 2018; **138**: 232-40.
8. Smith EE, Kent DM, Bulsara KR, et al. Accuracy of Prediction Instruments for Diagnosing Large Vessel Occlusion in Individuals With Suspected Stroke: A Systematic Review for the 2018 Guidelines for the Early Management of Patients With Acute Ischemic Stroke. *Stroke* 2018; **49**(3): e111-e22.
9. Krebs W, Sharkey-Toppen TP, Cheek F, et al. Prehospital Stroke Assessment for Large Vessel Occlusions: A Systematic Review. *Prehosp Emerg Care* 2018; **22**(2): 180-8.
10. Hacke W, Donnan G, Fieschi C, et al. Association of outcome with early stroke treatment: pooled analysis of ATLANTIS, ECASS, and NINDS rt-PA stroke trials. *Lancet* 2004; **363**(9411): 768-74.
11. Lakomkin N, Dhamoon M, Carroll K, et al. Prevalence of large vessel occlusion in patients presenting with acute ischemic stroke: a 10-year systematic review of the literature. *J Neurointerv Surg* 2019; **11**(3): 241-5.
12. Goyal M, Menon BK, van Zwam WH, et al. Endovascular thrombectomy after large-vessel ischaemic stroke: a meta-analysis of individual patient data from five randomised trials. *Lancet* 2016; **387**: 1723-31.
13. Weinstein MC, Stason WB. Foundations of cost-effectiveness analysis for health and medical practices. *N Engl J Med* 1977; **296**(13): 716-21.
14. Dijkland SA, Voormolen DC, Venema E, et al. Utility-Weighted Modified Rankin Scale as Primary Outcome in Stroke Trials: A Simulation Study. *Stroke* 2018; **49**: 965-71.
15. Samsa GP, Reutter RA, Parmigiani G, et al. Performing cost-effectiveness analysis by integrating randomized trial data with a comprehensive decision model: application to treatment of acute ischemic stroke. *J Clin Epidemiol* 1999; **52**(3): 259-71.
16. Emberson J, Lees KR, Lyden P, et al. Effect of treatment delay, age, and stroke severity on the effects of intravenous thrombolysis with alteplase for acute ischaemic stroke: a meta-analysis of individual patient data from randomised trials. *Lancet* 2014; **384**(9958): 1929-35.
17. Chaisinanunkul N, Adeoye O, Lewis RJ, et al. Adopting a Patient-Centered Approach to Primary Outcome Analysis of Acute Stroke Trials Using a Utility-Weighted Modified Rankin Scale. *Stroke* 2015; **46**(8): 2238-43.
18. Nogueira RG, Silva GS, Lima FO, et al. The FAST-ED App: A Smartphone Platform for the Field Triage of Patients With Stroke. *Stroke* 2017; **48**(5): 1278-84.

19. Schlemm E, Ebinger M, Nolte CH, Endres M, Schlemm L. Optimal Transport Destination for Ischemic Stroke Patients With Unknown Vessel Status: Use of Prehospital Triage Scores. *Stroke* 2017; **48**: 2184-91.
20. Benoit JL, Khatri P, Adeoye OM, et al. Prehospital Triage of Acute Ischemic Stroke Patients to an Intravenous tPA-Ready versus Endovascular-Ready Hospital: A Decision Analysis. *Prehosp Emerg Care* 2018; **22**: 722-33.
21. Holodinsky JK, Williamson TS, Demchuk AM, et al. Modeling Stroke Patient Transport for All Patients With Suspected Large-Vessel Occlusion. *JAMA Neurol* 2018; **75**(12): 1477-86.
22. del Zoppo GJ, Poeck K, Pessin MS, et al. Recombinant tissue plasminogen activator in acute thrombotic and embolic stroke. *Ann Neurol* 1992; **32**(1): 78-86.
23. Demchuk AM, Goyal M, Yeatts SD, et al. Recanalization and clinical outcome of occlusion sites at baseline CT angiography in the Interventional Management of Stroke III trial. *Radiology* 2014; **273**(1): 202-10.
24. Mistry EA, Mistry AM, Nakawah MO, et al. Mechanical Thrombectomy Outcomes With and Without Intravenous Thrombolysis in Stroke Patients: A Meta-Analysis. *Stroke* 2017; **48**(9): 2450-6.
25. Kaesmacher J, Mordasini P, Arnold M, et al. Direct mechanical thrombectomy in tPA-ineligible and -eligible patients versus the bridging approach: a meta-analysis. *J Neurointerv Surg* 2019; **11**(1): 20-7.
26. Nogueira RG, Jadhav AP, Haussen DC, et al. Thrombectomy 6 to 24 Hours after Stroke with a Mismatch between Deficit and Infarct. *N Engl J Med* 2018; **378**(1): 11-21.
27. Albers GW, Marks MP, Kemp S, et al. Thrombectomy for Stroke at 6 to 16 Hours with Selection by Perfusion Imaging. *N Engl J Med* 2018; **378**(8): 708-18.

Supplementary material

Supplementary Table 9.1. Overview of the model parameters used in the probabilistic sensitivity analysis.

Model parameter	Estimated value	SE	Distribution	Source
Treatment characteristics				
Probability of receiving IVT if presenting <4.5h with an ischemic stroke	0.55	±10%	β	Expert opinion
Effect of IVT, beta at time 0	0.56	0.18	Normal	Hacke et al. ¹ (n=2763)
Time-dependent decrease in effect of IVT, beta per minute	-0.0019	0.001	Normal	Hacke et al. ¹ (n=2763)
Probability of early reperfusion after IVT	0.11	0.008	β	Tsivgoulis et al. ² (n=1561)
Probability of receiving EVT if presenting <6h with an LVO	0.85	±10%	β	Expert opinion
Effect of EVT, beta at time 0	1.35	0.29	Normal	Saver et al. ³ (n=1275)
Time-dependent decrease in effect of EVT, beta per minute	-0.0026	0.001	Normal	Saver et al. ³ (n=1275)
Outcome parameters				
Utility values				Dijkland et al. ⁴
mRS scores 0	0.95	0.08	β	(n=7)
mRS scores 1	0.93	0.04	β	(n=36)
mRS scores 2	0.83	0.04	β	(n=84)
mRS scores 3	0.62	0.05	β	(n=87)
mRS scores 4	0.42	0.04	β	(n=133)
mRS scores 5	0.11	0.05	β	(n=45)
Death hazard rate ratios				Samsa et al. ⁵
mRS scores 0-1	1.00	NA	NA	
mRS scores 2	1.11	1.0-1.5	Triangular	
mRS scores 3	1.27	1.2-1.4	Triangular	
mRS scores 4	1.71	1.3-2.0	Triangular	
mRS scores 5	2.37	1.5-4.0	Triangular	

EVT, endovascular treatment; IVT, treatment with intravenous thrombolytics; mRS, modified Rankin Scale; NA, not applicable; SE, standard error.

Corresponding references

¹ Lancet 2004;363:768-774.

² Stroke 2018;49:232-235.

³ JAMA 2016;316:1279-1288.


⁴ Stroke 2018;49:965-971.

⁵ J Clin Epidemiol 1999;52:259-271.


Supplementary Table 9.2. Results of the probabilistic sensitivity analysis using 10,000 Monte Carlo simulations.

Probability of large vessel occlusion	Optimal transportation strategy (percentage of simulations)			Median benefit of direct transportation to intervention center, QALYs (95% credible interval)
	Primary stroke center	Indifferent	Intervention center	
Base case scenario				
Low risk (14%)	60%	35%	6%	-0.03 (-0.09 to 0.03)
Average risk (30%)	20%	43%	37%	0.01 (-0.06 to 0.08)
High risk (66%)	1%	7%	92%	0.09 (-0.01 to 0.20)
Urban scenario				
Low risk (14%)	8%	82%	10%	0.00 (-0.03 to 0.03)
Average risk (30%)	1%	33%	66%	0.03 (-0.01 to 0.08)
High risk (66%)	<1%	4%	96%	0.09 (0.01 to 0.19)
Rural scenario				
Low risk (14%)	87%	9%	4%	-0.10 (-0.25 to 0.03)
Average risk (30%)	68%	18%	14%	-0.05 (-0.19 to 0.07)
High risk (66%)	10%	16%	74%	0.06 (-0.07 to 0.20)


The percentage of simulations in which transportation to the primary stroke center or direct transportation to the intervention center was preferred and the median difference in expected outcome between the two strategies is shown for different scenarios and different likelihood of large vessel occlusion. The percentages may not add up to 100% due to rounding. QALYs, quality-adjusted life years.




Supplementary Figure 9.1. Distribution of the baseline modified Rankin Scale (mRS) scores. Outcome for untreated ischemic stroke patients without large vessel occlusion (A) and untreated ischemic stroke patients with large vessel occlusion (B).
 A. Hacke et al. placebo group, n=1384 (Lancet 2004;363:768-774).
 B. Goyal et al. control group, n=644 (Lancet 2016;387:1723-1731).



Supplementary Figure 9.2. The time-dependent decrease in treatment effect as used in the model. EVT, endovascular treatment; IVT, treatment with intravenous thrombolytics.



Supplementary Figure 9.3. Tornado-plot with the effect of changes in model parameters on the optimal transportation strategy. The bars illustrate the effect of changes in the model parameter estimates, within the indicated ranges, on the optimal transportation strategy. The bars are ordered according to their impact on the difference in outcome. EVT, endovascular treatment; IVT, treatment with intravenous thrombolytics; LVO, large vessel occlusion.



Supplementary Figure 9.4. Results of the sensitivity analyses. The bars show the optimal transportation strategy for different likelihood of large vessel occlusion in the base case scenario (primary stroke center at 20 minutes and intervention center at 45 minutes); the urban scenario (primary stroke center at 10 minutes and intervention center at 20 minutes); and the rural scenario (primary stroke center at 30 minutes and intervention center at 90 minutes).

0. Base case analysis.
1. Sensitivity analysis with increased workflow times in primary stroke center (door-to-needle time 60 minutes and door-in-door-out time 90 minutes).
2. Sensitivity analysis with female patient.
3. Sensitivity analysis with contra-indications for treatment with intravenous thrombolytics.
4. Sensitivity analysis with absent effect of treatment with intravenous thrombolysis for patients with a large vessel occlusion.
5. Sensitivity analysis with utility weights as defined in the study of Chaisinanunkul et al (Stroke 2015;46:2238-2243).



Chapter 10

Prehospital triage strategies for the transportation of suspected stroke patients in the United States

Accepted for publication

Esmee Venema, James F. Burke, Bob Roozenbeek, Jason Nelson, Hester F. Lingsma, Diederik W.J. Dippel, and David M. Kent



Chapter 11

Prehospital triage of patients with suspected stroke symptoms (PRESTO): protocol of a prospective observational study

BMJ Open, 2019

Esmee Venema, Martijne H.C. Duvekot, Hester F. Lingsma, Anouk D. Rozeman, Walid Moudrous, Frederique H. Vermeij, Marileen Biekart, Aad van der Lugt, Henk Kerkhoff, Diederik W.J. Dippel, and Bob Roozenbeek, on behalf of the PRESTO investigators

Abstract

Introduction

The efficacy of both intravenous treatment (IVT) and endovascular treatment (EVT) for patients with acute ischemic stroke strongly declines over time. Only a subset of patients with ischemic stroke caused by an intracranial large vessel occlusion (LVO) in the anterior circulation can benefit from EVT. Several prehospital stroke scales were developed to identify patients that are likely to have an LVO, which could allow for direct transportation of EVT eligible patients to an endovascular-capable center without delaying IVT for the other patients. We aim to prospectively validate these prehospital stroke scales simultaneously to assess their accuracy in predicting LVO in the prehospital setting.

Methods and analysis

Prehospital triage of patients with suspected stroke symptoms (PRESTO) is a prospective multicenter observational cohort study in the southwest of the Netherlands including adult patients with suspected stroke in the ambulance. The paramedic will assess a combination of items from five prehospital stroke scales, without changing the normal workflow. Primary outcome is the clinical diagnosis of ischemic stroke with an intracranial LVO in the anterior circulation. Additional hospital data concerning the diagnosis and provided treatment will be collected by chart review. Logistic regression analysis will be performed and performance of the prehospital stroke scales will be expressed as sensitivity, specificity and area under the receiver operator curve (AUC).

Ethics and dissemination

The Institutional Review Board of the Erasmus MC University Medical Center has reviewed the study protocol and confirmed that the Dutch Medical Research Involving Human Subjects Act (WMO) is not applicable. The findings of this study will be disseminated widely through peer-reviewed publications and conference presentations. The best performing scale, or the simplest scale in case of clinical equipoise, will be integrated in a decision model with other clinical characteristics and real-life driving times to improve prehospital triage of suspected stroke patients.

Trial registration number NTR7595 (www.trialregister.nl)

Box 11.1 Strengths and limitations of this study.

- *Prospective simultaneous validation of several prehospital stroke scales allows for direct comparison of their accuracy.*
- *In contrast to previous studies based on in-hospital assessment by experienced physicians, assessment of the prehospital stroke scales will be performed by paramedics in daily clinical practice.*
- *The results of this study will provide unique insight in the characteristics of an unselected group of patients with suspected stroke in the prehospital setting.*
- *The best performing scale will be integrated in a prehospital decision tool with other clinical characteristics and real-life driving times to select those patients that benefit from direct transportation to an endovascular-capable center.*
- *Performance will be measured with the area under the receiver operator curve (AUC), which does not always relate directly to the clinical usefulness of these scales.*

Introduction

Rapid treatment with intravenous thrombolytics (IVT) is effective for patients with an ischemic stroke of less than 4.5 hours after onset.^{1,2} However, the effect of IVT is limited for ischemic stroke caused by an intracranial large vessel occlusion (LVO) in the anterior circulation, which accounts for approximately 30% of the patients.³ These patients can benefit from endovascular treatment (EVT), preferably started within 6 hours after onset of symptoms, but this treatment can only be performed in specialized intervention centers.⁴ The effect of both treatments strongly declines over time.⁵⁻⁷ In current clinical practice, most suspected stroke patients are transported by ambulance to the nearest hospital for immediate treatment with IVT. Patients can subsequently be transferred to an endovascular-capable center, if eligible for EVT. This is one of the main causes of treatment delay and is associated with worse functional outcomes after EVT.^{8,9}

Several prehospital stroke scales were developed to identify patients that are likely to have an LVO, which could allow for direct transportation of EVT eligible patients to an endovascular-capable center without delaying IVT for the other patients.^{10,11} Most of these scales were derived from the National Institute of Health Stroke Severity (NIHSS) score and external validation was often attempted by retrospective assessment of the items based on the NIHSS score completed by the treating physician at the emergency department.¹²⁻¹⁴ The results of existing prehospital validation studies are limited due to small sample sizes, selected populations or the exclusion of stroke mimics.¹⁵⁻¹⁸ Further prospective validation

is therefore required to assess and compare the accuracy of these scales when used by emergency medical services (EMS) personnel in a broad population of suspected stroke patients under circumstances that reflect usual care.

Objective

The primary objective of this study is to prospectively validate several prehospital stroke scales simultaneously to assess their accuracy in predicting the likelihood of ischemic stroke caused by an intracranial LVO in the prehospital setting.

Methods and analysis

Study design

Prehospital triage of patients with suspected stroke symptoms (PRESTO) is a prospective multicenter observational cohort study. Patients will be recruited in the ambulance and a combination of items from different prehospital stroke scales will be assessed by the paramedic. The normal workflow will not be affected and there is no intervention. Additional hospital data will be collected by chart review. Routinely performed neuro-imaging will be collected and centrally assessed. Follow-up will only be performed in patients with a final diagnosis of ischemic stroke.

Study population

We will include patients in the southwest of the Netherlands, a region with approximately 2 million inhabitants. Participating paramedics have ample experience with the initial management of patients with acute neurological deficits and they received additional training before the start of the study with regards to the study procedures and the use of the prehospital stroke scales. Additional to the prior training, an instruction video is available for all paramedics. Also, during the duration of the study, regular visits are paid to all ambulance stations to provide feedback and address uncertainty or questions of the paramedics. All adult patients with acute neurologic deficit, defined as at least 1 point on the Face-Arm-Speech-Test (FAST), and a suspected diagnosis of stroke by the paramedic, will be included. Patients with a blood glucose level below 2.5 mmol/L will be excluded.

Prehospital stroke scales

We choose five well known prehospital stroke scales to validate: the Los Angeles Motor Scale (LAMS),^{19,20} the Rapid Arterial occlusion Evaluation (RACE),¹⁸ the Cincinnati Stroke Triage Assessment Tool (C-STAT),²¹ the Prehospital Acute Stroke Severity scale (PASS),²² and the Gaze-Face-Arm-Speech-Test (G-FAST).²³ These scales have many similarities in the items that are being used, but there are differences in the scoring systems and the degree of complexity of these scores. In the PRESTO study, we will assess a combination of the items used in these five scales (Table 11.1).

Table 11.1. Overview of the items and corresponding scores used in the prehospital stroke scales.

	LAMS	RACE	C-STAT	PASS	G-FAST	Items collected in this study
Answering questions (age and current month)						
Correctly answers both questions			0	0		0
Correctly answers one question			1*	1		1
Does not correctly answer either question						
Following commands ('close your eyes, 'make a fist')						
Correctly performs both tasks		0†	0			0
Correctly performs one task		1†	1*			1
Does not correctly perform either task		2†				2
Head and gaze deviation						
Normal; able to follow pen or finger to both sides		0	0	0	0	0
Gaze palsy or deviation (total or partial)		1	2	1	1	1
Facial palsy						
Normal and symmetrical movement	0	0			0	0
Mild palsy (flattened nasolabial fold or minor asymmetry in smile)		1			1	1
Moderate to severe palsy	1	2				2
Grip strength						
Normal grip strength	0					0
Weak grip strength	1					1
No grip possible	2					2
Motor function arm						
Normal	0	0	0	0	0	0
Drift (minimal drift with closed eyes)	1	1	1	1	1	1
Mild palsy (arm drifts down within 10 seconds)						
Severe palsy (not able to lift arm)	2	2				2
Motor function leg						
Normal		0				0
Drift (minimal drift with closed eyes)						
Mild palsy (leg drifts down within 5 seconds)		1				1
Severe palsy (not able to lift leg)		2				2
Language						
Normal speech					0	0
Speech problems (dysarthria, language abnormality, or unable to speak)					1	1
Agnosia						
Patient recognises his/her arm and the impairment		0‡				0‡
Does not recognise his/her arm or the impairment		1‡				1‡
Does not recognise his/her arm nor the impairment		2‡				2‡

*1 point if the patient answers at least one question incorrect and does not follow at least one command. †Only scored if right hemiparesis. ‡Only scored if left hemiparesis.

Data collection

Eligible patients presenting with suspected stroke symptoms will be recruited in the ambulance. The items from the prehospital stroke scales will be assessed by the paramedic and entered in a web-based database (LimeSurvey GmbH/Carsten Schmitz, www.limesurvey.org). The paramedic will also enter the transportation number (to link with EMS data and hospital data), the time of symptom onset or last known well (according to patient or bystander), the side of the hemiparesis (if applicable), and the presence of a known neurological deficit on the symptomatic side. Data concerning demographics, vital functions, general neurological examination and transportation times will be collected from the EMS databases.

After arrival in the hospital, patients will receive the usual care. A non-contrast CT scan and additional imaging (eg, CT angiography (CTA), digital subtraction angiography (DSA) and/or CT perfusion) can be performed as part of the regular workup of a suspected stroke. No additional imaging will be performed in the context of this study. Clinical data concerning the medical history, medication use, laboratory results, physical examination, and diagnosis will be collected by chart review. All diagnostic neuro-imaging data and radiology reports will be collected. If applicable, we will also collect information on the given treatment and corresponding treatment times (eg, the door-to-needle time, the door-to-groin time, the imaging-to-treatment time, and the door-in-door-out time of transferred patients).

Follow-up will only be collected for patients with a final diagnosis of ischemic stroke. We will use the outcome registration of the hospitals to collect length of hospital stay, discharge destination, and the modified Rankin Scale (mRS) score after 90 days.

Outcome measures

Primary outcome will be the clinical diagnosis of ischemic stroke with an intracranial LVO in the anterior circulation, defined as an occlusion of the internal carotid artery, the middle cerebral artery segment M1 or M2, or the anterior cerebral artery segment A1 or A2 (assessed on CTA or DSA). Secondary outcome measures include the presence of an LVO in the posterior circulation (vertebral artery or basilar artery); the final diagnosis at hospital discharge; the given treatment (IVT, EVT, or both) and corresponding treatment times; and the functional outcome, measured with the 90-day mRS.

Sample size calculation

At least 100 events (ie, intracranial LVOs) are required for the external validation of predictive models.^{24,25} The annual incidence of suspected ischemic stroke within 6 hours after onset of symptoms is estimated to be 50 per 100,000 people, based on an earlier cohort study.¹⁴ In the catchment area of the participating EMS (approximately 2 million inhabitants), this would imply 1,000 patients every year presenting with stroke symptoms within the 6-hour time window. Of these 1,000 patients, approximately 15% are assumed

to have an ischemic stroke due to an LVO; 31% an ischemic stroke without the presence of an LVO; 9% a transient ischemic attack (TIA); 10% an intracerebral hemorrhage; and 35% a stroke mimic.¹⁴ To reach the required number of 100 stroke patients with an LVO, we will have to include at least (number of cases / prevalence = $100 / 0.15$) 667 patients with stroke symptoms of less than 6 hours. To allow for a 5% loss of follow up, we will aim for a sample size of 700 patients.

After inclusion of the first 500 patients, we will perform an interim analysis to calculate the percentage of LVO in our study population. If necessary, the required sample size will be adjusted based on this information. Although patients presenting after 6 hours will be included in the study, they will not count for the required sample size.

Data analysis plan

After completion of the last inclusion, the data will be checked and the database will be locked for statistical analyses. We will report the absolute numbers and percentages of patients based on the final diagnosis (eg, ischemic stroke, hemorrhagic stroke, TIA or stroke mimic) and, if applicable, the location of the intracranial LVO. For ischemic stroke patients, we will report the given treatment (IVT, EVT, or both) and corresponding treatment times, the number of inter-hospital transfers, and the functional outcome after 90 days. Missing values will be imputed with simple imputation based on the mean or mode (if less than 5% missing) or multiple imputation based on relevant covariates and outcome (if more than 5% missing).

The different prehospital stroke scales will be reconstructed based on the items assessed in the ambulance (Table 11.1). We will validate the prehospital stroke scales for patients presented within 6 hours after symptom onset using a logistic regression model with the presence of an LVO in the anterior circulation as outcome measure. We will analyze the scores both continuously and dichotomised, based on the previously reported cut points in the original studies. Sensitivity and specificity of all cut points will be reported separately. The global performance of the prehospital stroke scales will be expressed as the area under the receiver operator curve (AUC).

Prespecified sensitivity analyses will be performed for patients that presented more than 6 hours after symptom onset, for the separate occlusion locations, and for the presence of an LVO in the posterior circulation. We will also assess the original outcome definitions as defined in each prehospital stroke scale instead of our own primary outcome and we will analyse the correlation between the prehospital stroke scales and the NIHSS assessed at the emergency department. Additional analyses will be performed to predict the probability of treatment with EVT based on the prehospital stroke scales and relevant factors in the medical history, medication use or vital signs.

Patient and public involvement

Patients and public were not involved in the development of the research questions or the design of this study. All study participants and every interested person in the public will have the possibility to read regular project updates on the project website (www.presto-studie.nl).

Duration and current status of the study

The study was registered in The Netherlands Trial Register on November 11, 2018 under number NTR7595 (www.trialregister.nl). The study started on August 13, 2018 in the region Zuid-Holland Zuid and on September 1, 2018 in the region Rotterdam-Rijnmond. Recruitment of patients is ongoing and at the time of submission, April, 2019, 665 patients have been included in the study within 6 hours of symptom onset. In anticipation of a formal interim analysis, first raw data analysis shows a prevalence of 8% LVO in our study population. Based on this information, we increased our sample size to 1250 patients. With the current inclusion rate, we expect to reach the required sample size of 1250 patients by September 2019.

Ethics and dissemination

Ethical aspects and informed consent

This study will be conducted in accordance with the principles of Good Clinical Practice, the Dutch Agreement on Medical Treatment Act (WGBO), and the European General Data Protection Regulation (GDPR). The Institutional Review Board of the Erasmus MC University Medical Center has reviewed the study protocol and confirmed that the Dutch Medical Research Involving Human Subjects Act (WMO) is not applicable.

Acquiring informed consent can be very challenging in the prehospital inclusion of suspected stroke patients. Many patients suffer from a language deficit, anosognosia, or other cognitive symptoms that impede an informed consent procedure, and often there is no (legal) representative of the patient present in the prehospital setting. Furthermore, an adequate informed consent procedure takes time, which is not available in the prehospital setting. Sometimes a deferred consent procedure can be used, but in the context of the WGBO this should be done by the treating physician. Since our unselected population of patients, including many stroke mimics, will spread towards different directions after presentation in the hospital, a disproportionate number of health care providers from a variety of specialisms (eg, neurologists, emergency physicians, internists, cardiologists) should be involved in the research to enable a deferred consent procedure.

The extent of the effort by a large number of health care providers needed to obtain permission from the participating patients is disproportionate to the relatively limited sensitivity of the collected and linked personal data and the related limited intrusion to the

personal privacy. We will therefore use an opt-out procedure in this study. The including paramedic will provide a leaflet with information about the study to the patient or their relatives. In this leaflet, we will explain that some routinely collected data can be collected from the EMS databases and the hospital charts for further analysis. Patients or their relatives are offered the opportunity to object to the use of these data in this study. When a patient or relative objects to study participation, all data will be destroyed and the patient will be excluded from the study.

Dissemination plan

The main study results will be disseminated via publication in an international peer-reviewed journal and presentation at international conferences for stroke and emergency medicine experts. Representatives of the EMS providers and participating hospitals will be given the opportunity to comment on the manuscript and to participate as co-author, following the recommendations of the International Committee of Journal Editors (ICMJE). We plan to disseminate the results of the planned secondary analyses in one or more separate papers.

The best performing scale, or the simplest scale in case of clinical equipoise, will be integrated in a decision model with other clinical characteristics and real-life driving times.²⁶ This model can be implemented in an online tool to improve prehospital triage of patients with suspected stroke symptoms without harming those patients that benefit from rapid IVT in the nearest hospital. Patients eligible for EVT will be directly transported to an endovascular-capable center, which will lead to an increased number of treated patients, reduced treatment times and improved patient outcomes. Moreover, avoiding unnecessary inter-hospital transfers will lead to more efficient use of EMS resources.

References

1. National Institute of Neurological D, Stroke rt PASSG. Tissue plasminogen activator for acute ischemic stroke. *N Engl J Med* 1995; **333**(24): 1581-7.
2. Hacke W, Kaste M, Bluhmki E, et al. Thrombolysis with alteplase 3 to 4.5 hours after acute ischemic stroke. *N Engl J Med* 2008; **359**(13): 1317-29.
3. Lakomkin N, Dharmoon M, Carroll K, et al. Prevalence of large vessel occlusion in patients presenting with acute ischemic stroke: a 10-year systematic review of the literature. *J Neurointerv Surg* 2019; **11**(3): 241-5.
4. Goyal M, Menon BK, van Zwam WH, et al. Endovascular thrombectomy after large-vessel ischaemic stroke: a meta-analysis of individual patient data from five randomised trials. *Lancet* 2016; **387**(10029): 1723-31.
5. Emberson J, Lees KR, Lyden P, et al. Effect of treatment delay, age, and stroke severity on the effects of intravenous thrombolysis with alteplase for acute ischaemic stroke: a meta-analysis of individual patient data from randomised trials. *Lancet* 2014; **384**(9958): 1929-35.
6. Mulder M, Jansen IGH, Goldhoorn RB, et al. Time to Endovascular Treatment and Outcome in Acute Ischemic Stroke: MR CLEAN Registry Results. *Circulation* 2018; **138**(3): 232-40.
7. Saver JL, Goyal M, van der Lugt A, et al. Time to Treatment With Endovascular Thrombectomy and Outcomes From Ischemic Stroke: A Meta-analysis. *JAMA* 2016; **316**(12): 1279-88.
8. Venema E, Boodt N, Berkhemer OA, et al. Workflow and factors associated with delay in the delivery of intra-arterial treatment for acute ischemic stroke in the MR CLEAN trial. *J Neurointerv Surg* 2018; **10**(5): 424-8.
9. Froehler MT, Saver JL, Zaidat OO, et al. Interhospital Transfer Before Thrombectomy Is Associated With Delayed Treatment and Worse Outcome in the STRATIS Registry (Systematic Evaluation of Patients Treated With Neurothrombectomy Devices for Acute Ischemic Stroke). *Circulation* 2017; **136**(24): 2311-21.
10. Smith EE, Kent DM, Bulsara KR, et al. Accuracy of Prediction Instruments for Diagnosing Large Vessel Occlusion in Individuals With Suspected Stroke: A Systematic Review for the 2018 Guidelines for the Early Management of Patients With Acute Ischemic Stroke. *Stroke* 2018; **49**(3): e111-e22.
11. Krebs W, Sharkey-Toppen TP, Cheek F, et al. Prehospital Stroke Assessment for Large Vessel Occlusions: A Systematic Review. *Prehosp Emerg Care* 2018; **22**(2): 180-8.
12. Heldner MR, Hsieh K, Broeg-Morvay A, et al. Clinical prediction of large vessel occlusion in anterior circulation stroke: mission impossible? *J Neurol* 2016; **263**(8): 1633-40.
13. Turc G, Maier B, Naggara O, et al. Clinical Scales Do Not Reliably Identify Acute Ischemic Stroke Patients With Large-Artery Occlusion. *Stroke* 2016; **47**(6): 1466-72.
14. Zhao H, Coote S, Pesavento L, et al. Large Vessel Occlusion Scales Increase Delivery to Endovascular Centers Without Excessive Harm From Misclassifications. *Stroke* 2017; **48**(3): 568-73.
15. Kim JT, Chung PW, Starkman S, et al. Field Validation of the Los Angeles Motor Scale as a Tool for Paramedic Assessment of Stroke Severity. *Stroke* 2017; **48**(2): 298-306.
16. McMullan JT, Katz B, Broderick J, Schmit P, Sucharew H, Adeoye O. Prospective Prehospital Evaluation of the Cincinnati Stroke Triage Assessment Tool. *Prehosp Emerg Care* 2017; **21**(4): 481-8.
17. Noorian AR, Sanossian N, Shkirkova K, et al. Los Angeles Motor Scale to Identify Large Vessel Occlusion: Prehospital Validation and Comparison With Other Screens. *Stroke* 2018; **49**(3): 565-72.
18. Perez de la Ossa N, Carrera D, Gorchs M, et al. Design and validation of a prehospital stroke scale to predict large arterial occlusion: the rapid arterial occlusion evaluation scale. *Stroke* 2014; **45**(1): 87-91.
19. Llanes JN, Kidwell CS, Starkman S, Leary MC, Eckstein M, Saver JL. The Los Angeles Motor Scale (LAMS): a new measure to characterize stroke severity in the field. *Prehosp Emerg Care* 2004; **8**(1): 46-50.

20. Nazliel B, Starkman S, Liebeskind DS, et al. A brief prehospital stroke severity scale identifies ischemic stroke patients harboring persisting large arterial occlusions. *Stroke* 2008; **39**(8): 2264-7.
21. Katz BS, McMullan JT, Sucharew H, Adeoye O, Broderick JP. Design and validation of a prehospital scale to predict stroke severity: Cincinnati Prehospital Stroke Severity Scale. *Stroke* 2015; **46**(6): 1508-12.
22. Hastrup S, Damgaard D, Johnsen SP, Andersen G. Prehospital Acute Stroke Severity Scale to Predict Large Artery Occlusion: Design and Comparison With Other Scales. *Stroke* 2016; **47**(7): 1772-6.
23. Scheitz JF, Abdul-Rahim AH, MacIsaac RL, et al. Clinical Selection Strategies to Identify Ischemic Stroke Patients With Large Anterior Vessel Occlusion: Results From SITS-ISTR (Safe Implementation of Thrombolysis in Stroke International Stroke Thrombolysis Registry). *Stroke* 2017; **48**(2): 290-7.
24. Vergouwe Y, Steyerberg EW, Eijkemans MJ, Habbema JD. Substantial effective sample sizes were required for external validation studies of predictive logistic regression models. *J Clin Epidemiol* 2005; **58**(5): 475-83.
25. Collins GS, Ogundimu EO, Altman DG. Sample size considerations for the external validation of a multivariable prognostic model: a resampling study. *Stat Med* 2016; **35**(2): 214-26.
26. Venema E, Lingsma HF, Chalos V, et al. Personalized Prehospital Triage in Acute Ischemic Stroke. *Stroke* 2019; **50**(2): 313-20.



Chapter 12

General discussion

The overall aim of this thesis was to increase the benefit of endovascular treatment for ischemic stroke by optimizing prediction of outcome and treatment effect, reducing treatment delay, and improving prehospital triage strategies. In this final chapter, I present the main findings, followed by recommendations for clinical practice and future research.

Treat the right patient

Endovascular treatment (EVT) greatly improves the overall functional outcome of patients with ischemic stroke due to a proximal, intracranial large vessel occlusion (LVO) in the anterior circulation.¹ However, to maximize the effect and reduce potential harms or costs, it is important to identify those patients who are likely to benefit and those who are not. Although subgroup analyses are valuable to explore potential differences in the relative treatment effect, they are insufficient for individual treatment decisions.² Patients are placed in a specific subgroup based on a single feature, such as old age or poor collateral score, without taking into account the large variation between individual patients. Also, subgroup analyses are mostly underpowered and prone to false-negative and false-positive results.^{3,4} Therefore, we developed the MR PREDICTS decision tool which combines the effect of multiple clinical and imaging characteristics on outcome and treatment effect simultaneously (Chapters 2 and 3). Included in the model were age, baseline National Institute of Health Stroke Severity (NIHSS) score, systolic blood pressure, history of ischemic stroke, history of diabetes mellitus, degree of pre-stroke disability, prior intravenous treatment with alteplase (IVT), Alberta Stroke Program Early Computed Tomography Score (ASPECTS), location of the occlusion, collateral score, and time from onset to start of treatment, defined as groin puncture. We found large differences in the predicted benefit between individual patients, but almost no harm was predicted. Even in subgroups where previous subgroup analyses showed absence of treatment effect (ie, patients with no collaterals or poor ASPECTS), the model identified a substantial proportion of patients who would benefit.

External validation of MR PREDICTS with data from the HERMES trials and the MR CLEAN Registry showed reasonable to good model performance for the prediction of outcome (Chapter 4). However, prediction of treatment benefit, defined as the difference between the probability of functional independence with and without EVT, is more complex. The observed treatment benefit in HERMES was systematically larger than predicted, especially in the quintile of patients with the lowest predicted benefit. This might be explained by the strict selection of patients who were included in the HERMES trials; patients with unfavorable imaging characteristics were only randomized when a large treatment effect was expected based on other characteristics. Patient selection in the MR CLEAN Registry was less strict, but treatment benefit could not be assessed directly due to the lack of a control group. However, patients with small predicted treatment benefit had low rates of functional independence at 3 months, irrespective of their reperfusion status after the intervention, which might suggest absence of treatment benefit.

Box 12.1 Overview of the main findings per research question***Can we reliably and accurately predict outcome and treatment benefit of endovascular treatment for individual patients?***

We developed and validated a prediction model (MR PREDICTS) that combines multiple baseline clinical and radiological characteristics to predict outcome with and without endovascular treatment. This model performed reasonably well in several validation cohorts and can support clinical judgement in distinguishing between patients who are likely to benefit from endovascular treatment and those who are not. Individual outcome predictions can be improved one day after the intervention using our post-procedural model (MR PREDICTS@24H), which showed excellent discriminative ability.

What are the main causes of prehospital and in-hospital delay of endovascular treatment?

Inter-hospital transfer proved to be the main determinant of treatment delay and a significant cause of poor outcomes even in the Netherlands, where the between-center distances are relatively small. In-hospital workflow processes in the emergency department and the angiosuite should be streamlined to ensure rapid treatment.

How do workflow improvements effect treatment delay and outcome?

Workflow interventions aimed at reducing any delay in the time between symptom onset and treatment, lead to a higher chance of good functional outcome for individual patients, especially when multiple interventions are combined. Effective interventions concern prehospital triage and transportation, in-hospital patient transfers, parallel workflow processes, anesthetic management, teamwork, and feedback on target times.

Which factors should influence the decision to transport individual patients directly to an intervention center?

Prehospital triage decisions should be based on the individual likelihood of large vessel occlusion, transportation times to the nearest primary stroke center and nearest intervention center, and in-hospital workflow characteristics such as the door-in-door-out time and door-to-groin time.

What is the optimal prehospital triage strategy for suspected stroke patients?

The use of a prehospital stroke scale to directly transport patients with suspected large vessel occlusion to an intervention center can improve outcomes, but the optimal triage strategy depends on geographic and organizational characteristics. We constructed a decision model that can be used to inform personalized triage decisions and to evaluate triage policies for specific regions.

Outcome predictions are more accurate when taking into account post-procedural variables such as the degree of reperfusion achieved and the National Institute of Health Stroke Scale (NIHSS) score after 24 hours. We used data available within one day after the intervention to develop MR PREDICTS@24H, a simple tool that combines nine variables to predict functional outcome following EVT (Chapter 5). The severity of stroke symptoms after one day, measured with the NIHSS, was the strongest predictor for functional outcome after 3 months. External validation with MR CLEAN Registry data showed excellent discriminative performance, but suboptimal calibration. The model was therefore updated with a separate intercept to reflect the baseline risk of the population treated in clinical practice. Although this model does not affect the decision to provide EVT, it may guide physicians in personalizing a patients' treatment and rehabilitation plan based on the probability of full recovery.

Several other pre-procedural models for the prediction of functional outcome following EVT exist, with c-statistics ranging between 0.60 and 0.80, but none of these predict outcome with and without EVT to assess the absolute treatment effect.⁵ The most frequently used predictors are age, baseline NIHSS score, and some kind of imaging characteristic. The addition of collateral score improves the predictive performance of existing models, which strengthens the hypothesis that collateral blood flow is a major predictor of outcome and treatment effect.⁶ Two models found potential subgroups of patients with small treatment benefit, based on a poor outcome irrespective of the reperfusion status, but these results need further validation in larger samples.^{7,8} Of the previously published post-procedural models, only three were externally validated and these are all developed with data from the period before the proven effectiveness of EVT.⁹⁻¹¹

Overall, MR PREDICTS and MR PREDICTS@24H showed large variations in anticipated outcome and treatment benefit of individual patients based on multiple characteristics. Both models were developed and validated using recent high-quality datasets of patients treated within 6 hours after stroke onset. The updated models are available online to aid patient selection and outcome prediction in daily clinical practice (www.mrpredicts.com).

Treatment benefit in perspective

With a number needed to treat of 2.6 to reduce disability with one level on the modified Rankin Scale (mRS), EVT ranks among the most effective treatments in the medical field.¹ However, it is important to realize that the total benefit on population-level is not only affected by the size of the treatment effect, but also to a large extent by the number of eligible patients. Treatments with a small effect for the individual patient, might have a large effect on the whole population when applicable to a larger proportion of patients. Organized stroke unit care by a multidisciplinary team, for example, has a small absolute treatment benefit, but leads to a high increase in the number of patients achieving functional independence. EVT has the largest effect on individual patients, but the effect size on

Table 12.1 An illustration of the annual effect of ischemic stroke interventions in the Netherlands, based on 29,082 hospitalizations in 2018.¹²

	Target population	Estimated number of eligible patients (% of total)	Percentage of patients achieving functional independence (mRS 0-2)	Absolute risk difference	Additional risk number of patients achieving functional independence (% of total)
Stroke unit care	All ischemic stroke patients admitted to the hospital	29,082 (100%)	Treatment 49.6% Control 39.8% ¹³	9.8% (NNT = 10)	2,850 (9.8%)
IV treatment with alteplase	Patients with ischemic stroke presenting within 4.5 hours after onset of symptoms or last seen well	5,917 ¹² (20.3%)	Treatment 45.9% Control 39.5% ¹⁴	6.4% (NNT = 16)	333 (1.1%)
Endovascular treatment	Patients with ischemic stroke due to an intracranial LVO in the anterior circulation, presenting within 6 hours after onset of treatment or last seen well	1,906 ¹² (6.6%)	Treatment 46.0% Control 26.5% ¹	19.5% (NNT = 5)	372 (1.3%)
Decompressive hemicraniectomy	Ischemic stroke patients under 60 years with malignant middle cerebral artery infarction	87 (0.3% ¹⁵)	Treatment 26.7% Control 13.9% ¹⁶ (mRS 0-3)	12.8% (NNT = 8)	11 (0.04%)
Early antiplatelet therapy	All ischemic stroke patients presenting within 2 weeks after stroke onset	29,082 (100%)	Treatment 54.3% Control 53.0% ¹⁷	1.3% (NNT = 79)	378 (1.3%)

IV, intravenous; LVO, large vessel occlusion; mRS, modified Rankin Scale; NNT, number needed to treat.

population-level is comparable to the effect of IVT or early treatment with antiplatelet agents (Table 12.1).

Strict selection of patients with a high expected treatment benefit, for example based on favorable imaging characteristics, will increase the treatment effect found in a study, but decreases the total number of patients treated in clinical practice and thereby the overall impact on the stroke population.¹⁸ A clear example is the large treatment effect that was shown in the recent randomized controlled trials (RCTs) on EVT for patients presenting between 6 and 24 hours since symptom onset or last seen well.¹⁹⁻²¹ The reported absolute risk difference of 28-36% for achieving functional independence suggests an enormous effect of treatment in the late time window, but the overall impact is limited due to the small percentage of eligible patients.^{22,23} Further studies on late treatment should therefore focus on expanding the number of treatable patients. Within the 6 hour time window, benefit of EVT is considerable within all subgroups. MR PREDICTS should not be used to select patients with the highest expected benefit, but only to identify patients for whom treatment would likely be futile. Referral of these patients to an intervention center is costly for the health system and unnecessarily inconvenient for their families.

Treat at the right time

Time is brain and rapid initiation of EVT is of great importance to maximize outcomes of ischemic stroke patients with intracranial LVO.^{24,25} Any delay between onset of symptoms and achieving reperfusion directly translates to a decreased chance of a favorable outcome. Early recognition of stroke symptoms and public awareness of the importance of rapid treatment are necessary to avoid patient delay in seeking medical attention.²⁶⁻²⁸ In-hospital determinants of delay in the MR CLEAN trial were presentation during off-hours and the use of general anesthesia (Chapter 6), which was confirmed in several studies.²⁹⁻³³ Analyses of treatment delay in two other RCTs, Endovascular Treatment for Small Core and Proximal Occlusion Ischemic Stroke (ESCAPE) and Solitaire With the Intention for Thrombectomy as Primary Endovascular Treatment (SWIFT PRIME), did not find a significant association with off-hour treatment, but these trials were focused on improving workflow and participating centers were trained in the delivery of fast care and logistics.^{34,35} Prior IVT was not associated with treatment delay, which is likely explained by the streamlined protocols for the administration of alteplase while preparing for CT angiography and transferring the patient to the angiosuite, without waiting for a clinical response.

The most significant modifiable source of delay in the delivery of EVT is the transportation from a primary stroke center to an intervention center.³⁴⁻³⁷ In the MR CLEAN trial, 44% of all patients were transferred, and time from onset to treatment was increased with more than one hour in these patients (Chapter 6). The total delay that occurred in the primary stroke center was much larger, but part of this delay was compensated by a faster work-up in the intervention center. Similar results were found for transferred patients in the MR CLEAN

Registry, reflecting current clinical practice (Chapter 7). The probability of achieving functional independence decreases with 8.5% when a patient was first presented in a primary stroke center. In addition, there are probably also patients who were not transferred to an intervention center because the time window for EVT does not allow for the transfer time. On the other hand, patients without LVO may have benefited from early admission to a primary stroke center by receiving IVT as early as possible, but these patients were not included in the Registry.

Although transferred patients arrive later after stroke onset at the intervention center, they are treated more quickly after arrival than directly admitted patients (Chapters 6 and 7). This finding is probably explained by the fact that part of the diagnostic work-up and treatment, including IVT, is already performed in the primary stroke center. Also, pre-notification allows for the interventional team to prepare in advance. The short door-to-groin time of transferred patients suggests that in-hospital workflow times can also be improved for directly admitted patients by streamlining processes on the emergency department and in the angiosuite.

Workflow improvements

In a systematic review and meta-analysis, we showed that interventions in the workflow of EVT lead to better clinical outcomes, and that this effect increases when multiple interventions are combined (Chapter 8). Strategies that reduced transfer-related delay include requiring the emergency medical services to wait at the primary stroke center until the decision of EVT eligibility has been made, improving cloud-based image sharing with the intervention center, and transporting transfer patients directly to the angiosuite. In regions with very large between-center distances, the additional use of air ambulance services for inter-hospital transportation should also be considered. Another possibility would be to bypass the primary stroke center by using a mobile stroke unit. This is an ambulance with a CT scanner and a specialized team on board, allowing in-ambulance initiation and administration of IVT and direct transportation to the intervention center.³⁸⁻⁴¹ Despite its potential advantages, the clinical benefit and cost-effectiveness of this strategy are unclear and warrant further research.⁴²

Interventions that reduced in-hospital delay were focused on pre-notification by the emergency medical services, direct activation of the entire stroke team, rapid acquisition of brain imaging, and provision of feedback on workflow times to the entire team. Parallel workflow processes for the administration of IVT without delaying EVT might be improved by the use of novel thrombolytic agents such as IV tenecteplase, that do not depend on a 60-minute infusion duration.^{43,44} The 24/7 in-house presence of a vascular neurologist and an interventional team could further improve treatment times during off-hours, but this might be limited by the low number of stroke patients currently eligible for EVT.⁴⁵ It has also been suggested that increased experience, reflected in higher volumes of patients treated

with EVT, leads to shorter treatment times.⁴⁶ Centralization of stroke care could increase the experience of stroke teams and neuro-interventional teams, but should be weighed against the harm of longer transportation times that will emerge. Finally, procedural factors also affect the time between groin puncture and reperfusion, but this is beyond the scope of this thesis.

The significant effect of workflow interventions on time to treatment is consistent with previous studies related to IVT. Studies that successfully reduced treatment times used a combination of several interventions, a multidisciplinary approach, intensive training, and frequent evaluation. After implementation of a national quality improvement initiative and education program by the American Heart Association/American Stroke Association (AHA/ASA), the percentage of patients treated within 60 minutes increased from 29.6% to 53.3% in participating centers.⁴⁷ A Dutch multicenter study proved cost-effectiveness of an intensive implementation program to increase the proportion of patients treated with IVT, although it was lacking power to show an effect on functional outcome.^{48,49} Reported factors of delay in patients with door-to-needle times of >60 minutes include delayed diagnosis, inability to determine eligibility, and difficulties in the management of comorbidities.⁵⁰ A dedicated center in Helsinki systematically implemented multiple interventions over a period of more than 10 years and was able to reduce the median door-to-needle time to 20 minutes.⁵¹ In 2018, the median door-to-needle time in the Netherlands was 24 minutes (IQR 18–34) and the proportion of patients receiving IVT is currently among the highest of Europe.^{12,52}

Target times

Specified target times can be used to improve workflow and compare between centers, but what times are reasonable to achieve in clinical practice? Many different time metrics are proposed in the literature, but one should always keep in mind that the total time interval from symptom onset to reperfusion is relevant for patients.

The door-in-door-out time reflects the total workflow in the primary stroke center from arrival at the emergency department until departure to an intervention center. A recent study suggested a median target time of <45 minutes, which was the top 15th percentile of

Table 12.2 Proposed target times and real-world time metrics of Dutch intervention centers, in minutes.

	Target times		Time metrics in the Netherlands		
	AHA/ASA ⁵⁶	This thesis	MR CLEAN trial	MR CLEAN Registry	Dutch Acute Stroke Audit 2018 ¹²
Door-to-groin time transferred patients	60	25	96 (75–120)	47 (31–70)	26 (17–39)
Door-to-groin time directly admitted patients	90	60	170 (142–205)	104 (80–135)	69 (54–93)

their own door-in-door-out times.⁵³ This target might be too ambitious, considering the median door-in-door-out time of 88 minutes (IQR 68 to 117) in the MR CLEAN Registry and approximately 100 minutes in another large registry in the United States.³⁷ The so called picture-to-puncture time, defined as the time from first imaging to groin puncture, describes the total workflow for transferred patients in both centers, with a proposed target time of <90 minutes.^{54,55} Since this measure includes the driving time between the centers, which is a non-modifiable part of the workflow that depends on the local distribution of hospitals, it should only be used to describe regional workflow characteristics.

For the workflow in the intervention center, the HERMES collaborators proposed a median door-to-groin time of <75 minutes, which was achieved in only 11.5% of the patients treated in the MR CLEAN trial (Chapter 6). It seems more appropriate to determine specific target times for patients that arrive directly at the intervention center and patients that are referred from a primary stroke center, as proposed by the AHA/ASA Target: Stroke Phase III initiative (Table 12.2).

As a last remark, I would like to emphasize that these target times might not be sufficient to ensure rapid treatment of the majority of stroke patients by focusing on the median. By its definition, only 50% of the ischemic stroke population is treated within a median target time, while significant delay may occur in the other 50%. It is therefore always important to assess the interquartile range, and maybe even the 90th percentile, when evaluating treatment times.⁵⁷

Treat in the right place

Prehospital triage of suspected stroke patients involves the trade-off between allowing rapid IVT in the nearest hospital versus avoiding transfer-related delay by direct transportation to an intervention center. We constructed a robust decision tree model to predict the optimal strategy based on individual and regional characteristics (Chapter 9). Factors that influence optimal triage decisions are the likelihood of LVO, estimated driving times to the primary stroke center and the intervention center, and local in-hospital workflow times. This finding is consistent with the results of other (mathematical) models, which used fixed values for the probability of LVO.⁵⁸⁻⁶² Our model is based on the entire range from 0% to 100% likelihood of LVO, and can therefore be used with different prehospital stroke scales as input. It can also be used to assess which differences in sensitivity or specificity between different stroke scales would be clinically relevant.

In Chapter 10, the decision model was combined with a geographic information system analysis to estimate the effect of nationwide, state-level, and county-level implementation of several prehospital triage strategies in the US. We evaluated the current AHA Mission: Lifeline® Stroke algorithm, which recommends direct transportation of suspected LVO patients (based on any positive prehospital stroke scale) to an intervention center when the

additional driving time will not disqualify for IVT and the total driving time from scene to the nearest intervention center is less than 30 minutes.⁶³ The results showed that nationwide implementation of this AHA algorithm prevents death or severe disability in 692 patients per year compared to transportation of all patients to the nearest center, but is suboptimal compared to strategies that allow to bypass the primary stroke center even when leading to delays of 30 minutes or beyond. The current AHA algorithm is only preferred in scenarios with very short driving times, a low prevalence of LVO, very efficient workflow times in the primary stroke centers, or a low number of additional patients without LVO accepted in the intervention centers. A modified algorithm without a time limit for the transportation of LVO-suspected patients would be optimal in the majority of states and could greatly improve outcomes of the ischemic stroke population in the United States.

There is no randomized data available to support these findings, and although clinical data showed better outcomes for patients directly transported to an intervention center, these studies did not take into account the negative effect of bypassing the primary stroke center for patients without LVO.⁶⁴ An ongoing RCT in Catalonia, Spain, will be the first to directly compare standard transportation to the nearest hospital to prehospital triage with the Rapid Arterial Occlusion Evaluation (RACE) score.⁶⁵ This study might provide real-world evidence for the use of a prehospital stroke scale, but the results will only be applicable to regions with similar population density, geographic distribution of hospitals and in-hospital workflow times. Modeling studies will therefore always be required to translate trial results to regions with other geographic and organizational characteristics.

Prehospital stroke scales

Because the likelihood of LVO is one of the main drivers of triage decisions, early prehospital identification of patients at risk of having an LVO in the ambulance is of utmost importance. Numerous prehospital stroke scales have been proposed, but none of these scales predict LVO with both a high sensitivity and a high specificity.^{66,67} Most validation studies used retrospective reconstruction of the scores with data from the in-hospital assessment by experienced physicians, which is not representative of clinical practice. The RACE score was prospectively validated in a prehospital setting and showed a sensitivity of 84% and a specificity of 60% at a cutoff at ≥ 5 points.⁶⁸ The prevalence of LVO among suspected stroke patients was 20%, resulting in a positive predictive value of 34%. When transporting all patients with a positive RACE score to an intervention center, there will be two non-LVO patients (false positives) for every patient with LVO stroke (true positives).

Further prospective, prehospital validation of prehospital stroke scales is needed to compare their accuracy. Choosing the optimal prehospital stroke scale for triage of suspected stroke patients depends on the false positives (ie, patients without LVO that are transported to an intervention center) and false negatives (ie, patients with LVO that are transported to the primary stroke center). These numbers are based on the sensitivity and specificity of a scale,

but also on the prevalence of LVO in the local population. Previous studies reported a prevalence among ischemic stroke patients ranging from 7% to 60%.⁶⁹ Most physicians will prefer a scale with a high sensitivity to increase the number of patients with LVO stroke directly transported to an intervention center. However, the specificity of a score reflects the number of false positives, which is a very relevant measure when financial resources or the capacity of intervention centers are limited.

We designed the PRESTO study to prospectively evaluate and compare several prehospital stroke scales used by emergency medical services in a broad population of suspected stroke patients (Chapter 11). The simultaneous in-field validation of several prehospital stroke scales allows for direct comparison of their accuracy in an unselected population. Patient inclusion was finished in September 2019 and the final results of this study, including core lab assessed imaging data, are expected soon.

Strengths and limitations

The main analyses in this thesis were performed with data from the MR CLEAN trial, which is considered to have the broadest inclusion criteria of the recent trials, and the MR CLEAN Registry, one of the largest cohorts of consecutive patients treated with EVT. Both cohorts are representative of patients treated in routine clinical practice without mandatory clinical or imaging-based selection criteria, which makes them especially suitable for the development of clinical prediction models. Patient outcomes in MR CLEAN and the Registry were less favorable than in the HERMES cohort, but this might be explained by the strict selection criteria, the inclusion of high-quality centers with ample experience, and the fast workflow times of the studies included in HERMES, which may not correspond to real-life clinical practice. On the other hand, the relatively small sample size of the MR CLEAN trial (n=500) limits generalizability, especially for small subgroups (eg, patients with M2 occlusions or low ASPECTS). The Registry, although larger in sample size, is limited by the lack of a control group and the high number of missing values, including missing outcomes.

Experienced clinicians are able to consider multiple factors when predicting outcome of individual patients with ischemic stroke, but several cognitive biases may occur and physicians' predictive accuracy showed to be suboptimal compared to prediction models that combine the information of large datasets.⁷⁰⁻⁷³ Well developed and externally validated models can be used to aid the clinician when making treatment decisions or rehabilitation plans. MR PREDICTS showed modest predictive ability, especially in the clinical trial population, but the main predictor effects were comparable in the different cohorts, and the model was updated using data from daily clinical practice. The effect of MR PREDICTS on patient selection will be limited, due to the substantial treatment effect and small potential harm of EVT. More difficult treatment decision arise when patients present after 6 hours of symptom onset, but there is currently insufficient data to develop an accurate model for late window patients.

The analyses of treatment delay and prehospital transportation times were performed with data from the Netherlands, which is a small, densely populated country compared to other Western regions. The average transfer time in our study was less than 30 minutes, which limits the generalizability of our results to other regions. It is likely that the negative effect of inter-hospital transfer is even larger in countries with large between-center distances. Therefore, we used the wide variety of geographical areas in the United States to further evaluate triage strategies.

The driving force of our prehospital model was to optimize patient outcomes, but the effect of triage strategies extends far beyond functional outcome alone. Unnecessary transportations to a more distant intervention center are inconvenient for patients and their relatives, and may lead to problems with resources and crowding at the intervention centers. This can be reduced by choosing a higher cut point or a more specific prehospital stroke scale. The shift of patient volume will also have economic consequences for the primary stroke centers that will receive and treat less stroke patients. This will create a substantial incentive for these centers to improve their workflow times, which makes transportation to a primary stroke center less harmful.

Comparison with acute myocardial infarction

The current developments in the field of vascular neurology have many similarities with previous developments in cardiology due to the large similarity between ischemic stroke and myocardial infarction (MI). During MI, the blood flow in a coronary artery is suddenly blocked, causing ischemia of the heart muscle. An electrocardiogram, which records the electrical activity of the heart, is used to identify patients with an ST elevation MI, which indicates salvageable tissue in distress. These patients benefit from direct restoration of the blood flow by a percutaneous coronary intervention (PCI), while patients without ST elevation are initially treated with medication.

Treatment of MI is time-critical, just like ischemic stroke treatment, and delayed initiation of PCI is associated with increased mortality.⁷⁴ In 1999, the American Heart Association recommended a door-to-balloon time of <90 minutes to increase the chance of good outcome.^{75,76} This target time was achieved in only half of the American patients in 2005, which increased to 76% of the patients in 2008 due to the efforts of a national quality campaign implementing several workflow interventions.⁷⁷ A study on specific strategies showed a significant effect of rapid activation and availability of the catheterization laboratory, 24/7 availability of an attending cardiologist, and the provision of real-time feedback to the entire team.⁷⁸ Early emergency department activation of the catheterization team can reduce door-to-balloon times with >30 minutes.^{79,80}

As in ischemic stroke, the transfer of patients from a non-intervention center to an intervention center is associated with significant delay. Several strategies to improve workflow of the emergency medical services, emergency department and in-hospital care

Table 12.3 Potential interventions to reduce delay between symptom onset and start of treatment.

Time metric	Prehospital targets	In-hospital targets
Symptom onset to EMS notification	<ul style="list-style-type: none"> Increasing public awareness Training family doctors 	
Scene to first hospital	<ul style="list-style-type: none"> Improving stroke recognition of EMS staff 	
Scene to intervention center	<ul style="list-style-type: none"> Bypassing the primary stroke center when the likelihood of LVO is high Use of mobile stroke unit to identify patients with LVO 	
Door-to-imaging	<ul style="list-style-type: none"> Prehospital notification Direct EMS transfer to CT scanner 	<ul style="list-style-type: none"> Direct activation of the stroke team
Door-in-door-out	<ul style="list-style-type: none"> Enlisting EMS crew to remain at primary stroke center until the definitive treatment decision is made 	<ul style="list-style-type: none"> Rapid imaging acquisition Parallel treatment with IV alteplase Direct contact with intervention center Initiation of transfers without awaiting approval of the intervention center in patients clearly meeting well-defined criteria Providing time targets with frequent feedback to the entire team
Transfer time	<ul style="list-style-type: none"> Air transportation 	
Door-to-groin puncture (general)		<ul style="list-style-type: none"> Rapid imaging acquisition Parallel treatment with IV alteplase Extending the on-site presence of stroke team members during off-hours Early activation of the interventional team More experienced teams Providing time targets with frequent feedback to the entire team
Door-to-groin puncture (transferred patients)	<ul style="list-style-type: none"> Direct EMS transfer to the angiosuite 	<ul style="list-style-type: none"> Electronic transmission of CT images Early activation of the interventional team
Angiosuite to groin puncture		<ul style="list-style-type: none"> Protocols to preselect devices Ready-to-use equipment tray Early consultation of the anesthetist Avoiding general anesthesia if possible

CT, computed tomography; EMS, emergency medical services; IV, intravenous; LVO, large vessel occlusion.

were independently associated with decreased door-in-door-out times, but the prehospital processes were the most important factor.⁸¹ Despite the similarities between the treatment of MI and stroke, the prehospital triage of MI patients is less complex. MI patients that should receive rapid treatment with PCI are easily identified in the ambulance using an electrocardiogram. This enables direct transportation of these patients to a PCI center, while the triage of suspected stroke patients is complicated by the fact that the diagnosis can only be confirmed with a CT angiography in the hospital. Our efforts should therefore be aimed on identifying eligible EVT patients in the prehospital setting, for example using prehospital stroke scales or a mobile stroke unit.

Recommendations for clinical practice

The MR PREDICTS decision tool and MR PREDICTS@24H are online available for use in clinical practice (www.mrpredicts.com), and emphasize the importance of taking various factors into account when making treatment decisions. EVT should never be withheld based on a single characteristic, and patients directly presenting at the intervention center within 6 hours after symptom onset should be treated as soon as possible, considering the large treatment effect and the small potential harm. I suggest that the decision tool may be particularly relevant when patients have to be transferred to an intervention center and there is doubt whether EVT will be beneficial. Transferring patients with low expected benefit causes unnecessary health care costs and inconvenience for patients and their families. Clinical judgement remains important, especially when there are rare, but strongly predictive characteristics not captured by the model, for example an abnormal anatomy of the circle of Willis or fluctuations in the course of symptoms.

Further reorganization of acute stroke care should be aimed on structuring workflow processes to ensure rapid treatment independent of the time or day of presentation. Multiple interventions can be combined to maximize the potential effect (Table 12.3). Ambitious but feasible time targets should be determined to motivate hospital staff and policy makers to improve in-hospital workflow. These time metrics should be monitored in every hospital and reported back to the stroke team regularly. At this moment, I would suggest that primary stroke centers in the Netherlands aim for a median door-in-door-out time of <45 minutes, which will only be possible when the ambulance waits at the emergency department until the result of CT angiography. Intervention centers should strive for a door-to-groin time of <60 minutes for directly admitted patients and <25 minutes for transferred patients.

Prehospital triage of suspected stroke patients is essential to reduce the number of inter-hospital transfers. I would recommend to extend the current AHA triage algorithm for prehospital triage in the US and allow more delay when bypassing a primary stroke center. No universal time threshold can be given, because this strongly depends on geographic circumstances and organization of stroke care, but it seems beneficial to allow at least 30-45

minutes additional driving time for suspected LVO patients. Regional protocols should be further optimized based on local driving times, financial resources, and capacity of the intervention centers. The online tool, based on the decision model described in Chapter 9, can be used to inform these decisions.

Currently, there is insufficient evidence to recommend one prehospital stroke scale over another, but one should at least use a scale that was prospectively validated in the prehospital setting (eg, RACE or the Cincinnati Prehospital Stroke Severity Scale (CPSSS)), and that is easy to use for emergency medical services.^{68,82} The choice for a prehospital stroke scale should not depend on sensitivity alone, because adequate specificity is required to limit the number of patients unnecessarily transported to an intervention center. Assessment of the individual likelihood of LVO based on a continuous scale instead of one fixed threshold, may further improve the effectiveness of prehospital triage.

Recommendations for future research

Despite the huge impact of EVT on clinical outcome, still more than half of the treated patients are deceased or disabled at 3 months after stroke.¹ This emphasizes the devastating effect of the disease and warrants innovations for further improvement of the current treatment options.

Patient selection is especially important for patients presenting more than 6 hours after onset of symptoms or last seen well, because EVT might be futile or even harmful in many of these patients. However, the currently available trial data only includes a strictly selected patient population and is therefore insufficient to use for predictive modeling. Large databases of patients treated in the late time window are required to develop a robust model that can aid patient selection. Results of the ongoing MR CLEAN LATE trial (ISRCTN19922220), which assesses the use of more liberal selection criteria without restrictions on infarct core size, might be useful in identifying which late window patients benefit from treatment.

Machine learning methods, or artificial intelligence, are increasingly popular in the medical field and are suggested to improve personalized predictions by discovering data patterns and associations through automated algorithms.^{83,84} However, these algorithms require very large datasets for which clinical data sources are often not sufficient.⁸⁵ The relatively small number of variables available might explain why machine learning methods did not outperform conventional logistic regression in the MR CLEAN Registry.⁸⁶ Also, machine learning models do not provide further insight in the underlying pathophysiological mechanisms and cannot be adjusted for the use in different settings or populations. Therefore, I believe that machine learning is mostly valuable for pattern recognition tasks, for example for the automatic interpretation of neuroimaging. Prediction modeling should focus on the development of simple, robust models and adequate external validation of existing models.

It is important to acknowledge that patients with a poor prognosis might still benefit from treatment, and therefore MR PREDICTS estimates outcome with and without EVT. However, evaluation of treatment benefit brings a number of challenges. In Chapter 4, I used the c-for-benefit to evaluate the discriminative performance of the predicted treatment benefit. While the traditional c-statistic assesses whether a model can discriminate between patients with a low risk versus high risk of the outcome, this novel measure focuses on the whether the model can separate patients who are very likely to benefit from treatment and those who are not.⁸⁷ Because the real treatment benefit of a patient cannot be observed, the c-for-benefit uses the outcomes of patient pairs that were matched on predicted benefit but discordant for treatment assignment. This method needs further evaluation and new benchmark values should to be defined to guide the interpretation of this promising measure.

A large number of prehospital stroke scales have been developed, and researchers should now focus on thorough validation of these scales in the prehospital setting, rather than the development of new scales. The future results of the PRESTO study will allow direct comparison of several prehospital stroke scales to improve triage decisions. Further decision modeling combined with geographic information systems can provide guidance for triage decisions in different settings. It can also be used to determine in which regions the implementation of a mobile stroke unit could be beneficial. Integration of a decision model with a GPS-controlled navigation application will facilitate personalized triage based on real-time information about driving times and in-hospital workflow times.

Final remarks

Although the overall effect of EVT is overwhelming, there are many targets to further improve outcome of individual patients. In this thesis, I developed and validated prediction models to predict outcome and treatment benefit; identified determinants of treatment delay and interventions to improve workflow; and applied decision-analyses to evaluate and optimize prehospital triage strategies.

The MR PREDICTS decision tool is the only model that estimates individual benefit of EVT and thereby improves patient selection. It performed reasonably well in a an international trial population and a large cohort of patients treated in daily clinical practice. Treatment benefit is substantial in the vast majority of patients, but absence of treatment effect is suggested in a small number of patients with multiple unfavorable characteristics. Rapid initiation of EVT is essential to improve patient outcomes, and requires the implementation of various prehospital and in-hospital workflow interventions. Inter-hospital transfer is the main determinant of treatment delay and should be avoided by transporting suspected stroke patients directly to an intervention center, but the optimal triage strategy varies based on regional characteristics. The future results of the PRESTO study will provide more information on the real-world performance of prehospital stroke scales, allowing further improvement of prehospital decision-making.

Box 12.2 Overview of the main recommendations.***Treat the right patient***

Endovascular treatment should never be withheld from an individual patient based on a single characteristic. The MR PREDICTS decision tool can be used to evaluate the effect of multiple patient and imaging characteristics simultaneously, when there is uncertainty about the treatment indication or when patients need to be transferred to an intervention center. Patient selection is particularly important for patients presenting more than 6 hours after symptom onset, but large databases of not strictly selected patients are required to develop a robust model for the late time window.

Treat at the right time

Rapid initiation of endovascular treatment is essential to improve patient outcomes, and requires the implementation of various prehospital and in-hospital workflow interventions. Acute stroke care should be reorganized to reduce any delay between stroke onset and start of treatment. Every effort is needed to streamline workflow processes, improve team work and communication, and reduce the number of inter-hospital transfers. Door-in-door-out times and door-to-groin times should be monitored and reported back to the stroke team regularly.

Treat in the right place

A validated prehospital stroke scale should be used to identify patients that benefit from direct transportation to an intervention center. The choice of a specific prehospital stroke scale and the maximum allowed delay when bypassing the primary stroke center depends on local geographic characteristics, resources and preferences. Decision modeling will therefore be needed to translate results of clinical studies on prehospital triage to different settings and regions.

References

1. Goyal M, Menon BK, van Zwam WH, et al. Endovascular thrombectomy after large-vessel ischaemic stroke: a meta-analysis of individual patient data from five randomised trials. *Lancet* 2016; **387**(10029): 1723-31.
2. Dahabreh IJ, Hayward R, Kent DM. Using group data to treat individuals: understanding heterogeneous treatment effects in the age of precision medicine and patient-centered evidence. *Int J Epidemiol* 2016; **45**(6): 2184-93.
3. Brookes ST, Whitley E, Egger M, Smith GD, Mulheran PA, Peters TJ. Subgroup analyses in randomized trials: risks of subgroup-specific analyses; power and sample size for the interaction test. *J Clin Epidemiol* 2004; **57**(3): 229-36.
4. Brookes ST, Whitley E, Peters TJ, Mulheran PA, Egger M, Davey Smith G. Subgroup analyses in randomised controlled trials: quantifying the risks of false-positives and false-negatives. *Health Technol Assess* 2001; **5**(33): 1-56.
5. Raza SA, Rangaraju S. A Review of Pre-Intervention Prognostic Scores for Early Prognostication and Patient Selection in Endovascular Management of Large Vessel Occlusion Stroke. *Interv Neurol* 2018; **7**(3-4): 171-81.
6. Ryu CW, Kim BM, Kim HG, et al. Optimizing Outcome Prediction Scores in Patients Undergoing Endovascular Thrombectomy for Large Vessel Occlusions Using Collateral Grade on Computed Tomography Angiography. *Neurosurgery* 2019; **85**(3): 350-8.
7. Rangaraju S, Aghaebrahim A, Streib C, et al. Pittsburgh Response to Endovascular therapy (PRE) score: optimizing patient selection for endovascular therapy for large vessel occlusion strokes. *J Neurointerv Surg* 2015; **7**(11): 783-8.
8. Sarraj A, Albright K, Barreto AD, et al. Optimizing prediction scores for poor outcome after intra-arterial therapy in anterior circulation acute ischemic stroke. *Stroke* 2013; **44**(12): 3324-30.
9. Natarajan SK, Dandona P, Karmon Y, et al. Prediction of adverse outcomes by blood glucose level after endovascular therapy for acute ischemic stroke. *J Neurosurg* 2011; **114**(6): 1785-99.
10. Prabhakaran S, Jovin TG, Tayal AH, et al. Posttreatment variables improve outcome prediction after intra-arterial therapy for acute ischemic stroke. *Cerebrovasc Dis* 2014; **37**(5): 356-63.
11. Rangaraju S, Liggins JT, Aghaebrahim A, et al. Pittsburgh outcomes after stroke thrombectomy score predicts outcomes after endovascular therapy for anterior circulation large vessel occlusions. *Stroke* 2014; **45**(8): 2298-304.
12. Dutch Institute for Clinical Auditing. Dutch Acute Stroke Audit (DASA) Jaarrapportage 2018. <https://dica.nl/jaarrapportage-2018/dasa> (accessed May 1 2020).
13. Stroke Unit Trialists' Collaboration. Organised inpatient (stroke unit) care for stroke. *Cochrane Database Syst Rev* 2013; (9): CD000197.
14. Wardlaw JM, Murray V, Berge E, del Zoppo GJ. Thrombolysis for acute ischaemic stroke. *Cochrane Database Syst Rev* 2014; (7): CD000213.
15. Rahme R, Curry R, Kleindorfer D, et al. How often are patients with ischemic stroke eligible for decompressive hemicraniectomy? *Stroke* 2012; **43**(2): 550-2.
16. Alexander P, Heels-Ansdell D, Siemieniuk R, et al. Hemicraniectomy versus medical treatment with large MCA infarct: a review and meta-analysis. *BMJ Open* 2016; **6**(11): e014390.
17. Sandercock PA, Counsell C, Tseng MC, Cecconi E. Oral antiplatelet therapy for acute ischaemic stroke. *Cochrane Database Syst Rev* 2014; (3): CD000029.
18. Goyal M, Jadhav AP. Denominator fallacy revisited. *Journal of Neurointerventional Surgery* 2017; **9**(10): 915-6.
19. Albers GW, Marks MP, Kemp S, et al. Thrombectomy for Stroke at 6 to 16 Hours with Selection by Perfusion Imaging. *N Engl J Med* 2018; **378**(8): 708-18.
20. Nogueira RG, Jadhav AP, Haussen DC, et al. Thrombectomy 6 to 24 Hours after Stroke with a Mismatch between Deficit and Infarct. *N Engl J Med* 2018; **378**(1): 11-21.

21. Albers GW. Late Window Paradox. *Stroke* 2018; **49**(3): 768-71.
22. Nannoni S, Strambo D, Sirimarco G, et al. Eligibility for late endovascular treatment using DAWN, DEFUSE-3, and more liberal selection criteria in a stroke center. *J Neurointerv Surg* 2019.
23. Jadhav AP, Desai SM, Kenmuir CL, et al. Eligibility for Endovascular Trial Enrollment in the 6- to 24-Hour Time Window: Analysis of a Single Comprehensive Stroke Center. *Stroke* 2018; **49**(4): 1015-7.
24. Mulder M, Jansen IGH, Goldhoorn RB, et al. Time to Endovascular Treatment and Outcome in Acute Ischemic Stroke: MR CLEAN Registry Results. *Circulation* 2018; **138**(3): 232-40.
25. Saver JL, Goyal M, van der Lugt A, et al. Time to Treatment With Endovascular Thrombectomy and Outcomes From Ischemic Stroke: A Meta-analysis. *JAMA* 2016; **316**(12): 1279-88.
26. Fladt J, Meier N, Thilemann S, et al. Reasons for Prehospital Delay in Acute Ischemic Stroke. *J Am Heart Assoc* 2019; **8**(20): e013101.
27. Lacy CR, Suh DC, Bueno M, Kostis JB. Delay in presentation and evaluation for acute stroke: Stroke Time Registry for Outcomes Knowledge and Epidemiology (S.T.R.O.K.E.). *Stroke* 2001; **32**(1): 63-9.
28. Jin H, Zhu S, Wei JW, et al. Factors associated with prehospital delays in the presentation of acute stroke in urban China. *Stroke* 2012; **43**(2): 362-70.
29. Vukasinovic I, Darcourt J, Guenego A, et al. "Real life" impact of anesthesia strategy for mechanical thrombectomy on the delay, recanalization and outcome in acute ischemic stroke patients. *J Neuroradiol* 2019; **46**(4): 238-42.
30. Powers CJ, Dornbos D, 3rd, Mlynash M, et al. Thrombectomy with Conscious Sedation Compared with General Anesthesia: A DEFUSE 3 Analysis. *AJNR Am J Neuroradiol* 2019; **40**(6): 1001-5.
31. Hoepner R, Weber R, Reimann G, et al. Stroke admission outside daytime working hours delays mechanical thrombectomy and worsens short-term outcome. *Int J Stroke* 2019; **14**(5): 517-21.
32. Kim JG, Choi JC, Kim DJ, et al. Effect of the Number of Neurointerventionalists on Off-Hour Endovascular Therapy for Acute Ischemic Stroke Within 12 Hours of Symptom Onset. *J Am Heart Assoc* 2019; **8**(21): e011933.
33. Hinsenveld WH, de Ridder IR, van Oostenbrugge RJ, et al. Workflow Intervals of Endovascular Acute Stroke Therapy During On- Versus Off-Hours: The MR CLEAN Registry. *Stroke* 2019; **50**(10): 2842-50.
34. Goyal M, Jadhav AP, Bonafe A, et al. Analysis of Workflow and Time to Treatment and the Effects on Outcome in Endovascular Treatment of Acute Ischemic Stroke: Results from the SWIFT PRIME Randomized Controlled Trial. *Radiology* 2016; **279**(3): 888-97.
35. Menon BK, Sajobi TT, Zhang Y, et al. Analysis of Workflow and Time to Treatment on Thrombectomy Outcome in the Endovascular Treatment for Small Core and Proximal Occlusion Ischemic Stroke (ESCAPE) Randomized, Controlled Trial. *Circulation* 2016; **133**(23): 2279-86.
36. Shah S, Xian Y, Sheng S, et al. Use, Temporal Trends, and Outcomes of Endovascular Therapy After Interhospital Transfer in the United States. *Circulation* 2019; **139**(13): 1568-77.
37. Froehler MT, Saver JL, Zaidat OO, et al. Interhospital Transfer Before Thrombectomy Is Associated With Delayed Treatment and Worse Outcome in the STRATIS Registry (Systematic Evaluation of Patients Treated With Neurothrombectomy Devices for Acute Ischemic Stroke). *Circulation* 2017; **136**(24): 2311-21.
38. Ebinger M, Winter B, Wendt M, et al. Effect of the use of ambulance-based thrombolysis on time to thrombolysis in acute ischemic stroke: a randomized clinical trial. *JAMA* 2014; **311**(16): 1622-31.
39. Rajan SS, Baraniuk S, Parker S, Wu TC, Bowry R, Grotta JC. Implementing a mobile stroke unit program in the United States: why, how, and how much? *JAMA Neurol* 2015; **72**(2): 229-34.
40. Kostopoulos P, Walter S, Haass A, et al. Mobile stroke unit for diagnosis-based triage of persons with suspected stroke. *Neurology* 2012; **78**(23): 1849-52.
41. Walter S, Kostopoulos P, Haass A, et al. Diagnosis and treatment of patients with stroke in a mobile stroke unit versus in hospital: a randomised controlled trial. *Lancet Neurol* 2012; **11**(5): 397-404.

42. Fassbender K, Grotta JC, Walter S, Grunwald IQ, Ragoschke-Schumm A, Saver JL. Mobile stroke units for prehospital thrombolysis, triage, and beyond: benefits and challenges. *Lancet Neurol* 2017; **16**(3): 227-37.
43. Parsons M, Spratt N, Bivard A, et al. A randomized trial of tenecteplase versus alteplase for acute ischemic stroke. *N Engl J Med* 2012; **366**(12): 1099-107.
44. Coutts SB, Dubuc V, Mandzia J, et al. Tenecteplase-tissue-type plasminogen activator evaluation for minor ischemic stroke with proven occlusion. *Stroke* 2015; **46**(3): 769-74.
45. Hill M, Glenn BA, Reese BJ, Morrow B. Recommendations for Endovascular Care of Stroke Patients. *Interv Neurol* 2018; **7**(1-2): 65-90.
46. Gupta R, Horev A, Nguyen T, et al. Higher volume endovascular stroke centers have faster times to treatment, higher reperfusion rates and higher rates of good clinical outcomes. *J Neurointerv Surg* 2013; **5**(4): 294-7.
47. Fonarow GC, Zhao X, Smith EE, et al. Door-to-needle times for tissue plasminogen activator administration and clinical outcomes in acute ischemic stroke before and after a quality improvement initiative. *JAMA* 2014; **311**(16): 1632-40.
48. Dirks M, Niessen LW, van Wijngaarden JD, et al. Promoting thrombolysis in acute ischemic stroke. *Stroke* 2011; **42**(5): 1325-30.
49. Dirks M, Baeten SA, Dippel DW, et al. Real-life costs and effects of an implementation program to increase thrombolysis in stroke. *Neurology* 2012; **79**(6): 508-14.
50. Kamal N, Sheng S, Xian Y, et al. Delays in Door-to-Needle Times and Their Impact on Treatment Time and Outcomes in Get With The Guidelines-Stroke. *Stroke* 2017; **48**(4): 946-54.
51. Meretoja A, Strbian D, Mustanoja S, Tatlisumak T, Lindsberg PJ, Kaste M. Reducing in-hospital delay to 20 minutes in stroke thrombolysis. *Neurology* 2012; **79**(4): 306-13.
52. Aguiar de Sousa D, von Martial R, Abilleira S, et al. Access to and delivery of acute ischaemic stroke treatments: A survey of national scientific societies and stroke experts in 44 European countries. *Eur Stroke J* 2019; **4**(1): 13-28.
53. Choi PMC, Tsoi AH, Pope AL, et al. Door-in-Door-Out Time of 60 Minutes for Stroke With Emergent Large Vessel Occlusion at a Primary Stroke Center. *Stroke* 2019; **50**(10): 2829-34.
54. Sun CH, Nogueira RG, Glenn BA, et al. "Picture to puncture": a novel time metric to enhance outcomes in patients transferred for endovascular reperfusion in acute ischemic stroke. *Circulation* 2013; **127**(10): 1139-48.
55. McTaggart RA, Ansari SA, Goyal M, et al. Initial hospital management of patients with emergent large vessel occlusion (ELVO): report of the standards and guidelines committee of the Society of NeuroInterventional Surgery. *J Neurointerv Surg* 2017; **9**(3): 316-23.
56. American Heart Association. Introducing Target: Stroke Phase III. 2020. <https://www.heart.org/en/professional/quality-improvement/target-stroke/introducing-target-stroke-phase-iii> (accessed April 1 2020).
57. Holodinsky JK, Kamal N, Wilson AT, Hill MD, Goyal M. Workflow in Acute Stroke: What Is the 90th Percentile? *Stroke* 2017; **48**(3): 808-12.
58. Milne MS, Holodinsky JK, Hill MD, et al. Drip 'n Ship Versus Mothership for Endovascular Treatment: Modeling the Best Transportation Options for Optimal Outcomes. *Stroke* 2017; **48**(3): 791-4.
59. Schlemm E, Ebinger M, Nolte CH, Endres M, Schlemm L. Optimal Transport Destination for Ischemic Stroke Patients With Unknown Vessel Status: Use of Prehospital Triage Scores. *Stroke* 2017; **48**(8): 2184-91.
60. Benoit JL, Khatri P, Adeoye OM, et al. Prehospital Triage of Acute Ischemic Stroke Patients to an Intravenous tPA-Ready versus Endovascular-Ready Hospital: A Decision Analysis. *Prehosp Emerg Care* 2018; **22**(6): 722-33.
61. Holodinsky JK, Williamson TS, Demchuk AM, et al. Modeling Stroke Patient Transport for All Patients With Suspected Large-Vessel Occlusion. *JAMA Neurol* 2018; **75**(12): 1477-86.
62. Xu Y, Parikh NS, Jiao B, Willey JZ, Boehme AK, Elkind MSV. Decision Analysis Model for Prehospital Triage of Patients With Acute Stroke. *Stroke* 2019; **50**(4): 970-7.
63. American Heart Association. Mission: Lifeline Stroke Severity-Based Stroke Triage Algorithm for EMS. <https://www.heart.org/en/professional/quality-improvement/mission->

- lifeline/missionlifeline-stroke 2020 (accessed April 1 2020).
64. Chartrain AG, Shoirah H, Jauch EC, Mocco J. A review of acute ischemic stroke triage protocol evidence: a context for discussion. *J Neurointerv Surg* 2018; **10**(11): 1047-52.
 65. Abilleira S, Perez de la Ossa N, Jimenez X, et al. Transfer to the Local Stroke Center versus Direct Transfer to Endovascular Center of Acute Stroke Patients with Suspected Large Vessel Occlusion in the Catalan Territory (RACECAT): Study protocol of a cluster randomized within a cohort trial. *Int J Stroke* 2019; **14**(7): 734-44.
 66. Smith EE, Kent DM, Bulsara KR, et al. Accuracy of Prediction Instruments for Diagnosing Large Vessel Occlusion in Individuals With Suspected Stroke: A Systematic Review for the 2018 Guidelines for the Early Management of Patients With Acute Ischemic Stroke. *Stroke* 2018; **49**(3): e111-e22.
 67. Krebs W, Sharkey-Toppen TP, Cheek F, et al. Prehospital Stroke Assessment for Large Vessel Occlusions: A Systematic Review. *Prehosp Emerg Care* 2018; **22**(2): 180-8.
 68. Carrera D, Gorchs M, Querol M, et al. Revalidation of the RACE scale after its regional implementation in Catalonia: a triage tool for large vessel occlusion. *J Neurointerv Surg* 2019; **11**(8): 751-6.
 69. Lakomkin N, Dhmoon M, Carroll K, et al. Prevalence of large vessel occlusion in patients presenting with acute ischemic stroke: a 10-year systematic review of the literature. *J Neurointerv Surg* 2019; **11**(3): 241-5.
 70. Geurts M, de Kort FAS, de Kort PLM, van Tuijl JH, Kappelle LJ, van der Worp HB. Predictive accuracy of physicians' estimates of outcome after severe stroke. *PLoS One* 2017; **12**(9): e0184894.
 71. Weimar C, König IR, Kraywinkel K, Ziegler A, Diener HC, German Stroke Study C. Age and National Institutes of Health Stroke Scale Score within 6 hours after onset are accurate predictors of outcome after cerebral ischemia: development and external validation of prognostic models. *Stroke* 2004; **35**(1): 158-62.
 72. Saposnik G, Cote R, Mamdani M, et al. JURASSIC: accuracy of clinician vs risk score prediction of ischemic stroke outcomes. *Neurology* 2013; **81**(5): 448-55.
 73. Fargen KM, Leslie-Mazwi TM, Chen M, Hirsch JA. Physician, know thyself: implicit and explicit decision-making for mechanical thrombectomy in stroke. *J Neurointerv Surg* 2020.
 74. Berger PB, Ellis SG, Holmes DR, Jr., et al. Relationship between delay in performing direct coronary angioplasty and early clinical outcome in patients with acute myocardial infarction: results from the global use of strategies to open occluded arteries in Acute Coronary Syndromes (GUSTO-IIb) trial. *Circulation* 1999; **100**(1): 14-20.
 75. Ryan TJ, Antman EM, Brooks NH, et al. 1999 update: ACC/AHA guidelines for the management of patients with acute myocardial infarction. A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee on Management of Acute Myocardial Infarction). *J Am Coll Cardiol* 1999; **34**(3): 890-911.
 76. Cannon CP, Gibson CM, Lambrew CT, et al. Relationship of symptom-onset-to-balloon time and door-to-balloon time with mortality in patients undergoing angioplasty for acute myocardial infarction. *JAMA* 2000; **283**(22): 2941-7.
 77. Bradley EH, Nallamothu BK, Herrin J, et al. National efforts to improve door-to-balloon time results from the Door-to-Balloon Alliance. *J Am Coll Cardiol* 2009; **54**(25): 2423-9.
 78. Bradley EH, Herrin J, Wang Y, et al. Strategies for reducing the door-to-balloon time in acute myocardial infarction. *N Engl J Med* 2006; **355**(22): 2308-20.
 79. Mahler SA, Chan HY, Carden DL, Wolcott C, Conrad SA. Emergency Department Activation of Interventional Cardiology to Reduce Door-to-Balloon Time. *West J Emerg Med* 2010; **11**(4): 363-6.
 80. Singer AJ, Shembekar A, Visram F, et al. Emergency department activation of an interventional cardiology team reduces door-to-balloon times in ST-segment-elevation myocardial infarction. *Ann Emerg Med* 2007; **50**(5): 538-44.
 81. Glickman SW, Lytle BL, Ou FS, et al. Care processes associated with quicker door-in-door-out times for patients with ST-elevation-myocardial infarction requiring transfer: results from a statewide regionalization program. *Circ Cardiovasc Qual Outcomes* 2011; **4**(4): 382-8.

82. Kummer BR, Gialdini G, Sevush JL, Kamel H, Patsalides A, Navi BB. External Validation of the Cincinnati Prehospital Stroke Severity Scale. *J Stroke Cerebrovasc Dis* 2016; **25**(5): 1270-4.
83. Kamal H, Lopez V, Sheth SA. Machine Learning in Acute Ischemic Stroke Neuroimaging. *Front Neurol* 2018; **9**: 945.
84. Rajkomar A, Dean J, Kohane I. Machine Learning in Medicine. *N Engl J Med* 2019; **380**(14): 1347-58.
85. van der Ploeg T, Austin PC, Steyerberg EW. Modern modelling techniques are data hungry: a simulation study for predicting dichotomous endpoints. *BMC Med Res Methodol* 2014; **14**: 137.
86. van Os HJA, Ramos LA, Hilbert A, et al. Predicting Outcome of Endovascular Treatment for Acute Ischemic Stroke: Potential Value of Machine Learning Algorithms. *Front Neurol* 2018; **9**: 784.
87. van Klaveren D, Steyerberg EW, Serruys PW, Kent DM. The proposed 'concordance-statistic for benefit' provided a useful metric when modeling heterogeneous treatment effects. *J Clin Epidemiol* 2018; **94**: 59-68.



Appendices

Summary | Samenvatting

Acknowledgments

Dankwoord

List of publications

PhD portfolio

About the author



Summary

The effect of endovascular treatment (EVT) for ischemic stroke due to a proximal, intracranial large vessel occlusion (LVO) varies between individual patients. Treatment benefit, defined as the difference between the chance of a good outcome with and without treatment, is affected by differences in clinical characteristics at baseline. For example, the treatment effect strongly declines over time because a larger part of the brain tissue is already irreversibly damaged. Early initiation of EVT is therefore associated with better functional outcome, and time to treatment should be reduced to improve chances of good recovery. A possibility to reduce delay is to directly transport patients with suspected LVO to an intervention center capable of performing EVT, instead of bringing them to a primary stroke center for the first evaluation. However, this strategy might be harmful for stroke patients without LVO, who only benefit from rapid treatment with intravenous thrombolytics in the nearest hospital. The likelihood of LVO in suspected stroke patients can be assessed in the ambulance using a prehospital stroke scale, but the optimal triage strategy requires a trade-off between the harm of delaying intravenous thrombolytics versus the potential benefit of rapid EVT.

The overall aim of this thesis was to increase the benefit of EVT by optimizing prediction of outcome and treatment effect (**Part I**), reducing treatment delay (**Part II**), and improving prehospital triage strategies (**Part III**).

The specific research questions were:

1. Which are the right patients to treat?
 - a. Can we reliably and accurately predict outcome and treatment benefit of EVT for individual patients?
2. How can we treat patients at the right time?
 - a. What are the main causes of prehospital and in-hospital delay of EVT?
 - b. How do workflow improvements effect treatment delay and outcome?
3. How to direct patients to the right place?
 - a. Which factors should influence the decision to transport individual patients directly to an intervention center?
 - b. What is the optimal prehospital triage strategy for suspected stroke patients?

Part I – Treat the right patient

The first part of this thesis concerns the development and external validation of a prediction model for functional outcome and treatment benefit of individual patients (MR PREDICTS), and a post-procedural model to improve outcome prediction for patients treated with EVT at one day after the intervention (MR PREDICTS@24H).

Chapters 2 and 3 describe the development of the MR PREDICTS decision tool with data from the MR CLEAN trial. Eleven baseline patient and imaging characteristics were included in the model, together with three interaction terms to account for differential relative treatment effects. An external validation with data from the IMS III trial showed a moderate ability of the model to distinguish between patients with low and high likelihood of a good outcome. Predicted treatment benefit varied substantially between individual patients, and clinically relevant benefit was predicted for some patients from subgroups in which no effect of EVT was expected, such as those with no or poor collaterals.

In **Chapter 4**, we performed external validation of MR PREDICTS with data from multiple international trials within the HERMES collaboration, including patients from different healthcare systems and countries. Based on these results, the model was updated and validated again in the MR CLEAN Registry, representing daily clinical practice in the Netherlands. The model performed reasonably well in both cohorts. Median predicted treatment benefit of routinely treated patients in the Registry was 10.6% (IQR 6.4% to 14.5%). A small group of patients in the Registry with low predicted treatment benefit had poor outcomes irrespective of reperfusion status, suggesting potential absence of treatment benefit.

MR PREDICTS@24H was developed with data from the HERMES collaboration, as described in **Chapter 5**. This prognostic model includes nine pre- and post-procedural characteristics, of which the National Institute of Health Stroke Severity scale at 24 hours after EVT was the most predictive of functional outcome at 3 months. External validation with data from the MR CLEAN Registry showed excellent discriminative performance, although the observed probability of survival was lower than predicted. This was resolved by providing separate intercepts that reflect the differences in outcome between the trial population and the broad group of routinely treated patients.

Part II – Treat at the right time

Chapter 6 describes the effect of workflow characteristics on the time intervals in the MR CLEAN trial. Patients admitted during off-hours had an average delay of 23 minutes before the start of EVT, and patients treated under general anesthesia were treated 19 minutes

later from arrival at the emergency department. Prior administration of intravenous thrombolytics was not related with any significant treatment delay. Most importantly, almost half of the patients (44%) were transferred from a primary stroke center, and those patients received EVT on average 65 minutes later than patients directly admitted to an intervention center.

To further explore the effect of inter-hospital transfer on time to treatment and functional outcome after EVT in daily clinical practice, we performed an analysis with data from the MR CLEAN Registry (**Chapter 6**). More than half of the routinely treated patients (54%) were transferred from a primary stroke center and these patients had an average delay of 57 minutes between first presentation and start of EVT. The negative impact of this delay on functional outcome of transferred patients was reflected in an absolute decrease of 8.5% in the probability of achieving functional independence after 3 months.

Chapter 7 consists of a systematic review and meta-analysis on the effect of workflow improvements for EVT, including 51 studies on different types of workflow interventions. The most frequently reported interventions were aimed at using local anesthesia or conscious sedation instead of general anesthesia, reducing in-hospital patient transfers, and optimizing the prehospital workflow. The mean time to treatment was 26 minutes shorter for patients in the intervention group, and simultaneous implementation of multiple interventions even reduced treatment delay with 50 minutes.

Part III – Treat in the right place

Because inter-hospital transfer is the most important factor of modifiable treatment delay, we constructed a decision tree model to determine the optimal prehospital transportation strategy of suspected stroke patients (**Chapter 8**). This model combines the likelihood of LVO in an individual patient with local driving times and in-hospital workflow times. Direct transportation to an intervention center was beneficial for patients with a high risk of having an LVO, especially in scenarios with short between-center distances. The optimal triage decision was also affected by the in-hospital workflow characteristics of both the primary stroke center (door-to-needle time and door-in-door-out time) and the intervention center (door-to-groin time).

In **Chapter 9**, our decision model was integrated in a geographic information system analysis to assess the effect of prehospital triage strategies for the United States. Evaluation of the current triage algorithm of the American Heart Association showed that this strategy is suboptimal for most states and counties. Modified versions of this algorithm, allowing for more time delay when bypassing a primary stroke center, showed large improvement of outcomes of the ischemic stroke population.

These triage strategies rely on prehospital stroke scales to assess the likelihood of LVO in suspected stroke patients, but prospective, prehospital validation of these scales is lacking. Therefore, I present the study protocol of PRESTO (prehospital triage of patients with suspected stroke symptoms) in **Chapter 10**. This prospective, multicenter, observational cohort study will evaluate the performance of several prehospital stroke scales based on the initial assessment in the ambulance. The results of this study will provide a unique insight in the characteristics of an unselected group of patients with suspected stroke in the prehospital setting.

Discussion

Although the overall effect of EVT is overwhelming, there are many targets to further improve outcome of individual patients. In this thesis, I developed and validated prediction models to predict outcome and treatment benefit; identified determinants of treatment delay and interventions to improve workflow; and applied decision-analyses to evaluate and optimize prehospital triage strategies.

Based on my findings, I recommend to never withheld EVT from an individual patient based on a single characteristic. The vast majority of patients will benefit when treated within 6 hours after stroke onset, but the MR PREDICTS decision tool can be used to evaluate the effect of multiple patient and imaging characteristics when there is uncertainty about the treatment indication or when patients need to be transferred to an intervention center. Patient selection is particularly important for patients presenting more than 6 hours after symptom onset, but large databases of not strictly selected patients are required to develop a robust model for the late time window.

Rapid initiation of EVT is essential to improve patient outcomes, and requires the implementation of various prehospital and in-hospital workflow interventions. Acute stroke care should be reorganized to reduce any delay between stroke onset and start of EVT. Every effort is needed to streamline workflow processes, improve team work and communication, and reduce the number of inter-hospital transfers. Ambitious target times should be set for the door-in-door-out time in the primary stroke center and the door-to-groin time in the intervention center. These time metrics should be monitored and reported back to the stroke team regularly.

To improve prehospital triage, a validated prehospital stroke scale should be used to identify patients that benefit from direct transportation to an intervention center. The choice of a specific prehospital stroke scale and the maximum allowed delay when bypassing the primary stroke center depends on local geographic characteristics, resources and preferences. The future results of the PRESTO study will provide more information on the

real-world performance of prehospital stroke scales, allowing further improvement of prehospital decision-making. However, decision modeling will always be needed to translate results of clinical studies to different settings and regions.



Samenvatting

Patiënten met een herseninfarct op basis van een proximale intracraniële occlusie in de voorste circulatie kunnen baat hebben bij intra-arteriële therapie waarbij het stolsel wordt verwijderd (trombectomie). De winst van deze behandeling, gedefinieerd als het verschil tussen de kans op een goed resultaat met en zonder behandeling, varieert tussen individuele patiënten en wordt beïnvloed door verschillen in klinische factoren. Zo neemt het effect van intra-arteriële therapie sterk af in de eerste uren na het ontstaan van de symptomen, doordat met het verstrijken van de tijd een groter deel van het hersenweefsel onomkeerbaar beschadigd raakt. Een snelle start van de behandeling wordt geassocieerd met een betere functionele uitkomst. De tijd tussen het ontstaan van de symptomen en het starten van intra-arteriële therapie moet daarom zoveel mogelijk worden gereduceerd om de kans op goed herstel te vergroten.

Een mogelijkheid om vertraging te voorkomen is om patiënten rechtstreeks naar een interventiecentrum te brengen waar intra-arteriële therapie kan worden uitgevoerd, in plaats van naar een primair stroke centrum voor de initiële opvang. Deze triage strategie kan echter schadelijk zijn voor patiënten met een herseninfarct zonder intracraniële occlusie, aangezien die patiënten alleen profiteren van snelle intraveneuze trombolysen in het dichtstbijzijnde ziekenhuis. Bij patiënten met de verdenking op een herseninfarct moet daarom worden ingeschat hoe waarschijnlijk het is dat zij een intracraniële occlusie hebben. Dit kan in de ambulance worden gedaan met behulp van een prehospital stroke score. Optimale prehospital triage vereist echter een afweging tussen de schade die kan ontstaan door het uitstellen van intraveneuze trombolysen versus het potentiële voordeel van snelle intra-arteriële therapie.

Het algemene doel van dit proefschrift was om de winst van intra-arteriële therapie voor patiënten met een herseninfarct te vergroten door voorspellingen van uitkomst en behandelingseffect te optimaliseren (**Deel I**), vertraging van de behandeling te verminderen (**Deel II**) en prehospital triage strategieën te verbeteren (**Deel III**).

De specifieke onderzoeksvragen waren:

1. Welke patiënten moeten worden behandeld?
 - a. Kunnen we uitkomsten en de winst van intra-arteriële therapie voor individuele patiënten betrouwbaar en nauwkeurig voorspellen?
2. Hoe kunnen we patiënten op het juiste moment behandelen?
 - a. Wat zijn de belangrijkste oorzaken van vertraging in de prehospital setting en in het ziekenhuis?
 - b. Hoe beïnvloeden workflow interventies de tijd tot behandeling en de uitkomsten van behandelde patiënten?

3. Hoe kunnen we patiënten naar de juiste plaats leiden?
 - a. Welke factoren zouden van invloed moeten zijn op de beslissing om individuele patiënten rechtstreeks naar een interventiecentrum te vervoeren?
 - b. Wat is de optimale prehospital triage strategie voor patiënten met de verdenking op een herseninfarct?

Deel I - Behandel de juiste patiënt

Het eerste deel van dit proefschrift betreft de ontwikkeling en externe validatie van een predictiemodel voor functionele uitkomst en behandelwinst voor individuele patiënten (MR PREDICTS), en een post-procedureel model om de voorspelling van uitkomsten te verbeteren één dag na de interventie (MR PREDICTS@24H).

Hoofdstukken 2 en 3 beschrijven de ontwikkeling van het MR PREDICTS model met gegevens van de MR CLEAN studie. Elf patiëntkarakteristieken en radiologische kenmerken werden in het model opgenomen, samen met drie interactietermen om rekening te houden met differentiële relatieve behandel-effecten. Een externe validatie met gegevens van de IMS III studie toonde een redelijk voorspellend vermogen van het model in het onderscheiden van patiënten met een lage en een hoge kans op een goede uitkomst. De voorspelde winst van de behandeling varieerde aanzienlijk tussen individuele patiënten. Er werd ook baat van de behandeling voorspeld voor sommige patiënten uit subgroepen waarin eerder geen effect van intra-arteriële therapie werd verwacht, zoals patiënten met een zeer slechte collaterale bloedvoorziening.

In **Hoofdstuk 4** heb ik een externe validatie van MR PREDICTS uitgevoerd met behulp van zes internationale, gerandomiseerde studies die zijn aangesloten bij de HERMES collaboration. Op basis van deze resultaten is het model geactualiseerd en opnieuw gevalideerd in de MR CLEAN Registry, een database waarin alle patiënten zijn opgenomen die in de dagelijkse praktijk in Nederland worden behandeld. Het model presteerde redelijk goed in beide cohorten. Voor de routinematig behandelde patiënten in de Registry werd een absolute toename van 10.6% voorspeld in de kans op goed functioneel herstel (interkwartiel range 6.4% tot 14.5%). Bij een kleine groep patiënten in de Registry werd geen baat van de behandeling voorspeld en een slechte uitkomst waargenomen ongeacht de reperfusiestatus, wat duidt op een mogelijk afwezig behandel-effect.

MR PREDICTS@24H is ontwikkeld met gegevens van de HERMES collaboration, zoals beschreven in **Hoofdstuk 5**. Dit prognostische model omvat 9 pre- en post-procedurele kenmerken, waarvan de National Institute of Health Stroke Severity score afgenomen 24 uur na de interventie de belangrijkste voorspellende factor was van de functionele uitkomst na 3 maanden. Externe validatie met data van de MR CLEAN Registry toonde een uitstekend

discriminerend vermogen, maar de waargenomen overlevingskans in dit cohort was lager dan voorspeld. Dit werd opgelost door twee afzonderlijke intercepts te presenteren die het verschil in uitkomst weerspiegelen tussen de studiepopulatie en de brede groep van patiënten die in de dagelijkse praktijk behandeld wordt.

Deel II - Behandel op het juiste moment

Hoofdstuk 6 beschrijft het effect van workflow factoren op de tijd tot behandeling in de MR CLEAN studie. Patiënten die buiten kantooruren werden opgenomen zijn gemiddeld 23 minuten later behandeld dan patiënten binnen kantooruren, en patiënten onder algehele narcose zijn gemiddeld 19 minuten later behandeld. Het toedienen van intraveneuze trombolysie was niet gerelateerd aan enige significante vertraging van de behandeling. De belangrijkste bevinding was dat bijna de helft van de patiënten (44%) werd overgeplaatst vanuit een primair stroke centrum. Deze overgeplaatste patiënten werden gemiddeld 65 minuten later behandeld dan patiënten die rechtstreeks werden opgenomen in een interventiecentrum.

Om het effect van overplaatsingen tussen ziekenhuizen op de tijd tot behandeling en functionele uitkomst na intra-arteriële therapie in de dagelijkse klinische praktijk te onderzoeken, heb ik ook een analyse uitgevoerd met data van de MR CLEAN Registry (**Hoofdstuk 6**). Meer dan de helft van de routinematig behandelde patiënten in Nederland (54%) werd overgebracht vanuit een primair stroke centrum en deze patiënten hadden een gemiddelde vertraging van 57 minuten tussen de eerste presentatie en het starten van de intra-arteriële therapie. Het negatieve effect hiervan kwam tot uiting in een absolute afname van 8.5% in de kans op goed functioneel herstel na 3 maanden.

Hoofdstuk 7 bestaat uit een systematische review en meta-analyse naar het effect van workflow interventies, waarin 51 studies van verschillende type interventies geïncorporeerd werden. De meest frequent gerapporteerde interventies waren gericht op het gebruik van lokale anesthesie of sedatie in plaats van algehele narcose, het verminderen van overplaatsingen van patiënten en het optimaliseren van de prehospital workflow. De gemiddelde tijd tot de behandeling was 26 minuten korter voor patiënten in de interventiegroep en gelijktijdige implementatie van meerdere interventies verkortte de tijd tot behandeling zelfs met 50 minuten.

Deel III - Behandel op de juiste plaats

Omdat overplaatsingen tussen ziekenhuizen de belangrijkste factor zijn in de vertraging van intra-arteriële therapie, heb ik een besliskundig model opgesteld om de optimale

prehospitale strategie te bepalen voor patiënten waarbij een herseninfarct vermoed wordt (**Hoofdstuk 8**). Dit model combineert de waarschijnlijkheid van een intracraniale occlusie bij een individuele patiënt met lokale reistijden en doorlooptijden in de regionale ziekenhuizen. Direct vervoer naar een interventiecentrum was gunstig voor patiënten waarbij een intracraniale occlusie waarschijnlijk is, vooral in scenario's met korte afstanden tussen de verschillende centra. De optimale triage strategie werd ook beïnvloed door de doorlooptijden in het primaire stroke centrum (deur-tot-naald tijd en deur-in-deur-uit tijd) en het interventiecentrum (deur-tot-lies tijd).

In **Hoofdstuk 9** werd dit besliskundige model geïntegreerd in een geografisch informatiesysteem om het effect van prehospital triage strategieën in de Verenigde Staten te beoordelen. Het huidige triage algoritme van de American Heart Association bleek onvoldoende effectief te zijn in de meerderheid van de staten en provincies. Aangepaste versies van dit algoritme, die meer vertraging toestaan bij het vervoer naar een interventiecentrum, hadden een groot positief effect op de uitkomsten van de Amerikaanse patiëntenpopulatie.

Prehospital stroke scores worden gebruikt om de waarschijnlijkheid van een intracraniale occlusie te beoordelen in de ambulance, maar prospectieve, prehospital validatie van deze scores ontbreekt. Daarom presenteer ik in **Hoofdstuk 10** het studieprotocol van PRESTO (prehospital triage van patiënten met de verdenking op een herseninfarct). Deze prospectieve, multicenter, observationele cohortstudie zal het voorspellend vermogen van verschillende prehospital stroke scores evalueren op basis van de initiële beoordeling in de ambulance. De resultaten van deze studie zullen tevens inzicht bieden in de ongeselecteerde groep van patiënten met de verdenking op een herseninfarct.

Discussie

Hoewel het algehele effect van intra-arteriële therapie overweldigend is, zijn er veel mogelijkheden om de uitkomsten van individuele patiënten met een herseninfarct verder te verbeteren. In dit proefschrift heb ik predictiemodellen ontwikkeld en gevalideerd om functionele uitkomst en behandelwinst te voorspellen; determinanten die leiden tot vertraging van de behandeling en interventies om de workflow te verbeteren geïdentificeerd; en besliskundige analyses toegepast om prehospital triage strategieën te evalueren en te optimaliseren.

Op basis van mijn bevindingen raad ik aan om nooit een individuele patiënt intra-arteriële therapie te onthouden op basis van één enkele eigenschap. Het MR PREDICTS model kan worden gebruikt om het effect van meerdere patiëntkarakteristieken en radiologische kenmerken tegelijkertijd te evalueren, wanneer er onzekerheid bestaat over de

behandelindicatie of wanneer patiënten moeten worden overgeplaatst naar een interventiecentrum. Patiëntselectie is vooral belangrijk voor patiënten die zich meer dan 6 uur na het begin van de symptomen presenteren, maar er zijn grote databases met niet strikt geselecteerde patiënten nodig om een robuust model voor het late tijdvenster te ontwikkelen.

Een snelle start van intra-arteriële therapie is essentieel om de uitkomsten van patiënten te verbeteren en vereist implementatie van verschillende workflow interventies in de prehospital setting en in het ziekenhuis. De acute beroertezorg moet worden gereorganiseerd om de vertraging tussen het begin van een herseninfarct en de start van de behandeling te verminderen. Er moet alles aan gedaan worden om de logistieke processen te stroomlijnen, samenwerking en communicatie te verbeteren en het aantal overplaatsingen tussen ziekenhuizen te verminderen. Het is raadzaam om de doorstroomtijden te monitoren en regelmatig aan het stroke team te rapporteren.

Om de prehospital triage te verbeteren, moet een gevalideerde prehospital stroke score worden gebruikt om patiënten te identificeren die baat hebben bij direct transport naar een interventiecentrum. De keuze voor een specifieke prehospital stroke score en de maximaal toegestane vertraging wanneer wordt omgereden naar een interventiecentrum hangt af van geografische kenmerken, beschikbare capaciteit van de ziekenhuizen en lokale voorkeuren. De toekomstige resultaten van de PRESTO studie zullen naar verwachting meer informatie verschaffen over de werkelijke prestaties van prehospital stroke scores, zodat de besluitvorming kan worden geoptimaliseerd. Besliskundige modellen zullen echter altijd nodig zijn om de resultaten van klinische studies te vertalen naar verschillende regionale omstandigheden.



Acknowledgments

MR CLEAN Registry Investigators – group authors

Executive committee

Diederik W.J. Dippel¹; Aad van der Lugt²; Charles B.L.M. Majoie³; Yvo B.W.E.M. Roos⁴; Robert J. van Oostenbrugge⁵; Wim H. van Zwam⁶; Jelis Boiten¹⁴; Jan Albert Vos⁸

Study coordinators

Ivo G.H. Jansen³; Maxim J.H.L. Mulder^{1,2}; Robert-Jan B. Goldhoorn^{5,6}; Kars C.J. Compagne²; Manon Kappelhof³; Josje Brouwer⁴; Sanne J. den Hartog^{1,2,40}; Wouter H. Hinsenveld^{5,6}

Local principal investigators

Diederik W.J. Dippel¹; Bob Roozenbeek¹; Aad van der Lugt²; Adriaan C.G.M. van Es²; Charles B.L.M. Majoie³; Yvo B.W.E.M. Roos⁴; Bart J. Emmer³; Jonathan M. Coutinho⁴; Wouter J. Schonewille⁷; Jan Albert Vos⁸; Marieke J.H. Wermer⁹; Marianne A.A. van Walderveen¹⁰; Julie Staals⁵; Robert J. van Oostenbrugge⁵; Wim H. van Zwam⁶; Jeannette Hofmeijer¹¹; Jasper M. Martens¹²; Geert J. Lycklama à Nijeholt¹³; Jelis Boiten¹⁴; Sebastiaan F. de Bruijn¹⁵; Lukas C. van Dijk¹⁶; H. Bart van der Worp¹⁷; Rob H. Lo¹⁸; Ewoud J. van Dijk¹⁹; Hieronymus D. Boogaarts²⁰; J. de Vries²²; Paul L.M. de Kort²¹; Julia van Tuijl²¹; Jo P. Peluso²⁶; Puck Fransen²²; Jan S.P. van den Berg²²; Boudewijn A.A.M. van Hasselt²³; Leo A.M. Aerden²⁴; René J. Dallinga²⁵; Maarten Uyttenboogaart²⁸; Omid Eschgi²⁹; Reinoud P.H. Bokkers²⁹; Tobien H.C.M.L. Schreuder³⁰; Roel J.J. Heijboer³¹; Koos Keizer³²; Lonneke S.F. Yo³³; Heleen M. den Hertog²²; Emiel J.C. Sturm³⁵; Paul J.A.M. Brouwers³⁴

Imaging assessment committee

Charles B.L.M. Majoie³(chair); Wim H. van Zwam⁶; Aad van der Lugt²; Geert J. Lycklama à Nijeholt¹³; Marianne A.A. van Walderveen¹⁰; Marieke E.S. Sprengers³; Sjoerd F.M. Jenniskens²⁷; René van den Berg³; Albert J. Yoo³⁸; Ludo F.M. Beenen³; Alida A. Postma⁶; Stefan D. Roosendaal³; Bas F.W. van der Kallen¹³; Ido R. van den Wijngaard¹³; Adriaan C.G.M. van Es²; Bart J. Emmer³; Jasper M. Martens¹²; Lonneke S.F. Yo³³; Jan Albert Vos⁸; Joost Bot³⁶; Pieter-Jan van Doormaal²; Anton Meijer²⁷; Elyas Ghariq¹³; Reinoud P.H. Bokkers²⁹; Marc P. van Proosdij³⁷; G. Menno Krietemeijer³³; Jo P. Peluso²⁶; Hieronymus D. Boogaarts²⁰; Rob Lo¹⁸; Dick Gerrits³⁵; Wouter Dinkelaar²; Auke P.A. Appelman²⁹; Bas Hammer¹⁶; Sjoert Pegge²⁷; Anouk van der Hoorn²⁹; Saman Vinke²⁰

Writing committee

Diederik W.J. Dippel¹(chair); Aad van der Lugt²; Charles B.L.M. Majoie³; Yvo B.W.E.M. Roos⁴; Robert J. van Oostenbrugge⁵; Wim H. van Zwam⁶; Geert J. Lycklama à Nijeholt¹³; Jelis Boiten¹⁴; Jan Albert Vos⁸; Wouter J. Schonewille⁷; Jeannette Hofmeijer¹¹; Jasper M. Martens¹²; H. Bart van der Worp¹⁷; Rob H. Lo¹⁸

Adverse event committee

Robert J. van Oostenbrugge⁵(chair);Jeannette Hofmeijer¹¹;H. Zwenneke Flach²³

Trial methodologist

Hester F. Lingsma⁴⁰

Research nurses / local trial coordinators

Naziha el Ghannouti¹;Martin Sterrenberg¹;Wilma Pellikaan⁷;Rita Sprengers⁴;Marjan Elfrink¹¹;Michelle Simons¹¹;Marjolein Vossers¹²;Joke de Meris¹⁴;Tamara Vermeulen¹⁴;Annet Geerlings¹⁹;Gina van Vemde²²;Tiny Simons³⁰;Gert Messchendorp²⁸;Nynke Nicolaij²⁸;Hester Bongenaar³²;Karin Bodde²⁴;Sandra Kleijn³⁴;Jasmijn Lodico³⁴; Hanneke Droste³⁴;Maureen Wollaert⁵;Sabrina Verheesen⁵;D. Jeurissen⁵;Erna Bos⁹;Yvonne Drabbe¹⁵;Michelle Sandiman¹⁵;Nicoline Aaldering¹¹;Berber Zweedijk¹⁷;Jocova Vervoort²¹;Eva Ponjee²²;Sharon Romviel¹⁹;Karin Kanselaar¹⁹;Denn Barning¹⁰

PhD / Medical students:

Esmee Venema⁴⁰; Vicky Chalos^{1,40}; Ralph R. Geuskens³; Tim van Straaten¹⁹;Saliha Ergezen¹; Roger R.M. Harmsma¹; Daan Muijres¹; Anouk de Jong¹;Olvert A. Berkhemer^{1,3,6};Anna M.M. Boers^{3,39}; J. Huguet³;P.F.C. Groot³;Marieke A. Mens³;Katinka R. van Kranendonk³;Kilian M. Treurniet³;Manon L. Tolhuisen^{3,39};Heitor Alves³;Annick J. Weterings³;Eleonora L.F. Kirkels³;Eva J.H.F. Voogd¹¹;Lieve M. Schupp³;Sabine Collette^{28,29};Adrien E.D. Groot⁴;Natalie E. LeCouffe⁴;Praneeta R. Konduri³⁹;Haryadi Prasetya³⁹;Nerea Arrarte-Terreros³⁹;Lucas A. Ramos³⁹

List of affiliations

Department of Neurology¹, Radiology², Public Health⁴⁰,
Erasmus MC University Medical Center;
Department of Radiology and Nuclear Medicine³, Neurology⁴,
Biomedical Engineering & Physics³⁹, Amsterdam UMC, University of Amsterdam, Amsterdam;
Department of Neurology⁵, Radiology⁶, Maastricht University Medical Center and
Cardiovascular Research Institute Maastricht (CARIM);
Department of Neurology⁷, Radiology⁸, Sint Antonius Hospital, Nieuwegein;
Department of Neurology⁹, Radiology¹⁰, Leiden University Medical Center;
Department of Neurology¹¹, Radiology¹², Rijnstate Hospital, Arnhem;
Department of Radiology¹³, Neurology¹⁴, Haaglanden MC, the Hague;
Department of Neurology¹⁵, Radiology¹⁶, HAGA Hospital, the Hague;
Department of Neurology¹⁷, Radiology¹⁸, University Medical Center Utrecht;
Department of Neurology¹⁹, Neurosurgery²⁰, Radiology²⁷,
Radboud University Medical Center, Nijmegen;
Department of Neurology²¹, Radiology²⁶, Elisabeth-TweeSteden ziekenhuis, Tilburg;
Department of Neurology²², Radiology²³, Isala Klinieken, Zwolle;

Department of Neurology²⁴, Radiology²⁵, Reinier de Graaf Gasthuis, Delft;
 Department of Neurology²⁸, Radiology²⁹, University Medical Center Groningen;
 Department of Neurology³⁰, Radiology³¹, Atrium Medical Center, Heerlen;
 Department of Neurology³², Radiology³³, Catharina Hospital, Eindhoven;
 Department of Neurology³⁴, Radiology³⁵, Medical Spectrum Twente, Enschede;
 Department of Radiology³⁶, Amsterdam UMC, Vrije Universiteit van Amsterdam, Amsterdam;
 Department of Radiology³⁷, Noordwest Ziekenhuisgroep, Alkmaar;
 Department of Radiology³⁸, Texas Stroke Institute, Texas, United States of America

PRESTO Investigators – group authors

Executive committee

Diederik W.J. Dippel¹; Bob Roozenbeek^{1,2}; Henk Kerkhoff³; Hester F. Lingsma⁴; Aad van der Lugt²; Adriaan C.G.M van Es²; Anouk D. Rozeman³; Walid Moudrous⁵; Frédérique H. Vermeij⁶

Study coordinators

Esmee Venema^{1,3}; Martijne H.C. Duvekot^{1,2,3,4}

Steering committee

Diederik W.J. Dippel¹; Bob Roozenbeek^{1,2}; Hester F. Lingsma⁴; Aad van der Lugt²; Adriaan C.G.M. van Es²; Henk Kerkhoff³; Anouk D. Rozeman³; Walid Moudrous⁵; Frédérique H. Vermeij⁶; Kees C.L. Alblas⁶; Laus J.M.M. Mulder⁷; Annemarie D. Wijnhoud⁸; Lisette Maasland⁹; Roeland P.J. van Eijkelenburg¹⁰; Marileen Biekart¹¹; M.L. Willeboer¹²; Bianca Buijck¹³

Imaging committee

Bob Roozenbeek^{1,2}; Henk Kerkhoff³; Aad van der Lugt²; Adriaan C.G.M. van Es²

Imaging core laboratory

Aad van der Lugt²; Adriaan C.G.M. van Es²; Pieter Jan van Doormaal²; Jeannette Bakker¹⁴; Jan-Hein Hensen¹⁵; Aarnout Plaisier¹⁶; Geert Lycklama à Nijeholt¹⁷

Local principal investigators

Diederik W.J. Dippel¹; Bob Roozenbeek^{1,2}; Aad van der Lugt²; Amber Hoek¹⁸; Henk Kerkhoff³; Anouk D. Rozeman³; Jeannette Bakker¹⁴; Erick Oskam¹⁹; Walid Moudrous⁵; Jan-Hein Hensen¹⁵; Frédérique H. Vermeij⁶; Mandy M.A. van der Zon²⁰; Egon D. Zwets²¹; Kees C.L. Alblas⁶; Laus J.M.M. Mulder⁷; Jan Willem Kuiper²²; Annemarie D. Wijnhoud⁸; Bruno J.M. van Moll⁸; Aarnout Plaisier¹⁶; Lisette Maasland⁹; Mirjam Woudenberg²³; Roeland P.J. van Eijkelenburg¹⁰; Arnoud M. de Leeuw²⁴

Local trial coordinators

Anja Noordam-Reijm¹¹; Timo Bevelander¹²

PhD / Medical students

Vicky Chalos^{1,2,3}; Eveline J.A. Wiegers⁴; Lennard Wolff²; Dennis C. van Kalker¹

List of affiliations

Department of Neurology¹, Radiology & Nuclear Medicine², Public Health⁴,
Emergency Medicine¹⁸, Erasmus MC University Medical Center;
Department of Neurology³, Radiology¹⁴, Emergency Medicine¹⁹,
Albert Schweitzer hospital, Dordrecht;
Department of Neurology⁵, Radiology¹⁵, Maasstad hospital, Rotterdam;
Department of Neurology⁶, Radiology²⁰, Emergency Medicine²¹,
Franciscus Gasthuis & Vlietland, Rotterdam;
Department of Neurology⁷, Radiology²², Ikazia hospital, Rotterdam;
Department of Neurology⁸, Radiology¹⁶; IJsselland hospital, Rotterdam;
Department of Neurology⁹, Radiology²³, van Weel Bethesda hospital, Dirksland;
Department of Neurology¹⁰, Radiology²⁴, Rivas Zorggroep Beatrix hospital, Gorinchem;
Department of Radiology¹⁷, Haaglanden Medical Center, the Hague;
Ambulance service Rotterdam-Rijnmond¹¹, Barendrecht;
Ambulance service Zuid-Holland Zuid¹², Papendrecht;
Rotterdam Stroke Service¹³, Rotterdam



In life, it's not where you go, it's who you travel with
Charles M. Schulz

Dankwoord

Het is eindelijk zover, mijn boekje is af! Er is hiermee een einde gekomen aan een mooie en bijzondere periode in mijn leven. Een promotieonderzoek doe je niet alleen en ik ben blij dat ik nu iedereen kan bedanken die mij de afgelopen jaren geholpen heeft. Ik heb het geluk gehad om met een groep geweldige mensen samen te mogen werken en daarnaast veel lieve mensen om me heen te hebben die het leven zo mooi maken.

Allereerst wil ik natuurlijk mijn promotoren, professor Ewout Steyerberg en professor Diederik Dippel, bedanken.

Beste Ewout, ik kan me onze eerste meeting nog goed herinneren. Nadat we eerst gezellig over hardlopen en familie zaken spraken, ging je plotseling in de tweede versnelling en had je in een mum van tijd alle pijnpunten van onze studieopzet blootgelegd. Ik kwam terug met een lijstje vol nieuwe ideeën en analyses waar ik nog nooit van gehoord had. In de loop van de tijd leerde ik je gelukkig steeds iets beter bijhouden en anders had ik altijd nog je boek om het terug te zoeken. Dank voor je kritische blik, waardevolle commentaar en indrukwekkende snelheid van reageren op mijn manuscripten.

Beste Diederik, het begon eigenlijk allemaal toen ik onder jouw begeleiding tijdens mijn keuze-onderzoek aan de MR CLEAN pre-trial mocht werken. Later gaf je me de mogelijkheid om dit promotieonderzoek te doen waarvoor ik je nog steeds erg dankbaar ben. Je bent een ontzettend fijne supervisor: enthousiast, gedreven en makkelijk benaderbaar. Ik heb veel van je geleerd, niet alleen medisch-inhoudelijk, maar ook over samenwerken, grammatica en het opzetten van studies en beslisbomen. Je bent een voorbeeld voor iedereen die gedegen methodologisch onderzoek wil combineren met de klinische praktijk.

Naast twee promotoren heb ik ook twee fantastische copromotoren. Hester, ontzettend bedankt voor alle kansen en de ruimte die je me hebt geboden om mezelf te ontwikkelen. Onze meetings zijn altijd gezellig maar ook heel productief. Ik heb met veel plezier met je kunnen sparren, zowel over lopende projecten als over mijn toekomstplannen. Ik hoop dat we onze samenwerking nog lang mogen voortzetten. Bob, je bent betrokken en ambitieus met een goed gevoel voor humor. Tijdens de langdurige procedures in het kader van de nieuwe AVG gaf ik de moed soms bijna op, maar jij wist altijd overal weer een oplossing voor. Bedankt voor je scherpe blik en nuttige suggesties bij het beoordelen van mijn manuscripten.

Beste leden van de promotiecommissie, bedankt voor het doornemen van mijn proefschrift en het deelnemen aan de verdediging. Ik hoop op een interessante gedachteswisseling. Professor Fiona Lecky and professor David Kent, thank you for your willingness to travel to Rotterdam to take place in the committee if the situation regarding COVID-19 allows.

Dear David, thanks for your guidance during my time in Boston and afterwards. You taught me a lot about research (and good food!) and I still enjoy working together.

De artikelen in dit proefschrift zouden er niet zijn gekomen zonder de waardevolle inbreng van alle coauteurs. Ik wil jullie allen bedanken voor de prettige samenwerking de afgelopen jaren. Speciale dank gaat uit naar Maxim, Vicky, Paula, Nikki en Adrien voor de papers die ik heb mogen gebruiken in mijn proefschrift. I would like to thank professor Joseph Broderick, professor Mayank Goyal, and the HERMES investigators for their cooperation on the MR PREDICTS project. Dank ook aan alle onderzoekers van de MR CLEAN trial en MR CLEAN Registry (pagina 259) voor het beschikbaar stellen van de door hen met veel toewijding verzamelde data.

Maxim, ooit werkten we al met veel plezier samen aan de MR CLEAN pre-trial en ik ben blij dat we dat tijdens onze promotietijd konden voortzetten. Het was heerlijk om samen te sleutelen aan MR PREDICTS en voor het eerst onze helden tegen te komen op congres. Je enthousiasme en doortastendheid heb ik wel eens gemist tijdens het afronden van dit boekje! Vicky, we hebben de afgelopen jaren lief en leed gedeeld. Urenlang bijpraten op de gang of lekker samen R'en, waarbij we meestal alleen maar nieuwe problemen tegenkomen (en weer oplossen!). Wat hebben we veel gelachen, zeker tijdens alle tripjes en de ESO summerschool. Bedankt dat je mijn paranimf wilt zijn!

Martijne, aan jou kon ik PRESTO met een gerust hart overdragen en daar ben ik je heel dankbaar voor! Je hebt een enorme hoeveelheid werk verricht en hopelijk kunnen we snel de eerste resultaten publiceren. Daarnaast natuurlijk dank aan alle PRESTO onderzoekers (pagina 261), met name Henk Kerkhoff, Anouk Rozeboom, Walid Moudrous, Frederique Vermeij, Ad van Es en professor Aad van der Lugt. Ook veel dank aan alle deelnemende ambulanceverpleegkundigen van regio Rotterdam-Rijnmond en Zuid-Holland Zuid.

Charlie, Eveline en Benjamin, waar moet ik beginnen? Jullie brachten een hoop leven, muziek en nieuwe ideeën in de toren. Daarnaast hebben jullie me te vaak uit de brand geholpen als mijn planning weer eens faalde (valet parking in Leiden of een visum halen in Brussel, ga zo maar door), waarvoor eeuwig dank. Met jullie enthousiasme en daadkracht hebben jullie veel bereikt in korte tijd!

De afgelopen jaren heb ik veel meer mensen leren kennen binnen en buiten het Erasmus MC. José en Veerle, kamergenootjes van het eerste uur. Collega's op MGZ en binnen de sectie Medische Besliskunde (onder andere Jilske, Ernest, Daphne, Nikki, Daan, David, Arvind, Isabel, Marzyeh, Ana). Commissieleden van Promeras en het JVO. De MR CLEAN predictie groep (Noor, Rob, Kars, Femke, Nikki, Sanne, Nadinda, Jim) en promovendi van de MR CLEAN Registry. Allen bedankt voor de gezellige tijd op de afdeling, de congressen, de borrels en de etentjes!

Many thanks to George Ntaios for organizing the fantastic ESO summerschool. I've made many friends within Europe and really enjoyed the trip with the Lisbon group (Vicky, Sebastian, Alex, and Mariana). All colleagues at the PACE center, especially Chrissy, Jinny, and Jenn, thanks for your hospitality in Boston and the pleasant collaboration.

SEH-artsen en verpleegkundigen uit Bergen op Zoom, jullie hebben me laten zien wat een mooi vak de spoedeisende geneeskunde is. Collega's en oud-collega's van de SEH, zowel in het Erasmus MC als het Albert Schweitzer ziekenhuis, ik ben blij om met jullie samen te kunnen werken.

Wat het leven buiten het werk om zo leuk maakt, zijn natuurlijk alle vrienden en familie waarmee ik mezelf gelukkig kan prijzen. Ik kan jullie niet genoeg bedanken voor alle steun en gezelligheid!

Van den Bergjes, al zo lang als ik me kan herinneren zijn jullie mijn tweede familie. Van horror nights, tuinhuisparty's en de Yellow tot aan Radewijk en de Dominicaanse republiek, de Tokkies hebben overal de boel op stelten gezet (vooral die paaltjes waar water uit komt!). Rox, Mies en Eef, hoelang we elkaar ook niet gezien hebben, het is altijd weer als vanouds! In de loop van de tijd is de groep langzaam uitgebreid en het wordt er alleen maar gezelliger op (zolang we maar niet gaan midgetgolfen!).

Meiden van de doc groep, het is alweer 14 jaar geleden dat we aan de geneeskundestudie begonnen. Hoe had ik het ooit zonder jullie kunnen volhouden? Lonneke, ik bewonder je gedrevenheid en hoop later samen met jou op een SEH te kunnen werken. Simone, we zijn al vriendinnen vanaf de basisschool, hoe bijzonder is dat! Natas, mijn huisgenootje tijdens de Eurekaweek, bij de nonnen in Mtinko en later op de Boezemlaan. We hebben zoveel gelachen en zoveel mooie herinneringen samen! Sher, je bent altijd vrolijk en deelt mijn enthousiasme voor fanatiek zingen/meeblèren in de auto. Samen met Ro en Will zijn we homies op de piste en ver daarbuiten. Ik hoop op nog vele gezellige avondjes!

Sigrid, je bent een heerlijke optimist en je kaartjes vrolijken me altijd weer op! We zijn samen de halve wereld over geweest en ik hoop dat we binnenkort weer onze beruchte Belgisch bier tripjes kunnen maken. Daarnaast horen de spelletjesmiddagen met Anne en Eva er inmiddels natuurlijk ook zeker bij!

Paardenvrouwen uit Oostvoorne, bedankt voor de gezelligheid en de heerlijke ritjes door het bos en over het strand. Nu mijn proefschrift af is kan ik hopelijk weer wat vaker langskomen. Martin, je bent de beste hoefsmid van Nederland! Mariska, ik denk nog vaak terug aan onze tijd op de Pruiwendijk en ben heel blij dat we nog steeds vriendinnen zijn. Rox, ik ben je eeuwig dankbaar voor het vertrouwen dat je in me had met Donna.

Lieve ooms en tantes, neven en nichten, bedankt voor jullie gezelligheid en belangstelling. Toen ik Romero leerde kennen werd ik met liefde verwelkomt in de familie Camagong, daar ben ik erg blij mee! Familie Uysal, birbirimizin ailesinin bir parçası olduğumuz için çok mutluyum. Tuğçe, speciaal voor jou een nieuwe Nederlandse uitdrukking: je bent een topper! Hans en Tonny, voor mij voelen jullie ook als familie! Manuéla, bedankt voor je oprechte interesse en ik hoop dat je erg gelukkig gaat zijn in Rotterdam. Willem, je maakt de beste foto's en lekkerste tiramisu's. Ik vind het altijd fijn als je erbij bent!

Mijn lieve oma's zijn altijd een groot voorbeeld voor me geweest. Zoveel doorzettingsvermogen en altijd positief, daar heb ik veel bewondering voor. Oma Door, ik ben heel blij dat u bij dit bijzondere moment kunt zijn en ik hoop dat u ervan geniet. Ik ben trots wanneer mensen zeggen dat ik op u lijk en kan alleen maar hopen dat ik op uw manier oud mag worden!

Timo, als broer en zus zijn we twee handen op één buik! Toen ik startte met dit promotietraject wist ik al dat jij m'n paranimf zou moeten worden en ik ben dan ook blij dat je tijdens mijn promotie naast me zal staan. Ik ben super trots op je, met alles wat je de afgelopen jaren hebt gedaan en hoe je in het leven staat. Als je weer terug in Nederland bent, hebben we nog heel wat avondjes in te halen bij Leo!

Mijn lieve ouders, de beste die ik me kan voorstellen! Jullie steunen me onvoorwaardelijk en staan altijd voor me klaar, daar ben ik heel dankbaar voor. Pap, we zijn allebei levensgenieters en delen volgens mij hetzelfde gen om altijd en overal als laatste weg te gaan. Mam, wat zijn we samen veel met de paarden op stap geweest, we genieten er altijd van! Je bent de liefste moeder die ik me kan wensen en kent me soms beter dan ik mezelf ken. Ik hou van jullie!

Romero, ik ben zo blij dat ik 12 jaar geleden Koninginnedag ging vieren in Amsterdam. Wie had toen kunnen bedenken dat we nu nog steeds zo gelukkig zouden zijn samen. Bedankt voor alles, ik zou het niet gekund hebben zonder jou! Je remt me af als ik weer eens teveel hooi op m'n vork heb genomen en bent er altijd voor me als het toch mis is gegaan. Ik kijk er elke dag weer naar uit om naar huis te gaan omdat ik weet dat jij er bent!





List of publications

In this thesis

- **Venema E**, Burke JF, Roozenbeek B, Nelson J, Lingsma HF, Dippel DWJ, Kent DM. Prehospital Triage Strategies for the Transportation of Suspected Stroke Patients in the United States. *Accepted for publication*.
- **Venema E**, Roozenbeek B, Mulder MJHL, Brown S, Majoie CBLM, Steyerberg EW, Demchuck AM, Muir KW, Dávalos A, Mitchell PJ, Bracard S, Berkhemer OA, Lycklama à Nijeholt GJ, van Oostenbrugge RJ, Roos YBWEM, van Zwam WH, van der Lugt A, Hill MD, White P, Campbell BCV, Guillemin F, Saver JL, Jovin TG, Goyal M, Dippel DWJ, Lingsma HF; HERMES collaborators and MR CLEAN Registry Investigators. Improving selection of patients for endovascular treatment of ischaemic stroke: validation and updating of the MR PREDICTS decision tool with data from 4,398 patients. *Submitted*.
- **Venema E**, Duvekot MHC, Lingsma HF, Rozeman AD, Moudrour W, Vermeij FH, Biekart M, van der Lugt A, Kerkhoff H, Dippel DWJ, Roozenbeek B; PRESTO investigators. Prehospital triage of patients with suspected stroke symptoms (PRESTO): protocol of a prospective observational study. *BMJ Open*. 2019;9:e028810.
- **Venema E***, Groot AE*, Lingsma HF, Hinsenveld W, Treurniet KM, Chalos V, Zinkstok SM, Mulder MJHL, de Ridder IR, Marquering HA, Schonewille WJ, Wermer MJH, Majoie CBLM, Roos YBWEM, Dippel DWJ, Coutinho JM*, Roozenbeek B*. Effect of inter-hospital transfer on endovascular treatment for acute ischemic stroke. *Stroke*. 2019;50:923–30.
- Janssen PM, **Venema E**, Dippel DWJ. Effect of Workflow Improvements in Endovascular Stroke Treatment: A Systematic Review and Meta-Analysis. *Stroke*. 2019;50:665–74.
- **Venema E**, Lingsma HF, Chalos V, Mulder MJHL, Lahr MMH, van der Lugt A, van Es ACGM, Steyerberg EW, Hunink MGM, Dippel DWJ, Roozenbeek B. Personalized Prehospital Triage in Acute Ischemic Stroke. *Stroke*. 2019;50:313–20.
- **Venema E***, Boedt N*, Berkhemer OA, Rood PPM, van Zwam WH, van Oostenbrugge RJ, van der Lugt A, Roos YBWEM, Majoie CBLM, Lingsma HF, Dippel DWJ; MR CLEAN investigators. Workflow and factors associated with delay in the delivery of intra-arterial treatment for acute ischemic stroke in the MR CLEAN trial. *J Neurointerv Surg*. 2018;10:424–8.
- **Venema E***, Mulder MJHL*, Roozenbeek B, Broderick JP, Yeatts SD, Khatri P, Berkhemer OA, Emmer BJ, Roos YBWEM, Majoie CBLM, van Oostenbrugge RJ, van Zwam WH, van der Lugt A, Steyerberg EW, Dippel DWJ, Lingsma HF. Selection of patients for intra-arterial treatment for acute ischaemic stroke: development and validation of a clinical decision tool in two randomised trials. *BMJ*. 2017;357: j1710.

- Mulder MJ*, **Venema E***, Roozenbeek B, Broderick JP, Yeatts SD, Khatri P, Berkhemer OA, Roos YB, Majoie CB, van Oostenbrugge RJ, van Zwam WH, van der Lugt A, Steyerberg EW, Dippel DW, Lingsma HF. Towards personalised intra-arterial treatment of patients with acute ischaemic stroke: a study protocol for development and validation of a clinical decision aid. *BMJ Open*. 2017;7:e013699.

Other publications

- Duvekot MHC, **Venema E**, Rozeman AD, Moudrour W, Vermeij FH, Biekart M, Lingsma HF, Maasland L, Wijnhoud AD, Mulder LJMM, van Eijkelenburg RPJ, Buijk B, Bakker J, Plaisier AS, Hensen, JH, Lyklama à Nijeholt GJ, van Doormaal PJ, van Es ACGM, van der Lugt A, Kerkhoff H, Dippel DWJ, Roozenbeek B; PRESTO Investigators. Prehospital triage of patients with suspected stroke (PRESTO): A prospective in-field validation of eight prehospital stroke scales to detect intracranial large vessel occlusion. *Submitted*.
- Kremers FCC, **Venema E**, Duvekot MHC, Yo LSF, Bokkers RPH, Lycklama à Nijeholt GJ, van Es ACGM, van der Lugt A, Majoie CBLM, Burke JF, Roozenbeek B, Lingsma HF, Dippel DWJ; MR CLEAN Registry Investigators. Outcome Prediction Models for Patients Undergoing Endovascular Treatment for Acute Ischemic Stroke: A Systematic Review And External Validation. *Submitted*.
- Gravesteijn BY, Sewalt CA, **Venema E**, Nieboer D, Steyerberg EW. Missing data in prediction research: A five step approach for multiple imputation, illustrated in CENTER-TBI. *Submitted*.
- Sewalt CA, **Venema E**, van Zwet E, van Ditshuizen J, Schuit SCE, Polinder S, Lingsma HF, den Hartog D. The relationship between hospital volume and in-hospital mortality of severely injured patients in Dutch Level-1 trauma centers. *Submitted*.
- Duvekot MHC, **Venema E**, Lingsma HF, Coutinho JM, van der Worp HB, Hofmeijer J, van der Lugt A, Kerkhoff H, Dippel DWJ, Roozenbeek B; MR CLEAN Registry Investigators. Early detection of large vessel occlusions: sensitivity of fourteen prehospital stroke scales for different occlusion locations in the MR CLEAN Registry. *Submitted*.
- Wieggers EJA, Compagne KCJ, Janssen PM, **Venema E**, Deckers JW, Schonewille WJ, Vos JA, Lycklama à Nijeholt GJ, Roozenbeek B, Martens JM, Hofmeijer J, van Oostenbrugge RJ, van Zwam WH, Majoie CBLM, van der Lugt A, Lingsma HF, YBWEM Roos, DWJ Dippel. Path From Clinical Research to Implementation: Endovascular Treatment of Ischemic Stroke in the Netherlands. *Stroke*. 2020;51:1941–50.

- Van Meenen LCC, Groot AE, **Venema E**, Emmer BJ, Smeekes MD, Kommer GJ, Majoie CBLM, Roos YBWEM, Schonewille WJ, Roozenbeek B, Coutinho JM; MR CLEAN Registry Investigators. Interhospital Transfer vs. Direct Presentation of Patients With a Large Vessel Occlusion Not Eligible for IV Thrombolysis. *J Neurol*. 2020;267:2142-50.
- Wiegers EJA, Mulder MJHL, Jansen IGH, **Venema E**, Compagne KCJ, Berkhemer OA, Emmer BJ, Marquering HA, van Es ACGM, Sprengers ME, van Zwam WH, van Oostenbrugger RJ, Roos YBWEM, Majoie CBLM, Roozenbeek B, Lingsma HF, Dippel DWJ, van der Lugt A; Clinical and Imaging Determinants of Collateral Status in Patients With Acute Ischemic Stroke in MR CLEAN Trial and Registry. *Stroke*. 2020;51:1493-1502.
- Sewalt CA, Wiegers EJA, Lecky FE, den Hartog D, Schuit SCE, **Venema E**, Lingsma HF. The volume-outcome relationship among severely injured patients admitted to English major trauma centers: a registry study. *Scand J Trauma Resusc Emerg Med*. 2020;28:18.
- Verboon C, van den Berg B, Cornblath DR, **Venema E**, Gorson KC, Lunn MP, Lingsma H, Van den Bergh P, Harbo T, Bateman K, Pereon Y, Sindrup SH, Kusunoki S, Miller J, Islam Z, Hartung HP, Chavada G, Jacobs BC, Hughes RAC, van Doorn PA; IGOS Consortium. Second IVIg course in Guillain-Barré syndrome with poor prognosis: the non-randomised ISID study. *J Neurol Neurosurg Psychiatry*. 2020;91:113-21.
- Chalos V, van der Ende NAM, Lingsma HF, Mulder MJHL, **Venema E**, Dijkland SA, Berkhemer OA, Yoo AJ, Broderick JP, Palesch YY, Yeatts SD, Roos YBWEM, van Oostenbrugger RJ, van Zwam WH, Majoie CBLM, van der Lugt A, Roozenbeek B, Dippel DWJ; MR CLEAN Investigators. National Institutes of Health Stroke Scale: An Alternative Primary Outcome Measure for Trials of Acute Treatment for Ischemic Stroke. *Stroke*. 2020;51:282-90.
- Sewalt CA, **Venema E**, Wiegers EJA, Lecky FE, Schuit SCE, den Hartog D, Steyerberg EW, Lingsma HF. Early Identification of Major Trauma: External Validation and Comparison of Prognostic Prehospital Trauma Models in 154,476 patients. *Br J Surg*. 2020;107:373-80.
- Oemrawsingh A, van Leeuwen N, **Venema E**, Limburg M, de Leeuw FE, Wijffels MP, de Groot AJ, Hilken PHE, Hazelzet JA, Dippel DWJ, Bakker CH, Voogdt-Pruis HR, Lingsma HF. Value-based healthcare in ischemic stroke care: case-mix adjustment models for clinical and patient-reported outcomes. *BMC Med Res Methodol*. 2019;19:229.
- Chalos V*, LeCouffe NE*, Uyttenboogaart M, Lingsma HF, Mulder MJHL, **Venema E**, Treurniet KM, Eshghi O, van der Worp HB, van der Lugt A, Roos YBWEM, Majoie CBLM, Dippel DWJ, Roozenbeek B*, Coutinho JM*; MR CLEAN Registry Investigators.. Endovascular Treatment With or Without Prior Intravenous Alteplase for Acute Ischemic Stroke. *J Am Heart Assoc*. 2019;8:e011592.

- Klompenhouwer AJ, Alblas M, van Rosmalen BV, Haring MPD, **Venema E**, Doukas M, Thomeer MGJ, Takkenberg RB, Verheij J, de Meijer VE, van Gulik TM, Lingsma HF, de Man RA, Ijzermans JNM. Development and Validation of a Model to Predict Regression of Large Size Hepatocellular Adenoma. *Am J Gastroenterol*. 2019;114:1292–8.
- Wiegers EJA*, Sewalt CA*, **Venema E**, Schep NWL, Verhaar JAN, Lingsma HF, Den Hartog D. The volume–outcome relationship for hip fractures: a systematic review and meta-analysis of 2,023,469 patients. *Acta Orthop*. 2019;90:26–32.
- Baggen VJM, **Venema E**, Živná R, van den Bosch AE, Eindhoven JA, Witsenburg M, Cuypers JAAE, Boersma E, Lingsma H, Popelová JR, Roos-Hesselink JW. Development and validation of a risk prediction model in patients with adult congenital heart disease. *Int J Cardiol*. 2019;276:87–92.
- van Os HJA, Ramos LA, Hilbert A, van Leeuwen M, van Walderveen MAA, Kruyt ND, Dippel DWJ, Steyerberg EW, van der Schaaf IC, Lingsma HF, Schonewille WJ, Majoie CBLM, Olabarriga SD, Zwinderman KH, **Venema E**, Marquering HA, Wermer MJH; MR CLEAN Registry Investigators. Predicting Outcome of Endovascular Treatment for Acute Ischemic Stroke: Potential Value of Machine Learning Algorithms. *Front Neurol*. 2018;9:784.
- Sewalt CA*, Wiegers EJA*, **Venema E**, Lecky FE, Schuit SCE, Den Hartog D, Lingsma HF. The volume-outcome relationship in severely injured patients: A systematic review and meta-analysis. *J Trauma Acute Care Surg*. 2018;85:810–9.
- Mulder MJHL, Jansen IGH, Goldhoorn RB, **Venema E**, Chalos V, Compagne KCJ, Roozenbeek B, Lingsma HF, Schonewille WJ, van den Wijngaard IR, Boiten J, Albert Vos J, Roos YBWE, van Oostenbrugge RJ, van Zwam WH, Majoie CBLM, van der Lugt A, Dippel DWJ; MR CLEAN Registry Investigators. Time to Endovascular Treatment and Outcome in Acute Ischemic Stroke: MR CLEAN Registry Results. *Circulation*. 2018;138:232–40.
- Dijkland SA, Voormolen DC, **Venema E**, Roozenbeek B, Polinder S, Haagsma JA, Nieboer D, Chalos V, Yoo AJ, Schreuders J, van der Lugt A, Majoie CBLM, Roos YBWE, van Zwam WH, van Oostenbrugge RJ, Steyerberg EW, Dippel DWJ, Lingsma HF; MR CLEAN Investigators. Utility-Weighted Modified Rankin Scale as Primary Outcome in Stroke Trials: A Simulation Study. *Stroke*. 2018;49:965–71.

Letters to the editor

- Chalos V, **Venema E**, Lingsma HF, Dippel DWJ. Letter to the Editor Regarding "Predicting Clinical Outcome After Mechanical Thrombectomy: The GADIS (Gender, Age, Diabetes Mellitus History, Infarct Volume, and Sex) Score". *World Neurosurg.* 2020;138:587–8.
- Klompenhouwer AJ, Alblas M, van Rosmalen BV, Haring MPD, **Venema E**, Doukas M, Thomeer MGJ, Takkenberg RB, Verheij J, de Meijer VE, van Gulik TM, Lingsma HF, de Man RA, Ijzermans JNM. Response to Chen et al. *Am J Gastroenterol.* 2020;115:147–8.
- **Venema E**, Groot AE, Coutinho JM. Response by Venema et al to Letter Regarding Article, "Effect of Interhospital Transfer on Endovascular Treatment for Acute Ischemic Stroke". *Stroke.* 2019;50:e260.
- Sewalt CA, Wiegers EJA, **Venema E**, Lingsma HF. Letter to the Editor regarding the article: "Identifying pre-hospital factors associated with outcome for major trauma patients in a regional trauma network: an exploratory study". *Scand J Trauma Resusc Emerg Med.* 2017;25:113.
- **Venema E**, Mulder M, Lingsma HF. Letter by Venema et al Regarding Article, "Validating a Predictive Model of Acute Advanced Imaging Biomarkers in Ischemic Stroke". *Stroke.* 2017;48:e225.
- **Venema E**, Lingsma HF, Dippel DW. Letter by Venema et al Regarding Article, "Analysis of Workflow and Time to Treatment on Thrombectomy Outcome in the Endovascular Treatment for Small Core and Proximal Occlusion Ischemic Stroke (ESCAPE) Randomized, Controlled Trial". *Circulation.* 2016;134:e404–e405.

*equal contribution



PhD portfolio

Name: Esmee Venema
 Erasmus MC Department: Public Health & Neurology
 Research School: Netherlands Institute for Health Sciences (NIHES)
 PhD period: November 2015 – November 2019
 Promotors: Prof.dr. E.W. Steyerberg & prof.dr. D.W.J. Dippel
 Copromotors: Dr. H.F. Lingsma & dr. B. Roozenbeek

	Year	Workload (ECTS)
1. PhD training		
<i>General academic courses</i>		
Master of Science – Clinical Epidemiology	2016-2017	39
Biomedical English Writing and Communication	2017	4.0
Research Integrity	2017	0.3
Basiscursus Regelgeving en Organisatie voor Klinisch onderzoekers (BROK)	2019	1.0
<i>Presentations at (inter)national conferences</i>		
European Stroke Organisation Conference; Barcelona (two oral presentations)	2016	2.0
European Stroke Organisation Conference; Prague (oral and poster presentation)	2017	1.5
1st Translational Cardiovascular Research Meeting; Utrecht (poster presentation)	2017	0.5
39 th Annual meeting of the Society for Medical Decision Making; Pittsburgh (poster presentation)	2017	0.5
Dutch Society for Neurology Science Days; Nunspeet (oral presentation)	2017	1.0
European Stroke Organisation Conference; Gothenburg (oral presentation)	2018	1.0
European Meeting of the Society for Medical Decision Making; Leiden (oral and poster presentation)	2018	1.5
Dutch Emergency Medicine Conference; Egmond aan Zee (two oral presentations)	2018	2.0
40th Annual meeting of the Society for Medical Decision Making; Montreal (oral and poster presentation)	2018	1.5
<i>Seminars and workshops</i>		
Various seminars and research meetings of the department of Public Health, the department of Neurology, the MR CLEAN Registry and the CONTRAST consortium	2016-2019	4.0

European Stroke Organisation Stroke Summer School; Larissa, Greece	2017	1.0
Sensitivity Analysis and Value of Information Analysis Using Regression Metamodeling in R	2017	0.2
2. Teaching activities		
<i>General teaching activities</i>		
Revising bachelor essays	2016	0.5
Supervising community projects of bachelor students	2016-2019	2.0
Lecturing 'VO Choices in healthcare'	2017-2018	0.5
Developing R assignments NIHES Biostatistical Methods I (CC02)	2018	0.2
Teaching-assistant NIHES Clinical Epidemiology (CE02)	2018	0.2
<i>Supervising research master students</i>		
Eveline Wiegers	2017-2018	2.0
Charlie Sewalt	2017-2018	2.0
Femke Kremers	2019	1.0
3. Other activities		
Member of the organizing committee of the Erasmus MC PhD day and Career Event	2016	1.0
Secretary of the Promeras PhD students association	2016-2017	1.5
Member of the Junior Representatives committee, department of Public Health	2016-2017	0.5
Training paramedics to use prehospital stroke scales (ambulance regions Rotterdam-Rijnmond, Zuid-Holland Zuid, and Zeeland)	2018	2.5
Visiting researcher at the Predictive Analytics and Comparative Effectiveness (PACE) Center, Boston, USA	2018	2.0
Peer reviews for several journals (European Journal of Neurology, Journal of NeuroInterventional Surgery, European Stroke Journal, Intensive Care Medicine)	2018-2019	0.6
4. Awards		
Netherlands Institute for Health Sciences (NIHES) award for the best paper written in the academic year 2016-2017	2017	
Dutch Emergency Medicine Conference award for the best abstract presentation	2018	
Lee B. Lusted Student Award for outstanding presentation of research at the Society for Medical Decision Making meeting	2018	

Total ECTS

77.5





About the author

Esmee Venema was born on March 1st, 1988 in Rotterdam, the Netherlands. After finishing secondary school at the Dalton Lyceum in Barendrecht, she returned to Rotterdam to study medicine at the Erasmus Medical Center. As part of her study, she conducted a research project on the effect of intravenous thrombolytic therapy given prior to endovascular treatment of ischemic stroke in the MR CLEAN pre-trial (under supervision of professor Diederik Dippel). In 2013, she finished her clinical rotations after working for 3 months at the St. Carolus hospital in Tanzania. She obtained her medical degree cum laude and started working as resident at the Emergency Department of the Bravis hospital in Bergen op Zoom. It is there that she discovered a great passion for acute care and emergency medicine.



In 2015, she was given the opportunity to start a PhD on personalized treatment of ischemic stroke at the Department of Public Health and the Department of Neurology in the Erasmus Medical Center, under supervision of professor Ewout Steyerberg and professor Diederik Dippel. She combined her PhD trajectory with a Master of Science in Clinical Epidemiology, which she finished in 2017, and was secretary of the Promeras PhD students association. In 2018, she worked for 2 months under the supervision of professor David Kent at the Predictive Analytics and Comparative Effectiveness Center of Tufts Medical Center in Boston (USA), a collaboration that continues to this day. She designed and coordinated the PRESTO study, a prospective multicenter study for the in-field validation of prehospital stroke scales. She returned to her clinical work in 2019. Currently, she works at the Emergency Department of the Albert Schweitzer hospital in Dordrecht and is pursuing a career as emergency physician and researcher. She lives in Rotterdam, together with her boyfriend Romero.



