


TR/ 55

JULY 1975

THE LINEAR DEPENDENCE RELATIONS AND
TRUNCATION ERRORS FOR INTERPOLATING
EVEN ORDER SPLINES

W.D. Hoskins and D. S.Meek.



(121)
1
CA

W 9252524

The Linear Dependence Relations and Truncation
Errors for Interpolating Even Order Splines

W. D. Hoskins and D. S. Meek

Abstract:

Linear dependence relations are derived for even order polynomial splines defined over the points midway between a uniformly spaced set of knots. The leading term of the truncation error is determined and is illustrated by an example.

1. Introduction

The linear dependence relations for odd order polynomial splines have been studied for small orders by a variety of workers, however the general relations for odd polynomial splines on a uniform partition were obtained by Fyfe in [6] and Swartz [11]. The local truncation error of the odd order polynomial splines defined in this way has been analysed by Swartz [11] and Albasiny and Hoskins [3]. Less work has been done with even order polynomial splines defined over points midway between the knots of a uniform partition as in [8] and [9]. Subbotin [10] has demonstrated the existence of such even degree splines with certain boundary conditions and now the linear dependence relations and the local truncation error will be described [7].

2. Linear Dependence Relations

Let $s(x)$ be a spline of degree N interpolating to the function $f(x)$ on $[x_0, x_n]$ and defined over points midway between the knots of a uniform partition, then $s(x)$ satisfies the following conditions:

- (a) $s(x)$ is a polynomial of degree at most N in each interval $[x_k - h/2, x_k + h/2]$, $k = 1, 2, \dots, n-1$, and in the intervals $[x_0, x_0 + h/2]$, $[x_n - h/2, x_n]$, $h = (x_n - x_0)/n$ and $x_k = x_0 + kh$,
 $k = 0, 1, 2, \dots, n$

(b) using the notation $s_k \equiv s(x_k)$ and $f_k \equiv f(x_k)$, $s_k = f_k$,

$k = 0, 1, 2, \dots, n$

and

(c) $s(x) \in C^{N-1}[x_0, x_n]$.

Although N even is the case of interest, all the formulae below also hold for odd N . It is convenient subsequently to refer to such a spline as a midpoint interpolating polynomial (m.i.p.) spline.

The linear dependence relations satisfied by an m.i.p. spline are

$$\frac{h^N}{N!} \sum_{i=0}^N C_{i+1/2, N}^{(0)} s_{k+i}^{(p)} = \frac{h^{N-p}}{(N-p)!} \sum_{i=0}^N C_{i+1/2, N}^{(p)} s_{k+i} \quad (1)$$

$$p = 0, 1, 2, \dots, N$$

where

$$C_{i+1/2, N}^{(p)} = (-1)^p (N-p)! Q_{N+1}^{(p)}(i+1/2)$$

and $Q_{N+1}^{(p)}(x)$ is the p th derivative with respect to x of the B-spline

$$Q_{N+1}(x) = \frac{1}{N!} \nabla_{x+}^{N+1} x_+^N$$

defined on the integer knots. Proof of the result (1) is exactly analogous to that found in Loscalzo and Talbot [6], Swartz [11], Fyfe [5] and is given in detail in Meek [7].

Several values of the coefficients $C_{i+1/2, N}^{(p)}$ are given in table 1.

N = 1

	i	0	1
p	0	1/2	1/2
	1	-1	1

N = 2

	i	0	1	2
p	0	1/4	6/4	1/4
	1	-1/2	0	1/2
	2	1	-2	1

N = 3

	i	0	1	2	3
p	0	1/8	23/8	23/8	1/8
	1	-1/4	-5/4	5/4	1/4
	2	1/2	-1/2	-1/2	1/2
	3	-1	3	-3	1

N = 4

	i	0	1	2	3	4
p	0	1/16	76/16	230/16	76/16	1/16
	1	-1/8	-22/8	0	22/8	1/8
	2	1/4	4/4	-10/4	4/4	1/4
	3	-1/2	2/2	0	-2/2	1/2
	4	1	-4	6	-4	1

Table 1

The coefficients $C_{i+1/2,N}^{(p)}$, $i = 0, 1, \dots, N$; $p = 0, \dots, N$

For example, a fourth degree m.i.p. spline satisfies the relation
($p=2$ in table 1.)

$$\frac{1}{16} \frac{h^4}{4!} (s_{i+2}'' + 76s_{i+1}'' + 230s_i'' + 76s_{i-1}'' + s_{i-2}'')$$

$$= \frac{1}{4} \frac{h^2}{2!} (s_{i+2} + 4s_{i+1} - 10s_i + 4s_{i-1} + s_{i-2}).$$

Various properties of the coefficients $C_{i+1/2,N}^{(p)}$ follow immediately from their definition in terms of B-spline [4].

A generating function for these coefficients is known [7] and is

$$\sum_{i=0}^N C_{i+1/2,N}^{(p)} z^{i+1/2} = (-1)^p (1-z)^{N+1} \left(z \frac{d}{dz} \right)^{N-p} \left(\frac{z^{1/2}}{1-z} \right), \quad (2)$$

$$p = 0, 1, 2, \dots, N.$$

3. Leading term of the Truncation Error

The leading term of the truncation error may be found in the following manner.

Define the coefficients $a_{0,N} = 1, a_{2,N}, a_{4,N} \dots$ by the equation

$$\sinh^{N+1} x = x^{N+1} (a_{0,N} + a_{2,N} x^2 + \dots)$$

and for $t = 0, 1, 2, \dots$ let

$$E_{t,N}^{(p)} = \frac{1}{t!} \sum_{i=0}^N C_{i+1/2,N}^{(p)} (i-N/2)^t, \quad (3)$$

with the indeterminate 0^0 taken to be unity. Then it follows that

$$E_{t,N}^{(p)} = 0 \quad \begin{cases} \text{if } t+p \text{ is odd} \\ \text{if } t < p \end{cases}, \quad (4)$$

and in addition if $q = [(N-p+2)/2]$ then

$$\text{and } \left. \begin{aligned} E_{p+2m,N}^{(p)} &= \frac{(N-p)!}{2^{2m}} a_{2m,N}, \quad m = 0, 1, \dots, q-1 \\ E_{p+2q,N}^{(p)} &= \frac{(N-p)!}{2^{2q}} a_{2q,N} + (-1)^{N-p} \frac{(2-2^{2q})}{2^{2q}} \frac{B_{2q}}{2q} \end{aligned} \right\} \quad (5)$$

with B_{2q} a Bernoulli number [1].

The proof is given from equation (3) by rewriting it as first

$$E_{t,N}^{(p)} = \frac{1}{t!} \left(z \frac{d}{dz} \right)^t \sum_{i=0}^N C_{i+1/2,N}^{(p)} z^{i-N/2} \Big|_{z=1}$$

then by using equation (2)

$$E_{t,N}^{(p)} = \frac{(-1)^p}{t!} \left(z \frac{d}{dz} \right)^t \left\{ \frac{(1-z)^{N+1}}{z^{(N+1)/2}} \left(z \frac{d}{dz} \right)^{N-p} \left(\frac{z^{1/2}}{1/z} \right) \right\} \Bigg|_{z=1}$$

Now substitution of $z=e^{2x}$ and performing the differentiation $N-p$ times gives

$$E_{t,N}^{(1)} = \frac{1}{2^{t-p} t!} \left(\frac{d}{dx} \right)^t \left\{ x^p \left(\sum_{i=0}^q e_{2i,N} x^{2i} + 0(x^{2q+2}) \right) \right\} \Bigg|_{x=0}$$

where

$$e_{2i,N} = \begin{cases} (N-p)! a_{2i,N} & \text{if } i = 0, 1, \dots, q-1 \\ (N-p)! a_{2i,N} + (-1)^{N-p} \frac{B_{2i}}{2i} (2-2^{2i}) & \text{if } i = q \end{cases}$$

Clearly the properties (4) are true and without loss of generality now let $t = p + 2m$, $m = 0, 1, 2, \dots$ and a

$$E_{p+2m,N}^{(p)} = \frac{1}{2^{2m}} e_{2m,N}$$

and equations (5) follow.

The local truncation error for m.i.p. splines is obtained in a manner similar to [3], is given in detail in [7], and brief summary now follows.

Since the m.i.p. spline satisfies equation (1), the appropriate equation giving the truncation error $e_{p,N,k+N/2}$ is

$$\frac{h^N}{N!} \sum_{i=0}^N C_{i+1/2,N}^{(0)} f_{k+i}^{(p)} = \frac{h^{N-p}}{(N-p)!} \sum_{i=0}^N C_{i+1/2,N}^{(p)} f_{k+i} + e_{p,N,k+N/2}. \quad (6)$$

The left-hand side of equation (6), expanded about the midpoint of the interval $[x_k, x_{k+N}]$ by Taylor's theorem, is

$$h^N \left(\sum_{m=0}^{j-1} \frac{h^{2m}}{2^{2m}} a_{2m,N} f_{k+N/2}^{(p+2m)} + \frac{h^{N+2j}}{N!} E_{2j,N}^{(0)} f_{k+N/2}^{(p+2j)} + o(h^{N+2j+2}) \right),$$

where $j = \left\lceil \frac{N+2}{2} \right\rceil$. Expansion of the right-hand side of (6) gives

$$h^N \left(\sum_{m=0}^{q-1} \frac{h^{2m}}{2^{2m}} a_{2m,N} f_{k+N/2}^{(p+2m)} + \frac{h^{N+2q}}{(N-p)!} E_{p+2q}^{(p)} f_{k+N/2}^{(p+2q)} + o(h^{N+2q+2}) \right) + e_{p,N,k+N/2},$$

where $q = \left\lceil \frac{N-p+2}{2} \right\rceil$. Now if $j > q$, then the truncation error is

$$e_{p,N,k+N/2} = (-1)^{N-p+1} \frac{h^{N+2q}}{(N-p)!} \frac{(2-2^{2q})}{2^{2q}} \frac{B_{2q}}{2q} f_{k+N/2}^{(p+2q)} + o(h^{N+2q+2}), \quad (7)$$

and it is easy to see that $j > q$ for all $p > 1$ and when $p=1$ and N is even. If $p=1$ and N is odd, then $j=q$ and the truncation error is given by

$$e_{1,N,k+N/2} = (-1)^N \frac{h^{2N+1}}{N!} \left(\frac{2-2^{N+1}}{2^{N+1}} \right) B_{N+1} f_{k+N/2}^{(N+2)} + o(h^{2N+3}) \quad (8)$$

For example, if $f(x)$ has sufficiently many continuous derivatives,

$$\begin{aligned} & \frac{1}{16} \frac{h^4}{4!} (f_{i+2}'' + 76f_{i+1}'' + 230f_i'' + 76f_{i-1}'' + f_{i-2}'') \\ &= \frac{1}{4} \frac{h^2}{2!} (f_{i+2} + 4f_{i+1} - 10f_i + 4f_{i-1} + f_{i-2}) - \frac{7}{1920} h^8 f_i^{vi} + o(h^{10}). \end{aligned}$$

References

- [1] M. Abramowitz and I.A. Stegun, "Handbook of Mathematical Functions". Dover Publications, Inc. New York, 1965.
- [2] J.H. Ahlberg, E.N. Nilson and J.L. Walsh. "The Theory of Splines and Their Applications", Academic Press, New York, 1967.
- [3] E.L. Albasiny and W.D. Hoskins. Explicit error bounds for periodic splines of odd order on a uniform mesh. *J.Inst.Math.Applies.* 12 (1973), pp 303-318.
- [4] C. deBoor. On calculating with B-splines. *J. Approx. Theory* 6(1973), pp. 50-62.
- [5] D.J. Fyfe. Linear dependence relations connecting equal interval N^{th} degree splines and their derivatives. *J. Inst. Math. Applies.* 7(1971), pp. 398-406.
- [6] F.R. Loscalzo and T.D. Talbot, Spline function approximation for solutions of ordinary differential equations. *SIAM J. Num. Anal.* 4(1967), pp. 433-445.
- [7] D.S. Meek. "On the the Numerical Construction and Approximation of Some Piecewise Polynomial Functions". Ph.D. Thesis, University of Manitoba (1973).
- [8] I.J. Schoenberg. Contributions to the problem of approximation of equidistant data by analytic functions. Part A.*Quart. Appl. Math.* 4(1946), pp. 45-99.

- [9] I.J. Schoenberg. Cardinal Interpolation and Spline Functions IV.
The Exponential Euler Splines. University of Wisconsin,
MRC Report No. 1153 (1971).
- [10] Y.N. Subbotin. Piecewise polynomial (spline) interpolation. Mat.
Zametki 1(1967), pp. 63-70 = Math. Notes 1(1967), pp. 41-45.
- [11] B. Swartz. $O(h^{2n+2-\ell})$ bounds on some spline interpolation errors.
Bull.Amer. Math. Soc. 74(1968), pp. 1072-1078.

**NOT TO BE
REMOVED**
FROM THE LIBRARY

XB 2356842 9

