




Practice recommendations for the use of sedation in routine hospital-based colonoscopy

Fahima Dossa ,^{1,2} Catherine Dubé,^{3,4} Jill Tinmouth,^{2,4,5} Anne Sorvari,⁶ Linda Rabeneck,^{4,5} Bronwen R McCurdy,⁴ Jason A Dominitz,⁷ Nancy N Baxter^{1,2,4,6}

To cite: Dossa F, Dubé C, Tinmouth J, *et al*. Practice recommendations for the use of sedation in routine hospital-based colonoscopy. *BMJ Open Gastro* 2020;**7**:e000348. doi:10.1136/bmjgast-2019-000348

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/bmjgast-2019-000348>).

Received 15 October 2019
Revised 10 December 2019
Accepted 11 December 2019

ABSTRACT

Objective Although sedation improves patient experience during colonoscopy, there is great jurisdictional variability in sedative practices. The objective of this study was to develop practice recommendations for the use of moderate and deep sedation in routine hospital-based colonoscopy to facilitate standardisation of practice.

Design We recruited 32 multidisciplinary panellists to participate in a modified Delphi process to establish consensus-based recommendations for the use of sedation in colonoscopy. Panel members participated in a values assessment survey followed by two rounds of anonymous online voting on preliminary practice recommendations. An inperson meeting was held between voting rounds to facilitate consensus-building. Consensus was defined as >60% agreement/disagreement with recommendation statements; >80% agreement/disagreement was considered indicative of strong consensus.

Results Twenty-nine panellists participated in the values assessment survey. Panellists ranked all factors presented as important to the development of practice recommendations. The factor considered most important was patient safety. Patient satisfaction, procedural efficiency, and cost were considered less important. Strong consensus was achieved for all nine practice recommendations presented to the panel. These recommendations included that all endoscopists be able to perform colonoscopy with moderate sedation, that an endoscopist and a single trained nurse are sufficient for performing colonoscopy with moderate sedation, and that anaesthesia-provided deep sedation be used for select patients.

Conclusion The recommendations presented in this study were agreed on by a multidisciplinary group and provide guidance for the use of sedation in routine hospital-based colonoscopy. Standardised sedation practices will promote safe, effective, and efficient colonoscopy for all patients.

INTRODUCTION

Colonoscopy is an essential tool for the diagnosis and treatment of many intestinal conditions, including colorectal cancer. Patient pain and discomfort experienced during colonoscopy can necessitate early termination of the procedure prior to complete colonic visualisation and can lead to reluctance to undergo future procedures.¹ Patient anxiety prior to and during the procedure may also be

Summary box

What is already known about this subject?

► Sedation improves patient experience during endoscopy; however, there is wide variation in sedation practices.

What are the new findings?

- Using consensus-based methods, an expert panel established recommendations for the use of sedation in colonoscopy; this study provides nine recommendations for sedation practice in colonoscopy.
- These include the recommendation that all endoscopists be capable of performing colonoscopy under moderate sedation, with the assistance of a single trained nurse; and that select patients, such as those with severe comorbidities, may benefit from the presence of an anaesthesia provider during colonoscopy, irrespective of the level of sedation.
- Further to this, the panel identified specific patient groups and contexts that should warrant consideration for deep sedation, including patients who are chronic opiate users, those who have previously not tolerated endoscopy under moderate sedation, and when a lengthy or complex procedure is anticipated.

How might it impact on clinical practice in the foreseeable future?

- This study provides consensus-based recommendations to guide the use of sedation in colonoscopy.
- These recommendations can be used to facilitate standardisation of sedation practices for routine, in-hospital colonoscopy, ensuring high-quality colonoscopy for all patients.

considerable and limit willingness to undergo colonoscopy. Therefore, strategies to optimise patient experience are important.

Sedation provides analgesia and anxiolysis during endoscopy and is recommended for most patients undergoing colonoscopy.² Moderate sedation, commonly provided by a combination of midazolam and fentanyl, is usually administered by an endoscopist and monitored by a nurse, with no additional healthcare providers required. Patients sedated to this level are rousable with minimal



© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Fahima Dossa;
fahima.dossa@mail.utoronto.ca




Figure 1 Consensus-building process.

tactile stimulation.³ In contrast, deep sedation with propofol can potentially alter cardiorespiratory function and even progress to general anaesthesia. Given this risk, in Ontario, propofol is delivered by individuals qualified to administer general anaesthesia—most commonly a trained anaesthesia professional who is responsible only for monitoring the patient^{3–6}—even if moderate sedation with propofol is intended. Deeply sedated patients require repeated or painful stimulation to be roused.³ Deep sedation can be appealing for endoscopy as

propofol has a quick onset and short plasma half-life,⁷ which can potentially enhance procedural efficiency,⁸ and is believed to result in greater patient satisfaction than moderate sedation.^{9,10}

There are wide variations in sedation practices globally and regionally.¹¹ In the UK, approximately 10% of colonoscopies are performed without sedation, <1% are performed with propofol, and the remainder are performed with moderate sedation.¹² In contrast, in 2009, 12% of Canadian endoscopists reported use of propofol,¹³ and importantly




Figure 2 Results of values assessment survey. Twenty-nine members of the expert panel participated in the values assessment survey. Panel members were provided with a list of 20 factors/outcomes and were asked to provide a score (1–10) indicating the importance of the factor in the generation of practice recommendations.

50% of gastroenterologists who were not using propofol expressed interest in its routine use,¹³ portending a rise in its popularity. In Ontario, Canada, the use of propofol has been quickly increasing. Nineteen per cent of colonoscopies performed in Ontario in 2005 included anaesthesia assistance¹⁴; by 2015, this estimate had increased to nearly 50%, although there was wide variability across institutions and regions (data available on request). This is important as the method of sedation delivery has potential quality and safety implications. Recent population-based studies report higher complication rates in colonoscopies performed with anaesthesia assistance, particularly aspiration.^{15 16} Additionally, use of anaesthesia services for colonoscopy comes at a significant monetary cost. In Ontario, colonoscopies performed with anaesthesia cost an additional \$152.27 per procedure, translating into a budgetary impact of over \$18 million in 2014/2015 (data available on request).

Given the variability in practices and the associated safety, satisfaction, and cost implications, guidance regarding sedation for colonoscopy is needed. The objective of this study was to develop consensus-based practice recommendations for the use of sedation in routine hospital-based colonoscopy to facilitate standardisation of practices.

METHODS

Overview

We used a modified Delphi technique^{17–19} to develop consensus-based practice recommendations guiding

the use of sedation for routine hospital-based colonoscopy (figure 1). We chose a consensus-based approach as a review of existing evidence revealed heterogeneous results and because this strategy allowed for incorporation of values from diverse perspectives. The Delphi technique is an iterative approach to gaining consensus from an expert panel. Participants are first asked to assess/rank statements through an anonymous survey. The group's results are tabulated and presented to the participants, who then rerank the statements. This process of receiving tabulated results and reranking statements is continued until consensus is reached. The anonymity of the Delphi approach attenuates the influence of dominant opinions and feedback promotes convergence to consensus.²⁰ The modified Delphi technique includes an inperson meeting, where additional information can be provided and clarifications can be made.

This study is reported using the CREDES (Conducting and Reporting Delphi Studies) recommendations.²¹

Multidisciplinary panel recruitment

We used purposive sampling to select a multidisciplinary group of 32 panellists. Our panel size reflects our desire to recruit diverse panellists who represented a range of hospital settings, practitioners, and administrators. We included physicians (gastroenterologists, general surgeons, anaesthesiologists); endoscopy nurses and managers; public representatives; experts in health economics; representatives from Cancer Care Ontario, the College of Physicians and Surgeons of Ontario, and the Ontario Ministry of Health and Long-Term Care; and international experts in endoscopy. All participants were required to disclose conflicts of interest prior to participation.

Evidence reviews

To inform preliminary recommendations, the study team first systematically reviewed existing guidelines to assess the degree of consistency among current recommendations. As there was little consistency found among guidelines, the team next reviewed and summarised published literature in the following areas: (1) whether the presence of an anaesthesiologist influenced the safety and effectiveness of colonoscopy (irrespective of sedative agent used), (2) the incidence of complications in colonoscopy performed with propofol versus traditional sedatives (opioids and/or benzodiazepines), and (3) safety, satisfaction, and efficiency outcomes between colonoscopies performed with propofol versus midazolam and fentanyl (irrespective of sedative provider). The results of a published meta-analysis comparing propofol with traditional sedatives were also reviewed.¹⁰ Briefly, there were no statistically significant differences in rates of hospital admission, major complications (death, aspiration, splenic injury, myocardial infarction, stroke), polyp detection, caecal intubation, or patient satisfaction between anaesthesia and non-anaesthesia provider-administered sedation; however,

**Table 1** Results of small group sessions**Who may need deep sedation for routine colonoscopy?***

Patient factors	Contextual factors
<ul style="list-style-type: none"> ▶ Chronic opiate users. ▶ Patients who have failed with moderate sedation in the past due to discomfort. ▶ Hypersensitivity to vomiting and nausea. ▶ History of sexual abuse. ▶ Selection of patients with irritable bowel syndrome, fibromyalgia, or previous diverticulitis. ▶ Patients with cognitive disabilities (eg, dementia). ▶ Paediatric patients. 	<ul style="list-style-type: none"> ▶ Anaesthesiologist on hand/readily available. <ul style="list-style-type: none"> – Smaller centres/hospitals may depend on an anaesthesiologist as a revenue stream. – May use anaesthesiologists because they need to provide them with enough cases to keep them employed. ▶ Patient preference. ▶ Lengthy/complex procedure (eg, endoscopic mucosal resection). ▶ Skill level of endoscopist. ▶ Patient is alone.

How do we enhance patient experience under moderate sedation?

Prior to the procedure

- ▶ Provide a friendly, clean environment for the patient.
- ▶ Educate the patient regardless of the sedation they will receive (standardised learning materials).
 - Information about what to expect before, during, and after procedure.
 - Start educating the patient as early as possible.
 - Train all staff in proper education protocols.
- ▶ Prepare the patient for possibility of pain/discomfort.
- ▶ Communicate and listen to the patient.
 - Address concerns and answer questions.
 - Build rapport.
 - Address language barriers, if necessary.
- ▶ Continual education/improvement for medical staff.
- ▶ Skill-enhancing courses for endoscopists and nurses.
- ▶ Ensure patients have a safe way home.

During the procedure

- ▶ Ensure the patient is comfortable (temperature of the room, choice in music, etc).
- ▶ ‘Time out’ before, during, and after the procedure where patient information (including relevant comorbidities and allergies), indications for the procedure, equipment required, findings, etc are reviewed.
- ▶ Use of anxiolytics to minimise recall of pain, when necessary.
- ▶ Communicate with the patient during the procedure (warn about any discomfort they might feel).
- ▶ Allow family member in the room with certain patients (eg, patients who are hearing impaired).
- ▶ Skilled intravenous (IV) placement.
- ▶ Patient-controlled sedation.
- ▶ Titration of sedation dosage.
- ▶ Start the patient lightly sedated and increase sedation, if necessary.
- ▶ Use of abdominal pressure and variation in patient positioning.
- ▶ Use carbon dioxide instead of air.
 - Carbon dioxide helps in the recovery phase.
 - Patients report less pain following procedure.
- ▶ Use of a scope guide.

After the procedure

- ▶ Confirm that patients have a safe way home.
- ▶ Provide next-day call or follow-up appointment.
 - Emphasise that patient feedback is important and how it is used to improve patient experience.
- ▶ Provide contact information for patients to contact with questions or concerns.
- ▶ Patient rating cards provided to endoscopist (the patient rates comfort level during the procedure).
- ▶ Use patient feedback to improve.

Members of the multidisciplinary panel participated in two small group sessions during an inperson consensus-building meeting. The first small group session explored patient and contextual factors that may warrant the use of deep sedation in specific circumstances. The second small group session focused on how to improve patient experience under moderate sedation.

*The patient and contextual factors listed here are not meant to imply that deep sedation should be used if any of these factors are present. Rather, if ≥ 1 of these factors are present, endoscopists should consider, on a case-by-case basis, whether deep sedation is necessary.

results of studies reporting aspiration and bleeding rates were conflicting. There were no differences in cardiorespiratory events between patients sedated with propofol versus traditional sedatives. There were small

improvements in patient satisfaction and recovery time with propofol versus midazolam and fentanyl. Members of the multidisciplinary panel received copies of the evidence summary.

Box 1 Final practice recommendations

Definitions

- ▶ Routine colonoscopy: colonoscopy provided in the elective, ambulatory setting to stable patients (ASA category I–III). Routine colonoscopy can include procedures such as biopsy and polypectomy, but does not include interventions such as planned resection of large polyps, placement of colonic stents and so on.
- ▶ Levels of sedation: we have defined moderate and deep sedation using categories from the ASA. Of note, the level of sedation refers to the targeted level of sedation, acknowledging that occasionally patients will become more deeply sedated than intended. Such occasions must be recognised and managed, and the patient returned to the intended level of sedation. In general, when ‘traditional agents’ such as midazolam and fentanyl are used, a moderate level of sedation is targeted. In general, when propofol is used, a deep level of sedation is targeted.
 - Moderate sedation is a drug-induced depression of consciousness during which patients respond purposefully to verbal commands, either alone or accompanied by light tactile stimulation. No interventions are required to maintain a patent airway, and spontaneous ventilation is adequate. Cardiovascular function is usually maintained.
 - Deep sedation is a drug-induced depression of consciousness during which patients cannot be easily roused but respond purposefully following repeated or painful stimulation. The ability to independently maintain ventilator function may be impaired. Patients may require assistance in maintaining a patent airway, and spontaneous ventilation may be inadequate. Cardiovascular function is usually maintained.
- ▶ Endoscopist: for these statements, an endoscopist is considered a practitioner (gastroenterologist, surgeon and so on) with sufficient training and skill to perform routine colonoscopy.
- ▶ Anaesthesia provider: the term ‘anaesthesia provider’ is used to denote a practitioner with additional training in the delivery of anaesthesia and airway management. Non-physician anaesthesia providers (anaesthesia assistants) provide care under the authority of medical directives with the direct supervision of an anaesthesiologist certified by the Royal College of Physicians and Surgeons of Canada.
- ▶ Brief interruptible task: for colonoscopy, brief interruptible tasks would include charting, providing pressure, as well as assistance with biopsies and simple polypectomies. If more technical assistance is required, a second assistant should be available to assist during the more demanding portion of the procedure.

Final practice recommendations

- ▶ All endoscopists performing colonoscopy should be able to complete colonoscopy safely and effectively (per accepted benchmarks) using moderate sedation or less. (93%)
- ▶ Endoscopists unable to complete colonoscopy safely and effectively (per accepted benchmarks) using moderate sedation should undergo additional training. (89%)
- ▶ For patients undergoing routine colonoscopy, endoscopists can safely administer moderate sedation with the assistance of a trained nurse. (89%)
- ▶ For routine in-hospital colonoscopy under moderate sedation, a single RN* can both monitor the patient and perform brief interruptible tasks. (86%)
- ▶ Select patients undergoing routine colonoscopy may benefit from deep sedation. (93%)
- ▶ Institutions will not mandate the use of deep sedation for routine colonoscopy. (93%)
- ▶ Deep sedation for colonoscopy should only be administered by an anaesthesia provider. (82%)
- ▶ For routine colonoscopy under deep sedation, an anaesthesia provider will be responsible for monitoring the patient and should not be responsible for additional tasks. (93%)
- ▶ Select patients undergoing colonoscopy, such as those with severe comorbidities, may benefit from having sedation administered and monitored by an anaesthesia provider, irrespective of level of sedation. (96%)

Level of agreement (agree/strongly agree) with each practice recommendation in the final round of voting is indicated in parentheses.

*Assumes this RN is an experienced, trained endoscopy nurse.

ASA, American Society of Anesthesiologists; RN, registered nurse.

Webinars and values assessment survey

Members of the multidisciplinary panel participated in 1.5-hour webinars where the evidence summary was discussed in detail. Additionally, the webinars were used to generate a list of factors and outcomes that panelists felt were important to consider in the development of practice recommendations for the use of sedation. Following the webinars, panellists received a link to an online, anonymous values assessment survey, where they were asked to indicate the importance, on a scale from 1 to 10, of each of 20 factors/outcomes that had emerged from the webinars.

Delphi process: round 1

Based on the results of the systematic reviews and values assessment survey, the study team drafted a set of nine

preliminary practice recommendations. In the first round of the Delphi process, panellists were asked to indicate their level of agreement with and provide feedback on the nine recommendations through an online survey using a 5-point Likert scale; response frequencies were grouped into three categories: (1) agreement (strongly agree and agree), (2) unsure, and (3) disagreement (strongly disagree and disagree). A priori, consensus for agreement or disagreement was set at 60%¹⁷; >80% agreement or disagreement was considered indicative of strong consensus.

Consensus-building meeting

Panellists attended a 1-day consensus-building meeting to review the results of the values assessment survey and the first round of the Delphi process. Panellists were

Table 2 Comparison of ASGE guidelines² and consensus-based practice recommendations

Category	ASGE guidelines	Consensus-based practice recommendations
Preprocedure risk assessment	'We recommend that all patients undergoing endoscopic procedures be evaluated to assess their risk of sedation related to pre-existing medical conditions'.	No recommendations provided.
Preferred sedative agent	'We recommend that the combination of an opioid and benzodiazepine is a safe and effective regimen for achieving minimal to moderate sedation for upper endoscopy and colonoscopy in patients without risk factors for sedation-related adverse events'. 'Minimal and/or moderate sedation can be delivered safely by endoscopists to patients who are ASA Class I, II, or III. Other candidates for minimal or moderate sedation include those with a history of previously successful procedures with moderate sedation and an expectation for moderate sedation as well as those undergoing a procedure that is expected to be uncomplicated or routine'. 'We suggest using an appropriate adjunctive agent (e.g., diphenhydramine, promethazine, or droperidol) in combination with conventional sedative drugs in select clinical circumstances'.	All endoscopists performing colonoscopy should be able to complete colonoscopy safely and effectively (per accepted benchmarks) <i>Deep sedation/propofol only in select circumstances (see below).</i>
Personnel capable of administering moderate sedation	'Minimal and/or moderate sedation can be delivered safely by endoscopists to patients who are ASA Class I, II, or III. Other candidates for minimal or moderate sedation include those with a history of previously successful procedures with moderate sedation and an expectation for moderate sedation as well as those undergoing a procedure that is expected to be uncomplicated or routine'. 'Medications targeting minimal and moderate sedation generally can be administered in an incremental fashion by an appropriately trained registered nurse (RN) under the supervision of an endoscopist'.	For patients undergoing routine colonoscopy, endoscopists can safely administer moderate sedation with the assistance of a trained nurse.
Skills and training required of endoscopists	'We recommend that providers undergo specific training in the administration of endoscopic sedation and possess the skills necessary for the diagnosis and management of sedation-related adverse events, including rescue from a level of sedation deep than that intended'.	All endoscopists performing colonoscopy should be able to complete colonoscopy safely and effectively (per accepted benchmarks) Using moderate sedation or less. Endoscopists unable to complete colonoscopy safely and effectively (per accepted benchmarks) using moderate sedation should undergo additional training.
Personnel responsible for monitoring moderately sedated patients	'For moderate sedation, the personnel assigned to monitoring the patient can be assigned brief and interruptible tasks (such as mucosal biopsy), provided that the patient has not reached a state of deep sedation'.	For routine in-hospital colonoscopy under moderate sedation, a single RN can both monitor the patient and perform brief interruptible tasks.
Equipment required for monitoring sedated patients	'We recommend that routine monitoring of blood pressure, oxygen saturation, and heart rate in addition to clinical observation for changes in cardiopulmonary status during all endoscopic procedures using sedation. Supplemental oxygen administration should be considered for moderate sedation and should be administered during deep sedation. Supplemental oxygen should be administered if hypoxemia is anticipated or develops'. 'We suggest that capnography monitoring be considered for patients undergoing endoscopy targeting deep sedation'.	No recommendations provided.
Techniques to enhance patient experience under moderate sedation	No recommendations provided.	<i>Preprocedure, intraprocedural and postprocedure techniques provided (table 1).</i>
Reasons to consider use of deep sedation/propofol	'We suggest that endoscopists use propofol-based sedation (endoscopist-directed or anesthesia-provider administered) when it is expected to improve patient safety, comfort, procedural efficiency, and/or successful procedure completion'.	Select patients undergoing routine colonoscopy may benefit from deep sedation. <i>Numerous patient and context factors to consider identified (table 1).</i> Institutions will not mandate the use of deep sedation for routine colonoscopy.

Continued

Table 2 Continued

Category	ASGE guidelines	Consensus-based practice recommendations
Reasons to involve an anaesthesia provider	<p>'We recommend anaesthesia provider-administered sedation be considered for complex endoscopic procedures or patients with multiple medical comorbidities or at risk for airway compromise'.</p> <p>'Patient risk factors include significant medical conditions such as extremes of age; severe pulmonary, cardiac, renal, or hepatic disease; pregnancy; the abuse of drugs or alcohol; uncooperative patients; a potentially difficult airway for positive-pressure ventilation; and individuals with anatomy that is associated with more difficult intubation. Additionally, an anaesthesia provider may be used to provide propofol-based sedation for settings in which regulations or policies do not allow endoscopist-administered propofol, but the treating physicians judge the benefits of a propofol regimen to outweigh the risks and costs'.</p>	<p>Select patients undergoing colonoscopy, such as those with severe comorbidities, may benefit from having sedation administered and monitored by an anaesthesia provider, irrespective of level of sedation (see table 1).</p>
Personnel capable of administering deep sedation/propofol	<p>'We suggest that endoscopists use propofol-based sedation (endoscopist-directed or anaesthesia-provider administered)...'.</p> <p>'Extensive data have demonstrated the safety and efficacy of non-anaesthesiologist-administered propofol sedation (NAAP)...NAAP requires specialized training, patient selection, and personnel dedicated to continuous physiologic monitoring. Regulations regarding administration of propofol are determined at the state, regional, and local levels regardless of targeted level of sedation. As a result, the practice of NAAP is quite limited nationally. Hence, propofol-based sedation for low-risk patients undergoing routine procedures often is administered by anaesthesia personnel'.</p>	<p>Deep sedation for colonoscopy should only be administered by an anaesthesia provider.</p>
Personnel responsible for monitoring deeply sedated patients	<p>'For deep sedation, personnel assigned to monitoring the patient must do so in a continuous and uninterrupted fashion'.</p>	<p>For routine colonoscopy under deep sedation, an anaesthesia provider will be responsible for monitoring the patient and should not be responsible for additional tasks.</p>

ASA, American Society of Anesthesiologists; ASGE, American Society for Gastrointestinal Endoscopy.

presented with data on the level of agreement/disagreement for each preliminary recommendation. Wording of recommendations was then refined through large group discussions. The meeting also included presentations by visiting experts, covering topics of discussion that had arisen during the webinars, and small group breakout sessions. The purpose of the small group sessions was to discuss (1) patients for whom deep sedation would be beneficial and (2) how to enhance the experience for patients undergoing routine colonoscopy with moderate sedation. Notes were taken during the small group and large group sessions to facilitate modifications to the practice recommendations based on the inperson discussion.

Delphi process: round 2

Following the consensus-building meeting, the practice recommendations were reworded and definitions of terms used were clarified. The multidisciplinary panel participated in a second round of online voting on the reworded practice recommendations, after which a set of final recommendations was drafted by the study team.

Patient and public involvement

We included public representatives in the expert panel to provide patient and public perspectives. These individuals participated in all aspects of the study involving the expert panel, including values assessment, webinars, Delphi process, and consensus-building meeting. We incorporated feedback received from these representatives into the recommendations provided.

RESULTS

Panel composition

The multidisciplinary panel consisted of 32 members (online supplementary table 1), including 5 (16%) anaesthesiologists, 9 (28%) endoscopists (7 gastroenterologists, 2 surgeons), 4 (13%) Cancer Care Ontario representatives, 3 (9%) public representatives, 3 (9%) administrators/funders, 2 (6%) health economics experts, 3 (9%) endoscopy nurses, 2 (6%) hospital endoscopy programme managers, and 1 (3%) individual from the College of Physicians and Surgeons of Ontario. Two of the endoscopists were international experts.

Values assessment

Twenty-nine panellists participated in the values assessment survey. The multidisciplinary panel deemed all 20 factors/outcomes included in the values assessment survey to be important considerations for the development of practice recommendations (figure 2). Factors judged as the most important (mean scores of 9–10 on a 10-point scale) were related to the safety of sedatives, including the outcomes of death, cardiorespiratory complications, and procedural complications. Patient-reported pain and factors related to colonoscopy quality (eg, caecal intubation rate, adenoma detection rate) were considered very important (mean score of 8); factors

related to procedural efficiency and cost were considered less important (mean scores of 6–7).

Delphi process round 1: preliminary practice recommendations

Based on the evidence review and results of the values assessment survey, nine preliminary practice recommendations were developed (online supplementary table 2) and presented to the multidisciplinary panel for ranking and comments. Although the safety of sedatives was considered the most important factor in the values assessment survey, preliminary practice recommendations did not include a statement favouring particular sedatives as the evidence review did not find any significant differences in safety outcomes between sedatives. Twenty-nine panellists participated in the first round of the Delphi process. Consensus was reached for all recommendations; strong consensus was observed for six of the nine recommendations (online supplementary table 2).

Consensus-building meeting

Thirty-one panellists attended the 1-day inperson meeting. As consensus was reached for all recommendations presented in the first round of online voting, the inperson meeting was used to present and discuss the results of voting, clarify wording of recommendations, discuss the recommendations for which strong consensus had not been achieved, and facilitate small group breakout sessions.

Small group session 1: deep sedation for routine colonoscopy

In the first small group session, participants identified patient groups that may warrant consideration for the use of deep sedation in specific circumstances, although it was recognised that these patient and clinical contexts do not necessarily mandate the use of deep sedation, that is, use should be considered on a case-by-case basis. Patient factors identified by panellists that may warrant consideration for use of deep selection in select cases included chronic opiate users, those who could not previously tolerate colonoscopy under moderate sedation due to discomfort, paediatric patients, and patients with hypersensitivities to traditional sedative agents, a history of sexual abuse, irritable bowel syndrome, fibromyalgia, cognitive disabilities (eg, dementia), or a history of diverticulitis. Contextual factors that could be considered when weighing the use of deep sedation included complex procedures (eg, endoscopic mucosal resection), lengthy procedures, and performance of colonoscopy at small centres where revenue generated for anaesthesiologists through endoscopy is required to maintain the anaesthesiologist workforce in the region (table 1).

Small group session 2: enhancing patient experience under moderate sedation

Several strategies for enhancing patient experience under moderate sedation were identified by panellists (table 1). Methods suggested to be undertaken prior to the procedure included skills enhancement courses and

setting patient expectations surrounding pain/discomfort. Suggested methods to be used during the procedure included appropriate titration of sedatives, use of abdominal pressure, variation in patient positioning, and use of carbon dioxide rather than air for insufflation. Postprocedure methods included follow-up phone calls or appointments for solicitation of feedback on the experience.

Delphi process round 2: final practice recommendations

Twenty-eight panellists participated in the second round of the Delphi, where they were presented with the nine reworded practice recommendations (box 1, online supplementary table 3). Strong consensus was achieved for all recommendations.

INTERPRETATION

Using a multidisciplinary panel and modified Delphi methodology, we developed nine consensus-based practice recommendations for the use of sedation in routine hospital-based colonoscopy. Participants agreed that all endoscopists should be able to perform colonoscopy under moderate sedation with the assistance of a single trained nurse. Select patients, including those with severe comorbidities, may benefit from deep sedation and monitoring by an anaesthesia provider; however, deep sedation should not be mandated by hospitals.

The recommendations developed in our study largely align with recent guidelines put forth by the American Society for Gastrointestinal Endoscopy (ASGE) (table 2).² These guidelines reinforce that moderate sedation can be administered by an endoscopist for most patients and that a trained nurse can both monitor a moderately sedated patient and perform brief, interruptible tasks. However, our recommendations differ from those of the ASGE in several key areas (table 2). Unique to our recommendations, participants agreed that endoscopists who are unable to perform colonoscopy with moderate sedation should undergo further training. Due to the high use of propofol at some hospitals, there may be endoscopists who have not performed colonoscopy under moderate sedation at all or for some time who would feel uncomfortable doing so. For these individuals, skills enhancement courses can be beneficial by providing strategies to minimise patient discomfort and improve caecal intubation and adenoma detection rates, such as by teaching torque steering and loop reduction, varying patient positioning to facilitate passage of the colonoscope,^{22–26} and use of carbon dioxide insufflation^{27–29} and water infusion techniques.^{30–32} Further recommendations made by the expert panel for improving patient experience under moderate sedation are provided in table 1. The ASGE document recommended that propofol be considered if it is expected to improve patient safety, comfort, procedural efficiency, or success, but suggested that whether an anaesthesiologist is involved in administration of deep sedation for low-risk cases be determined

by state, regional, and local regulations. In contrast, the expert panel recommended that deep sedation only be administered by anaesthesia personnel, and identified specific patient and clinical contexts, not discussed in the ASGE document, that warrant consideration for use of propofol. These include select patients with irritable bowel syndrome, fibromyalgia, or previous diverticulitis, as well as smaller hospitals where revenue from providing endoscopic sedation is required to maintain the anaesthesiologist workforce. Both groups agreed that when deep sedation is used, an individual dedicated to administering and monitoring deep sedation, without any other responsibilities, is required.

The systematic reviews that informed our consensus-building process also identified multiple areas in which evidence is currently lacking or conflicting. We found many discrepancies among current guidelines relating to the administration of sedation. Our review of the safety of non-anaesthesia provider-administered sedation found conflicting results for the outcomes of aspiration and bleeding. Additionally, we found that propofol was associated with small improvements in patient satisfaction, but it was unclear whether differences of this magnitude were clinically meaningful. Further research in these areas will be important to providing evidence-based guidance for sedation.

Although our study focused on the Ontario hospital context, high use of deep sedation and variability in practices are a growing issue in other jurisdictions as well. These increases cannot be attributed to a changing patient demographic or case complexity. In a study of the use of monitored anaesthesia care (MAC) in the Veterans Health Administration, facility-level factors had stronger associations with MAC use than patient-level factors.^{33 34} Similarly, in Canada, an analysis by Cancer Care Ontario found that large endoscopy units, where complex procedures are most likely to be performed, had high variability in the use of anaesthesia, indicating that procedural complexity was not the main driver of anaesthesia use (data available on request).

Our study has limitations. Although we selected a diverse panel, sedation practices vary considerably by jurisdiction and members of our panel may not reflect the breadth of practice in all clinical settings, although they do adequately reflect a range of hospital-based practice models. Additionally, as recommended by the Guidelines International Network,³⁵ the US Institute of Medicine/National Academy of Medicine,³⁶ and the UK National Institute for Health and Care Excellence,³⁷ we included public representatives in our panel to ensure incorporation of patient priorities and preferences. As most of our panellists were from Ontario, the recommendations developed in this study may be less applicable elsewhere; however, the panellists from outside of Ontario found the recommendations to be reflective of their experiences as well. Importantly, our reviews and recommendations did not evaluate long-term outcomes related to the effectiveness of colonoscopy.

Although our recommendations focused on hospital-based colonoscopy, the use of deep sedation has increased for other endoscopic procedures, including gastroscopy and flexible sigmoidoscopy performed in hospitals over the same period.^{38 39} Given that the role of deep sedation for these procedures is even more controversial,^{40 41} recommendations pertaining to other endoscopic procedures may also be helpful. Additionally, as consensus was built for our recommendations in the context of hospital-based colonoscopies, our recommendations may not be generalisable to endoscopy performed in the clinic-based setting, where other factors may necessitate the use of anaesthesia services. Further work will be required to create recommendations for sedation practices for colonoscopy performed in clinic-based facilities.

In conclusion, using a diverse panel and modified Delphi process, we developed nine consensus-based practice recommendations to guide use of sedation for routine hospital-based colonoscopy. The results can be used to inform hospital policy with respect to the use of sedation for in-hospital colonoscopy and can help facilitate the standardisation of sedation practices.

Author affiliations

¹Division of General Surgery, Department of Surgery, University of Toronto, Toronto, Ontario, Canada

²Institute of Health Policy, Management, and Evaluation, Dalla Lana School of Public Health, University of Toronto, Toronto, Ontario, Canada

³Department of Medicine, The Ottawa Hospital, University of Ottawa, Ottawa, Ontario, Canada

⁴Cancer Care Ontario, Toronto, Ontario, Canada

⁵Sunnybrook Health Sciences Centre, Toronto, Ontario, Canada

⁶Department of General Surgery, St Michael's Hospital, Toronto, Ontario, Canada

⁷Department of Medicine, Division of Gastroenterology, VA Puget Sound Health Care System and University of Washington School of Medicine, Seattle, Washington, USA

Twitter Fahima Dossa @fdossa and Nancy N Baxter @enenbee

Acknowledgements The authors would like to thank the members of the multidisciplinary panel for their time and contributions to this work. Additionally, the authors thank Anna Gagliardi for her work in planning and facilitating the multidisciplinary panel meeting, and Michelle Helm, Zahrah Khalid, Melissa Coulson, and Shamara Baidobonso for their involvement in the meeting.

Contributors Conception and design: FD, CD, JT, AS, LR, BM, NB. Analysis and interpretation of data: FD, AS, JAD, NB. Drafting of the article: FD. Critical revision of the article for important intellectual content: FD, CD, JT, AS, LR, BM, JAD, NB. Final approval of the article: FD, CD, JT, AS, LR, BM, JAD, NB.

Funding This study was funded by the Canadian Institutes of Health Research (CIHR) Foundation Grant (no 148470) and with the support of Cancer Care Ontario (CCO) through funding provided by the Government of Ontario. The opinions, results, view, and conclusions reported in this publication are those of the authors and do not necessarily reflect those of Cancer Care Ontario.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval This study was approved by the Research Ethics Board at St Michael's Hospital.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made

indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Fahima Dossa <http://orcid.org/0000-0002-4670-7445>

REFERENCES

- Harewood GC, Wiersema MJ, Melton LJ. A prospective, controlled assessment of factors influencing acceptance of screening colonoscopy. *Am J Gastroenterol* 2002;97:3186–94.
- Early DS, Lightdale JR, Vargo JJ, et al. Guidelines for sedation and anesthesia in GI endoscopy. *Gastrointest Endosc* 2018;87:327–37.
- American Society of Anesthesiologists. practice guidelines for sedation and analgesia by non-anesthesiologists. *Anesthesiology* 2002;96:1004–17.
- Baxter Corporation. Propofol product monograph, 2017. Available: https://www.baxter.ca/sites/g/files/ebysai1431/files/2018-11/Propofol_EN.pdf [Accessed 15 Feb 2019].
- DIPRIVAN 3 (propofol) Injectable Emulsion - FDA, 2008. Available: https://www.accessdata.fda.gov/drugsatfda_docs/label/2008/019627s046lbl.pdf [Accessed 16 Feb 2019].
- College of Physicians and Surgeons of Ontario. Out-Of-Hospital premises inspection program (OHPIP) program standards 2017.
- Chutkan R, Cohen J, Abedi M, et al. Training guideline for use of propofol in gastrointestinal endoscopy. *Gastrointest Endosc* 2004;60:167–72.
- Faulx AL, Vela S, Das A, et al. The changing landscape of practice patterns regarding unsedated endoscopy and propofol use: a national web survey. *Gastrointest Endosc* 2005;62:9–15.
- Grocott HP. Propofol sedation improves efficiency and optimizes patient satisfaction during colonoscopy. *Can Med Assoc J* 2018;190:E751–E51.
- Singh H, Poluha W, Cheang M, et al. Propofol for sedation during colonoscopy. *Cochrane Database Syst Rev* 2008;50.
- Ladas SD, Satake Y, Mostafa I, et al. Sedation practices for gastrointestinal endoscopy in Europe, North America, Asia, Africa and Australia. *Digestion* 2010;82:74–6.
- Gavin DR, Valori RM, Anderson JT, et al. The National colonoscopy audit: a nationwide assessment of the quality and safety of colonoscopy in the UK. *Gut* 2013;62:242–9.
- Porostocky P, Chiba N, Colacino P, et al. A survey of sedation practices for colonoscopy in Canada. *Can J Gastroenterol* 2011;25:255–60.
- Alharbi O, Rabeneck L, Paszat LF, et al. A population-based analysis of outpatient colonoscopy in adults assisted by an anesthesiologist. *Anesthesiology* 2009;111:734–40.
- Cooper GS, Kou TD, Rex DK. Complications following colonoscopy with anesthesia assistance: a population-based analysis. *JAMA Intern Med* 2013;173:551–6.
- Bielawska B, Hookey LC, Sutradhar R, et al. Anesthesia assistance in outpatient colonoscopy and risk of aspiration pneumonia, bowel perforation, and splenic injury. *Gastroenterology* 2018;154:77–85.
- Fink A, Kosecoff J, Chassin M, et al. Consensus methods: characteristics and guidelines for use. *Am J Public Health* 1984;74:979–83.
- Boukheld R, Abdoul H, Loustau M, et al. Using and reporting the Delphi method for selecting healthcare quality indicators: a systematic review. *PLoS One* 2011;6:e20476.
- Jones J, Hunter D. Consensus methods for medical and health services research. *BMJ* 1995;311:376–80.
- Jairath N, Weinstein J. The Delphi methodology (Part one): a useful administrative approach. *Can J Nurs Adm* 1994;7:29–42.
- Jünger S, Payne SA, Brine J, et al. Guidance on conducting and reporting Delphi studies (CREDES) in palliative care: recommendations based on a methodological systematic review. *Palliat Med* 2017;31:684–706.
- East JE, Bassett P, Arebi N, et al. Dynamic patient position changes during colonoscopy withdrawal increase adenoma detection: a randomized, crossover trial. *Gastrointest Endosc* 2011;73:456–63.
- Ball AJ, Johal SS, Riley SA. Position change during colonoscopy withdrawal increases polyp and adenoma detection in the right but not in the left side of the colon: results of a randomized controlled trial. *Gastrointest Endosc* 2015;82:488–94.
- Wilson A, Saunders BP. Position change during colonoscopy: the oldest and best trick in the book. *Gastrointest Endosc* 2015;82:495–6.
- Lee S-W, Chang JH, Ji J-S, et al. Effect of dynamic position changes on adenoma detection during colonoscopy withdrawal:

- a randomized controlled multicenter trial. *Am J Gastroenterol* 2016;111:63–9.
- 26 Vergis N, McGrath AK, Stoddart CH, *et al.* Right or left in colonoscopy (ROLCOL)? A randomized controlled trial of right- versus left-sided starting position in colonoscopy. *Am J Gastroenterol* 2015;110:1576–81.
 - 27 Wang WL, Wu ZH, Sun Q, *et al.* Meta-Analysis: the use of carbon dioxide insufflation vs. room air insufflation for gastrointestinal endoscopy. *Aliment Pharmacol Ther* 2012;35:1145–54.
 - 28 Sajid MS, Caswell J, Bhatti MI, *et al.* Carbon dioxide insufflation vs conventional air insufflation for colonoscopy: a systematic review and meta-analysis of published randomized controlled trials. *Colorectal Dis* 2015;17:111–23.
 - 29 Memon MA, Memon B, Yunus RM, *et al.* Carbon dioxide versus air insufflation for elective colonoscopy: a meta-analysis and systematic review of randomized controlled trials. *Surg Laparosc Endosc Percutan Tech* 2016;26:102–16.
 - 30 Hafner S, Zolk K, Radaelli F, *et al.* Water infusion versus air insufflation for colonoscopy. *Cochrane Database Syst Rev* 2015;56.
 - 31 Cadoni S, Hassan C, Frazzoni L, *et al.* Impact of water exchange colonoscopy on endoscopy room efficiency: a systematic review and meta-analysis. *Gastrointest Endosc* 2019;89:159–67.
 - 32 Fuccio L, Frazzoni L, Hassan C, *et al.* Water exchange colonoscopy increases adenoma detection rate: a systematic review with network meta-analysis of randomized controlled studies. *Gastrointest Endosc* 2018;88:589–97.
 - 33 Adams MA, Prenovost KM, Dornitz JA, *et al.* Predictors of use of monitored anesthesia care for outpatient gastrointestinal endoscopy in a capitated payment system. *Gastroenterology* 2017;153:1496–503.
 - 34 Adams MA, Prenovost KM, Dornitz JA, *et al.* National trends in use of monitored anesthesia care for outpatient gastrointestinal endoscopy in the Veterans health administration. *JAMA Intern Med* 2017;177:436–8.
 - 35 Qaseem A *et al.* Guidelines international network: toward international standards for clinical practice guidelines. *Ann Intern Med* 2012;156:525–31.
 - 36 Graham R, Mancher M, Miller Wolman D. Committee on Standards for Developing Trustworthy Clinical Practice Guidelines. In: *Clinical practice guidelines we can trust*. Washington, DC: The National Academies Press, 2011. <http://www.nationalacademies.org/hmd/Reports/2011/Clinical-Practice-Guidelines-We-Can-Trust.aspx>
 - 37 Jarrett L. *Patient involvement unit (PIU). A report on a study to evaluate patient/carer membership of the first NICE Guideline development groups*. London: National Institute for Clinical Excellence (NICE), 2004. <https://www.nice.org.uk/media/default/About/NICE-Communities/Public-involvement/Public-involvement-programme/PIU-GDG-evaluation-report-2004-1.pdf>
 - 38 Liu H, Waxman DA, Main R, *et al.* Utilization of anesthesia services during outpatient endoscopies and colonoscopies and associated spending in 2003–2009. *JAMA* 2012;307:1178–84.
 - 39 Predmore Z, Nie X, Main R, *et al.* Anesthesia service use during outpatient gastroenterology procedures continued to increase from 2010 to 2013 and potentially discretionary spending remained high. *Am J Gastroenterol* 2017;112:297–302.
 - 40 Garg PK, Singh AP, Jain BK, *et al.* Safety and acceptance of non-sedated upper gastrointestinal endoscopy: a prospective observational study. *J Laparoendosc Adv Surg Tech A* 2012;22:315–8.
 - 41 Ferrer Rosique Juan Ángel, Canaval Zuleta HJ, Cacho Acosta G. Patients' perspective on sedation during upper gastrointestinal endoscopy. systematic use of sedation or systematic prior information? *Rev Esp Enferm Dig* 2018;110:262–3.