

Title	On orientations of real algebraic curves determined by spin structures (The theory of transformation groups and its applications)
Author(s)	Nagami, Seiji
Citation	数理解析研究所講究録 = RIMS Kokyuroku (2019), 2135: 70-73
Issue Date	2019-11
URL	<a href="http://hdl.handle.net/2433/254824">http://hdl.handle.net/2433/254824</a>
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

# On orientations of real algebraic curves determined by spin structures

Seiji Nagami  
Academic Support Center,  
Setsunan University

## 1 Introduction

It is a classical problem to decide possible arrangements of ovals of real algebraic curves in the real projective plane  $\mathbf{R}P^2$ , which is known as the Hilbert's 16'th problem([2]). Harnack was the first to show that the number of components of ovals of a given non-singular real algebraic curve of degree  $m$  is less than or equal to  $g(m) = (m-1)(m-2)/2$  ([1]), which is known as Harnack's inequality.

In [4], Rokhlin introduced 'complex orientations' on the algebraic curves of even type, which turned out to be useful for this problem. So searching canonical orientations on a given algebraic curve seems interesting problem.

## 2 Orientation by Wang

In [5], by using spin structures, Wang gave orientations of fixed point set  $E_{\mathbf{R}}$  of a complex conjugate involution  $\sigma_E : E \rightarrow E$  on a complex vector bundle  $\pi : E \rightarrow X$  that covers an involution  $\sigma : X \rightarrow X$  on a closed smooth manifold  $X$ . Note that, in case  $X = \mathbf{C}A = \{[z_0; z_1; z_2] \in \mathbf{C}P^2 | A(z_0, z_1, z_2) = 0\}$ ,  $E = T\mathbf{C}A$ , and  $\sigma_E$  is the differential of the involution  $\sigma : X \rightarrow X$  given by  $\sigma([z_0; z_1; z_2]) = [\bar{z}_0; \bar{z}_1; \bar{z}_2]$ ,  $E_{\mathbf{R}}$  is the total space of real vector bundle of the real algebraic curve  $\mathbf{R}A = \{[z_0; z_1; z_2] \in \mathbf{R}P^2 | A(z_0, z_1, z_2) = 0\}$ , where  $A$  is a real non-singular homogenous polynomial of three variables.

In this section, we see the construction given in [5]. Let  $\mathbf{U}(E)$  denote the  $\mathbf{U}(r)$ -frame bundle for  $E$  and  $\sigma' : \mathbf{U}(E) \rightarrow \mathbf{U}(E)$  its induced involution. Set  $P = \mathbf{U}(E) \times_i \mathbf{SO}(2r)$ , where  $i : \mathbf{U}(r) \rightarrow \mathbf{SO}(2r)$  is given by  $i(X + \sqrt{-1}Y) = \begin{pmatrix} X & -Y \\ Y & X \end{pmatrix}$ . Note that  $i(\overline{X + \sqrt{-1}Y}) = \begin{pmatrix} X & Y \\ -Y & X \end{pmatrix} = T \begin{pmatrix} X & -Y \\ Y & X \end{pmatrix} T^{-1}$ , where  $T = \begin{pmatrix} E & O \\ O & -E \end{pmatrix}$ .

**Definition 2.1** We define the involution  $\sigma_P : P \rightarrow P$  by  $\sigma_P([V, M]) = [\sigma_E(V), TMT^{-1}]$  for  $V \in \mathbf{U}(r)$  and  $M \in \mathbf{SO}(2r)$ .

Suppose that the principal  $\mathbf{SO}(2r)$  bundle  $P \rightarrow X$  admits a spin structure  $\xi \in H^1(P; \mathbf{Z}_2)$ , i.e., a double cover  $\tilde{P} \rightarrow P$  such that the composite  $\tilde{P} \rightarrow P \rightarrow X$  is a principal  $\mathbf{Spin}(2r)$  bundle whose restriction to a point  $* \in X$  is the non-trivial one.

**Definition 2.2** The involution  $\sigma_E : E \rightarrow E$  is compatible with  $\xi \in H^1(P; \mathbf{Z}_2)$  if and only if there exists a bundle automorphism  $\tilde{\sigma}_P : \tilde{P} \rightarrow \tilde{P}$  such that  $\tilde{\sigma}_P(xg) = \tilde{\sigma}_P(x)\bar{g}$  holds for any  $g \in \mathbf{Spin}(2r) \subset \mathbf{Cl}(\mathbf{R}^{2r}) = \mathbf{Cl}(\mathbf{C}^r)$ , and that the following diagram commutes;

$$\begin{array}{ccc} \tilde{P} & \xrightarrow{\tilde{\sigma}_P} & \tilde{P} \\ \downarrow & & \downarrow \\ P & \xrightarrow{\sigma_P} & P. \end{array}$$

We say that  $\sigma_P$  is compatible with a spin structure  $\xi$ , and that  $\tilde{\sigma}_P$  a conjugate lift of  $\sigma_P$ .

With this notation, Wang showed in [5];

**Theorem 2.1** Suppose that  $\sigma_E : E \rightarrow E$  is compatible with a spin structure  $\xi \in H^1(P; \mathbf{Z}_2)$ . Then for each conjugate lift  $\tilde{\sigma}_E : P_\xi \rightarrow P_\xi$  there is a canonical orientation of real vector bundle  $E_{\mathbf{R}} \rightarrow X_{\mathbf{R}}$ , where  $X_{\mathbf{R}}$  denotes the fixed point set of the involution  $\sigma : X \rightarrow X$ .

(Case 1) First suppose that  $E \rightarrow X$  is a complex line bundle. Then  $P \rightarrow X$  is a principal  $\mathbf{U}(1)$  bundle. Let  $P^\sigma$  denote the fixed point set of  $\sigma_P$ . Then  $P^\sigma \rightarrow X_{\mathbf{R}}$  is a principal  $\mathbf{Z}_2$  bundle associated with  $E_{\mathbf{R}} \rightarrow X_{\mathbf{R}}$ . By our assumption, we have a spin structure  $P_\xi \rightarrow P$  and a conjugate morphism  $\sigma_\xi : P_\xi \rightarrow P_\xi$  such that the following diagram commutes;

$$\begin{array}{ccc} P_\xi & \xrightarrow{\sigma_\xi} & P_\xi \\ \downarrow & & \downarrow \\ P & \xrightarrow{\sigma_P} & P \\ \downarrow & & \downarrow \\ X & \xrightarrow{\sigma} & X. \end{array}$$

Since  $\sigma_\xi$  is conjugate,  $P_\xi^\sigma \rightarrow X_{\mathbf{R}}$  also is principal  $\mathbf{Z}_2$  bundle. Let  $x \in X_{\mathbf{R}}$ , and set  $\pi^{-1}(x) = \{a, b\}$ . Then we have that  $b = -a$ . Therefore any fiber of  $P_\xi^\sigma \rightarrow X_{\mathbf{R}}$  is sent to one point. This implies that  $P_\xi^\sigma$  determines a section of  $P^\sigma \rightarrow X_{\mathbf{R}}$ . Thus we obtain an orientation of  $E_{\mathbf{R}} = P^\sigma \times_{\mathbf{Z}_2} \mathbf{R} \rightarrow X_{\mathbf{R}}$ .

(Case 2) When the complex dimension of the complex vector bundle  $E \rightarrow X$  is greater than 1, we apply the method of Case 1 to the complex line bundle  $\det_{\mathbf{C}} E \rightarrow X$ .

### 3 Construction of orientations by using pin structures

Let  $L \rightarrow X$  be a complex line bundle and  $\sigma_L : L \rightarrow L$  a complex conjugate involution such that the following diagram commutes;

$$\begin{array}{ccc} L & \xrightarrow{\sigma_L} & L \\ \downarrow & & \downarrow \\ X & \xrightarrow{\sigma} & X. \end{array}$$

Let  $\mathbf{U}(1) \rightarrow \mathbf{U}(L) \rightarrow X$ ,  $\mathbf{SO}(2) \rightarrow \mathbf{SO}(L) \rightarrow X$ , and  $\mathbf{O}(2) \rightarrow \mathbf{O}(L) \rightarrow X$  denote the principal bundles associated with  $L \rightarrow X$ .

Then for each  $e \in \mathbf{U}(L)$ , set  $\sigma(e) = \alpha e$ , where  $\alpha = a + \sqrt{-1}b \in \mathbf{U}(1)$ . Then we have that  $\sigma(\sqrt{-1}e) = -\sqrt{-1}\sigma(e) = (b - \sqrt{-1}a)e$ . Thus we obtain that, by setting  $M_\alpha = \begin{pmatrix} a & b \\ b & -a \end{pmatrix}$ ,  $\sigma(e, \sqrt{-1}e) = (e, \sqrt{-1}e)M_\alpha = (e, \sqrt{-1}e)R_\alpha T$ , where  $R_\alpha = \begin{pmatrix} a & -b \\ b & a \end{pmatrix}$  and  $T = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ . Define  $i_P : P = \mathbf{U}(L) \times_\rho \mathbf{SO}(2) \rightarrow \mathbf{SO}(L)$  by  $i_P([e, A]) = (e, \sqrt{-1}e)A$ . Then we have the following commutative diagram, where  $R_T$  denotes the right multiplication by  $T$ ;

$$\begin{array}{ccc} P & \xrightarrow{\sigma_U} & P \\ i_P \downarrow & & i_P \downarrow \\ \mathbf{SO}(L) & \xrightarrow{R_T \circ \sigma_U} & \mathbf{SO}(L). \end{array}$$

Therefore the following diagram commutes between the induced homomorphisms of cohomology;

$$\begin{array}{ccc} H^1(P; \mathbf{Z}_2) & \xleftarrow{\sigma_U^*} & H^1(P; \mathbf{Z}_2) \\ i_P^* \uparrow & & i_P^* \uparrow \\ H^1(\mathbf{SO}(L); \mathbf{Z}_2) & \xleftarrow{\sigma_L^* \circ R_T^*} & H^1(\mathbf{SO}(L); \mathbf{Z}_2) \end{array}$$

**Definition 3.1** For a spin structure  $\xi \in H^1(\mathbf{SO}(L); \mathbf{Z}_2)$ , set  $\tilde{\xi} = \xi \oplus R_T^*(\xi) \in H^1(\mathbf{SO}(L); \mathbf{Z}_2) \oplus H^1(\mathbf{SO}(L)T; \mathbf{Z}_2) = H^1(\mathbf{O}(L); \mathbf{Z}_2)$ .

With this definition, we have the following;

**Proposition 3.1**  $i_P^*(\xi) \in H^1(P; \mathbf{Z}_2)$  is compatible with  $\sigma_U$  if and only if  $\tilde{\xi} \in H^1(\mathbf{O}(L); \mathbf{Z}_2)$  is a pin structure that is preserved by  $\sigma$ .

By a similar method in [3], we obtain a section  $\tilde{d}\sigma \in \Gamma(\text{Ad}(\mathbf{Pin}(L)|_{X_{\mathbf{R}}})) \subset \Gamma(\mathbf{Pin})(X)|_{X_{\mathbf{R}}} \times_{\text{Ad}} \mathbf{Cl}(2) \cong \Gamma(\wedge^* L|_F)$  via  $\tilde{\xi}$ .

Fix a point  $x \in X_{\mathbf{R}}$ . Then we may assume that  $L_x = \mathbf{R}\langle u, w \rangle$  and that  $\sigma_L|_x$  is the reflection determined by  $v$ . Thus we have that  $\tilde{d}\sigma|_x = \pm v$ .

Whether this section coincides with the section given in Section 2 or not should be investigated.

## References

- [1] Harnack. Uber vieltheiligkeit der ebenen algebraischen curven. *Math. Ann.*, 10:189–99, 1876.
- [2] David Hilbert. Mathematische probleme. *Nachrichten von der Koniglichen Gesellschaft der Wissenschaften zu Göttingen*, 1900.

- [3] Kaoru Ono et al. On a theorem of edmonds. In *Progress in differential geometry*, pages 243–245. Mathematical Society of Japan, 1993.
- [4] Vladimir Abramovich Rokhlin. Complex orientations of real algebraic curves. *Functional Analysis and its Applications*, 8(4):331–334, 1974.
- [5] Shuguang Wang. Orientability of real parts and spin structures. *JP J. Geom. Topol*, 7(1):159–174, 2007.