



TITLE:

Effective quantum kinetic theory for spin transport of fermions with collisional effects

AUTHOR(S):

Yang, Di-Lun; Hattori, Koichi; Hidaka, Yoshimasa

CITATION:

Yang, Di-Lun ...[et al]. Effective quantum kinetic theory for spin transport of fermions with collisional effects. Journal of High Energy Physics 2020, 2020(7): 70.

ISSUE DATE:

2020-07

URL:

<http://hdl.handle.net/2433/255130>

RIGHT:

© The Authors. This article is distributed under the terms of the Creative Commons Attribution License (CC-BY 4.0), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited



PUBLISHED FOR SISSA BY SPRINGER

RECEIVED: February 15, 2020

REVISED: May 28, 2020

ACCEPTED: June 18, 2020

PUBLISHED: July 10, 2020

Effective quantum kinetic theory for spin transport of fermions with collisional effects

Di-Lun Yang,^a Koichi Hattori^b and Yoshimasa Hidaka^{c,d}

^a*Faculty of Science and Technology, Keio University,
Yokohama 223-8522, Japan*

^b*Yukawa Institute for Theoretical Physics, Kyoto University,
Kyoto 606-8502, Japan*

^c*RIKEN Nishina Center, RIKEN,
Wako, Saitama 351-0198, Japan*

^d*RIKEN iTHEMS, RIKEN,
Wako, Saitama 351-0198, Japan*

E-mail: dilunyang@gmail.com, koichi.hattori@yukawa.kyoto-u.ac.jp,
hidaka@post.kek.jp

ABSTRACT: We systematically derive the collision term for the axial kinetic theory, a quantum kinetic theory delineating the coupled dynamics of the vector/axial charges and spin transport carried by the massive spin-1/2 fermions traversing a medium. We employ the Wigner-function approach and propose a consistent power-counting scheme where the axial-charge distribution function, a non-conserved quantity for massive particles, is accounted as the first-order quantity in the \hbar expansion, while the vector-charge distribution function the zeroth-order quantity. This specific power-counting scheme allows us to organize a reduced \hbar expansion for the collision term and to formally identify the spin-diffusion effect and the spin-polarization effect at the same order. We confirm that the obtained collisional axial kinetic theory smoothly reduces to the chiral kinetic theory in the massless limit, serving as a consistency check. In the absence of electromagnetic fields, we further present the simplified axial kinetic equations suitable for tracking dynamical spin polarization of heavy and light fermions, respectively. As an application to the weakly coupled quark-gluon plasma at high temperature, we compute the spin-diffusion term for massive quarks within the leading-log approximation. The formal expression for the first-order terms provides a path toward evaluation of the spin polarization effect in quantum chromodynamics.

KEYWORDS: Heavy Ion Phenomenology, QCD Phenomenology

ARXIV EPRINT: [2002.02612](https://arxiv.org/abs/2002.02612)

Contents

| | | |
|----------|---|-----------|
| 1 | Introduction | 1 |
| 2 | Wigner-function approach with collisions | 3 |
| 2.1 | Full master equations | 3 |
| 2.2 | Setting the power-counting scheme | 7 |
| 2.3 | Effective master equations and constrains | 9 |
| 3 | Effective axial kinetic theory with collisions | 11 |
| 3.1 | Axial kinetic equation with general frame vector and its massless limit | 11 |
| 3.2 | Rest-frame expression | 14 |
| 3.3 | \hbar sorting with the present order counting | 15 |
| 3.4 | Discussions for the collision terms | 16 |
| 4 | Example: spin diffusion of quarks in weakly coupled QGP | 17 |
| 4.1 | Scattering between massive fermions and a medium | 17 |
| 4.2 | Weakly coupled QGP and hard-thermal-loop approximation | 19 |
| 4.3 | SKE and AKE with diffusion effects in the leading-log approximation | 21 |
| 4.3.1 | Results | 22 |
| 4.3.2 | Nonrelativistic limit | 23 |
| 4.3.3 | Consistency checks | 23 |
| 5 | Concluding remarks and outlook | 24 |
| A | Derivation of the master equations | 26 |
| A.1 | Spinor decomposition | 26 |
| A.2 | Eliminating \mathcal{S} , \mathcal{P} and $\mathcal{S}_{\mu\nu}$ | 30 |
| B | Angular-momentum decomposition for fermions and spin polarization | 31 |
| C | Decomposition of the collision term in AKE | 32 |
| D | Derivation of the spin diffusion for SKE | 35 |
| E | Derivation of the spin diffusion for AKE | 38 |
| E.1 | Half of the terms: $[Q_2^{\mu\nu} \tilde{a}_{q'\nu} - Q_1 \tilde{a}_q^\mu]$ | 38 |
| E.1.1 | Evaluating $Q_{\text{cl}(a)}^\mu$ | 40 |
| E.1.2 | Evaluating $Q_{\text{cl}(b)}^\mu$ | 40 |
| E.1.3 | Evaluating $Q_{\text{cl}(c)}^\mu$ | 42 |
| E.2 | Rest of the terms: $[Q_2^{\mu\nu} f_{Vq} \tilde{a}_{q'\nu} - Q_1(1 - f_{Vq'}) \tilde{a}_q^\mu]$ | 42 |
| E.3 | Assembling all the pieces | 44 |

JHEP07(2020)070

1 Introduction

The observations of global polarization of Λ hyperons in heavy ion collisions (HIC) have triggered intensive studies for the spin polarization of relativistic fermions [1, 2]. In particular, the experimental measurements have been motivated by theoretical proposals upon the polarization led by non-head-on scattering of hard partons [3] and by the thermal vorticity from the statistical model in equilibrium [4]. Although the simulations based on the modified Cooper-Frye formula for spin polarization [5, 6] have shown remarkable agreements with global polarization of Λ [7, 8], more recent observations for local polarization caused the new tension between experiments and theories [9, 10]. Further phenomenological studies have alluded to an essential role for non-equilibrium corrections [11, 12], where the spin polarization is no longer just dictated by thermal vorticity and equilibrium distribution functions for Λ . Also, the feed-down effects are analyzed in refs. [13, 14], which show only minor corrections upon the local polarization. It is hence imperative to understand the dynamics of the spin polarization for not only hadrons but also quarks (and even gluons) in quark gluon plasmas (QGP). In general, the current studies in theory for the dynamical spin polarization in heavy ion physics may be divided into two directions: one is to construct the so-called hydrodynamics of spin as a macroscopic effective theory incorporating spin as a hydrodynamic variable obeying angular momentum conservation [15–17]. Alternatively, non-equilibrium spin transport can be studied through quantum kinetic theory as a microscopic theory having a direct connection to the underlying quantum field theory.

In fact, the construction for quantum kinetic theory of massless fermions, known as chiral kinetic theory (CKT), was initiated by refs. [18, 19] from Berry connection and by refs. [20, 21] from the Wigner-function approach based on quantum field theory with the motivation to explore non-equilibrium transport in chiral matter beyond the renown anomalous phenomena in equilibrium such as the chiral magnetic/vortical effects [22–24]. There have been also plenty of followup studies for extension and applications [25–43]. In particular, the issue for Lorentz covariance associated with the side-jump phenomenon was addressed in refs. [26, 27] and further refined in ref. [30] from the Wigner-function approach with systematic inclusion of background electromagnetic fields and collisions in CKT. The side-jump effect is also shown to contribute to anti-symmetric part of the canonical energy momentum tensor responsible for angular-momentum transfer between spin and orbital angular momentum in chiral fluids [44] (see also ref. [45] for a related study), which further manifests the origin of side jumps in connection to spin-orbit interaction. Overall, the CKT can be regarded as a modified Boltzmann(Vlasov) equation involving quantum corrections such as the chiral anomaly, magnetic-moment coupling, and spin-orbit interaction. Moreover, the CKT has also been recently applied to the study of the Λ polarization in heavy ion collisions [46, 47].

However, in order to consistently investigate the spin transport of Λ or strange quarks as the seed for Λ in QGP before hadronization, it is inevitable to consider the mass corrections. Unlike massless fermions of which the spin orientation is enslaved to the momentum direction, the spin is now a new dynamical degree of freedom coupled with the charge transport for massive fermions. There have been recent studies on the construction of the

“free-streaming” quantum kinetic theory, without the collisional effects, delineating the entangled dynamics between the vector/axial charges and the spin polarization for massive fermions [48–53]. In particular, the axial kinetic theory, derived from the Wigner-function approach up to $\mathcal{O}(\hbar)$ in ref. [51], successfully cover both the massless and massive cases, which is made possible by maintaining the general spin frame vector allowed in the solution of the constraint equations. The set of quantum kinetic theories consists of a scalar kinetic equation (SKE) for the vector-charge and an axial-vector kinetic equation (AKE) for the spin degrees of freedom.

The next key ingredient for describing the spin diffusion/polarization is the collisional effects, which have not been incorporated in the above studies. On the other hand, in ref. [54], the spin-diffusion term at $\mathcal{O}(\hbar^0)$ in collisions was computed in perturbative QCD (see also refs. [55, 56] for other studies of collisions). Nonetheless, the inclusion of $\mathcal{O}(\hbar^1)$ corrections to the collisional effects, which is responsible for generating the spin polarization, has never been achieved. Therefore, we investigate the field-theoretical framework to systematically include the collisional effects to the axial kinetic theory. As will be explained in this paper, one will confront with tough technical problems. The full master equations directly obtained from the Kadanoff-Byam equations are quite involved with the quantum corrections up to $\mathcal{O}(\hbar^1)$. On the other hand, similar to the previous construction for the free-streaming part [51], we also have to ensure the smooth connection between the collisional AKE and the CKT in the massless limit. In spite of the potential importance of finite-mass effects in the collisions terms for the spin rotations and kinematics, the basic framework for the collisional effects has not been touched thus far. We, therefore, intend to establish the collisional SKE and AKE for arbitrary mass within a plausible approximation for physical systems of our interest.

In this paper, we propose a \hbar -counting scheme to circumvent the aforementioned difficulty and construct the quantum kinetic theory with collisions and background electromagnetic fields for spin transport of spin-1/2 fermions with arbitrary mass. Our power counting entails that the axial-charge distribution function f_A is at $\mathcal{O}(\hbar^1)$ as opposed to the vector-charge distribution function f_V at $\mathcal{O}(\hbar^0)$ in the \hbar expansion applied in the Wigner-function formalism. However, this power counting may be applicable in most of physical systems such as HIC. The consequent quantum kinetic theory can be regarded as an “effective” axial kinetic theory with collisions, for which the free-streaming part has been established in ref. [51]. Once deriving the full master equations with the self-energy terms in the Kadanoff-Baym equations, we apply our power-counting scheme and solve the reduced master equations. At the end of the day, we obtain the quantum kinetic equations with collisions for the spin polarization and axial-charge evolution. The collision term therein can be decomposed into the classical part responsible for the spin diffusion and the quantum part contributing to the spin polarization, which importantly stems from the spin-orbit interaction through the entangled dynamics of vector-charge distributions.

Applications of our general framework are not limited to QCD although we emphasize the motivation for the quark transport in QGP. As an example, we further evaluate the classical part of collisions for the spin-diffusion dynamics in the weakly-coupled QGP with specific forms of the self-energies from the hard thermal loop approximation. Computation

of the quantum corrections demand the so-far unknown self-energies with polarized gluons, which needs to be further investigated in the future.

The paper is organized as follows: in section 2, we briefly review the Wigner-function approach and derive the master equations led by Kadanoff-Baym equations with collisions. Then, we introduce our power-counting scheme in \hbar . Based on this scheme, we find the perturbative solutions for Wigner functions. In section 3, we then derive the SKE and AKE in an effective axial kinetic theory with collisions and background electromagnetic fields as a generic formalism for studying quantum transport of fermions. Moreover, we highlight the simplified version suitable for the application to heavy-ion physics and make more detailed discussions upon the collision terms. As a concrete example for applications, in section 4, we further investigate the spin-diffusion term in weakly-coupled QGP by utilizing our formalism. Finally, we make concluding remarks and outlook in section 5. For completeness, we present critical steps in the derivations/computations in appendices.

2 Wigner-function approach with collisions

In this section, we first derive the “master equations” by applying the spinor decomposition and the \hbar expansion to the Kadanoff-Baym equation. Those master equations are subsequently used to derive the quantum kinetic theories. We systematically include the collisional effects that are necessary for describing the relaxation dynamics.

The derivation of quantum kinetic theories in this framework has been investigated for massless fermions [30] and for massive fermions without collisions, that is, the “free-streaming” case [51]. Below, we follow the basic flow of the derivation established in those preceding studies and summarize the crucial intermediate steps below. One will find that the master equations (2.12) with the collisional effects have quite involved structures originating from the spinor structures of fermions. Those involved structures yield versatile transport phenomena that need to be investigated with appropriate strategies for further developments. Therefore, we propose an \hbar counting scheme in section 2.2 that enables one to cure the pathological complication of the collisional effects and to extract the entangled dynamics between the vector charge and the spin polarization.

2.1 Full master equations

We shall start with the Wigner transformation applied to quantum expectation values of correlation functions of fermionic fields,

$$S_{\alpha\beta}^{(>)}(q, X) = \int d^4Y e^{\frac{iq \cdot Y}{\hbar}} \tilde{S}_{\alpha\beta}^{(>)}(x, y), \quad (2.1)$$

where $X = (x + y)/2$ and $Y = x - y$ and we work in the Minkowski spacetime with the mostly negative spacetime metric. Here, $\tilde{S}_{\alpha\beta}^{<}(x, y) = \langle \bar{\psi}_\beta(y) U(y, x) \psi_\alpha(x) \rangle$ and $\tilde{S}_{\alpha\beta}^{>}(x, y) = \langle \psi_\alpha(x) U^\dagger(x, y) \bar{\psi}_\beta(y) \rangle$ are lesser and greater propagators, respectively. To maintain the gauge invariance, we also insert the gauge link, e.g., $U(y, x) = \exp(-i \int_x^y dz^\rho A_\rho(z))$ for QED with A_ρ denoting the U(1) electromagnetic gauge field. Note that q^μ thus represents the kinetic momentum. Hereafter, we focus on $S^{<}(q, X)$ and suppress the indices of spinors.

After the Wigner transformation, the lesser propagator obeys the Kadanoff-Baym equations derived from the Schwinger-Dyson equation,

$$\begin{aligned} (\not{M} - m)S^< + \gamma^\mu i \frac{\hbar}{2} \nabla_\mu S^< &= \frac{i\hbar}{2} \left(\Sigma^< \star S^> - \Sigma^> \star S^< \right), \\ S^< (\not{M} - m) - i \frac{\hbar}{2} \nabla_\mu S^< \gamma^\mu &= -\frac{i\hbar}{2} \left(S^> \star \Sigma^< - S^< \star \Sigma^> \right), \end{aligned} \quad (2.2)$$

where $\Sigma^{<(>)}$ represents the lesser (greater) self-energy. Since we only focus on the scattering process, here we drop the real parts of the retarded and advanced self-energies and of the retarded propagators. See ref. [30] for the same setup to derive the equations for Weyl fermions. The symbol of \star represents the Moyal product incorporating higher-order corrections in \hbar . The star product of two functions $A(q, X)$ and $B(q, X)$ are expanded as

$$A \star B = AB + \frac{i\hbar}{2} \{A, B\}_{\text{P.B.}} + \mathcal{O}(\hbar^2), \quad (2.3)$$

where we define the Poisson bracket as $\{A, B\}_{\text{P.B.}} \equiv (\partial_q^\mu A)(\partial_\mu B) - (\partial_\mu A)(\partial_q^\mu B)$. The sum and difference of eq. (2.2) read

$$\begin{aligned} \{(\not{M} - m), S^<\} + \frac{i\hbar}{2} \left([\gamma^\mu, \nabla_\mu S^<] - [\Sigma^<, S^>]_\star + [\Sigma^>, S^<]_\star \right) &= 0, \\ [(\not{M} - m), S^<] + \frac{i\hbar}{2} \left(\{\gamma^\mu, \nabla_\mu S^<\} - \{\Sigma^<, S^>\}_\star + \{\Sigma^>, S^<\}_\star \right) &= 0. \end{aligned} \quad (2.4)$$

Here, we introduced $\{F, G\} \equiv FG + GF$, $[F, G] \equiv FG - GF$, $\{F, G\}_\star \equiv F \star G + G \star F$ and $[F, G]_\star \equiv F \star G - G \star F$, where F and G are arbitrary matrix-valued functions.

The notations and conventions in the above equations are as follows. First, the derivative operators are given as [57]

$$\nabla_\mu = \partial_\mu + j_0(\square) F_{\nu\mu} \partial_q^\nu, \quad \Pi_\mu = q_\mu + \frac{\hbar}{2} j_1(\square) F_{\nu\mu} \partial_q^\nu, \quad \square = \frac{\hbar}{2} \partial_\rho \partial_q^\rho. \quad (2.5)$$

We will hereafter use $\partial_\mu \equiv \partial/\partial X^\mu$ for convenience. Here $j_0(\square), j_1(\square)$ are spherical Bessel functions and ∂_ρ in \square only act on the field strength $F_{\nu\mu}$ when having spacetime-dependent background fields. Making the \hbar expansion, which corresponds to the gradient expansion for $\partial_\mu \ll q_\mu$, one finds

$$\begin{aligned} \nabla_\mu &= \partial_\mu + F_{\nu\mu} \partial_q^\nu - \frac{\hbar^2}{24} (\partial_\rho \partial_\lambda F_{\nu\mu}) \partial_q^\rho \partial_q^\lambda \partial_q^\nu + \mathcal{O}(\hbar^4), \\ \Pi_\mu &= q_\mu + \frac{\hbar^2}{12} (\partial_\rho F_{\nu\mu}) \partial_q^\rho \partial_q^\nu + \mathcal{O}(\hbar^4). \end{aligned} \quad (2.6)$$

By using the complete basis for the Clifford algebra [57, 58], we may decompose the Wigner functions into

$$\begin{aligned} S^< &= \mathcal{S} + i\mathcal{P}\gamma^5 + \mathcal{V}_\mu \gamma^\mu + \mathcal{A}_\mu \gamma^5 \gamma^\mu + \frac{\mathcal{S}_{\mu\nu}}{2} \sigma^{\mu\nu}, \\ S^> &= \bar{\mathcal{S}} + i\bar{\mathcal{P}}\gamma^5 + \bar{\mathcal{V}}_\mu \gamma^\mu + \bar{\mathcal{A}}_\mu \gamma^5 \gamma^\mu + \frac{\bar{\mathcal{S}}_{\mu\nu}}{2} \sigma^{\mu\nu}, \end{aligned} \quad (2.7)$$

where $\sigma^{\mu\nu} = i[\gamma^\mu, \gamma^\nu]/2$ and $\gamma^5 = i\gamma^0\gamma^1\gamma^2\gamma^3$. We shall then focus on \mathcal{V}^μ and \mathcal{A}^μ , which give rise to the vector-charge and axial-charge currents through $J_V^\mu = 4 \int_q \mathcal{V}^\mu$ and $J_5^\mu = 4 \int_q \mathcal{A}^\mu$, where $\int_q \equiv \int d^4q/(2\pi)^4$. In fact, from field theory, the axial-charge current can be regarded as a spin current for fermions. One may also establish a direct connection between \mathcal{A}^μ and the momentum spectrum of spin polarization from the modified Cooper-Frye formula [5]. See e.g., refs. [6, 44] and appendix B for references. It is worthwhile to note that the axial-charge currents engendered by magnetic fields and vorticity in equilibrium, known as the chiral separation effect and axial vortical effect, could be thus pertinent to the spin polarization particularly in the case with mass corrections [59–61]. See also refs. [62–65] for axial-charge currents triggered by electric fields. Similarly, it is useful to carry out the same spinor-basis decomposition for the self-energies,

$$\begin{aligned}\Sigma^< &= \Sigma_S + i\Sigma_P\gamma^5 + \Sigma_{V\mu}\gamma^\mu + \Sigma_{A\mu}\gamma^5\gamma^\mu + \frac{\Sigma_{T\mu\nu}\sigma^{\mu\nu}}{2}, \\ \Sigma^> &= \bar{\Sigma}_S + i\bar{\Sigma}_P\gamma^5 + \bar{\Sigma}_{V\mu}\gamma^\mu + \bar{\Sigma}_{A\mu}\gamma^5\gamma^\mu + \frac{\bar{\Sigma}_{T\mu\nu}\sigma^{\mu\nu}}{2}.\end{aligned}\quad (2.8)$$

From the Kadanoff-Baym equations and decomposition of the Wigner functions and of the self-energies, one can derive the master equations leading to the derivation of axial kinetic theory.

Now, inserting the spinor decompositions (2.7) and (2.8) into the Kadanoff-Baym equation (2.4), we find ten equations according to the orthogonal property of the spinor basis. This calculation is tedious mostly due to the manipulation of the spinor structures. Nevertheless, it is a straightforward substitution and further decomposition of the products of the gamma matrices. For brevity, we provide the intermediate steps in appendix A.1 and here show the results that serve as a set of the master equations for the derivation of quantum kinetic theories [same as eqs. (A.17) and (A.18)]:

$$m\mathcal{S} = \Pi^\mu \mathcal{V}_\mu - \frac{\hbar^2}{4} \left[\widehat{\Sigma_S \mathcal{S}} - \widehat{\Sigma_P \mathcal{P}} + \widehat{\Sigma_{V\mu} \mathcal{V}^\mu} - \widehat{\Sigma_{A\mu} \mathcal{A}^\mu} + \frac{1}{2} \widehat{\Sigma_{T\mu\nu} \mathcal{S}^{\mu\nu}} \right]_{\text{P.B.}} + \mathcal{O}(\hbar^3), \quad (2.9a)$$

$$m\mathcal{P} = -\frac{\hbar}{2} (\tilde{\mathcal{D}}_\mu \mathcal{A}^\mu - \widehat{\Sigma_{A\mu} \mathcal{V}^\mu}) - \frac{\hbar^2}{4} \left[\widehat{\Sigma_S \mathcal{P}} + \widehat{\Sigma_P \mathcal{S}} + \frac{1}{4} \epsilon_{\mu\nu\alpha\beta} \widehat{\Sigma_T^{\mu\nu} \mathcal{S}^{\alpha\beta}} \right]_{\text{P.B.}} + \mathcal{O}(\hbar^3), \quad (2.9b)$$

$$\begin{aligned}2\Pi_\alpha \mathcal{S} - \hbar \tilde{\mathcal{D}}^\nu \mathcal{S}_{\nu\alpha} - 2m\mathcal{V}_\alpha - \hbar (\widehat{\Sigma_P \mathcal{A}_\alpha} - \widehat{\Sigma_{A\alpha} \mathcal{P}} - \widehat{\Sigma_{T\mu\alpha} \mathcal{V}^\mu}) \\ = \frac{\hbar^2}{2} \left[\widehat{\Sigma_S \mathcal{V}_\alpha} + \widehat{\Sigma_{V\alpha} \mathcal{S}} + \frac{1}{2} \epsilon_{\mu\nu\lambda\alpha} (\widehat{\Sigma_A^\mu \mathcal{S}^{\nu\lambda}} + \widehat{\Sigma_T^{\mu\nu} \mathcal{A}^\lambda}) \right]_{\text{P.B.}} + \mathcal{O}(\hbar^3),\end{aligned}\quad (2.9c)$$

$$\begin{aligned}\hbar \tilde{\mathcal{D}}_\alpha \mathcal{P} - \epsilon_{\alpha\nu\rho\sigma} \Pi^\sigma \mathcal{S}^{\nu\rho} - 2m\mathcal{A}_\alpha - \hbar (\widehat{\Sigma_P \mathcal{V}_\alpha} + \widehat{\Sigma_A^\mu \mathcal{S}_{\mu\alpha}} - \widehat{\Sigma_{T\mu\alpha} \mathcal{A}^\mu}) \\ = \frac{\hbar^2}{2} \left[\widehat{\Sigma_S \mathcal{A}_\alpha} + \widehat{\Sigma_{A\alpha} \mathcal{S}} + \frac{1}{2} \epsilon_{\mu\nu\lambda\alpha} (\widehat{\Sigma_V^\mu \mathcal{S}^{\nu\lambda}} + \widehat{\Sigma_T^{\mu\nu} \mathcal{V}^\lambda}) \right]_{\text{P.B.}} + \mathcal{O}(\hbar^3),\end{aligned}\quad (2.9d)$$

$$\begin{aligned}m\mathcal{S}_{\alpha\beta} + \epsilon_{\alpha\beta\rho\sigma} \Pi^\rho \mathcal{A}^\sigma - \frac{\hbar}{2} (\tilde{\mathcal{D}}_{[\alpha} \mathcal{V}_{\beta]} - \widehat{\Sigma_{A[\alpha} \mathcal{A}_{\beta]}} + \widehat{\Sigma_{T\mu[\alpha} \mathcal{S}^\mu_{\beta]}}) \\ = -\frac{\hbar^2}{4} \left[\widehat{\Sigma_S \mathcal{S}_{\alpha\beta}} + \widehat{\Sigma_{T\alpha\beta} \mathcal{S}} - \epsilon_{\mu\nu\alpha\beta} \widehat{\Sigma_V^\mu \mathcal{A}^\nu} + \epsilon_{\mu\nu\alpha\beta} \widehat{\Sigma_A^\mu \mathcal{V}^\nu} - \frac{1}{2} \epsilon_{\mu\nu\alpha\beta} (\widehat{\Sigma_P \mathcal{S}^{\mu\nu}} + \widehat{\Sigma_T^{\mu\nu} \mathcal{P}}) \right]_{\text{P.B.}} + \mathcal{O}(\hbar^3),\end{aligned}\quad (2.9e)$$

and

$$\tilde{\mathcal{D}}^\mu \mathcal{V}_\mu = -\widehat{\Sigma_S \mathcal{S}} + \widehat{\Sigma_P \mathcal{P}} + \widehat{\Sigma_{A\mu} \mathcal{A}^\mu} - \frac{1}{2} \widehat{\Sigma_{T\mu\nu} \mathcal{S}^{\mu\nu}} + \mathcal{O}(\hbar^3), \quad (2.10a)$$

$$2\Pi^\nu \mathcal{A}_\nu = -\hbar \left(\widehat{\Sigma_S \mathcal{P}} + \widehat{\Sigma_P \mathcal{S}} + \frac{1}{4} \epsilon_{\mu\nu\alpha\beta} \widehat{\Sigma_T^{\mu\nu} \mathcal{S}^{\alpha\beta}} \right) - \frac{\hbar^2}{2} [-\widehat{\Sigma_{V\alpha} \mathcal{A}^\alpha} + \widehat{\Sigma_{A\alpha} \mathcal{V}^\alpha}]_{\text{P.B.}} + \mathcal{O}(\hbar^3), \quad (2.10b)$$

$$\begin{aligned} 2\Pi^\nu \mathcal{S}_{\nu\alpha} + \hbar \tilde{\mathcal{D}}_\alpha \mathcal{S} + \hbar \left(\widehat{\Sigma_S \mathcal{V}_\alpha} + \frac{1}{2} \epsilon_{\mu\nu\lambda\alpha} (\widehat{\Sigma_A^\mu \mathcal{S}^{\nu\lambda}} + \widehat{\Sigma_T^{\mu\nu} \mathcal{A}^\lambda}) \right) \\ = \frac{\hbar^2}{2} [\widehat{\Sigma_P \mathcal{A}_\alpha} + \widehat{\Sigma_V^\mu \mathcal{S}_{\mu\alpha}} - \widehat{\Sigma_{A\alpha} \mathcal{P}} + \widehat{\Sigma_{T\alpha\beta} \mathcal{V}^\beta}]_{\text{P.B.}} + \mathcal{O}(\hbar^3), \end{aligned} \quad (2.10c)$$

$$\begin{aligned} 2\Pi_\alpha \mathcal{P} + \frac{\hbar}{2} \epsilon_{\alpha\nu\rho\sigma} (\tilde{\mathcal{D}}^\nu \mathcal{S}^{\rho\sigma} + \widehat{\Sigma_T^{\nu\rho} \mathcal{V}^\sigma}) - \hbar (\widehat{\Sigma_S \mathcal{A}_\alpha} + \widehat{\Sigma_{A\alpha} \mathcal{S}}) \\ = 1 - \frac{\hbar^2}{2} [\widehat{\Sigma_P \mathcal{V}_\alpha} - \widehat{\Sigma_{V\alpha} \mathcal{P}} + \widehat{\Sigma_A^\mu \mathcal{S}_{\mu\alpha}} + \widehat{\Sigma_{T\alpha\beta} \mathcal{A}^\beta}]_{\text{P.B.}} + \mathcal{O}(\hbar^3), \end{aligned} \quad (2.10d)$$

$$\begin{aligned} \Pi_{[\alpha} \mathcal{V}_{\beta]} + \frac{\hbar}{2} \epsilon_{\alpha\beta\mu\nu} (\tilde{\mathcal{D}}^\mu \mathcal{A}^\nu - \widehat{\Sigma_A^\mu \mathcal{V}^\nu}) - \frac{\hbar}{2} (\widehat{\Sigma_S \mathcal{S}_{\alpha\beta}} + \widehat{\Sigma_{T\alpha\beta} \mathcal{S}}) + \frac{\hbar}{4} \epsilon_{\mu\nu\alpha\beta} (\widehat{\Sigma_P \mathcal{S}^{\mu\nu}} + \widehat{\Sigma_T^{\mu\nu} \mathcal{P}}) \\ = \frac{\hbar^2}{4} [\widehat{\Sigma_{V[\alpha} \mathcal{V}_{\beta]}} - \widehat{\Sigma_{A[\alpha} \mathcal{A}_{\beta]}} + \widehat{\Sigma_{T\mu[\alpha} \mathcal{S}^{\mu}_{\beta]}}]_{\text{P.B.}} + \mathcal{O}(\hbar^3), \end{aligned} \quad (2.10e)$$

where $\epsilon_{\mu\nu\rho\sigma}$ is the totally antisymmetric tensor with $\epsilon_{0123} = -1$. In the above, we introduced a few shorthand notations. $[AB]_{\text{P.B.}} = \{A(q, X), B(q, X)\}_{\text{P.B.}}$ is a shorthand notation for the Poisson bracket. We use $\widehat{XY} = \bar{X}Y - X\bar{Y}$, where X and Y are the coefficients of the Clifford decomposition of the propagators and self-energies. We also use $\tilde{\mathcal{D}}_\mu \mathcal{M} = \nabla_\mu \mathcal{M} + \widehat{\Sigma_{V\mu} \mathcal{M}} + \mathcal{O}(\hbar^2)$.¹

As has been known in light of the derivation of the free-streaming case without the collision term [51], one can first reduce the number of variables to solve the above master equations. Namely, one can replace \mathcal{S} , \mathcal{P} and $\mathcal{S}_{\mu\nu}$ by \mathcal{V}^μ and \mathcal{A}^μ by the use of eqs. (2.9a), (2.9b) and (2.9e). This procedure is explained in appendix A.2. Since $\mathcal{S}^{\mu\nu}$ is contracted with $\Sigma_{T\mu\nu}$ and $\bar{\Sigma}_{T\mu\nu}$ on the right-hand side, one may not express $\mathcal{S}_{\mu\nu}$ as an explicit function of \mathcal{V}_μ and \mathcal{A}_μ only. Nevertheless, assuming that the interaction is sufficiently weak, we may drop the nonlinear terms in the self-energy. Maintaining the linear terms in the self-energies and the explicit \hbar dependence up to $\mathcal{O}(\hbar^1)$, \mathcal{S} , \mathcal{P} , and $\mathcal{S}_{\mu\nu}$ are expressed as [cf. eq. (A.19)]

$$\mathcal{S}_{\alpha\beta} = -\frac{1}{m} \epsilon_{\alpha\beta\rho\sigma} q^\rho \mathcal{A}^\sigma + \frac{\hbar}{2m} \left\{ \mathcal{D}_{[\alpha} \mathcal{V}_{\beta]} - \widehat{\Sigma_{A[\alpha} \mathcal{A}_{\beta]}} + \frac{q_\mu}{m} \epsilon^{\mu\rho\sigma} \widehat{\Sigma_{T[\beta] \rho} \mathcal{A}_\sigma} \right\}, \quad (2.11a)$$

$$\mathcal{S} = \frac{q^\mu}{m} \mathcal{V}_\mu, \quad (2.11b)$$

$$\mathcal{P} = -\frac{\hbar}{2m} (\mathcal{D}_\mu \mathcal{A}^\mu - \widehat{\Sigma_{A\mu} \mathcal{V}^\mu}), \quad (2.11c)$$

where we define $\mathcal{D}_\mu \mathcal{M} = \nabla_\mu \mathcal{M} + \widehat{\Sigma_{V\mu} \mathcal{M}}$ and the antisymmetrization $T_{[\mu\nu]} = T_{\mu\nu} - T_{\nu\mu}$. After elimination of those variables, the other master equations up to the linear order in

¹The original definition before the \hbar bar expansion is given below eq. (A.16).

the self-energies and $\mathcal{O}(\hbar)$ read [same as eqs. (A.20) and (A.21)]

$$q^\mu(q \cdot \mathcal{V}) - m^2 \mathcal{V}^\mu = \frac{\hbar}{2} \left[m(\widehat{\Sigma_P \mathcal{A}^\mu} + \widehat{\Sigma_T^{\mu\alpha} \mathcal{V}_\alpha}) - 2 \left(\tilde{F}^{\mu\beta} \mathcal{A}_\beta + \frac{1}{2} \epsilon^{\mu\alpha\beta\gamma} q_\alpha \Delta_\beta \mathcal{A}_\gamma \right) - \epsilon^{\mu\alpha\beta\gamma} q_\alpha \widehat{\Sigma_{V\beta} \mathcal{A}_\gamma} \right], \quad (2.12a)$$

$$q^2 \mathcal{A}^\mu - q^\mu q \cdot \mathcal{A} - m^2 \mathcal{A}^\mu = \frac{1}{2} \hbar \left[m(\widehat{\Sigma_P \mathcal{V}^\mu} + \widehat{\Sigma_T^{\mu\alpha} \mathcal{A}_\alpha}) + \epsilon^{\mu\alpha\beta\gamma} q_\alpha \mathcal{D}_\beta \mathcal{V}_\gamma - 2\epsilon^{\mu\alpha\beta\gamma} q_\alpha \widehat{\Sigma_{A\beta} \mathcal{A}_\gamma} - \frac{1}{m} \epsilon^\mu_{\alpha\beta\gamma} \epsilon^{\beta\lambda\rho\sigma} q^\alpha q_\lambda \widehat{\Sigma_{T\rho}^\gamma \mathcal{A}_\sigma} \right], \quad (2.12b)$$

$$\mathcal{D}^\mu \mathcal{V}_\mu = \widehat{\Sigma_{A\mu} \mathcal{A}^\mu} - \frac{1}{2m} \left[2q_\mu \widehat{\Sigma_S \mathcal{V}^\mu} - \epsilon_{\mu\nu\alpha\beta} q^\alpha \widehat{\Sigma_T^{\mu\nu} \mathcal{A}^\beta} + \hbar \widehat{\Sigma_P (\nabla \cdot \mathcal{A})} + \frac{\hbar}{2} \widehat{\Sigma_T^{\mu\nu} \nabla_{[\mu} \mathcal{V}_{\nu]}} \right], \quad (2.12c)$$

$$q \cdot \mathcal{A} = -\frac{\hbar}{2m} q_\mu (\widehat{\Sigma_P \mathcal{V}^\mu} + \widehat{\Sigma_T^{\mu\nu} \mathcal{A}_\nu}), \quad (2.12d)$$

$$\begin{aligned} \mathcal{D}_\mu(q \cdot \mathcal{V}) - q^\nu \mathcal{D}_{[\mu} \mathcal{V}_{\nu]} &= -m \widehat{\Sigma_S \mathcal{V}_\mu} + q_{[\mu} \widehat{\Sigma_{A\alpha]} \mathcal{A}^\alpha + \frac{1}{m} q^\nu q_\alpha \epsilon^{\alpha\beta\gamma} \widehat{\Sigma_{T\nu] \beta} \mathcal{A}_\gamma} \\ &\quad + \frac{1}{2} \epsilon_{\mu\nu\rho\sigma} \left\{ m \widehat{\Sigma_T^{\nu\rho} \mathcal{A}^\sigma} - \hbar (\partial^\nu F^{\rho\beta}) \partial_{q\beta} \mathcal{A}^\sigma + \hbar \widehat{\Sigma_A^\nu (\nabla^\rho \mathcal{V}^\sigma)} \right\} \\ &\quad - \frac{\hbar q^\nu}{2m} (\widehat{\Sigma_{T\gamma} [\nu \Delta^\gamma \mathcal{V}_\mu]} - \widehat{\Sigma_{T\gamma} [\nu \Delta_\mu] \mathcal{V}^\gamma}) \\ &\quad + \frac{\hbar q^\nu}{2} \left[\frac{1}{m} \widehat{\Sigma_{T\nu\mu} (q \cdot \mathcal{V})} - \epsilon_{\nu\mu\rho\sigma} \widehat{\Sigma_V^\rho \mathcal{A}^\sigma} + \epsilon_{\nu\mu\rho\sigma} \widehat{\Sigma_A^\rho \mathcal{V}^\sigma} - \frac{1}{m} \widehat{\Sigma_P (q_{[\nu} \mathcal{A}_{\mu]})} \right]_{\text{P.B.}} \\ &\quad + \frac{\hbar m}{2} \left[\widehat{\Sigma_P \mathcal{A}_\mu} - \frac{\epsilon_{\nu\mu\rho\sigma}}{m} \widehat{\Sigma_V^\nu (q^\rho \mathcal{A}^\sigma)} + \widehat{\Sigma_{T\mu\beta} \mathcal{V}^\beta} \right]_{\text{P.B.}}, \end{aligned} \quad (2.12e)$$

$$\begin{aligned} F_{\mu\nu} \mathcal{A}^\nu - q \cdot \mathcal{D} \mathcal{A}_\mu + \frac{\hbar}{4} \epsilon_{\mu\nu\rho\sigma} [\mathcal{D}^\nu, \mathcal{D}^\rho] \mathcal{V}^\sigma - \frac{\hbar}{2m} q_\mu \left[\widehat{\Sigma_P (q \cdot \mathcal{V})} + \frac{1}{2} \widehat{\Sigma_T^{\rho\sigma} (q_{[\rho} \mathcal{A}_{\sigma]})} \right]_{\text{P.B.}} \\ = m \left[\widehat{\Sigma_S \mathcal{A}_\mu} - \frac{1}{2} \epsilon_{\mu\nu\rho\sigma} \widehat{\Sigma_T^{\nu\rho} \mathcal{V}^\sigma} \right] + q_\alpha \widehat{\Sigma_{A\mu} \mathcal{V}^\alpha} - q_\mu \widehat{\Sigma_{A\alpha} \mathcal{V}^\alpha} + \frac{\hbar}{2} \epsilon_{\mu\nu\rho\sigma} \Delta^\nu \left[\widehat{\Sigma_A^\rho \mathcal{A}^\sigma} - \frac{1}{m} q^\alpha \epsilon_{\alpha\beta\gamma} \widehat{\Sigma_T^{\sigma\beta} \mathcal{A}_\gamma} \right] \\ - \frac{\hbar m}{2} \left[\widehat{\Sigma_P \mathcal{V}_\mu} + \frac{1}{m} \epsilon_{\mu\nu\rho\sigma} \widehat{\Sigma_A^\nu (q^\rho \mathcal{A}^\sigma)} + \widehat{\Sigma_{T\mu\nu} \mathcal{A}^\nu} \right]_{\text{P.B.}}, \end{aligned} \quad (2.12f)$$

$$q^{[\mu} \mathcal{V}^{\nu]} + \frac{\hbar}{2} \epsilon^{\mu\nu\alpha\beta} (\mathcal{D}_\alpha \mathcal{A}_\beta - \widehat{\Sigma_{A\alpha} \mathcal{V}_\beta}) = -\frac{\hbar}{2m} \left[\epsilon^{\mu\nu\alpha\beta} q_\alpha \widehat{\Sigma_S \mathcal{A}_\beta} - q_\alpha \widehat{\Sigma_T^{\mu\nu} \mathcal{V}^\alpha} + q^{[\mu} \widehat{\Sigma_P \mathcal{A}^{\nu]} \right]. \quad (2.12g)$$

As opposed to the free-streaming case, the inclusion of the collision terms cause the drastic complication in the master equations as seen in the above expressions. This hinders finding the perturbative solution for the Wigner functions and deriving the consequent quantum kinetic equations. In the subsequent sections, we tackle this problem that needs to be addressed at both formal and practical levels before describing any kinds of the relaxation dynamics. Our main proposal is an appropriate \hbar counting scheme that greatly simplifies the collision terms and may work in some applications such as the spin transport in the quark-gluon plasma.

2.2 Setting the power-counting scheme

Thus far, we have not specified the orders of \mathcal{V}^μ and \mathcal{A}^μ in the \hbar expansion. In general, they could be both comparable and different in magnitudes. In most of practical situations such as in heavy ion collisions, the axial-charge current is usually smaller than the vector-charge current since the spin polarization is basically generated by quantum effects. This observation motivates us to introduce the power counting:

$$\mathcal{V}^\mu \sim \mathcal{O}(\hbar^0) \quad \text{and} \quad \mathcal{A}^\mu \sim \mathcal{O}(\hbar). \quad (2.13)$$

In light of eq. (2.11b) and eq. (2.11a), these assignments lead to the order counting:

$$\mathcal{S} \sim \mathcal{O}(\hbar^0) \quad \text{and} \quad \mathcal{S}^{\mu\nu} \sim \mathcal{O}(\hbar). \quad (2.14)$$

Also, in the free case, from eq. (2.11c), we find

$$\mathcal{P} \sim \mathcal{O}(\hbar^2). \quad (2.15)$$

Consequently, similar power counting will be applied to $\Sigma^{<(>)}$. From eq. (2.12c), it is clear that the ordinary Boltzmann equation at $\mathcal{O}(\hbar^0)$ comprises only Σ_S and $\Sigma_{V\mu}$ in collisions when $\mathcal{V}^\mu \sim \mathcal{O}(\hbar^0)$ and $\mathcal{A}^\mu \sim \mathcal{O}(\hbar)$. This implies

$$\Sigma_S \sim \mathcal{O}(\hbar^0) \quad \text{and} \quad \Sigma_V^\mu \sim \mathcal{O}(\hbar^0), \quad (2.16)$$

and the same counting applies to $\bar{\Sigma}_S$ and $\bar{\Sigma}_V^\mu$. Those self-energies are responsible for the “classical” collision term. On the contrary, other components in $\Sigma^{<(>)}$ come from quantum corrections. In light of eq. (2.12f), one finds that at most

$$\Sigma_A^\mu \sim \mathcal{O}(\hbar^1) \quad \text{and} \quad \Sigma_T^{\mu\nu} \sim \mathcal{O}(\hbar^1). \quad (2.17)$$

This is required to balance the orders of free-streaming and collision parts, and is also consistent with eq. (2.11c).

Physically, the “classical” Boltzmann equation only incorporates the vector-charge conservation. In order to have nonzero Σ_A^μ or $\Sigma_T^{\mu\nu}$, either the scattered fermion or gluon should carry nonzero chirality imbalance or spin (angular-momentum), which has to come from quantum corrections at least at $\mathcal{O}(\hbar)$ since our power counting has implied the suppression of spin currents compared to the vector-charge currents. Albeit there exists no explicit restriction for Σ_P from generic master equations, it is expected that the presence of Σ_P has to be induced by nonzero pseudo-scalar condensate, which should be at $\mathcal{O}(\hbar^2)$ from the consistency with the anomaly equation (mass correction upon the chiral anomaly). Nonetheless, due to the lack of a rigorous proof for the order of Σ_P from Kadanoff-Baym equations, we will naively take $\Sigma_P \sim \mathcal{O}(\hbar^0)$ for completeness. We hence establish our power counting in that we have

$$\Sigma_S \sim \mathcal{O}(\hbar^0) \quad \text{and} \quad \Sigma_V^\mu \sim \mathcal{O}(\hbar^0), \quad (2.18)$$

while

$$\Sigma_A^\mu \sim \mathcal{O}(\hbar^1), \quad \Sigma_T^{\mu\nu} \sim \mathcal{O}(\hbar^1) \quad \text{and} \quad \Sigma_P \sim \mathcal{O}(\hbar^0). \quad (2.19)$$

All components certainly can also have higher-order corrections in \hbar depending on the details of collisions for different systems.

2.3 Effective master equations and constrains

By implementing the power counting and the results shown in eq. (2.12), the master equations obtained from the Kadanoff-Baym equations at “the leading order $\mathcal{O}(\hbar^0)$ ” now read

$$\mathcal{D}^\mu \mathcal{V}_\mu = -\frac{1}{m} q_\mu \widehat{\Sigma_S \mathcal{V}^\mu}, \quad (2.20a)$$

$$q^{[\mu} \mathcal{V}^{\nu]} = 0, \quad (2.20b)$$

$$(q^2 - m^2) \mathcal{V}_\mu = 0, \quad (2.20c)$$

$$q \cdot \mathcal{A} = -\frac{\hbar}{2m} q_\mu \widehat{\Sigma_P \mathcal{V}^\mu}, \quad (2.20d)$$

$$(q^2 - m^2) \mathcal{A}^\mu = \frac{\hbar}{2} \epsilon^{\mu\alpha\beta\gamma} q_\alpha D_\beta \mathcal{V}_\gamma, \quad (2.20e)$$

$$\begin{aligned} q \cdot \mathcal{D} \mathcal{A}_\mu - F_{\mu\nu} \mathcal{A}^\nu + \frac{\hbar}{2m} q_\mu [\widehat{\Sigma_P (q \cdot \mathcal{V})}]_{\text{P.B.}} - \frac{\hbar m}{2} [\widehat{\Sigma_P \mathcal{V}_\mu}]_{\text{P.B.}} \\ = \frac{\hbar}{4} \epsilon_{\mu\nu\rho\sigma} [\mathcal{D}^\nu, \mathcal{D}^\rho] \mathcal{V}^\sigma - m \left(\widehat{\Sigma_S \mathcal{A}_\mu} - \frac{1}{2} \epsilon_{\mu\nu\rho\sigma} \widehat{\Sigma_T^{\nu\rho} \mathcal{V}^\sigma} \right) - q_\alpha \widehat{\Sigma_{A\mu} \mathcal{V}^\alpha} + q_\mu \widehat{\Sigma_{A\alpha} \mathcal{V}^\alpha}, \end{aligned} \quad (2.20f)$$

$$q \cdot \mathcal{D} \mathcal{V}_\mu - F_{\mu\nu} \mathcal{V}^\nu = -m \widehat{\Sigma_S \mathcal{V}_\mu}. \quad (2.20g)$$

Note that we have to keep the $\mathcal{O}(\hbar)$ terms linear to \mathcal{V}^μ in eqs. (2.20e) and (2.20f) since $\mathcal{A}^\mu \sim \mathcal{O}(\hbar)$. In addition, we consider weakly coupled systems and thus drop the $\mathcal{O}(\Sigma^2)$ terms. One can further implement

$$\frac{\hbar}{2m} q_\mu [\widehat{\Sigma_P (q \cdot \mathcal{V})}]_{\text{P.B.}} - \frac{\hbar m}{2} [\widehat{\Sigma_P \mathcal{V}_\mu}]_{\text{P.B.}} = \frac{\hbar}{2m} (\partial_\mu \widehat{\Sigma_P}) q \cdot \mathcal{V} \quad (2.21)$$

to simplify eq. (2.20f).

From eqs. (2.20b) and (2.20c), one immediately obtains the leading-order solution for the vector part,

$$\mathcal{V}^\mu = 2\pi\delta(q^2 - m^2) q^\mu f_V, \quad (2.22)$$

where $f_V(q, X)$ denotes the vector-charge distribution function. For completeness, we should, in principle, multiply \mathcal{V}^μ by the sign function for energy to include antiparticles. For notational simplicity, we will mostly focus on just the positive-energy solution throughout the paper. For \bar{V}^μ , we simply replace f_V by $\bar{f}_V = 1 - f_V$ as the vector-charge distribution function for an outgoing fermion. We may also easily show that eq. (2.20g) is a redundant equation, which can be derived from eqs. (2.20a)–(2.20c). Plugging the leading-order solution for \mathcal{V}^μ into eq. (2.20a), one acquires the SKE as the usual Boltzmann equation

$$\delta(q^2 - m^2) \left(q \cdot \Delta f_V + q_\mu \widehat{\Sigma_V^\mu} f_V + m \widehat{\Sigma_S} f_V \right) + \mathcal{O}(\hbar) = 0. \quad (2.23)$$

For the axial part \mathcal{A}^μ , we have to solve eqs. (2.20d) and (2.20e) up to $\mathcal{O}(\hbar)$. With the leading-order solution (2.22) constrained above, one however finds that the collisional term vanishes on the left-hand side of eq. (2.20e). This means that the collisional effects do not modify the dispersion relation for \mathcal{A}^μ at the leading order. Nevertheless, the same

as the collisionless case [51], we are not able to uniquely determine the magnetization-current term at the next-to-leading order from the master equations. In the absence of a background field, one can determine the corresponding term by explicitly solving the free Dirac equation [51]. In addition, we refer to the correspondence between the cases with and without a background field in the massless limit, where the latter is given by the former with the simple replacement of the partial derivative by the derivative operator with a background field. Based on those observations, we could generalize the free-theory form to the case with background fields and collisions.

Namely, we “postulate”²

$$\mathcal{A}^\mu = 2\pi \left[\delta(q^2 - m^2) \left(a^\mu f_A + \hbar S_{m(n)}^{\mu\nu} \mathcal{D}_\nu f_V + \frac{\hbar}{2m} q^\mu C_P[f_V] \right) + \hbar \tilde{F}^{\mu\nu} q_\nu \delta'(q^2 - m^2) f_V \right], \quad (2.24)$$

where $C_P[f_V] = -\widehat{\Sigma_P f_V}$ and

$$S_{m(n)}^{\mu\nu} = \frac{\epsilon^{\mu\nu\alpha\beta} q_\alpha n_\beta}{2a \cdot n} = \frac{\epsilon^{\mu\nu\alpha\beta} q_\alpha n_\beta}{2(q \cdot n + m)}, \quad (2.25)$$

which is generalization of the \mathcal{A}^μ found in ref. [51] with the replacement of $\Delta_\nu f_V$ by $\mathcal{D}_\nu f_V$. Here, a^μ represents a spin four-vector satisfying $q \cdot a = q^2 - m^2$ and f_A denotes the axial-charge distribution function.

At $\mathcal{O}(\hbar)$, the dispersion relation is modified by, e.g., the magnetic-moment coupling from the last term in eq. (2.24) with $\delta'(q^2 - m^2) = d\delta(q^2 - m^2)/dq^2$. On the other hand, the $S_{m(n)}^{\mu\nu}$ term corresponds to the so-called magnetization-current term led by spin-orbit interaction, which depends on a frame vector n^μ specifying the spin basis. The presence of such a term implies the frame dependence of f_V because of the frame invariance of full \mathcal{A}^μ . In the massless limit, $a^\mu = q^\mu$ according to the spin enslavement specified by the helicity. The expression in eq. (2.24) then agrees with the solution directly solved from Kadanoff-Baym equations of Weyl fermions [30]. Furthermore, for the solution of $\tilde{\mathcal{A}}^\mu$, we simply replace f_A and f_V in eq. (2.24) by \tilde{f}_A and \tilde{f}_V , respectively. Nonetheless, unlike $\tilde{f}_V = 1 - f_V$, we have $\tilde{f}_A = -f_A$ due to its origin from the expectation value of the fermionic density operator in spinor space (see ref. [51] for a detailed definition of f_A in field theory). As a consequence of our power counting, eq. (2.24) corresponds to the leading-order solution for \mathcal{A}^μ starting at $\mathcal{O}(\hbar)$. Albeit the power counting we apply, we will still dub the terms with \hbar prefactors as the “quantum corrections” at $\mathcal{O}(\hbar)$ throughout this paper for convenience. That is, in the rest part of the paper, \hbar is simply a parameter indicating the quantum origin of certain terms, while these terms are in the same order of magnitudes as the classical terms without \hbar prefactors. More precisely, one should distinguish the difference between \hbar terms and $\mathcal{O}(\hbar)$ terms.

²Thanks to the power-counting scheme, the generalization of the magnetization-current term with collisions here is rather natural since Σ_V^μ and $\tilde{\Sigma}_V^\mu$ are only vectors at $\mathcal{O}(\hbar^0)$ in self-energies, which can then be coupled to $S_{m(n)}^{\mu\nu}$.

3 Effective axial kinetic theory with collisions

In this section, we elaborate the collision terms for the effective axial kinetic theory arising from our \hbar counting scheme introduced in section 2.2. Assembling the ingredients obtained in section 2.3, the general expression will be given in eq. (3.2) with the collision terms $\hat{\mathcal{C}}_1^{(n)\mu}$ and $\hbar\hat{\mathcal{C}}_2^{(n)\mu}$. Here, we maintain the general frame vector n^μ .

We provide a simpler form in eq. (3.9) by choosing a specific frame vector $n^\mu = (1, 0)$ and dropping the background electromagnetic field, which will be useful for applications. A further alternative expression (3.12) enables us to take the massless limit and to confirm the agreement to the (massless) chiral kinetic theory obtained in ref. [30]. In section 3.2, we take another frame vector $n^\mu = n_r^\mu(q) = q^\mu/m$ specifying the rest frame of a massive fermion. $\hat{\mathcal{C}}_{1,2}^{(n_r)\mu}$ in eq. (3.14) adds collisional effects to the free-streaming part investigated in refs. [49, 50], and leads to a useful expression of the effective axial kinetic theory (3.17).

Besides the explicit \hbar dependences originating from those in the Kadanoff-Baym equation (2.2), there are implicit \hbar dependences contained in the self-energies Σ . In section 3.3, we therefore sort the \hbar dependences in the actual order of our counting scheme, which results in the form of $\hat{\mathcal{C}}_{\text{cl}}^\mu$ and $\hbar\hat{\mathcal{C}}_{\text{Q}}^{(n)\mu}$ in eq. (3.18). Finally, in section 3.4, we discuss the physical meaning of the quantum corrections to the collision terms and the conditions to make efficient applications of our effective axial kinetic theories. As noted in the end of previous section, one should bear in mind that $\hat{\mathcal{C}}_{\text{cl}}^\mu$ and $\hbar\hat{\mathcal{C}}_{\text{Q}}^{(n)\mu}$ have the same order of magnitude and we hereafter use \hbar as just a parameter indicating the quantum origin of attached terms unless specified.

3.1 Axial kinetic equation with general frame vector and its massless limit

We now utilize eqs. (2.20f) and (2.24) to derive the effective AKE. Given that the collisionless (free-streaming) part up to $\mathcal{O}(\hbar)$ has been obtained in ref. [51], we only need to further work out the collisional part. This can be carried out by inserting the solution (2.24) of the constraint equations into eq. (2.20f). By the use of an identity

$$\epsilon_{\mu\nu\rho\sigma}[\mathcal{D}^\nu, \mathcal{D}^\rho]\mathcal{V}^\sigma = \epsilon_{\mu\nu\rho\sigma} \left([\Delta^\nu, \Delta^\rho]\mathcal{V}^\sigma + 2(\Delta^\nu \widehat{\Sigma_V^\rho})\mathcal{V}^\sigma \right), \quad (3.1)$$

the AKE with the collisional effects and the general frame vector $n^\mu = n^\mu(X)$ takes the form

$$\square^{(n)}\mathcal{A}^\mu = \hat{\mathcal{C}}_1^{(n)\mu} + \hbar\hat{\mathcal{C}}_2^{(n)\mu}. \quad (3.2)$$

Here the free-streaming part is given by [51]

$$\begin{aligned} \square^{(n)}\mathcal{A}^\mu &= \delta(q^2 - m^2) \left(q \cdot \Delta (a^\mu f_A) + F^{\nu\mu} a_\nu f_A \right) \\ &+ \hbar q^\mu \left\{ \delta(q^2 - m^2) \left[(\partial_\alpha S_{m(n)}^{\alpha\nu}) \Delta_\nu + \frac{S_{m(n)}^{\alpha\nu} F_{\alpha\beta} n^\beta \Delta_\nu}{q \cdot n + m} \right. \right. \\ &+ \left. \left. S_{m(n)}^{\rho\nu} (\partial_\rho F_{\beta\nu}) \partial_q^\beta \right] - \delta'(q^2 - m^2) \frac{q^\alpha \tilde{F}_{\alpha\beta} n^\beta}{q \cdot n + m} q \cdot \Delta \right\} f_V \\ &+ \hbar m \left\{ \frac{\delta(q^2 - m^2) \epsilon^{\mu\nu\alpha\beta}}{2(q \cdot n + m)} \left[m(\partial_\alpha n_\beta) \Delta_\nu + (mn_\beta + q_\beta) \left(\frac{(F_{\alpha\rho} n^\rho - \partial_\alpha(q \cdot n))}{q \cdot n + m} \Delta_\nu \right. \right. \right. \\ &\left. \left. \left. - (\partial_\nu F_{\rho\alpha}) \partial_q^\rho \right) \right] + \delta'(q^2 - m^2) \frac{(mn_\beta + q_\beta) \tilde{F}^{\mu\beta}}{q \cdot n + m} q \cdot \Delta \right\} f_V. \end{aligned} \quad (3.3)$$

Taking $\hbar = 0$, the equation above reproduces the so-called Bargmann-Michel-Telegdi (BMT) equation as a classical kinetic equation for spin transport [66]. On the other hand, the collision terms are

$$\hat{C}_1^{(n)\mu} = \delta(q^2 - m^2) \left[-a^\mu q_\nu \widehat{\Sigma_V^\nu f_A} - m^2 \widehat{\Sigma_A^\mu f_V} + q^\mu q_\nu \widehat{\Sigma_A^\nu f_V} - m \left(a^\mu \widehat{\Sigma_S f_A} - \frac{1}{2} \epsilon^{\mu\nu\rho\sigma} q_\nu \widehat{\Sigma_{T\rho\sigma} f_V} \right) \right], \quad (3.4)$$

and

$$\begin{aligned} \hat{C}_2^{(n)\mu} = & \frac{\delta(q^2 - m^2)}{2} \left[\epsilon^{\mu\nu\rho\sigma} q_\nu (\Delta_\rho \widehat{\Sigma_{V\sigma} f_V}) - m (\partial^\mu \widehat{\Sigma_P}) f_V - 2 \widehat{\Sigma_{V\nu} f_V} (q \cdot \Delta S_{m(n)}^{\mu\nu} - F_\lambda^\mu S_{m(n)}^{\lambda\nu}) \right. \\ & \left. - 2 S_{m(n)}^{\mu\nu} \left((q \cdot \Sigma_V \widehat{\Delta_\nu f_V}) + m \Sigma_S \widehat{\Delta_\nu f_V} + (q \cdot \Delta \widehat{\Sigma_{V\nu} f_V}) + \frac{1}{m} q^\mu (q \cdot \Delta \widehat{\Sigma_P} f_V) \right) \right] \\ & - S_{m(n)}^{\mu\nu} \Delta_\nu (\delta(q^2 - m^2) q \cdot \Delta f_V) - \delta(q^2 - m^2) (\Delta_\nu S_{m(n)}^{\mu\nu}) q \cdot \Delta f_V \\ & - \tilde{F}^{\mu\nu} q_\nu \delta'(q^2 - m^2) (q \cdot \widehat{\Sigma_V f_V} + m \widehat{\Sigma_S f_V}). \end{aligned} \quad (3.5)$$

Note that $\hat{C}_1^{(n)\mu}$ implicitly contains the \hbar terms from, e.g., Σ_A^μ and $\Sigma_{T\rho\sigma}$, which hence implicitly depends on the frame choice.

Implementing the leading-order SKE in eq. (2.23), the following term in $\hat{C}_2^{(n)\mu}$ can be also written as

$$\begin{aligned} & - S_{m(n)}^{\mu\nu} \Delta_\nu (\delta(q^2 - m^2) q \cdot \Delta f_V) \\ & = - S_{m(n)}^{\mu\nu} \left[\delta(q^2 - m^2) \Delta_\nu + 2q^\lambda F_{\lambda\nu} \delta'(q^2 - m^2) \right] (C_V[f_V] + m C_S[f_V]), \end{aligned} \quad (3.6)$$

where $C_V[f_V] = q \cdot \Sigma_V \bar{f}_V - q \cdot \bar{\Sigma}_V f_V$ and $C_S[f_V] = \Sigma_S \bar{f}_V - \bar{\Sigma}_S f_V$. Moreover, one may rewrite

$$\begin{aligned} & - (q \cdot \Sigma_V \widehat{\Delta_\nu f_V}) + m \Sigma_S \widehat{\Delta_\nu f_V}) \\ & = \Delta_\nu C_V[f_V] + q^\rho (\Delta_\nu \widehat{\Sigma_{V\rho} f_V}) + F_\nu^\rho \widehat{\Sigma_{V\rho} f_V} + m (\Delta_\nu \widehat{f_V}) \Sigma_S. \end{aligned} \quad (3.7)$$

Then we may re-express $\hat{C}_2^{(n)\mu}$ as

$$\begin{aligned} \hat{C}_2^{(n)\mu} = & \frac{\delta(q^2 - m^2)}{2} \left\{ \epsilon^{\mu\nu\rho\sigma} q_\nu (\Delta_\rho \widehat{\Sigma_{V\sigma} f_V}) + 2 S_{m(n)}^{\mu\nu} \left[m (\Delta_\nu \widehat{\Sigma_S} f_V) + F_\nu^\rho \widehat{\Sigma_{V\rho} f_V} + q^\rho (\Delta_\nu \widehat{\Sigma_{V\rho} f_V}) \right. \right. \\ & \left. \left. - (q \cdot \Delta \widehat{\Sigma_{V\nu} f_V}) \right] + 2 \widehat{f_V} \widehat{\Sigma_{V\nu}} (q \cdot \Delta S_{m(n)}^{\mu\nu} - F_\lambda^\mu S_{m(n)}^{\lambda\nu}) \right\} \\ & - \left[(2 S_{m(n)}^{\mu\nu} q^\lambda F_{\lambda\nu} - \tilde{F}^{\mu\nu} q_\nu) \delta'(q^2 - m^2) + \delta(q^2 - m^2) (\Delta_\nu S_{m(n)}^{\mu\nu}) \right] (C_V[f_V] + m C_S[f_V]) \\ & + \frac{\delta(q^2 - m^2)}{2m} (q^\mu (q \cdot \Delta \widehat{\Sigma_P} f_V) - m^2 (\partial^\mu \widehat{\Sigma_P} f_V)). \end{aligned} \quad (3.8)$$

In fact, we can further decompose $\hat{C}_2^{(n)\mu}$ into the piece proportional to q^μ , which survives in the massless limit and reproduces the collision term in CKT, and another piece proportional

to m , which stems from the purely finite-mass correction. This decomposition can be used as a consistency check in the massless limit. Note that the procedure is simply to rewrite $\hat{C}_2^{(n)\mu}$ in the form with a more apparent connection to the CKT despite some of the computational complexity. A similar procedure and comparison are also performed for the free-streaming AKE in ref. [51]. We hence present such a complicated yet straightforward check in appendix C (see eq. (C.15)), while we will later show such an alternative expression for $\hat{C}_2^{(n)\mu}$ in a simpler case suitable for the application in HIC.

Although $\hat{C}_1^{(n)\mu} + \hbar\hat{C}_2^{(n)\mu}$ given by eq. (3.4) and eq. (3.8) serves as the generic collision term in the presence of spacetime-dependent background electromagnetic fields and an arbitrary spacetime-dependent frame vector $n^\mu(X)$, we may drop unnecessary terms for practical applications in HIC. First, it is more convenient to work with a constant frame vector such that $\partial_\mu n^\nu = 0$. More precisely, we could simply take $n^\mu = (1, \mathbf{0})$, which also corresponds to the frame choice for the CKT presented in early works obtained from the Berry phase [18, 19]. Note that the choice of a frame vector is analogous to the choice of a gauge, which does not affect the physics in the end (see e.g. refs. [30, 51] for comprehensive discussions). Second, the background electromagnetic fields and particularly the magnetic field may only exist in HIC for a rather short period in the pre-equilibrium phase although the finite electric conductivity of QGP may slightly mitigate the decrease of magnetic fields in time [67, 68]. It is thus more practical to drop the contributions from electromagnetic fields in the QGP phase. In such a case, the explicit form of $\hat{C}_1^{(n)\mu}$ remains unchanged. On the other hand, not only $\hbar\hat{C}_2^{(n)\mu}$ but also the free-streaming part in the AKE become much simpler. The AKE now is given by

$$\delta(q^2 - m^2) \left\{ q \cdot \partial (a^\mu f_A) + a^\mu q_\nu \widehat{\Sigma_V^\nu f_A} + m^2 \widehat{\Sigma_A^\mu f_V} + q^\mu q_\nu \widehat{\Sigma_A^\nu f_V} + m \left(a^\mu \widehat{\Sigma_S f_A} + \frac{1}{2} \epsilon^{\mu\nu\rho\sigma} q_\nu \widehat{\Sigma_{T\rho\sigma} f_V} \right) - \frac{\hbar}{2} \left[\epsilon^{\mu\nu\rho\sigma} q_\nu (\partial_\rho \widehat{\Sigma_{V\sigma}}) f_V + 2S_{m(n)}^{\mu\nu} \left(m (\partial_\nu \widehat{\Sigma_S}) f_V + q^\rho (\partial_\nu \widehat{\Sigma_{V\rho}}) f_V - (q \cdot \partial \widehat{\Sigma_{V\nu}}) f_V \right) \right] \right\} = 0, \quad (3.9)$$

where we also took $\Sigma_P = \bar{\Sigma}_P = 0$ as expected higher-order contributions. Note that the \hbar terms shown above only come from $\hbar\hat{C}_2^{(n)\mu}$.

By employing the Schouton identity

$$\eta_\mu^\lambda \epsilon_{\rho\nu\alpha\beta} - \eta_\rho^\lambda \epsilon_{\mu\nu\alpha\beta} - \eta_\nu^\lambda \epsilon_{\rho\mu\alpha\beta} - \eta_\alpha^\lambda \epsilon_{\rho\nu\mu\beta} - \eta_\beta^\lambda \epsilon_{\rho\nu\alpha\mu} = 0, \quad (3.10)$$

one finds

$$S_{m(n)}^{\mu\nu} q^\alpha = q^\mu S_{m(n)}^{\alpha\nu} + q^\nu S_{m(n)}^{\mu\alpha} + \frac{\epsilon^{\mu\nu\rho\alpha} q_\rho}{2} - \frac{\epsilon^{\mu\nu\rho\alpha}}{2(q \cdot n + m)} (mq_\rho + q^2 n_\rho). \quad (3.11)$$

Given the relation above, the AKE in eq. (3.9) can be alternatively written as

$$\delta(q^2 - m^2) \left\{ q \cdot \partial (a^\mu f_A) + a^\mu q_\nu \widehat{\Sigma_V^\nu f_A} + m^2 \widehat{\Sigma_A^\mu f_V} + q^\mu q_\nu \widehat{\Sigma_A^\nu f_V} + m \left(a^\mu \widehat{\Sigma_S f_A} + \frac{1}{2} \epsilon^{\mu\nu\rho\sigma} q_\nu \widehat{\Sigma_{T\rho\sigma} f_V} \right) + \hbar \left[q^\mu S_{m(n)}^{\rho\nu} (\partial_\rho \widehat{\Sigma_{V\nu}}) f_V - m \left(S_{m(n)}^{\mu\nu} (\partial_\nu \widehat{\Sigma_S}) f_V + \frac{\epsilon^{\mu\nu\rho\sigma} (q_\rho + mn_\rho)}{2(q \cdot n + m)} (\partial_\sigma \widehat{\Sigma_{V\nu}}) f_V \right) \right] \right\} = 0, \quad (3.12)$$

where we rearrange the explicit \hbar corrections in the collision term based on the decomposition for the terms proportional to q^μ and to m , respectively. One can now more easily check that eq. (3.12) reduces to the CKT with a constant frame vector in the absence of electromagnetic fields [30] in the massless limit by taking $m = 0$ and $a^\mu = q^\mu$. We shall discuss later about the physical interpretations of the explicit \hbar corrections in eq. (3.9) and in eq. (3.12) as two mathematically equivalent expressions.

3.2 Rest-frame expression

Notably, when focusing on massive fermions with mass much greater than the gradient scale, we can set the frame vector at their rest frame $n^\mu = n_r^\mu(q) = q^\mu/m$ to simplify both the Wigner functions and AKE. Such a frame choice is also applied in refs. [49, 50]. Nonetheless, this frame choice is rather different from the previous one when $n^\mu(X)$ only depends on spacetime coordinates. In such a case, the magnetization-current term in \mathcal{A}^μ vanishes and the \mathcal{A}^μ reduces to

$$\mathcal{A}^\mu = 2\pi \left[\delta(q^2 - m^2) a^\mu f_A + \hbar \tilde{F}^{\mu\nu} q_\nu \delta'(q^2 - m^2) f_V + \frac{\hbar \delta(q^2 - m^2)}{2m} q^\mu C_P[f_V] \right]. \quad (3.13)$$

Accordingly, the AKE from eq. (2.20f) becomes

$$\square^{(n_r)} \mathcal{A}^\mu = \hat{\mathcal{C}}_1^{(n_r)\mu} + \hbar \hat{\mathcal{C}}_2^{(n_r)\mu}, \quad (3.14)$$

where $\hat{\mathcal{C}}_1^{(n_r)\mu}$ is the same as eq. (3.4) by taking $n^\mu = n_r^\mu$, while

$$\begin{aligned} \square^{(n_r)} \mathcal{A}^\mu &= \delta(q^2 - m^2) \left(q \cdot \Delta(a^\mu f_A) + F^{\nu\mu} a_\nu f_A - \frac{1}{2} \hbar \epsilon^{\mu\nu\rho\sigma} q_\rho (\partial_\sigma F_{\beta\nu}) \partial_q^\beta f_V \right) \\ &\quad + \hbar \tilde{F}^{\mu\nu} q_\nu \delta'(q^2 - m^2) q \cdot \Delta f_V \end{aligned} \quad (3.15)$$

and

$$\begin{aligned} \hat{\mathcal{C}}_2^{(n_r)\mu} &= \frac{\delta(q^2 - m^2)}{2} \left(\epsilon^{\mu\nu\rho\sigma} q_\nu (\Delta_\rho \widehat{\Sigma_{V\sigma}}) f_V + \frac{1}{m} q^\mu (q \cdot \Delta \widehat{\Sigma_P}) f_V - m (\partial^\mu \widehat{\Sigma_P}) f_V \right) \\ &\quad - \tilde{F}^{\mu\nu} q_\nu \delta'(q^2 - m^2) (q \cdot \widehat{\Sigma_V} f_V + m \widehat{\Sigma_S} f_V). \end{aligned} \quad (3.16)$$

Finally, when considering the application to a heavy quark traveling in QGP, the AKE in the rest frame could be simplified as

$$\begin{aligned} \delta(q^2 - m^2) \left\{ q \cdot \partial(a^\mu f_A) + a^\mu q_\nu \widehat{\Sigma_V^\nu} f_A + m^2 \widehat{\Sigma_A^\mu} f_V + q^\mu q_\nu \widehat{\Sigma_A^\nu} f_V \right. \\ \left. + m \left(a^\mu \widehat{\Sigma_S} f_A + \frac{1}{2} \epsilon^{\mu\nu\rho\sigma} q_\nu \widehat{\Sigma_{T\rho\sigma}} f_V \right) - \frac{\hbar}{2} \epsilon^{\mu\nu\rho\sigma} q_\nu (\partial_\rho \widehat{\Sigma_{V\sigma}}) f_V \right\} = 0, \end{aligned} \quad (3.17)$$

by taking $\Sigma_P = \bar{\Sigma}_P = 0$ and $F_{\mu\nu} = 0$.

In general, when involving also the \hbar corrections in \mathcal{V}^μ , such a frame choice is only valid when m is much larger than the gradient and electromagnetic scales in the system. The magnetization-current term in \mathcal{V}^μ explicitly reveals the breakdown for the choice of a

rest frame away from the aforementioned regime. Although we do not explicitly include the \hbar term in \mathcal{V}^μ based on our power counting, it is still essential to be aware of the valid regime for the frame choice $n^\mu = n_r^\mu$. In heavy-ion phenomenology, one may assume the validity is held, which could be “somewhat applicable” for the spin transport of strange quarks in QGP. For up and down quarks or other applications, it is inevitable to maintain the general frame vector $n^\mu = n^\mu(X)$. Note that the quantum correction on collisions in eq. (3.16) is also presented in eq. (3.5). We may regard other terms in eq. (3.5) as the $\mathcal{O}(|\mathbf{q}|/m)$ corrections on top of eq. (3.16), where \mathbf{q} here denotes the spatial momentum.

3.3 \hbar sorting with the present order counting

In order to explicitly disentangle the (classical) spin-diffusion and (quantum) spin-polarization parts in collisions, we have to retrieve the \hbar terms in $\Sigma_{V\rho}$, Σ_S , $\Sigma_{A\rho}$ and $\Sigma_{T\rho\sigma}$. That is, we have to further make the decomposition $\bar{\Sigma}_{V\rho} = \bar{\Sigma}_{V\rho}^{\text{cl}} + \hbar\bar{\Sigma}_{V\rho}^{\text{Q}(n)}$, $\bar{\Sigma}_S = \bar{\Sigma}_S^{\text{cl}} + \hbar\bar{\Sigma}_S^{\text{Q}(n)}$, $\bar{\Sigma}_{A\rho} = \bar{\Sigma}_{A\rho}^{\text{cl}} + \hbar\bar{\Sigma}_{A\rho}^{\text{Q}(n)}$ and $\bar{\Sigma}_{T\rho\sigma} = \bar{\Sigma}_{T\rho\sigma}^{\text{cl}} + \hbar\bar{\Sigma}_{T\rho\sigma}^{\text{Q}}$, where their explicit forms depend on the details of collisions in systems. Nonetheless, $\hbar\bar{\Sigma}_{V\rho}^{\text{Q}(n)}$ and $\hbar\bar{\Sigma}_S^{\text{Q}(n)}$ are coupled to f_A in $\hat{\mathcal{C}}_1^{(n)\mu}$, which actually contribute to $\mathcal{O}(\hbar^2)$ (in the order of magnitude) corrections from our power counting. Consequently, we only have to retain the quantum corrections from $\hbar\bar{\Sigma}_{A\rho}^{\text{Q}(n)}$ and $\hbar\bar{\Sigma}_{T\rho\sigma}^{\text{Q}}$. One can then rewrite the collision term in the AKE as

$$\hat{\mathcal{C}}_1^{(n)\mu} + \hbar\hat{\mathcal{C}}_2^{(n)\mu} = \hat{\mathcal{C}}_{\text{cl}}^\mu + \hbar\hat{\mathcal{C}}_{\text{Q}}^{(n)\mu}, \quad (3.18)$$

where

$$\hat{\mathcal{C}}_{\text{cl}}^\mu = \delta(q^2 - m^2) \left[q^\mu q_\alpha \widehat{\Sigma_A^{\text{cl}\alpha} f_V} - m^2 \widehat{\Sigma_A^{\text{cl}\mu} f_V} - a^\mu (q_\alpha \widehat{\Sigma_V^{\text{cl}\alpha} f_A} + m \widehat{\Sigma_S^{\text{cl}} f_A}) + m \frac{\epsilon^{\mu\nu\rho\sigma}}{2} q_\nu \widehat{\Sigma_{T\rho\sigma}^{\text{cl}} f_V} \right] \quad (3.19)$$

and

$$\begin{aligned} \hbar\hat{\mathcal{C}}_{\text{Q}}^{(n)\mu} = & \hbar q^\mu \left(\hat{\mathcal{C}}_{q_2}^{(n)} + \delta(q^2 - m^2) q_\nu \widehat{\Sigma_A^{\text{Q}(n)\nu} f_V} \right) + \hbar m \left[\hat{\mathcal{C}}_{m_2}^{(n)\mu} + \delta(q^2 - m^2) \left(\frac{1}{2} \epsilon^{\mu\nu\rho\sigma} q_\nu \widehat{\Sigma_{T\rho\sigma}^{\text{Q}(n)} f_V} \right. \right. \\ & \left. \left. - m \widehat{\Sigma_A^{\text{Q}(n)\mu} f_V} \right) \right]. \end{aligned} \quad (3.20)$$

Analogously, in the rest frame, it is found

$$\hbar\hat{\mathcal{C}}_{\text{Q}}^{(n_r)\mu} = \hbar\hat{\mathcal{C}}_2^{(n_r)\mu} + \hbar\delta(q^2 - m^2) \left[q^\mu q_\alpha \widehat{\Sigma_A^{\text{Q}(n_r)\alpha} f_V} - m^2 \widehat{\Sigma_A^{\text{Q}(n_r)\mu} f_V} + m \frac{\epsilon^{\mu\nu\rho\sigma}}{2} q_\nu \widehat{\Sigma_{T\rho\sigma}^{\text{Q}(n_r)} f_V} \right]. \quad (3.21)$$

Note that the classical part $\hat{\mathcal{C}}_{\text{cl}}^\mu$ is explicitly frame independent. Now, all the \hbar terms are collected into $\hbar\hat{\mathcal{C}}_{\text{Q}}^{(n)\mu}$. Despite complication, one finds that $\hat{\mathcal{C}}_{\text{cl}}^\mu$ is proportional to $a^\mu f_A$; such a term hence results in the diffusion of spin.³ On the contrary, $\hat{\mathcal{C}}_{\text{Q}}^{(n)\mu}$ is instead proportional to f_V and \bar{f}_V . Even when initial spin ($\sim a^\mu f_A$) is zero, such a term can

³Here $\Sigma_A^{\text{cl}\alpha}$ and $\Sigma_{T\rho\sigma}^{\text{cl}}$ are in principle proportional to the axial-charge part of the Wigner functions for outgoing fermions. Therefore, the terms coupled to f_V in $\hat{\mathcal{C}}_{\text{cl}}^\mu$ are also proportional to the spin four vector. One may see an explicit example for application to QGP in the following section.

lead to the spin polarization from the entangled vector-charge transport. Similarly, when applying eqs. (3.9), (3.12), and (3.17), one should recall these extra \hbar corrections from $\hbar\bar{\Sigma}_{A\rho}^{Q(n)}$ and $\hbar\bar{\Sigma}_{T\rho\sigma}^{Q(n)}$ or from $\hbar\bar{\Sigma}_{A\rho}^{Q(n_r)}$ $\hbar\bar{\Sigma}_{T\rho\sigma}^{Q(n_r)}$.

3.4 Discussions for the collision terms

In this subsection, we would like to make a comparison between the AKE in different frame choices and discuss about their physical interpretations. However, to avoid complications, we focus on the simplified versions in eqs. (3.9), (3.12), and (3.17), which are the primary results in connection to the spin polarization of quarks in QGP. For preciseness, we will hereafter dub the terms proportional to \hbar in eqs. (3.9), (3.12), and (3.17) as explicit quantum (\hbar) corrections. In contrast, we refer the \hbar corrections encoded in self-energies discussed in section 3.3 as implicit quantum (\hbar) corrections.

First of all, eqs. (3.9) and (3.12) as mathematically equivalent expressions both work for an arbitrary mass of fermions. Nevertheless, the expression of eq. (3.9) could be more useful in the large-mass regime. When comparing eq. (3.9) with eq. (3.17) as an effective AKE with large-mass fermions, one finds they both incorporate the term $\frac{\hbar}{2}\epsilon^{\mu\nu\rho\sigma}q_\nu(\partial_\rho\widehat{\Sigma_{V\sigma}})f_V$ as an explicit quantum correction in collisions. In the non-relativistic condition, such a term further dominates over the rest of explicit \hbar corrections coupled with $S_{m(n)}^{\mu\nu}$ in eq. (3.9). When $m \rightarrow \infty$, one finds $q^\mu \rightarrow n^\mu m$ and accordingly $S_{m(n)}^{\mu\nu} \rightarrow 0$. It turns out that eq. (3.9) and eq. (3.14) coincide in the heavy-quark (fermion) limit. Consequently, as briefly mentioned in section 3.2, eq. (3.9) further incorporates the $\mathcal{O}(|\mathbf{q}|/m)$ corrections on top of eq. (3.17). It is thus more practical to utilize eq. (3.9) for exploring heavy-quark transport with the inclusion of non-relativistic corrections. On the contrary, eq. (3.12) has a more explicit connection to the CKT in the massless limit. The term $\hbar q^\mu S_{m(n)}^{\rho\nu}(\partial_\rho\widehat{\Sigma_{V\nu}})f_V$ in explicit \hbar corrections of collisions therein matches the \hbar correction in the collision term of CKT up to a prefactor q^μ . In fact, the prefactor q^μ further manifests that such a term is pertinent to the side-jump phenomena associated with the spin polarization enslaved by the momentum and chirality, which forces a^μ in the free-streaming part to align with q^μ . On the other hand, the rest of \hbar terms proportional to m in eq. (3.12) stem from the finite-mass effect suppressed by $\mathcal{O}(m/|\mathbf{q}|)$ in the relativistic limit, which could modify the orientation of spin characterized by the direction of a^μ . Therefore, the expression in eq. (3.12) could be more suitable for analyzing the axial-charge diffusion and spin polarization of light quarks in QGP.

In addition, similar to the case for CKT in the massless limit, the quantum corrections in the AKE now only come from collisions when choosing a constant frame vector in the absence of electromagnetic fields. As already mentioned in section 3.3 for further separation of the classical and quantum parts in the collision term, the classical part in eqs. (3.9), (3.12), and (3.17) with the same expression will yield the spin diffusion. However, it is believe that the spin polarization in HIC is led by the local vorticity of QGP. It is obvious to see the explicit \hbar corrections in the collision term now originate from the inhomogeneity of self-energies, from which the self-energies gradients could incorporate such vortical corrections in collisions. Also, these terms manifest the spin-orbit interactions through collisions, which entangle the dynamical evolution between $a^\mu f_A$ and f_V .

Nevertheless, the implicit \hbar corrections encoded in self-energies should also be taken into account, where at least the \hbar correction upon Wigner functions for outgoing quarks will contribute to the vortical corrections as well.

In a recent study for the application of CKT on the chiral radiation transport theory for neutrinos in core-collapse supernovae, it is shown the similar \hbar corrections of CKT give rise to the vorticity corrections associated with fluid helicity in the collision term for the neutrino absorption process [43]. One may expect a similar scenario when considering a strange quark probing the QGP near equilibrium yet with local vortical fields. Nonetheless, unlike the weak interaction governed the Weinberg-Salam model, the details of collisions in QGP are more sophisticated due to the interacting gluons. Even though the implicit \hbar corrections encoded in self-energies from light quarks in equilibrium could be derived from equilibrium Wigner functions shown in e.g. refs. [31, 51], how to include analogous corrections led by vorticity from polarized gluons is currently unknown. For the future application on the dynamical spin polarization of strange quarks traversing QGP, we will have to work out the Wigner functions for polarized gluons up to $\mathcal{O}(\hbar)$ with both classical and quantum contributions at least in equilibrium as one of essential ingredients. Such a development is beyond the scope of this work and left as the future research direction. It is however worthwhile to note that even the classical part of the collision term in AKE is an innovation. We will further apply such a theoretical framework to derive an explicit expression of the spin diffusion term for massive quarks traversing weakly-coupled QGP in the next section.

4 Example: spin diffusion of quarks in weakly coupled QGP

4.1 Scattering between massive fermions and a medium

We now apply the formalism established in the previous sections to investigate the collision term for massive quarks traversing weakly-coupled QGP in relativistic heavy ion collisions. For simplicity, we will just focus on the spin-diffusion term such as $\hat{\mathcal{C}}_{\text{cl}}^\mu$ in the AKE and leave the $\hbar\hat{\mathcal{C}}_{\text{Q}}^{(n)\mu}$ for future study. Recently, a related study for spin diffusion has been presented in ref. [54] with a different approach. We will mostly follow the theoretical setup therein. In the following, we call fermions quarks and gauge bosons gluons interchangeably, and include the color-group factors. Here, the color degrees of freedom do not play crucial roles (like in the color conductivity), and the same computation holds for QED with simple replacements of the relevant degrees of freedom. Furthermore, we consider the massive quarks with quark mass much greater than the scale of thermal mass in QGP and accordingly neglect the Compton scattering with gluons as the subleading effects analogous to the study of heavy-quark transport in heavy ion collisions (See e.g. ref. [69] and the same approximation in ref. [54]). Note that the $\mathcal{O}(f_A^2)$ terms are not dropped a priori in the calculations, whereas shall see that only the terms linear to f_A remain in the final result, which thus agrees with our power-counting scheme.

The gluon-exchange processes between a massive fermion and the medium constitutes are written down as

$$\Sigma^{>(<)}(q, X) = \lambda_c \int_{q'} \gamma^\mu S^{>(<)}(q', X) \gamma^\nu G_{\mu\nu}^{>(<)}(q - q', X), \quad (4.1)$$

where λ_c denotes an overall coefficient including the coupling and we drop $\mathcal{O}(\hbar^2)$ and higher-order correction in our order counting. The gluon propagator $G_{\mu\nu}^{>(<)}$ contains the information of the spectral functions which depend on the scatterers. Here, we assume a dilute population of the massive fermions in the medium and neglect the contributions of the massive scatterers, and retain the massless-fermion and gluon scatterers. Having assumed the weakly coupled system, we focus on the lowest-order contributions in the coupling constant g_c , i.e., the 2-to-2 scatterings between a massive quark and a massless quark/gluon.

Inserting eq. (2.7), one can straightforwardly decompose the gamma structures in eq. (4.1) as

$$\begin{aligned}\chi_{q'}^{>\mu\nu} &\equiv \gamma^\mu \mathcal{S}^>(q') \gamma^\nu \\ &= \bar{\mathcal{S}}(\eta^{\mu\nu} - i\sigma^{\mu\nu}) - i\bar{\mathcal{P}}\left(\eta^{\mu\nu}\gamma^5 + \frac{\epsilon^{\mu\nu\alpha\beta}}{2}\sigma_{\alpha\beta}\right) + \bar{\mathcal{V}}_\rho(\eta^{\mu\rho}\gamma^\nu + \eta^{\rho\nu}\gamma^\mu - \eta^{\mu\nu}\gamma^\rho + i\epsilon^{\mu\nu\rho\sigma}\gamma^5\gamma_\sigma) \\ &\quad + \bar{\mathcal{A}}_\rho(\eta^{\mu\nu}\gamma^5\gamma^\rho - \eta^{\rho\nu}\gamma^5\gamma^\mu - \eta^{\mu\rho}\gamma^5\gamma^\nu + i\epsilon^{\mu\rho\nu\sigma}\gamma_\sigma) \\ &\quad + \bar{\mathcal{S}}_{\alpha\beta}\left(\eta^{\mu\alpha}\sigma^{\beta\nu} + \eta^{\nu\alpha}\sigma^{\beta\mu} + \frac{\eta^{\mu\nu}}{2}\sigma^{\alpha\beta} + i\eta^{\mu\alpha}\eta^{\beta\nu} + \frac{\epsilon^{\mu\nu\alpha\beta}}{2}\gamma^5\right).\end{aligned}\quad (4.2)$$

Thus, contracted with the gluon propagator, we have

$$\begin{aligned}\chi_{q'}^{>\mu\nu} G_{\mu\nu}^> &= (\bar{\mathcal{S}}G_{\mu\nu}^> + i\bar{\mathcal{S}}^{\mu\nu}G_{\mu\nu}^>) + i\gamma^5\left(-\bar{\mathcal{P}}G_\mu^{>\mu} - i\bar{\mathcal{S}}_{\alpha\beta}G_{\mu\nu}^>\frac{\epsilon^{\mu\nu\alpha\beta}}{2}\right) \\ &\quad + \gamma^\rho(\bar{\mathcal{V}}^\mu(G_{(\mu\rho)}^> - \bar{\mathcal{V}}_\rho G_\mu^{>\mu} - i\epsilon_{\mu\nu\sigma\rho}\bar{\mathcal{A}}^\sigma G^{>\mu\nu})) \\ &\quad + \gamma^5\gamma^\rho(-\bar{\mathcal{A}}^\mu G_{(\mu\rho)}^> + \bar{\mathcal{A}}_\rho G_\mu^{>\mu} + i\epsilon_{\mu\nu\sigma\rho}\bar{\mathcal{V}}^\sigma G^{>\mu\nu}) \\ &\quad + \frac{1}{2}\sigma^{\rho\sigma}\left(2\bar{\mathcal{S}}_\rho^\mu G_{(\mu\sigma)}^> + \bar{\mathcal{S}}_{\rho\sigma}G_\mu^{>\mu} - 2i\bar{\mathcal{S}}G_{\rho\sigma}^> - i\bar{\mathcal{P}}\epsilon_{\mu\nu\rho\sigma}G^{>\mu\nu}\right),\end{aligned}\quad (4.3)$$

where $A_{(\mu}B_{\nu)} = A_\mu B_\nu + A_\nu B_\mu$. We consider gluon propagators $G_{\mu\nu}^{<(>)}$ symmetric in the Lorentz indices, and then find that the imaginary terms in the above vanish in the contractions:

$$\begin{aligned}\Sigma^>(q) &= \lambda_c \int_{q'} \left[\bar{\mathcal{S}}G_\mu^{>\mu} - i\bar{\mathcal{P}}G_\mu^{>\mu}\gamma^5 + \gamma^\rho(2\bar{\mathcal{V}}^\mu G_{\mu\rho}^> - \bar{\mathcal{V}}_\rho G_\mu^{>\mu}) \right. \\ &\quad \left. + \gamma^5\gamma^\rho(-2\bar{\mathcal{A}}^\mu G_{\mu\rho}^> + \bar{\mathcal{A}}_\rho G_\mu^{>\mu}) + \frac{\sigma^{\rho\sigma}}{2}(2\bar{\mathcal{S}}_{\mu[\rho}G_{\sigma]}^{>\mu} + \bar{\mathcal{S}}_{\rho\sigma}G_\mu^{>\mu}) \right],\end{aligned}\quad (4.4)$$

where $A_{[\mu}B_{\nu]} = A_\mu B_\nu - A_\nu B_\mu$. In general, $G_{\mu\nu}^{<(>)}$ possibly contains anti-symmetric components led by scatterings with spin-polarized scatterers in the medium. For example, when considering the scattering with massless quarks, such anti-symmetric components can arise from the side-jump terms, whereas $\chi_{q'}^{>\mu\nu}G_{\mu\nu}^>$ should still remain real. Such quantum corrections from a polarized medium will not be considered in the present work.

One can now read out the corresponding terms up to $\mathcal{O}(\hbar)$ between eqs. (2.8) and (4.4) as

$$\bar{\Sigma}_S = \lambda_c \int_{q'} \bar{S}_{q'} G_\alpha^{>\alpha} = \frac{\lambda_c}{m} \int_{q'} (q' \cdot \bar{V}_{q'}) G_\alpha^{>\alpha}, \quad (4.5a)$$

$$\bar{\Sigma}_P = \lambda_c \int_{q'} (-\bar{P}_{q'}) G_\alpha^{>\alpha}, \quad (4.5b)$$

$$\bar{\Sigma}_{V\mu} = \lambda_c \int_{q'} \bar{V}_{q'}^\alpha (2G_{\alpha\mu}^{>} - G_\beta^{>\beta} \eta_{\alpha\mu}), \quad (4.5c)$$

$$\bar{\Sigma}_{A\mu} = \lambda_c \int_{q'} \bar{A}_{q'}^\alpha (-2G_{\alpha\mu}^{>} + G_\beta^{>\beta} \eta_{\alpha\mu}), \quad (4.5d)$$

$$\bar{\Sigma}_{T\mu\nu} = \lambda_c \int_{q'} (2\bar{S}_{\alpha[\mu} G_{\nu]}^{>\alpha} + G_\alpha^{>\alpha} \bar{S}_{\mu\nu}) = \frac{\lambda_c}{m} \int_{q'} (2\bar{S}_{\alpha[\mu} G_{\nu]}^{>\alpha} + G_\alpha^{>\alpha} \bar{S}_{\mu\nu}). \quad (4.5e)$$

The rightmost sides are obtain by using eqs. (A.19a)–(A.19c) up to the linear orders in Σ 's and \hbar . In the present case, we confirm that $\bar{\Sigma}_P$ is at $\mathcal{O}(\hbar^2)$ as anticipated earlier.

Now, given explicit forms of $G_{\mu\nu}^{>}$, \bar{V}^μ , and \bar{A}^μ , we can directly evaluate $\Sigma_\mu^{>}$ from eq. (4.5) and $\Sigma_\mu^{<}$ in the same fashion. Inserting these expressions into the collision terms in the SKE (2.23), we have

$$(q \cdot \bar{\Sigma}_V + m\bar{\Sigma}_S) = \lambda_c \int_{q'} 2\pi\delta(q'^2 - m^2) \left(2q'^\mu G_{\mu\nu}^{>} q^\nu - p \cdot q' G_\mu^{>\mu} \right) \bar{f}_{Vq'}, \quad (4.6)$$

where $p^\mu = q^\mu - q'^\mu$. The other term $(q \cdot \Sigma_V + m\Sigma_S)$ takes a similar form. In addition, by making the decompositions $\bar{\Sigma}_{A\rho} = \bar{\Sigma}_{A\rho}^{\text{cl}} + \hbar\bar{\Sigma}_{A\rho}^{\text{Q}(n)}$ and $\bar{\Sigma}_{T\rho\sigma} = \bar{\Sigma}_{T\rho\sigma}^{\text{cl}} + \hbar\bar{\Sigma}_{T\rho\sigma}^{\text{Q}}$, we identify the classical and quantum parts. From the classical part in eq. (2.24), eqs. (4.5d) and (4.5e) yield

$$\bar{\Sigma}_{A\rho}^{\text{cl}} = \lambda_c \int_{q'} 2\pi\delta(q'^2 - m^2) (a_{q'\rho} G_\mu^{>\mu} - 2a_{q'}^\mu G_{\mu\rho}^{>}) \bar{f}_A(q'), \quad (4.7)$$

$$\bar{\Sigma}_{T\rho\sigma}^{\text{cl}} = -\frac{\lambda_c}{m} \int_{q'} 2\pi\delta(q'^2 - m^2) \left(2G^{>\mu}{}_{[\sigma} \epsilon_{\mu\rho]\alpha\beta} + G_\mu^{>\mu} \epsilon_{\rho\sigma\alpha\beta} \right) q'^\alpha a_{q'}^\beta \bar{f}_A(q'). \quad (4.8)$$

Those terms are further investigated with a specific gluon propagator provided by the hard-thermal loop approximation in the next section. The quantum parts are also identified in the same way. However, computation of those quantum corrections with specific gluon propagators are left as open issues. Note also that we have dropped possible antisymmetric parts of the gluon propagator in eq. (4.4).

4.2 Weakly coupled QGP and hard-thermal-loop approximation

Although eqs. (4.5a)–(4.5e) work for even non-equilibrium media, we now focus on the spin transport in equilibrium QGP as a concrete example. In such a case, we can make further simplification for the self-energy in eq. (4.1). For simplicity, we will only evaluate the spin-diffusion terms in the SKE and AKE, while the claculation for the spin-polarization term in the AKE is more involved, which will be presented in the followup work. We will implement the hard-thermal-loop (HTL) approximation, which allows us to derive the

leading-logarithmic result in weakly coupled QCD as in the derivation shown in ref. [54] from a distinct approach. Recall that $p^\mu = (q - q')^\mu$ and g_c denotes the coupling constant for strong interaction. We then apply the cut-gluon propagator in the Coulomb gauge and the fluid-rest frame, which gives rise to

$$G_{\mu\nu}^{<(>)}(q, q') \approx g_{\text{eqp}}^{<(>)} \left[\rho_L(p) P_{\mu\nu}^C + \rho_T(p) P_{\mu\nu}^T \right] \quad (4.9)$$

with

$$P_{\mu\nu}^C \equiv u_\mu u_\nu, \quad P_{\mu\nu}^T \equiv -\Theta_{\mu\alpha} \Theta_{\nu\beta} \left(\eta^{\alpha\beta} + \frac{p^\alpha p^\beta}{|\mathbf{p}|^2} \right) = - \left(\Theta_{\mu\nu} + \frac{p_\perp^\mu p_\perp^\nu}{|\mathbf{p}|^2} \right), \quad (4.10)$$

where $g_{\text{eqp}}^{<} = g_{0p} = 1/(e^{\beta p \cdot u} - 1)$, $g_{\text{eqp}}^{>} = 1 + g_{0p}$ and $\Theta^{\mu\nu} \equiv \eta^{\mu\nu} - u^\mu u^\nu$. Here, u^μ and $\beta = 1/T$ denote the fluid four velocity and the inverse of temperature in local equilibrium, respectively. We have introduced notations:

$$V^0 \equiv V \cdot u, \quad V_\perp^\mu \equiv V^\mu - V^0 u^\mu = \Theta^{\mu\nu} V_\nu, \quad (4.11a)$$

$$\mathbf{V}^i \equiv V_\perp^i, \quad \hat{\mathbf{V}}^i = \hat{V}_\perp^i \equiv V^i / |\mathbf{V}|, \quad (4.11b)$$

for an arbitrary vector V^μ . Then, we have $V_\perp \cdot k_\perp = -\mathbf{V} \cdot \mathbf{k}$ and, especially, $V_\perp^2 = -|\mathbf{V}|^2$ when $k^\mu = V^\mu$.

In our setup, the HTL approximation is more precisely applied to $g_c T \ll |p^\mu| \ll T$. On the other hand, $\rho_{L/T}(p)$ correspond to the HTL gluon spectral densities, which take explicit forms as (e.g., see ref. [70])

$$\rho_L(p) \approx \frac{\pi m_D^2 p_0}{|\mathbf{p}|^5}, \quad \rho_T(p) \approx \frac{\pi m_D^2 p_0}{2|\mathbf{p}|^5 \left(1 - \left(\frac{p_0}{|\mathbf{p}|} \right)^2 \right)}, \quad (4.12)$$

where $m_D \sim g_c T$ corresponds to the Debye mass. The explicit form from the gluons in $SU(N_c)$ color group and the N_f -flavored massless quarks is given by

$$m_D^2 = \frac{g_c^2 T^2 (2N_c + N_f)}{6}, \quad \lambda_c = g_c^2 C_2(F) = \frac{g_c^2 (N_c^2 - 1)}{2N_c}. \quad (4.13)$$

Moreover, one should keep in mind the relation

$$G_{\mu\nu}^{>}(p) = (1 + g_{0p}^{-1}) G_{\mu\nu}^{<}(p) \quad (4.14)$$

from detailed balance. In light of the theoretical frameworks constructed in the previous section, we may now write down the SKE and AKE in the HTL approximation. It is interesting that one can linearize the kinetic equations in terms of the distribution functions by taking $g_{0p}^{-1} \rightarrow 0$ and $G_{\mu\nu}^{>}(p) \approx G_{\mu\nu}^{<}(p)$. However, in the practical calculation, we have to at least approximate $g_{0p} \approx T/p_0 - 1/2 + p_0/(12T) + \mathcal{O}(p_0^3/T^3)$ to keep all relevant terms contributing to the leading logarithmic order. In the following, we will append subindices to $f_{V/A}$ and a^μ for specifying their momentum dependence.

For the SKE in eq. (2.23), it is found

$$0 = \delta(q^2 - m^2) \left[q \cdot \partial f_{Vq} + \lambda_c \int_p Q_1(q, p) \{ (1 + g_{0p}^{-1}) \bar{f}_{Vq'} f_{Vq} - \bar{f}_{Vq} f_{Vq'} \} \right], \quad (4.15)$$

where

$$Q_1(q, p) = 2\pi\delta(q'^2 - m^2) \left(2q'^\mu G_{\mu\nu}^< q^\nu - p \cdot q' G_\mu^< \right). \quad (4.16)$$

By using $\bar{f}_{Vq} = 1 - f_{Vq}$, we may further rewrite eq. (4.15) into

$$0 = \delta(q^2 - m^2) \left[q \cdot \partial f_{Vq} + \lambda_c \int_p \left\{ Q_1(f_{Vq} - f_{Vq-p}) + \tilde{Q}_1 f_{Vq}(1 - f_{Vq'}) \right\} \right], \quad (4.17)$$

where $\tilde{Q}_1 = g_{0p}^{-1} Q_1$. One can then further approximate $f_{Vq} - f_{Vq-p} \approx p^\mu \partial_{q^\mu} f_{Vq} + \mathcal{O}(|p|/|q|)$ assuming the small-momentum transfer $|p| \ll |q|$.

Next, we can simplify the AKE with the same approximation. Inserting the rightmost sides of eq. (4.5) into eq. (3.19), we obtain

$$\delta(q^2 - m^2) [q \cdot \partial \tilde{a}_q^\mu + F^{\nu\mu} \tilde{a}_{\nu q} - \hat{C}_{\text{cl}}^\mu] = 0, \quad (4.18a)$$

$$\hat{C}_{\text{cl}}^\mu = \lambda_c \int_p \left(-Q_1 \tilde{a}_q^\mu - \tilde{Q}_1 (1 - f_{Vq-p}) \tilde{a}_q^\mu + Q_2^{\mu\nu} \tilde{a}_{q-p\nu} + \tilde{Q}_2^{\mu\nu} f_{Vq} \tilde{a}_{q-p\nu} \right), \quad (4.18b)$$

where $\tilde{a}_q^\mu \equiv a_q^\mu f_{Aq}$, $\tilde{Q}_2^{\mu\nu} = g_{0p}^{-1} Q_2^{\mu\nu}$, and

$$Q_2^{\mu\nu} = -2\pi\delta(q'^2 - m^2) \left[\left(p^\mu q^\nu - \eta^{\mu\nu} q \cdot p \right) G_\rho^< \rho - 2 \left(p^\mu G_\rho^< \nu q^\rho - q \cdot p G^< \nu\mu \right) \right. \\ \left. + 2 \left(q^\nu G^< \rho\mu q'_\rho - \eta^{\mu\nu} q'_\sigma G^< \sigma\rho q_\rho \right) \right]. \quad (4.19)$$

Here we also use $\bar{f}_{Aq} = -f_{Aq}$. One can similarly approximate $\tilde{a}_{q-p\nu} \approx \tilde{a}_{q\nu} - p^\beta \partial_{q\beta} \tilde{a}_{q\nu}$. Such a diffusion term in collisions of the AKE has also been constructed in ref. [54] from a distinct approach, in which different parameterization of the spin vector is applied. Nonetheless, it may be more practical to adopt our parameterization for the spin vector, which has a direct connection to the axial-charge current equivalent to the spin polarization through the Wigner functions and the combination with the free-streaming part of the AKE. Note also that eqs. (4.17) and (4.18) contain nonlinear terms in distribution functions. While those nonlinear terms are not included in ref. [54], they are imperative to preserve the quantum statistics for fermions. For example, as will be shown, f_{Vq} follows the Fermi-Dirac distribution instead of just the Boltzmann distribution in equilibrium with the vanishing collision term in the SKE. In addition, as shown in eq. (4.18), the nonlinear terms further reveal the entangled dynamics between the vector/axial charges and spin diffusion.

4.3 SKE and AKE with diffusion effects in the leading-log approximation

We now explicitly compute the collision terms in axial kinetic theory with the HTL approximation. Notations have been introduced in eq. (4.11). The basic strategy is to collect all the terms up to $\mathcal{O}(|\mathbf{p}|^{-3})$ in the integrand. When combined with the integral measure, they give rise to the leading logarithmic results in g_c with the cut-offs provided by the HTL resummation. Moreover, we will consider the onshell kinetic equations. We hence take $f_{Vq} = f_{Vq}(\mathbf{q}, X)$ as just a function of \mathbf{q} and X by using $q_0 = E_q = \sqrt{|\mathbf{q}|^2 + m^2}$ for fermions (here we neglect anti-fermions) in the Wigner functions. Similarly, we take $u \cdot \tilde{a}_q = -q_\perp \cdot \tilde{a}_q / q_0$ for $\tilde{a}_{q\mu} = \tilde{a}_{q\mu}(\mathbf{q}, X)$.

4.3.1 Results

The computations for the diffusion terms in SKE and AKE are complicated yet straightforward. We hence present the details of computations for eqs. (4.15) and (4.18) in appendices D and E, respectively. In the following, we just summarize the final results. Up to the leading logarithmic order in g_c , the SKE takes the form

$$0 = \delta(q^2 - m^2) \left[q \cdot \partial - \kappa_{\text{LL}} \left\{ 2(1 - f_{Vq}) + \mathcal{S}^{(1)} \hat{q}_\perp^\beta \partial_{q_\perp^\beta} + \mathcal{S}^{(2)} \hat{q}_\perp^\alpha \hat{q}_\perp^\beta \partial_{q_\perp^\alpha} \partial_{q_\perp^\beta} + \mathcal{S}^{(3)} \eta^{\alpha\beta} \partial_{q_\perp^\alpha} \partial_{q_\perp^\beta} \right\} \right] f_{Vq}, \quad (4.20)$$

where we denote the coefficient of the leading log result $\kappa_{\text{LL}} \equiv [g_c^2 C_2(F) m_D^2 / (8\pi)] \ln(1/g_c)$ and the Minkowski metric $\eta^{\alpha\beta}$. We also introduce the four “velocity” $v^\mu = (v^0, \mathbf{v}^i) \equiv q^\mu / m$, which has the normalization $v^\mu v_\mu = 1$ under the delta function, and then the rapidity $\eta_q \equiv 2^{-1} \ln[(E_q + |\mathbf{q}|) / (E_q - |\mathbf{q}|)] = 2^{-1} \ln[(v^0 + |\mathbf{v}|) / (v^0 - |\mathbf{v}|)]$. The coefficient in each term is given as

$$\mathcal{S}^{(1)} = \frac{m v_0^2 \theta_{-1}}{|\mathbf{v}|^2} (1 - 2f_{Vq}), \quad \mathcal{S}^{(2)} = \frac{m T v_0^2}{2|\mathbf{v}|^3} \left(\frac{|\mathbf{v}|^2 \eta_q}{v_0^2} + \frac{3\theta_1}{v_0} \right), \quad \mathcal{S}^{(3)} = \frac{m T}{2} \left(\frac{v_0^3 \theta_{-3}}{|\mathbf{v}|^3} - 3v_0 \right), \quad (4.21)$$

where $\theta_n \equiv |\mathbf{v}| - v_0^n \eta_q$. Note that eq. (4.20) agrees with the result in ref. [54] except for additional nonlinear terms in f_{Vq} coming from Fermi-Dirac statistics.

Carrying out similar yet more sophisticated computations for eq. (4.18), we also derive the AKE with the spin-diffusion term up to leading-logarithmic order. Combining with the classical free-streaming part dictated by the BMT equation, the AKE reads

$$0 = \delta(q^2 - m^2) \left[q \cdot \partial \tilde{a}_q^\mu - \frac{\kappa_{\text{LL}} T}{E_q} \left(\tilde{a}_q^\mu \dot{\mathcal{Q}}_{\text{cl}}^{(1)} + u^\mu \dot{\mathcal{Q}}_{\text{cl}}^{(2)} + \hat{q}_\perp^\mu \dot{\mathcal{Q}}_{\text{cl}}^{(3)} \right) \right. \\ \left. + \dot{\mathcal{Q}}_{\text{cl}}^{(4)} \hat{q}_\perp^\nu \partial_{q_\perp^\nu} \tilde{a}_{q\nu} + \dot{\mathcal{Q}}_{\text{cl}}^{(5)} \hat{q}_\perp^\nu \partial_{q_\perp^\nu} \tilde{a}_q^\mu + \dot{\mathcal{Q}}_{\text{cl}}^{(6)} \eta^{\nu\rho} \partial_{q_\perp^\nu} \partial_{q_\perp^\rho} \tilde{a}_q^\mu + \dot{\mathcal{Q}}_{\text{cl}}^{(7)} \hat{q}_\perp^\nu \hat{q}_\perp^\rho \partial_{q_\perp^\nu} \partial_{q_\perp^\rho} \tilde{a}_q^\mu \right], \quad (4.22)$$

$$\dot{\mathcal{Q}}_{\text{cl}}^{(1)} = \frac{2m}{T} \left[\left(v_0(1 - 2f_{Vq}) - \frac{T}{m} \right) - \frac{v_0^3}{|\mathbf{v}|^2} m \theta_{-1} \hat{q}_\perp^\rho \partial_{q_\perp^\rho} f_{Vq} \right], \quad (4.23a)$$

$$\dot{\mathcal{Q}}_{\text{cl}}^{(2)} = m \frac{v_0}{|\mathbf{v}|^3} \left((\theta_1 - |\mathbf{v}|^3) \partial_{q_\perp^\nu} \tilde{a}_q^\nu + (3\theta_1 + |\mathbf{v}|^3) \hat{q}_\perp^\nu \hat{q}_\perp^\rho \partial_{q_\perp^\rho} \tilde{a}_{q\nu} \right) \\ + \frac{m}{v_0 |\mathbf{v}|^2 T} \left(v_0^3 (1 - 2f_{Vq}) \theta_{-1} - \frac{2T}{m} (\theta_{-1} + 2|\mathbf{v}|^3) \right) \hat{q}_\perp \cdot \tilde{a}_q, \quad (4.23b)$$

$$\dot{\mathcal{Q}}_{\text{cl}}^{(3)} = \frac{m}{|\mathbf{v}|^2} \left(v_0^2 \theta_{-1} \partial_{q_\perp^\nu} \tilde{a}_q^\nu + (3\theta_1 + |\mathbf{v}|^3) \hat{q}_\perp^\nu \hat{q}_\perp^\rho \partial_{q_\perp^\rho} \tilde{a}_{q\nu} \right) \\ + \frac{m}{|\mathbf{v}| T} \theta_{-1} \left(v_0(1 - 2f_{Vq}) - \frac{2T}{m} \right) \hat{q}_\perp \cdot \tilde{a}_q, \quad (4.23c)$$

$$\dot{\mathcal{Q}}_{\text{cl}}^{(4)} = -2m|\mathbf{v}|, \quad (4.23d)$$

$$\dot{\mathcal{Q}}_{\text{cl}}^{(5)} = \frac{m^2 v_0^3}{|\mathbf{v}|^2 T} \theta_{-1} (1 - 2f_{Vq}), \quad (4.23e)$$

$$\dot{\mathcal{Q}}_{\text{cl}}^{(6)} = -\frac{m^2 v_0^2}{2|\mathbf{v}|^3} (3|\mathbf{v}|^3 - v_0^2 \theta_{-3}), \quad (4.23f)$$

$$\dot{\mathcal{Q}}_{\text{cl}}^{(7)} = \frac{m^2 v_0}{2|\mathbf{v}|^3} (3v_0 \theta_1 + \eta_q |\mathbf{v}|^2). \quad (4.23g)$$

JHEP07(2020)070

4.3.2 Nonrelativistic limit

In the non-relativistic limit ($m \gg T, |\mathbf{q}|$), we have $v_0 \sim 1 + |\mathbf{v}|^2/2$ and $\theta_n \sim -(3n+2)|\mathbf{v}|^3/6$. We immediately find that the SKE and AKE reduce to similar forms,

$$0 = \left(\partial_0 + \frac{q_{\perp}^{\nu} \partial_{\perp\nu}}{m} \right) f_{Vq} + \frac{2}{3} \kappa_{LL} \left[T \eta^{\nu\rho} \partial_{q_{\perp}^{\nu}} \partial_{q_{\perp}^{\rho}} - \frac{1}{m} \left(3(1-f_{Vq}) + (1-2f_{Vq}) q_{\perp}^{\nu} \partial_{q_{\perp}^{\nu}} \right) \right] f_{Vq}, \quad (4.24a)$$

$$0 = \left(\partial_0 + \frac{q_{\perp}^{\nu} \partial_{\perp\nu}}{m} \right) \tilde{a}_q^{\mu} + \frac{2}{3} \kappa_{LL} \left[T \eta^{\nu\rho} \partial_{q_{\perp}^{\nu}} \partial_{q_{\perp}^{\rho}} \tilde{a}_q^{\mu} + \frac{1}{m} \left\{ \left(2(q_{\perp}^{\nu} \partial_{q_{\perp}^{\nu}} f_{Vq}) - 3(1-2f_{Vq}) \right) \tilde{a}_q^{\mu} + 2T(\partial_{q_{\perp}^{\nu}} \tilde{a}_q^{\nu}) u^{\mu} - (1-2f_{Vq}) q_{\perp}^{\nu} \partial_{q_{\perp}^{\nu}} \tilde{a}_q^{\mu} \right\} \right], \quad (4.24b)$$

where we further retain the terms up to $\mathcal{O}(1/m)$. It turns out that the orientation of spin for heavy quarks is fixed yet the “spin” (axial-charge) density characterized by f_{Aq} undergoes the diffusive process same as the vector-charge density led by f_{Vq} when $m \rightarrow \infty$. Nonetheless, the modification upon the spin orientation by, e.g., the fluid velocity could emerge at higher orders suppressed by the mass of heavy quarks. Note that the Compton scattering, neglected in this work, also give rise to $1/m$ corrections, that, however, do not come with the logarithm enhancement $\sim \log(1/g_c)$. Therefore, the above $1/m$ corrections provide the consistent results within the leading-log approximation.

4.3.3 Consistency checks

In thermal equilibrium, the vector-charge distribution function takes the Fermi-Dirac form $f_{Vq} = 1/(e^{(E_q - \mu)/T} + 1)$ such that

$$\begin{aligned} \partial_{q_{\perp}^{\beta}} f_{qV} &= f_{Vq} (1 - f_{Vq}) \frac{q_{\perp\beta}}{E_q T}, \\ \partial_{q_{\perp}^{\alpha}} \partial_{q_{\perp}^{\beta}} f_{qV} &= f_{Vq} (1 - 3f_{Vq} + 2f_{Vq}^2) \frac{q_{\perp\alpha} q_{\perp\beta}}{E_q^2 T^2} + \frac{f_{Vq} (1 - f_{Vq})}{E_q T} \left(\Theta_{\alpha\beta} + \frac{q_{\perp\alpha} q_{\perp\beta}}{E_q^2} \right), \end{aligned} \quad (4.25)$$

and hence

$$\begin{aligned} \hat{q}_{\perp}^{\beta} \partial_{q_{\perp}^{\beta}} f_{qV} &= -f_{Vq} (1 - f_{Vq}) \frac{|\mathbf{q}|}{E_q T}, \\ \eta^{\alpha\beta} \partial_{q_{\perp}^{\alpha}} \partial_{q_{\perp}^{\beta}} f_{qV} &= -f_{Vq} (1 - 3f_{Vq} + 2f_{Vq}^2) \frac{|\mathbf{q}|^2}{E_q^2 T^2} + \frac{f_{Vq} (1 - f_{Vq})}{E_q T} \left(3 - \frac{|\mathbf{q}|^2}{E_q^2} \right), \\ \hat{q}_{\perp}^{\alpha} \hat{q}_{\perp}^{\beta} \partial_{q_{\perp}^{\alpha}} \partial_{q_{\perp}^{\beta}} f_{qV} &= f_{Vq} (1 - 3f_{Vq} + 2f_{Vq}^2) \frac{|\mathbf{q}|^2}{E_q^2 T^2} - \frac{m^2 f_{Vq} (1 - f_{Vq})}{E_q^3 T}. \end{aligned} \quad (4.26)$$

Using eqs. (4.25) and (4.26), one can explicitly show that the Fermi-Dirac distribution satisfies the SKE in eq. (4.20).

As for the AKE, although each coefficient $\hat{\mathcal{Q}}_{cl}^{(i)}$ takes a complicated form, we can make a cross check with the SKE in the massless limit. Generically, to consider the spin diffusion

for massless or light quarks, it is inevitable to further incorporate the gluon Compton scattering. Nevertheless, taking the massless limit here is just to scrutinize the consistency of our results. In the massless limit, $\tilde{a}_q^\mu = q^\mu f_A$ and thus

$$\begin{aligned}\partial_{q_\perp^\beta} \tilde{a}_q^\mu &= (\Theta_\beta^\mu - u^\mu \hat{q}_{\perp\beta}) f_A + q^\mu \partial_{q_\perp^\beta} f_A, & \partial_{q_\perp^\beta} \tilde{a}_q^\beta &= 3f_A + q_\perp^\beta \partial_{q_\perp^\beta} f_A, \\ \hat{q}_\perp^\beta \partial_{q_\perp^\beta} \tilde{a}_q^\mu &= (\hat{q}_\perp^\mu + u^\mu) f_A + q^\mu \hat{q}_\perp^\beta \partial_{q_\perp^\beta} f_A, \\ \partial_{q_\perp^\alpha} \partial_{q_\perp^\beta} \tilde{a}_q^\mu &= (\Theta_\beta^\mu - u^\mu \hat{q}_{\perp\beta}) \partial_{q_\perp^\alpha} f_A + (\Theta_\alpha^\mu - u^\mu \hat{q}_{\perp\alpha}) \partial_{q_\perp^\beta} f_A + q^\mu \partial_{q_\perp^\alpha} \partial_{q_\perp^\beta} f_A - \frac{u^\mu}{|\mathbf{q}|} (\Theta_{\alpha\beta} + \hat{q}_{\perp\alpha} \hat{q}_{\perp\beta}) f_A, \\ \partial_{q_\perp^\alpha} \partial_{q_\perp^\beta} \tilde{a}_q^\mu &= 2(\partial_{q_\perp^\mu} - u^\mu \hat{q}_\perp^\beta \partial_{q_\perp^\beta}) f_A + q^\mu \partial_{q_\perp^\beta} \partial_{q_\perp^\alpha} f_A - \frac{2u^\mu}{|\mathbf{q}|} f_A,\end{aligned}\quad (4.27)$$

where $\partial_{q_\perp^\alpha} \hat{q}_{\perp\beta} = (\Theta_{\alpha\beta} + \hat{q}_{\perp\alpha} \hat{q}_{\perp\beta})/|\mathbf{q}|$. One can show that eq. (4.22) then reduces to

$$\begin{aligned}0 &= q^\mu \delta(q^2) \left[q \cdot \partial f_{Aq} - \kappa_{\text{LL}} \left(2f_{Aq} (1 - 2f_{Vq} - q_\perp^\rho \partial_{q_\perp^\rho} f_{Vq}) \right. \right. \\ &\quad \left. \left. + (1 - 2f_{Vq}) q_\perp^\rho \partial_{q_\perp^\rho} f_{Aq} - |\mathbf{q}| T \eta^{\rho\nu} \partial_{q_\perp^\rho} \partial_{q_\perp^\nu} f_{Aq} \right) \right].\end{aligned}\quad (4.28)$$

We may check the consistency with the SKE in the massless limit by dropping the nonlinear terms since the f_A contributions are ignored in the SKE based on our power counting, which could cause discrepancies from the chirality-mixing terms. By further taking $f_{Vq} = (f_{Rq} + f_{Lq})/2$ and $f_{Aq} = f_{Rq} - f_{Lq}$, one finds that eq. (4.28) is consistent with the SKE in eq. (4.20) in the massless limit. Both the linearized eqs. (4.28) and (4.20) result in

$$0 = \delta(q^2 - m^2) \left[q \cdot \partial - \kappa_{\text{LL}} \left(2 + q_\perp^\beta \partial_{q_\perp^\beta} - |\mathbf{q}| T \eta^{\alpha\beta} \partial_{q_\perp^\alpha} \partial_{q_\perp^\beta} \right) \right] f_{R/Lq}.\quad (4.29)$$

Such a remarkable check should support the correctness of eq. (4.22). The check is also performed in ref. [54].

5 Concluding remarks and outlook

In this paper, we have derived the effective axial kinetic theory with background fields and collisions in the cases when the vector charge is more dominant than the axial charge (or more precisely the spin current) as natural conditions in most of physical systems. It is found that the AKE as a kinetic equation dictating the spin transport not only embraces the spin-diffusion term but also quantum corrections responsible for spin polarization, which reveals nontrivial entanglement of vector/axial-charge and spin transport through collisions. Our SKE and AKT work for an arbitrary mass and they reproduce the CKT in the massless limit. For the case of massive quarks, we have shown how our formalism reproduces the spin-diffusion term up to the leading-logarithmic order in a weakly coupled QGP and the leading of \hbar expansion.

Here we further summarize the detailed achievements in a short list:

1. In our power counting, the leading-order SKE remains the same as a classical Boltzmann equation, whereas the AKE governing the dynamics of spin polarization characterized by an axial-vector component \mathcal{A}^μ in Wigner functions can be written as

$\square^{(n)}\mathcal{A}^\mu = \hat{\mathcal{C}}_{\text{cl}}^\mu + \hbar\hat{\mathcal{C}}_{\text{Q}}^{(n)\mu}$. Here $\square^{(n)}\mathcal{A}^\mu$ denotes the free-streaming AKE, while $\hat{\mathcal{C}}_{\text{cl}}^\mu$ and $\hbar\hat{\mathcal{C}}_{\text{Q}}^{(n)\mu}$ correspond to the “classical” and “quantum” parts of collisions. Such separation is explicitly shown in a generic spacetime-dependent frame and in the rest frame for massive fermions as well.

2. It is found $\hat{\mathcal{C}}_{\text{cl}}^\mu$ and $\hbar\hat{\mathcal{C}}_{\text{Q}}^{(n)\mu}$ are proportional to f_A and f_V , respectively. Consequently, $\hat{\mathcal{C}}_{\text{cl}}^\mu$ serves as a spin-diffusion term, which vanishes when $f_A = 0$. On the contrary, the “quantum” correction $\hbar\hat{\mathcal{C}}_{\text{Q}}^{(n)\mu}$, which survives when $f_A = 0$, is dubbed as the spin-polarization term and responsible for polarizing spin via the intertwined dynamics of vector-charge transport due to spin-orbit interaction.
3. Similar to the free-streaming case [51], for a generic spacetime-dependent frame, $\hbar\hat{\mathcal{C}}_{\text{Q}}^{(n)\mu}$ can be separated into the term proportional to the four momentum and to the mass, which establishes a smooth connection to the CKT with collisions [30, 31] and manifests spin enslavement by chirality in the massless limit. Also, we present the simplified versions of the AKE in eq. (3.9) [or eq. (3.17)] and eq. (3.12) suitable for tracking the spin polarization of heavy and light quarks in QGP, respectively.
4. The spin diffusion term $\hat{\mathcal{C}}_{\text{cl}}^\mu$ for massive quarks in weakly-coupled QGP is obtained up to the leading logarithmic order, which incorporates nonlinear terms in distribution functions as a consequence of quantum statistics for fermions. It turns out that even the spin diffusion is affected by entangled dynamics between f_V and f_A as opposed to the previous study with only the linearized collision terms in distribution functions [54].

Although we have explicitly evaluated the spin-diffusion term for massive quarks, which are sufficiently heavy for dropping the gluon Compton scattering, the quantum correction could be calculated in a similar fashion as the follow-up work. Moreover, when considering the spin transport for light quarks, it is inevitable to further incorporate the Compton scattering even for just spin diffusion. On the other hand, as already mentioned in the context, in rotating QGP, it is expected that both light quarks and gluons are polarized, which should be involved as quantum corrections in the self-energies. Recently, there have been some relevant studies for the quantum corrections upon polarized photons [71, 72]. Albeit the validity of our formalism is held even in the presence of such corrections, it could be challenging to systematically include the polarization of scattered gluons or even other quarks and to obtain an analytic form of the collision term. Nevertheless, to understand the dynamical evolution of the spin polarization for peculiarly strange quarks associated with the local polarization of Λ hyperons, it will be essential to carry out the aforementioned studies in the future.

On the other hand, our formalism is rather generic, which may have potential applications not only in heavy ion collisions but also other physical systems. For instance, it is proposed in refs. [73–75] that the electron and neutrino transport with anomalous effects led by chirality imbalance (axial charge) could influence the macroscopic hydrodynamic evolution of matter in core collapse supernovae. However, the chirality imbalance of elec-

trons produced by the electron capture process may be compensated by elastic electron scattering with the effect of nonzero electron mass [76–78]. Our formalism could be applied to track the axial-charge evolution in such a scenario. In such a case, eq. (3.12) will be useful for capturing the small-electron-mass effect.

Acknowledgments

Y.H. was partially supported by Japan Society of Promotion of Science (JSPS), Grants-in-Aid for Scientific Research(KAKENHI) Grants No. 15H03652, 16K17716, and 17H06462. Y.H. was also partially supported by RIKEN iTHES Project and iTHEMS Program. K.H. is supported in part by Japan Society of Promotion of Science (JSPS), Grants-in-Aid for Scientific Research (KAKENHI) under grant No. 20K03948 and also in part by Yukawa International Program for Quark-hadron Sciences (YIPQS). D.-L.Y. is supported by Keio Institute of Pure and Applied Sciences (KiPAS) project in Keio University and JSPS KAKENHI grant No. 20K14470. The authors also would like to thank participants of the molecular-type Yukawa Institute for Theoretical Physics (YITP) workshop “Quantum kinetic theories in magnetic and vortical fields (YITP-T-19-06)” for fruitful and useful discussions.

A Derivation of the master equations

A.1 Spinor decomposition

Based on the spinor decomposition of the propagators (2.7) and the self-energies (2.8), we perform the decomposition of the Kadanoff-Baym equations (2.4). One may use some useful relations such as

$$\gamma^\mu \gamma^\alpha \gamma^\beta = \eta^{\mu\alpha} \gamma^\beta - \eta^{\mu\beta} \gamma^\alpha + \eta^{\alpha\beta} \gamma^\mu - i\epsilon^{\mu\alpha\beta\lambda} \gamma^5 \gamma_\lambda, \quad \gamma^5 \sigma_{\mu\nu} = \frac{i}{2} \epsilon_{\mu\nu\kappa\sigma} \sigma^{\kappa\sigma}.$$

Here, $\epsilon_{\mu\nu\rho\sigma}$ is the totally antisymmetric tensor with $\epsilon_{0123} = -1$. The Kadanoff-Baym equations (2.4) contain the commutators $[\gamma^\mu, \nabla_\mu S^<]$ and $[(\not{M} - m), S^<]$ and their counterparts with the anticommutation relations. They can be decomposed with the following relations

$$[\gamma^\mu, S^<] = -2\mathcal{A}^\mu \gamma^5 + 2i\mathcal{S}^\mu{}_\alpha \gamma^\alpha - 2i\mathcal{P} \gamma^5 \gamma^\mu - 2i\mathcal{V}_\nu \sigma^{\mu\nu}, \quad (\text{A.1a})$$

$$\{\gamma^\mu, S^<\} = 2\mathcal{V}^\mu + 2\mathcal{S} \gamma^\mu + \epsilon^\mu{}_{\nu\rho\alpha} \mathcal{S}^{\nu\rho} \gamma^5 \gamma^\alpha - \epsilon^\mu{}_{\nu\alpha\beta} \mathcal{A}^\nu \sigma^{\alpha\beta}. \quad (\text{A.1b})$$

Based on the decompositions above, we now have

$$\frac{i}{2} [\gamma^\mu, \nabla_\mu S^<] = -i\nabla_\mu \mathcal{A}^\mu \gamma^5 - \nabla^\nu \mathcal{S}_{\nu\mu} \gamma^\mu + \nabla_\mu \mathcal{P} \gamma^5 \gamma^\mu + \nabla_\mu \mathcal{V}_\nu \sigma^{\mu\nu}, \quad (\text{A.1c})$$

$$\frac{i}{2} \{\gamma^\mu, \nabla_\mu S^<\} = i\nabla^\mu \mathcal{V}_\mu + i\nabla_\mu \mathcal{S} \gamma^\mu - \frac{i}{2} \epsilon^{\mu\nu\rho\sigma} \nabla_\sigma \mathcal{S}_{\nu\rho} \gamma^5 \gamma^\mu - \frac{i}{2} \epsilon_{\mu\nu\rho\sigma} \nabla^\rho \mathcal{A}^\sigma \sigma^{\mu\nu}, \quad (\text{A.1d})$$

$$[(\not{M} - m), S^<] = -2\Pi^\nu \mathcal{A}_\nu \gamma^5 + 2i\Pi^\nu \mathcal{S}_{\nu\mu} \gamma^\mu - 2i\Pi_\mu \mathcal{P} \gamma^5 \gamma^\mu - 2i\Pi_\mu \mathcal{V}_\nu \sigma^{\mu\nu}, \quad (\text{A.1e})$$

$$\begin{aligned} \{(\not{M} - m), S^<\} &= 2(\Pi^\mu \mathcal{V}_\mu - m\mathcal{S}) - 2im\mathcal{P} \gamma^5 + 2(\Pi_\mu \mathcal{S} - m\mathcal{V}_\mu) \gamma^\mu \\ &\quad - (\epsilon^{\mu\nu\rho\sigma} \Pi_\sigma \mathcal{S}_{\nu\rho} + 2m\mathcal{A}^\mu) \gamma^5 \gamma_\mu - (m\mathcal{S}_{\mu\nu} + \epsilon_{\mu\nu\rho\sigma} \Pi^\rho \mathcal{A}^\sigma) \sigma^{\mu\nu}. \end{aligned} \quad (\text{A.1f})$$

Assuming the following decomposition $G(q, X) = \hat{K}^\mu G_\mu(q, X)$ and $F(q, X) = \hat{Q}^\mu F_\mu(q, X)$ with \hat{K}^μ and \hat{Q}^μ being arbitrary matrices, it is found

$$\begin{aligned} \{G, F\}_\star &= \frac{1}{2} \{\hat{K}^\mu, \hat{Q}^\nu\} \{G_\mu(q, X), F_\nu(q, X)\}_\star + \frac{1}{2} [\hat{K}^\mu, \hat{Q}^\nu] [G_\mu(q, X), F_\nu(q, X)]_\star, \\ [G, F]_\star &= \frac{1}{2} [\hat{K}^\mu, \hat{Q}^\nu] \{G_\mu(q, X), F_\nu(q, X)\}_\star + \frac{1}{2} \{\hat{K}^\mu, \hat{Q}^\nu\} [G_\mu(q, X), F_\nu(q, X)]_\star. \end{aligned} \quad (\text{A.2})$$

Accordingly, the self-energy parts can be decomposed in the same and straightforward way as

$$\begin{aligned} [\Sigma^>, S^<]_\star &= i(\{\bar{\Sigma}_{V\alpha}, \mathcal{A}^\alpha\}_\star - \{\bar{\Sigma}_{A\alpha}, \mathcal{V}^\alpha\}_\star) i\gamma^5 \\ &\quad + i(\{\bar{\Sigma}_P, \mathcal{A}_\alpha\}_\star + \{\bar{\Sigma}_V^\mu, \mathcal{S}_{\mu\alpha}\}_\star - \{\bar{\Sigma}_{A\alpha}, \mathcal{P}\}_\star + \{\bar{\Sigma}_{T\alpha\beta}, \mathcal{V}^\beta\}_\star) \gamma^\alpha \\ &\quad + i(\{\bar{\Sigma}_P, \mathcal{V}_\alpha\}_\star - \{\bar{\Sigma}_{V\alpha}, \mathcal{P}\}_\star + \{\bar{\Sigma}_A^\mu, \mathcal{S}_{\mu\alpha}\}_\star + \{\bar{\Sigma}_{T\alpha\beta}, \mathcal{A}^\beta\}_\star) \gamma^5 \gamma^\alpha \\ &\quad + i(-\{\bar{\Sigma}_{V[\alpha}, \mathcal{V}_{\beta]}\}_\star + \{\bar{\Sigma}_{A[\alpha}, \mathcal{A}_{\beta]}\}_\star - \{\bar{\Sigma}_{T\mu[\alpha}, \mathcal{S}^{\mu}_{\beta]}\}_\star) \frac{\sigma^{\alpha\beta}}{2} \\ &\quad + [\bar{\Sigma}_S, \mathcal{S}]_\star - [\bar{\Sigma}_P, \mathcal{P}]_\star + [\bar{\Sigma}_{V\mu}, \mathcal{V}^\mu]_\star - [\bar{\Sigma}_{A\mu}, \mathcal{A}^\mu]_\star + \frac{1}{2} [\bar{\Sigma}_{T\mu\nu}, \mathcal{S}^{\mu\nu}]_\star \\ &\quad + \left([\bar{\Sigma}_S, \mathcal{P}]_\star + [\bar{\Sigma}_P, \mathcal{S}]_\star + \frac{1}{4} \epsilon_{\mu\nu\alpha\beta} [\bar{\Sigma}_T^{\mu\nu}, \mathcal{S}^{\alpha\beta}]_\star \right) i\gamma^5 \\ &\quad + \left([\bar{\Sigma}_S, \mathcal{V}_\alpha]_\star + [\bar{\Sigma}_{V\alpha}, \mathcal{S}]_\star + \frac{1}{2} \epsilon_{\mu\nu\lambda\alpha} ([\bar{\Sigma}_A^\mu, \mathcal{S}^{\nu\lambda}]_\star + [\bar{\Sigma}_T^{\mu\nu}, \mathcal{A}^\lambda]_\star) \right) \gamma^\alpha \\ &\quad + \left([\bar{\Sigma}_S, \mathcal{A}_\alpha]_\star + [\bar{\Sigma}_{A\alpha}, \mathcal{S}]_\star + \frac{1}{2} \epsilon_{\mu\nu\lambda\alpha} ([\bar{\Sigma}_V^\mu, \mathcal{S}^{\nu\lambda}]_\star + [\bar{\Sigma}_T^{\mu\nu}, \mathcal{V}^\lambda]_\star) \right) \gamma^5 \gamma^\alpha \\ &\quad + \left([\bar{\Sigma}_S, \mathcal{S}_{\alpha\beta}]_\star + [\bar{\Sigma}_{T\alpha\beta}, \mathcal{S}]_\star + \epsilon_{\mu\nu\alpha\beta} (-[\bar{\Sigma}_V^\mu, \mathcal{A}^\nu]_\star + [\bar{\Sigma}_A^\mu, \mathcal{V}^\nu]_\star) \right. \\ &\quad \left. - \frac{1}{2} \epsilon_{\mu\nu\alpha\beta} ([\bar{\Sigma}_P, \mathcal{S}^{\mu\nu}]_\star + [\bar{\Sigma}_T^{\mu\nu}, \mathcal{P}]_\star) \right) \frac{\sigma^{\alpha\beta}}{2}, \end{aligned} \quad (\text{A.3})$$

and

$$\begin{aligned} \{\Sigma^>, S^<\}_\star &= \{\bar{\Sigma}_S, \mathcal{S}\}_\star - \{\bar{\Sigma}_P, \mathcal{P}\}_\star + \{\bar{\Sigma}_{V\mu}, \mathcal{V}^\mu\}_\star - \{\bar{\Sigma}_{A\mu}, \mathcal{A}^\mu\}_\star + \frac{1}{2} \{\bar{\Sigma}_{T\mu\nu}, \mathcal{S}^{\mu\nu}\}_\star \\ &\quad + \left(\{\bar{\Sigma}_S, \mathcal{P}\}_\star + \{\bar{\Sigma}_P, \mathcal{S}\}_\star + \frac{1}{4} \epsilon_{\mu\nu\alpha\beta} \{\bar{\Sigma}_T^{\mu\nu}, \mathcal{S}^{\alpha\beta}\}_\star \right) i\gamma^5 \\ &\quad + \left(\{\bar{\Sigma}_S, \mathcal{V}_\alpha\}_\star + \{\bar{\Sigma}_{V\alpha}, \mathcal{S}\}_\star + \frac{1}{2} \epsilon_{\mu\nu\lambda\alpha} (\{\bar{\Sigma}_A^\mu, \mathcal{S}^{\nu\lambda}\}_\star + \{\bar{\Sigma}_T^{\mu\nu}, \mathcal{A}^\lambda\}_\star) \right) \gamma^\alpha \\ &\quad + \left(\{\bar{\Sigma}_S, \mathcal{A}_\alpha\}_\star + \{\bar{\Sigma}_{A\alpha}, \mathcal{S}\}_\star + \frac{1}{2} \epsilon_{\mu\nu\lambda\alpha} (\{\bar{\Sigma}_V^\mu, \mathcal{S}^{\nu\lambda}\}_\star + \{\bar{\Sigma}_T^{\mu\nu}, \mathcal{V}^\lambda\}_\star) \right) \gamma^5 \gamma^\alpha \\ &\quad + \left(\{\bar{\Sigma}_S, \mathcal{S}_{\alpha\beta}\}_\star + \{\bar{\Sigma}_{T\alpha\beta}, \mathcal{S}\}_\star + \epsilon_{\mu\nu\alpha\beta} (-\{\bar{\Sigma}_V^\mu, \mathcal{A}^\nu\}_\star + \{\bar{\Sigma}_A^\mu, \mathcal{V}^\nu\}_\star) \right. \\ &\quad \left. - \frac{1}{2} \epsilon_{\mu\nu\alpha\beta} (\{\bar{\Sigma}_P, \mathcal{S}^{\mu\nu}\}_\star + \{\bar{\Sigma}_T^{\mu\nu}, \mathcal{P}\}_\star) \right) \frac{1}{2} \sigma^{\alpha\beta} \\ &\quad + i([\bar{\Sigma}_{V\alpha}, \mathcal{A}^\alpha]_\star - [\bar{\Sigma}_{A\alpha}, \mathcal{V}^\alpha]_\star) i\gamma^5 \\ &\quad + i([\bar{\Sigma}_P, \mathcal{A}_\alpha]_\star + [\bar{\Sigma}_V^\mu, \mathcal{S}_{\mu\alpha}]_\star - [\bar{\Sigma}_{A\alpha}, \mathcal{P}]_\star + [\bar{\Sigma}_{T\alpha\beta}, \mathcal{V}^\beta]_\star) \gamma^\alpha \\ &\quad + i([\bar{\Sigma}_P, \mathcal{V}_\alpha]_\star + [\bar{\Sigma}_A^\mu, \mathcal{S}_{\mu\alpha}]_\star - [\bar{\Sigma}_{V\alpha}, \mathcal{P}]_\star + [\bar{\Sigma}_{T\alpha\beta}, \mathcal{A}^\beta]_\star) \gamma^5 \gamma^\alpha \\ &\quad + i(-[\bar{\Sigma}_{V[\alpha}, \mathcal{V}_{\beta]}\}_\star + [\bar{\Sigma}_{A[\alpha}, \mathcal{A}_{\beta]}\}_\star - [\bar{\Sigma}_{T\mu[\alpha}, \mathcal{S}^{\mu}_{\beta]}\}_\star) \frac{1}{2} \sigma^{\alpha\beta}, \end{aligned} \quad (\text{A.4})$$

where we defined the antisymmetrization $T_{[\mu\nu]} = T_{\mu\nu} - T_{\nu\mu}$. Plugging those decompositions back to the Kadanoff-Baym equations (2.4), we find

$$0 = \{(\not{M} - m), S^<\} + \frac{i\hbar}{2} \left([\gamma^\mu, \nabla_\mu S^<] - [\Sigma^<, S^>]_\star + [\Sigma^>, S^<]_\star \right) \quad (\text{A.5})$$

$$= \mathcal{K}_S + \mathcal{K}_5 i\gamma^5 + \mathcal{K}_{V\mu} \gamma^\mu + \mathcal{K}_{5\mu} \gamma^\mu \gamma^5 + \mathcal{K}_{T\mu\nu} \frac{1}{2} \sigma^{\mu\nu},$$

$$0 = [(\not{M} - m), S^<] + \frac{i\hbar}{2} \left(\{\gamma^\mu, \nabla_\mu S^<\} - \{\Sigma^<, S^>\}_\star + \{\Sigma^>, S^<\}_\star \right) \quad (\text{A.6})$$

$$= \mathcal{Q}_S + \mathcal{Q}_5 i\gamma^5 + \mathcal{Q}_{V\mu} \gamma^\mu + \mathcal{Q}_{5\mu} \gamma^\mu \gamma^5 + \mathcal{Q}_{T\mu\nu} \frac{1}{2} \sigma^{\mu\nu},$$

where

$$\mathcal{K}_S = 2\Pi^\mu \mathcal{V}_\mu - 2m\mathcal{S} + \frac{i\hbar}{2} \left([\widehat{\Sigma}_S, \mathcal{S}]_\star - [\widehat{\Sigma}_P, \mathcal{P}]_\star + [\widehat{\Sigma}_{V\mu}, \mathcal{V}^\mu]_\star - [\widehat{\Sigma}_{A\mu}, \mathcal{A}^\mu]_\star + \frac{1}{2} [\widehat{\Sigma}_{T\mu\nu}, \mathcal{S}^{\mu\nu}]_\star \right), \quad (\text{A.7})$$

$$\mathcal{K}_5 = -2m\mathcal{P} - \hbar\tilde{\mathcal{D}}\mathcal{A}^\mu + \frac{\hbar}{2} \{\widehat{\Sigma}_{A\mu}, \mathcal{V}^\mu\}_\star + \frac{i\hbar}{2} \left([\widehat{\Sigma}_S, \mathcal{P}]_\star + [\widehat{\Sigma}_P, \mathcal{S}]_\star + \frac{1}{4} \epsilon_{\mu\nu\alpha\beta} [\widehat{\Sigma}_T^{\mu\nu}, \mathcal{S}^{\alpha\beta}]_\star \right), \quad (\text{A.8})$$

$$\mathcal{K}_{V\alpha} = 2\Pi_\alpha \mathcal{S} - 2m\mathcal{V}_\alpha - \hbar\tilde{\mathcal{D}}^\nu \mathcal{S}_{\nu\alpha} - \frac{\hbar}{2} \left(\{\widehat{\Sigma}_P, \mathcal{A}_\alpha\}_\star - \{\widehat{\Sigma}_{A\alpha}, \mathcal{P}\}_\star + \{\widehat{\Sigma}_{T\alpha\mu}, \mathcal{V}^\mu\}_\star \right) + \frac{i\hbar}{2} \left([\widehat{\Sigma}_S, \mathcal{V}_\alpha]_\star + [\widehat{\Sigma}_{V\alpha}, \mathcal{S}]_\star + \frac{1}{2} \epsilon_{\mu\nu\lambda\alpha} ([\widehat{\Sigma}_A^\mu, \mathcal{S}^{\nu\lambda}]_\star + [\widehat{\Sigma}_T^{\mu\nu}, \mathcal{A}^\lambda]_\star) \right), \quad (\text{A.9})$$

$$\mathcal{K}_{5\alpha} = -\epsilon_{\alpha\nu\rho\sigma} \Pi^\sigma \mathcal{S}^{\nu\rho} - 2m\mathcal{A}_\alpha + \hbar\tilde{\mathcal{D}}_\alpha \mathcal{P} - \frac{\hbar}{2} \left(\{\widehat{\Sigma}_P, \mathcal{V}_\alpha\}_\star + \{\widehat{\Sigma}_A^\mu, \mathcal{S}_{\mu\alpha}\}_\star + \{\widehat{\Sigma}_{T\alpha\mu}, \mathcal{A}^\mu\}_\star \right) + \frac{i\hbar}{2} \left([\widehat{\Sigma}_S, \mathcal{A}_\alpha]_\star + [\widehat{\Sigma}_{A\alpha}, \mathcal{S}]_\star + \frac{1}{2} \epsilon_{\mu\nu\lambda\alpha} ([\widehat{\Sigma}_V^\mu, \mathcal{S}^{\nu\lambda}]_\star + [\widehat{\Sigma}_T^{\mu\nu}, \mathcal{V}^\lambda]_\star) \right), \quad (\text{A.10})$$

$$\mathcal{K}_{T\alpha\beta} = -2m\mathcal{S}_{\alpha\beta} - 2\epsilon_{\alpha\beta\rho\sigma} \Pi^\rho \mathcal{A}^\sigma + \hbar\tilde{\mathcal{D}}_{[\alpha} \mathcal{V}_{\beta]} - \frac{\hbar}{2} \left(\{\widehat{\Sigma}_{A[\alpha}, \mathcal{A}_{\beta]}\}_\star - \{\widehat{\Sigma}_{T\mu[\alpha}, \mathcal{S}^\mu_{\beta]}\}_\star \right) + \frac{i\hbar}{2} \left([\widehat{\Sigma}_S, \mathcal{S}_{\alpha\beta}]_\star + [\widehat{\Sigma}_{T\alpha\beta}, \mathcal{S}]_\star + \epsilon_{\mu\nu\alpha\beta} (-[\widehat{\Sigma}_V^\mu, \mathcal{A}^\nu]_\star + [\widehat{\Sigma}_A^\mu, \mathcal{V}^\nu]_\star) - \frac{1}{2} \epsilon_{\mu\nu\alpha\beta} ([\widehat{\Sigma}_P, \mathcal{S}^{\mu\nu}]_\star + [\widehat{\Sigma}_T^{\mu\nu}, \mathcal{P}]_\star) \right), \quad (\text{A.11})$$

and

$$\mathcal{Q}_S = i\hbar\tilde{\mathcal{D}}_\mu \mathcal{V}^\mu + \frac{i\hbar}{2} \left(\{\widehat{\Sigma}_S, \mathcal{S}\}_\star - \{\widehat{\Sigma}_P, \mathcal{P}\}_\star - \{\widehat{\Sigma}_{A\mu}, \mathcal{A}^\mu\}_\star + \frac{1}{2} \{\widehat{\Sigma}_{T\mu\nu}, \mathcal{S}^{\mu\nu}\}_\star \right), \quad (\text{A.12})$$

$$\mathcal{Q}_5 = 2i\Pi^\nu \mathcal{A}_\nu + \frac{i\hbar}{2} \left(\{\widehat{\Sigma}_S, \mathcal{P}\}_\star + \{\widehat{\Sigma}_P, \mathcal{S}\}_\star + \frac{1}{4} \epsilon_{\mu\nu\alpha\beta} \{\widehat{\Sigma}_T^{\mu\nu}, \mathcal{S}^{\alpha\beta}\}_\star \right) - \frac{\hbar}{2} ([\widehat{\Sigma}_{V\alpha}, \mathcal{A}^\alpha]_\star - [\widehat{\Sigma}_{A\alpha}, \mathcal{V}^\alpha]_\star), \quad (\text{A.13})$$

$$\mathcal{Q}_{V\alpha} = 2i\Pi^\nu \mathcal{S}_{\nu\alpha} + i\hbar\tilde{\mathcal{D}}_\alpha \mathcal{S} + \frac{i\hbar}{2} \left(\{\widehat{\Sigma}_S, \mathcal{V}_\alpha\}_\star + \frac{1}{2} \epsilon_{\mu\nu\lambda\alpha} (\{\widehat{\Sigma}_A^\mu, \mathcal{S}^{\nu\lambda}\}_\star + \{\widehat{\Sigma}_T^{\mu\nu}, \mathcal{A}^\lambda\}_\star) \right) - \frac{\hbar}{2} ([\widehat{\Sigma}_P, \mathcal{A}_\alpha]_\star + [\widehat{\Sigma}_V^\mu, \mathcal{S}_{\mu\alpha}]_\star - [\widehat{\Sigma}_{A\alpha}, \mathcal{P}]_\star + [\widehat{\Sigma}_{T\alpha\beta}, \mathcal{V}^\beta]_\star), \quad (\text{A.14})$$

$$\mathcal{Q}_{5\alpha} = -2i\Pi_\alpha \mathcal{P} - \frac{i\hbar}{2} \epsilon_{\alpha\nu\rho\sigma} \tilde{\mathcal{D}}^\nu \mathcal{S}^{\rho\sigma} + \frac{i\hbar}{2} \left(\{\widehat{\Sigma}_S, \mathcal{A}_\alpha\}_\star + \{\widehat{\Sigma}_{A\alpha}, \mathcal{S}\}_\star - \frac{1}{2} \epsilon_{\alpha\nu\rho\sigma} \{\widehat{\Sigma}_T^{\nu\rho}, \mathcal{V}^\sigma\}_\star \right) - \frac{\hbar}{2} ([\widehat{\Sigma}_P, \mathcal{V}_\alpha]_\star + [\widehat{\Sigma}_A^\mu, \mathcal{S}_{\mu\alpha}]_\star - [\widehat{\Sigma}_{V\alpha}, \mathcal{P}]_\star + [\widehat{\Sigma}_{T\alpha\mu}, \mathcal{A}^\mu]_\star), \quad (\text{A.15})$$

$$\mathcal{Q}_{T\alpha\beta} = -2i\Pi_{[\alpha} \mathcal{V}_{\beta]} - i\hbar\epsilon_{\alpha\beta\mu\nu} \tilde{\mathcal{D}}^\mu \mathcal{A}^\nu + \frac{i\hbar}{2} \left(\{\widehat{\Sigma}_S, \mathcal{S}_{\alpha\beta}\}_\star + \{\widehat{\Sigma}_{T\alpha\beta}, \mathcal{S}\}_\star + \epsilon_{\alpha\beta\mu\nu} \{\widehat{\Sigma}_A^\mu, \mathcal{V}^\nu\}_\star - \frac{1}{2} \epsilon_{\alpha\beta\mu\nu} (\{\widehat{\Sigma}_P, \mathcal{S}^{\mu\nu}\}_\star + \{\widehat{\Sigma}_T^{\mu\nu}, \mathcal{P}\}_\star) \right) - \frac{\hbar}{2} (-[\widehat{\Sigma}_{V[\alpha}, \mathcal{V}_{\beta]}\}_\star + [\widehat{\Sigma}_{A[\alpha}, \mathcal{A}_{\beta]}\}_\star - [\widehat{\Sigma}_{T\mu[\alpha}, \mathcal{S}^\mu_{\beta]}\}_\star). \quad (\text{A.16})$$

Here we introduce a shorthand notation $\widehat{XY} = \bar{X}Y - X\bar{Y}$, where X and Y are the coefficients of the Clifford decomposition of the propagators and self-energies, e.g., $\{\widehat{\Sigma_{V\mu}}, \mathcal{V}_\nu\}_* = \{\bar{\Sigma}_{V\mu}, \mathcal{V}_\nu\}_* - \{\Sigma_{V\mu}, \bar{\mathcal{V}}_\nu\}_*$. We also introduced $\tilde{\mathcal{D}}_\mu \mathcal{M} = \nabla_\mu \mathcal{M} + \{\widehat{\Sigma_{V\mu}}, \mathcal{M}\}_*/2$. $\mathcal{K}_S = \mathcal{K}_5 = \mathcal{K}_{V\alpha} = \mathcal{K}_{5\alpha} = \mathcal{Q}_{T\alpha\beta} = \mathcal{Q}_S = \mathcal{Q}_5 = \mathcal{Q}_{V\alpha} = \mathcal{Q}_{5\alpha} = \mathcal{Q}_{T\alpha\beta} = 0$ give the set of full quantum master equations. Explicitly, they read

$$m\mathcal{S} = \Pi^\mu \mathcal{V}_\mu - \frac{\hbar^2}{4} \left[\widehat{\Sigma_S \mathcal{S}} - \widehat{\Sigma_P \mathcal{P}} + \widehat{\Sigma_{V\mu} \mathcal{V}^\mu} - \widehat{\Sigma_{A\mu} \mathcal{A}^\mu} + \frac{1}{2} \widehat{\Sigma_{T\mu\nu} \mathcal{S}^{\mu\nu}} \right]_{\text{P.B.}} + \mathcal{O}(\hbar^3), \quad (\text{A.17a})$$

$$m\mathcal{P} = -\frac{\hbar}{2} (\tilde{\mathcal{D}}_\mu \mathcal{A}^\mu - \widehat{\Sigma_{A\mu} \mathcal{V}^\mu}) - \frac{\hbar^2}{4} \left[\widehat{\Sigma_S \mathcal{P}} + \widehat{\Sigma_P \mathcal{S}} + \frac{1}{4} \epsilon_{\mu\nu\alpha\beta} \widehat{\Sigma_T^{\mu\nu} \mathcal{S}^{\alpha\beta}} \right]_{\text{P.B.}} + \mathcal{O}(\hbar^3), \quad (\text{A.17b})$$

$$\begin{aligned} & 2\Pi_\alpha \mathcal{S} - \hbar \tilde{\mathcal{D}}^\nu \mathcal{S}_{\nu\alpha} - 2m\mathcal{V}_\alpha - \hbar (\widehat{\Sigma_P \mathcal{A}_\alpha} - \widehat{\Sigma_{A\alpha} \mathcal{P}} - \widehat{\Sigma_{T\mu\alpha} \mathcal{V}^\mu}) \\ &= \frac{\hbar^2}{2} \left[\widehat{\Sigma_S \mathcal{V}_\alpha} + \widehat{\Sigma_{V\alpha} \mathcal{S}} + \frac{1}{2} \epsilon_{\mu\nu\lambda\alpha} (\widehat{\Sigma_A^\mu \mathcal{S}^{\nu\lambda}} + \widehat{\Sigma_T^{\mu\nu} \mathcal{A}^\lambda}) \right]_{\text{P.B.}} + \mathcal{O}(\hbar^3), \end{aligned} \quad (\text{A.17c})$$

$$\begin{aligned} & \hbar \tilde{\mathcal{D}}_\alpha \mathcal{P} - \epsilon_{\alpha\nu\rho\sigma} \Pi^\sigma \mathcal{S}^{\nu\rho} - 2m\mathcal{A}_\alpha - \hbar (\widehat{\Sigma_P \mathcal{V}_\alpha} + \widehat{\Sigma_A^\mu \mathcal{S}_{\mu\alpha}} - \widehat{\Sigma_{T\mu\alpha} \mathcal{A}^\mu}) \\ &= \frac{\hbar^2}{2} \left[\widehat{\Sigma_S \mathcal{A}_\alpha} + \widehat{\Sigma_{A\alpha} \mathcal{S}} + \frac{1}{2} \epsilon_{\mu\nu\lambda\alpha} (\widehat{\Sigma_V^\mu \mathcal{S}^{\nu\lambda}} + \widehat{\Sigma_T^{\mu\nu} \mathcal{V}^\lambda}) \right]_{\text{P.B.}} + \mathcal{O}(\hbar^3), \end{aligned} \quad (\text{A.17d})$$

$$\begin{aligned} & m\mathcal{S}_{\alpha\beta} + \epsilon_{\alpha\beta\rho\sigma} \Pi^\rho \mathcal{A}^\sigma - \frac{\hbar}{2} (\tilde{\mathcal{D}}_{[\alpha} \mathcal{V}_{\beta]} - \widehat{\Sigma_{A[\alpha} \mathcal{A}_{\beta]}} + \widehat{\Sigma_{T\mu[\alpha} \mathcal{S}^\mu_{\beta]}}) \\ &= -\frac{\hbar^2}{4} \left[\widehat{\Sigma_S \mathcal{S}_{\alpha\beta}} + \widehat{\Sigma_{T\alpha\beta} \mathcal{S}} - \epsilon_{\mu\nu\alpha\beta} \widehat{\Sigma_V^\mu \mathcal{A}^\nu} + \epsilon_{\mu\nu\alpha\beta} \widehat{\Sigma_A^\mu \mathcal{V}^\nu} - \frac{1}{2} \epsilon_{\mu\nu\alpha\beta} (\widehat{\Sigma_P \mathcal{S}^{\mu\nu}} + \widehat{\Sigma_T^{\mu\nu} \mathcal{P}}) \right]_{\text{P.B.}} + \mathcal{O}(\hbar^3), \end{aligned} \quad (\text{A.17e})$$

and

$$\tilde{\mathcal{D}}^\mu \mathcal{V}_\mu = -\widehat{\Sigma_S \mathcal{S}} + \widehat{\Sigma_P \mathcal{P}} + \widehat{\Sigma_{A\mu} \mathcal{A}^\mu} - \frac{1}{2} \widehat{\Sigma_{T\mu\nu} \mathcal{S}^{\mu\nu}} + \mathcal{O}(\hbar^3), \quad (\text{A.18a})$$

$$\begin{aligned} 2\Pi^\nu \mathcal{A}_\nu &= -\hbar \left(\widehat{\Sigma_S \mathcal{P}} + \widehat{\Sigma_P \mathcal{S}} + \frac{1}{4} \epsilon_{\mu\nu\alpha\beta} \widehat{\Sigma_T^{\mu\nu} \mathcal{S}^{\alpha\beta}} \right) - \frac{\hbar^2}{2} [-\widehat{\Sigma_{V\alpha} \mathcal{A}^\alpha} + \widehat{\Sigma_{A\alpha} \mathcal{V}^\alpha}]_{\text{P.B.}} + \mathcal{O}(\hbar^3), \\ & \quad (\text{A.18b}) \end{aligned}$$

$$\begin{aligned} & 2\Pi^\nu \mathcal{S}_{\nu\alpha} + \hbar \tilde{\mathcal{D}}_\alpha \mathcal{S} + \hbar \left(\widehat{\Sigma_S \mathcal{V}_\alpha} + \frac{1}{2} \epsilon_{\mu\nu\lambda\alpha} (\widehat{\Sigma_A^\mu \mathcal{S}^{\nu\lambda}} + \widehat{\Sigma_T^{\mu\nu} \mathcal{A}^\lambda}) \right) \\ &= \frac{\hbar^2}{2} [\widehat{\Sigma_P \mathcal{A}_\alpha} + \widehat{\Sigma_V^\mu \mathcal{S}_{\mu\alpha}} - \widehat{\Sigma_{A\alpha} \mathcal{P}} + \widehat{\Sigma_{T\alpha\beta} \mathcal{V}^\beta}]_{\text{P.B.}} + \mathcal{O}(\hbar^3), \end{aligned} \quad (\text{A.18c})$$

$$\begin{aligned} & 2\Pi_\alpha \mathcal{P} + \frac{\hbar}{2} \epsilon_{\alpha\nu\rho\sigma} (\tilde{\mathcal{D}}^\nu \mathcal{S}^{\rho\sigma} + \widehat{\Sigma_T^{\nu\rho} \mathcal{V}^\sigma}) - \hbar (\widehat{\Sigma_S \mathcal{A}_\alpha} + \widehat{\Sigma_{A\alpha} \mathcal{S}}) \\ &= 1 - \frac{\hbar^2}{2} [\widehat{\Sigma_P \mathcal{V}_\alpha} - \widehat{\Sigma_{V\alpha} \mathcal{P}} + \widehat{\Sigma_A^\mu \mathcal{S}_{\mu\alpha}} + \widehat{\Sigma_{T\alpha\beta} \mathcal{A}^\beta}]_{\text{P.B.}} + \mathcal{O}(\hbar^3), \end{aligned} \quad (\text{A.18d})$$

$$\begin{aligned} & \Pi_{[\alpha} \mathcal{V}_{\beta]} + \frac{\hbar}{2} \epsilon_{\alpha\beta\mu\nu} (\tilde{\mathcal{D}}^\mu \mathcal{A}^\nu - \widehat{\Sigma_A^\mu \mathcal{V}^\nu}) - \frac{\hbar}{2} (\widehat{\Sigma_S \mathcal{S}_{\alpha\beta}} + \widehat{\Sigma_{T\alpha\beta} \mathcal{S}}) + \frac{\hbar}{4} \epsilon_{\mu\nu\alpha\beta} (\widehat{\Sigma_P \mathcal{S}^{\mu\nu}} + \widehat{\Sigma_T^{\mu\nu} \mathcal{P}}) \\ &= \frac{\hbar^2}{4} [\widehat{\Sigma_{V[\alpha} \mathcal{V}_{\beta]}} - \widehat{\Sigma_{A[\alpha} \mathcal{A}_{\beta]}} + \widehat{\Sigma_{T\mu[\alpha} \mathcal{S}^\mu_{\beta]}}]_{\text{P.B.}} + \mathcal{O}(\hbar^3), \end{aligned} \quad (\text{A.18e})$$

where $[AB]_{\text{P.B.}} = \{A(q, X), B(q, X)\}_{\text{P.B.}}$ is a shorthand notation for the Poisson bracket.

A.2 Eliminating \mathcal{S} , \mathcal{P} and $\mathcal{S}_{\mu\nu}$

Equations (A.17a), (A.17b) and (A.17e) can be used to eliminate \mathcal{S} , \mathcal{P} , and $\mathcal{S}_{\mu\nu}$ from the other master equations. Since $\mathcal{S}^{\mu\nu}$ is contracted with $\Sigma_{T\mu\nu}$ and $\widehat{\Sigma}_{T\mu\nu}$ on the right-hand side, one may not express $\mathcal{S}_{\mu\nu}$ as an explicit function of \mathcal{V}_μ and \mathcal{A}_μ only. However, assuming that the interaction is sufficiently weak, we may drop the nonlinear terms in the self-energy. Within this assumption, we may rewrite eqs. (A.17a), (A.17b) and (A.17e) as

$$\begin{aligned} \mathcal{S}_{\alpha\beta} \approx & -\frac{1}{m}\epsilon_{\alpha\beta\rho\sigma}\Pi^\rho\mathcal{A}^\sigma + \frac{\hbar}{2m}\left\{\mathcal{D}_{[\alpha}\mathcal{V}_{\beta]} - \widehat{\Sigma}_{A[\alpha}\widehat{\mathcal{A}}_{\beta]} + \frac{q_\mu}{m}\epsilon^{\mu\rho\sigma}{}_{[\alpha}\widehat{\Sigma}_{T\beta]\rho}\widehat{\mathcal{A}}_\sigma\right\} \\ & + \frac{\hbar^2}{4m^2}\left(\widehat{\Sigma}_{T\mu[\alpha}\widehat{\nabla}^\mu\mathcal{V}_{\beta]} - \widehat{\Sigma}_{T\mu[\alpha}\widehat{\nabla}^\mu\mathcal{V}_{\beta]}\right) - \frac{\hbar^2}{4m}\left[\frac{\epsilon_{\alpha\beta\mu\nu}}{m}\widehat{\Sigma}_S(q^\nu\mathcal{A}^\mu) + \frac{1}{m}\widehat{\Sigma}_{T\alpha\beta}(q\cdot\mathcal{V})\right. \\ & \left. - \epsilon_{\mu\nu\alpha\beta}\widehat{\Sigma}_V^\mu\mathcal{A}^\nu + \epsilon_{\mu\nu\alpha\beta}\widehat{\Sigma}_A^\mu\mathcal{V}^\nu - \frac{1}{m}\widehat{\Sigma}_P(q[\alpha\widehat{\mathcal{A}}_{\beta]})\right]_{\text{P.B.}} + \mathcal{O}(\hbar^3), \end{aligned} \quad (\text{A.19a})$$

$$\mathcal{S} = \frac{\Pi^\mu}{m}\mathcal{V}_\mu - \frac{\hbar^2}{4m}\left[\frac{\widehat{\Sigma}_S(q\cdot\mathcal{V})}{m} + \widehat{\Sigma}_{V\mu}\mathcal{V}^\mu - \widehat{\Sigma}_{A\mu}\mathcal{A}^\mu - \frac{1}{2m}\epsilon^{\mu\nu\alpha\beta}\widehat{\Sigma}_{T\mu\nu}(q_\alpha\mathcal{A}_\beta)\right]_{\text{P.B.}} + \mathcal{O}(\hbar^3), \quad (\text{A.19b})$$

$$\mathcal{P} = -\frac{\hbar}{2m}(\widehat{\mathcal{D}}_\mu\mathcal{A}^\mu - \widehat{\Sigma}_{A\mu}\mathcal{V}^\mu) - \frac{\hbar^2}{4m}\left[\frac{\widehat{\Sigma}_P(q\cdot\mathcal{V})}{m} + \frac{1}{2m}\widehat{\Sigma}_T^{\mu\nu}(q_{[\mu}\mathcal{A}_{\nu]})\right]_{\text{P.B.}} + \mathcal{O}(\hbar^3). \quad (\text{A.19c})$$

Inserting eqs. (A.19b), (A.19c) and (A.19a) and maintaining the linear terms in the self-energies and the explicit \hbar dependence up to $\mathcal{O}(\hbar)$ (or $\mathcal{O}(\hbar^2)$ for involving at least the next-leading-order corrections), the master equations (A.17c) and (A.17d) up to $\mathcal{O}(\hbar)$ are reduced as

$$\begin{aligned} q^\mu(q\cdot\mathcal{V}) - m^2\mathcal{V}^\mu & \approx \frac{\hbar}{2}\left[m(\widehat{\Sigma}_P\mathcal{A}^\mu + \widehat{\Sigma}_T^{\mu\alpha}\mathcal{V}_\alpha) - 2\left(\widehat{F}^{\mu\beta}\mathcal{A}_\beta + \frac{1}{2}\epsilon^{\mu\alpha\beta\gamma}q_\alpha\Delta_\beta\mathcal{A}_\gamma\right) - \epsilon^{\mu\alpha\beta\gamma}q_\alpha\widehat{\Sigma}_{V\beta}\mathcal{A}_\gamma\right], \\ q^2\mathcal{A}^\mu - q^\mu q\cdot\mathcal{A} - m^2\mathcal{A}^\mu & \approx \frac{1}{2}\hbar\left[m(\widehat{\Sigma}_P\mathcal{V}^\mu + \widehat{\Sigma}_T^{\mu\alpha}\mathcal{A}_\alpha) + \epsilon^{\mu\alpha\beta\gamma}q_\alpha\mathcal{D}_\beta\mathcal{V}_\gamma - 2\epsilon^{\mu\alpha\beta\gamma}q_\alpha\widehat{\Sigma}_{A\beta}\mathcal{A}_\gamma - \frac{1}{m}\epsilon^\mu{}_{\alpha\beta\gamma}\epsilon^{\beta\lambda\rho\sigma}q^\alpha q_\lambda\widehat{\Sigma}_{T\rho}^\gamma\mathcal{A}_\sigma\right]. \end{aligned} \quad (\text{A.20a})$$

Similarly, the master equations (A.18a)–(A.18e) up to $\mathcal{O}(\hbar)$ are reduced as

$$\mathcal{D}^\mu\mathcal{V}_\mu \approx \widehat{\Sigma}_{A\mu}\mathcal{A}^\mu - \frac{1}{2m}\left[2q_\mu\widehat{\Sigma}_S\mathcal{V}^\mu - \epsilon_{\mu\nu\alpha\beta}q^\alpha\widehat{\Sigma}_T^{\mu\nu}\mathcal{A}^\beta + \hbar\widehat{\Sigma}_P(\nabla\cdot\mathcal{A}) + \frac{\hbar}{2}\widehat{\Sigma}_T^{\mu\nu}\widehat{\nabla}_{[\mu}\mathcal{V}_{\nu]}\right], \quad (\text{A.21a})$$

$$q\cdot\mathcal{A} \approx -\frac{\hbar}{2m}q_\mu(\widehat{\Sigma}_P\mathcal{V}^\mu + \widehat{\Sigma}_T^{\mu\nu}\mathcal{A}_\nu), \quad (\text{A.21b})$$

$$\begin{aligned} \mathcal{D}_\mu(q\cdot\mathcal{V}) - q^\nu\mathcal{D}_{[\mu}\mathcal{V}_{\nu]} & \approx -m\widehat{\Sigma}_S\mathcal{V}_\mu + q_{[\mu}\widehat{\Sigma}_{A\alpha]}\mathcal{A}^\alpha + \frac{1}{m}q^\nu q_\alpha\epsilon^{\alpha\beta\gamma}{}_{[\mu}\widehat{\Sigma}_{T\nu]\beta}\mathcal{A}_\gamma \\ & + \frac{1}{2}\epsilon_{\mu\nu\rho\sigma}\left\{m\widehat{\Sigma}_T^{\nu\rho}\mathcal{A}^\sigma - \hbar(\partial^\nu F^{\rho\beta})\partial_{q\beta}\mathcal{A}^\sigma + \hbar\widehat{\Sigma}_A^\nu(\nabla^\rho\mathcal{V}^\sigma)\right\} - \frac{\hbar q^\nu}{2m}(\widehat{\Sigma}_{T\gamma[\nu}\widehat{\Delta}^\gamma\mathcal{V}_{\mu]} - \widehat{\Sigma}_{T\gamma[\nu}\widehat{\Delta}_{\mu]}\mathcal{V}^\gamma) \\ & + \frac{\hbar q^\nu}{2}\left[\frac{1}{m}\widehat{\Sigma}_{T\nu\mu}(q\cdot\mathcal{V}) - \epsilon_{\nu\mu\rho\sigma}\widehat{\Sigma}_V^\rho\mathcal{A}^\sigma + \epsilon_{\nu\mu\rho\sigma}\widehat{\Sigma}_A^\rho\mathcal{V}^\sigma - \frac{1}{m}\widehat{\Sigma}_P(q_{[\nu}\mathcal{A}_{\mu]})\right]_{\text{P.B.}} \\ & + \frac{\hbar m}{2}\left[\widehat{\Sigma}_P\mathcal{A}_\mu - \frac{\epsilon_{\nu\mu\rho\sigma}}{m}\widehat{\Sigma}_V^\nu(q^\rho\mathcal{A}^\sigma) + \widehat{\Sigma}_{T\mu\beta}\mathcal{V}^\beta\right]_{\text{P.B.}}, \end{aligned} \quad (\text{A.21c})$$

$$F_{\mu\nu}\mathcal{A}^\nu - q\cdot\mathcal{D}\mathcal{A}_\mu + \frac{\hbar}{4}\epsilon_{\mu\nu\rho\sigma}[\mathcal{D}^\nu, \mathcal{D}^\rho]\mathcal{V}^\sigma - \frac{\hbar}{2m}q_\mu\left[\widehat{\Sigma}_P(q\cdot\mathcal{V}) + \frac{1}{2}\widehat{\Sigma}_T^{\rho\sigma}(q_{[\rho}\mathcal{A}_{\sigma]})\right]_{\text{P.B.}} \quad (\text{A.21d})$$

$$\begin{aligned} & = m\left[\widehat{\Sigma}_S\mathcal{A}_\mu - \frac{1}{2}\epsilon_{\mu\nu\rho\sigma}\widehat{\Sigma}_T^{\nu\rho}\mathcal{V}^\sigma\right] + q_\alpha\widehat{\Sigma}_{A\mu}\mathcal{V}^\alpha - q_\mu\widehat{\Sigma}_{A\alpha}\mathcal{V}^\alpha + \frac{\hbar}{2}\epsilon_{\mu\nu\rho\sigma}\Delta^\nu\left[\widehat{\Sigma}_A^\rho\mathcal{A}^\sigma - \frac{1}{m}q^\alpha\epsilon_{\alpha\beta\gamma}{}^\rho\widehat{\Sigma}_T^{\sigma\beta}\mathcal{A}^\gamma\right] \\ & - \frac{\hbar m}{2}\left[\widehat{\Sigma}_P\mathcal{V}_\mu + \frac{1}{m}\epsilon_{\mu\nu\rho\sigma}\widehat{\Sigma}_A^\nu(q^\rho\mathcal{A}^\sigma) + \widehat{\Sigma}_{T\mu\nu}\mathcal{A}^\nu\right]_{\text{P.B.}}, \\ q^{[\mu}\mathcal{V}^{\nu]} + \frac{\hbar}{2}\epsilon^{\mu\nu\alpha\beta}(\mathcal{D}_\alpha\mathcal{A}_\beta - \widehat{\Sigma}_{A\alpha}\mathcal{V}_\beta) & \approx -\frac{\hbar}{2m}\left[\epsilon^{\mu\nu\alpha\beta}q_\alpha\widehat{\Sigma}_S\mathcal{A}_\beta - q_\alpha\widehat{\Sigma}_T^{\mu\nu}\mathcal{V}^\alpha + q^{[\mu}\widehat{\Sigma}_P\mathcal{A}^{\nu]}\right], \end{aligned} \quad (\text{A.21e})$$

where we defined $\mathcal{D}_\mu \mathcal{M} = \nabla_\mu \mathcal{M} + \widehat{\Sigma_{V\mu} \mathcal{M}}$, and we used the following relation

$$2\epsilon_{\nu\mu\rho\sigma}(\Pi^\nu \Pi^\rho) \mathcal{A}^\sigma = \frac{\hbar^2}{6} \epsilon_{\nu\mu\rho\sigma} (\partial^\rho F^{\beta\nu} + \partial^\beta F^{\rho\nu}) \partial_{q\beta} \mathcal{A}^\sigma = \frac{\hbar^2}{2} \epsilon_{\nu\mu\rho\sigma} (\partial^\rho F^{\beta\nu}) \partial_{q\beta} \mathcal{A}^\sigma. \quad (\text{A.22})$$

The master equations (A.20) and (A.21) are in general the full master equations with collisional effects up to $\mathcal{O}(\hbar^1)$. In our power counting $\mathcal{V}^\mu \sim \mathcal{O}(\hbar^0)$ and $\mathcal{A}^\mu \sim \mathcal{O}(\hbar^1)$, these equations reduce to eqs. (2.20a)–(2.20g).

B Angular-momentum decomposition for fermions and spin polarization

We briefly review the derivation of the gauge-invariant angular-momentum decomposition for fermions proposed in ref. [79] as a covariant version of the Ji’s decomposition [80]. Such a derivation is more explicitly shown in the review paper [81] (See also ref. [82]). Here we just summarize the derivation therein. The starting point is the Belinfante angular momentum for “on-shell” fermions,

$$M_B^{\mu\nu\rho} = \frac{x^\nu}{4} \bar{\psi} (\gamma^\mu i \overleftrightarrow{D}^\rho + \gamma^\rho i \overleftrightarrow{D}^\mu) \psi - (\nu \leftrightarrow \rho), \quad (\text{B.1})$$

where $\bar{\psi} \overleftrightarrow{D}_\mu \psi = \bar{\psi} (\overrightarrow{D}_\mu - \overleftarrow{D}_\mu^\dagger) \psi$ and $D_\mu = \partial_\mu + ieA_\mu$ denotes the covariant derivative. One may rewrite $M_B^{\mu\nu\rho}$ as

$$M_B^{\mu\nu\rho} = \frac{x^\nu}{2} \bar{\psi} \gamma^\mu i \overleftrightarrow{D}^\rho \psi - \frac{x^\nu}{4} \bar{\psi} (\gamma^\mu i \overleftrightarrow{D}^\rho - \gamma^\rho i \overleftrightarrow{D}^\mu) \psi - (\nu \leftrightarrow \rho). \quad (\text{B.2})$$

Next, one has to employ the relations,

$$\begin{aligned} \sigma^{\mu\nu} i \overrightarrow{D} &= \gamma^\nu \overrightarrow{D}^\mu - \gamma^\mu \overrightarrow{D}^\nu - i\epsilon^{\mu\nu\alpha\beta} \gamma_\beta \gamma_5 \overrightarrow{D}_\alpha, \\ i \overleftarrow{D} \sigma^{\mu\nu} &= \overleftarrow{D}^\mu \gamma^\nu - \overleftarrow{D}^\nu \gamma^\mu - i\epsilon^{\mu\nu\alpha\beta} \gamma_\beta \gamma_5 \overleftarrow{D}_\alpha, \end{aligned} \quad (\text{B.3})$$

with $\sigma^{\mu\nu} = i[\gamma^\mu, \gamma^\nu]/2$ and the equations of motion $i \overrightarrow{D} \psi = m\psi$ and $i \overleftarrow{D} \bar{\psi} = -m\bar{\psi}$, to obtain a useful identity,

$$\bar{\psi} (\gamma^\mu i \overleftrightarrow{D}^\rho - \gamma^\rho i \overleftrightarrow{D}^\mu) \psi = -\epsilon^{\rho\mu\alpha\beta} \partial_\alpha (\bar{\psi} \gamma_\beta \gamma_5 \psi). \quad (\text{B.4})$$

By using the identity, it is found

$$M_B^{\mu\nu\rho} = \frac{i}{2} \bar{\psi} \gamma^\mu (x^\nu \overleftrightarrow{D}^\rho - x^\rho \overleftrightarrow{D}^\nu) \psi - \frac{1}{2} \epsilon^{\mu\nu\rho\beta} \bar{\psi} \gamma_\beta \gamma_5 \psi - \frac{1}{4} \partial_\alpha \left((x^\nu \epsilon^{\mu\rho\alpha\beta} - x^\rho \epsilon^{\mu\nu\alpha\beta}) \bar{\psi} \gamma_\beta \gamma_5 \psi \right) \quad (\text{B.5})$$

and thus yields the gauge-invariant angular-momentum decomposition by dropping the surface term above,

$$M_C^{\mu\nu\rho} = \frac{i}{2} \bar{\psi} \gamma^\mu (x^\nu \overleftrightarrow{D}^\rho - x^\rho \overleftrightarrow{D}^\nu) \psi - \frac{1}{2} \epsilon^{\mu\nu\rho\beta} \bar{\psi} \gamma_\beta \gamma_5 \psi, \quad (\text{B.6})$$

where the first term above is regarded as the orbital angular momentum and the second term proportional to an axial-charge current is responsible for spin. This gauge-invariant decomposition introduced in ref. [79] is defined as the “canonical angular momentum” in ref. [44] though it actually agrees with the usual canonical angular momentum obtained from the Noether’s theorem only in the absence of gauge fields.

C Decomposition of the collision term in AKE

The expression of eq. (3.8) can be equivalently written as

$$\begin{aligned} \hat{C}_2^{(n)\mu} = & \frac{\delta(q^2 - m^2)}{2} \left\{ \epsilon^{\mu\nu\rho\sigma} q_\nu \left((\Delta_\rho \bar{\Sigma}_{V\sigma}) f_V - (\Delta_\rho \Sigma_{V\sigma}) \bar{f}_V \right) + 2S_{m(n)}^{\mu\nu} \left(m(\Sigma_S \Delta_\nu \bar{f}_V - \bar{\Sigma}_S \Delta_\nu f_V) \right. \right. \\ & - F_\nu^\rho (\Sigma_{V\rho} \bar{f}_V - \bar{\Sigma}_{V\rho} f_V) - q^\rho (\bar{f}_V \Delta_\nu \Sigma_{V\rho} - f_V \Delta_\nu \bar{\Sigma}_{V\rho}) + \bar{f}_V (q \cdot \Delta \Sigma_{V\nu}) - f_V (q \cdot \Delta \bar{\Sigma}_{V\nu}) \\ & \left. \left. + 2(\Sigma_{V\nu} \bar{f}_V - \bar{\Sigma}_{V\nu} f_V) \left(q \cdot \Delta S_{m(n)}^{\mu\nu} - F_\lambda^\mu S_{m(n)}^{\lambda\nu} \right) \right\} - m\delta(q^2 - m^2) S_{m(n)}^{\mu\nu} \Delta_\nu C_S[f_V] \\ & - 2S_{m(n)}^{\mu\nu} q^\lambda F_{\lambda\nu} \delta'(q^2 - m^2) (C_V[f_V] + mC_S[f_V]) - \delta(q^2 - m^2) (\Delta_\nu S_{m(n)}^{\mu\nu}) \\ & \times (C_V[f_V] + mC_S[f_V]) + \tilde{F}^{\mu\nu} q_\nu \delta'(q^2 - m^2) (C_V[f_V] + mC_S[f_V]) \\ & + \frac{\delta(q^2 - m^2)}{2m} \left[q^\mu (f_V q \cdot \Delta \bar{\Sigma}_P - \bar{f}_V q \cdot \Delta \Sigma_P) - m^2 ((\partial^\mu \bar{\Sigma}_P) f_V - (\partial^\mu \Sigma_P) \bar{f}_V) \right]. \end{aligned} \quad (C.1)$$

By analogy with the collisionless case, we would like to decompose $\hat{C}_2^{(n)\mu}$ above into the piece proportional to q^μ , which survives in the massless limit and reproduces the collision term in CKT, and another piece proportional to m , which stems from the purely finite-mass correction.

The Schouton identity gives rise to

$$S_{m(n)}^{\mu\nu} q^\alpha = q^\mu S_{m(n)}^{\alpha\nu} + q^\nu S_{m(n)}^{\mu\alpha} + \frac{\epsilon^{\mu\nu\rho\alpha} q_\rho}{2} - \frac{\epsilon^{\mu\nu\rho\alpha}}{2(q \cdot n + m)} (mq_\rho + q^2 n_\rho). \quad (C.2)$$

We thus find

$$\begin{aligned} & 2S_{m(n)}^{\mu\nu} \left(\bar{f}_V (q \cdot \Delta \Sigma_{V\nu}) - f_V (q \cdot \Delta \bar{\Sigma}_{V\nu}) \right) \\ & = 2 \left[(q^\mu S_{m(n)}^{\rho\nu} + q^\nu S_{m(n)}^{\mu\rho}) (\bar{f}_V \Delta_\rho \Sigma_{V\nu} - f_V \Delta_\rho \bar{\Sigma}_{V\nu}) \right] - \epsilon^{\mu\nu\rho\sigma} q_\nu \left((\Delta_\rho \bar{\Sigma}_{V\sigma}) f_V - (\Delta_\rho \Sigma_{V\sigma}) \bar{f}_V \right) \\ & \quad - \frac{\epsilon^{\mu\nu\rho\sigma} (mq_\rho + q^2 n_\rho)}{(q \cdot n + m)} (\bar{f}_V \Delta_\sigma \Sigma_{V\nu} - f_V \Delta_\sigma \bar{\Sigma}_{V\nu}) \end{aligned} \quad (C.3)$$

and $\hat{C}_2^{(n)\mu}$ becomes

$$\begin{aligned} & \hat{C}_2^{(n)\mu} \\ & = \frac{\delta(q^2 - m^2)}{2} \left\{ 2q^\mu S_{m(n)}^{\rho\nu} (\bar{f}_V \Delta_\rho \Sigma_{V\nu} - f_V \Delta_\rho \bar{\Sigma}_{V\nu}) + 2S_{m(n)}^{\mu\nu} \left(m(\Sigma_S \Delta_\nu \bar{f}_V - \bar{\Sigma}_S \Delta_\nu f_V) \right. \right. \\ & \quad \left. \left. - F_\nu^\rho (\Sigma_{V\rho} \bar{f}_V - \bar{\Sigma}_{V\rho} f_V) \right) - \frac{\epsilon^{\mu\nu\rho\sigma} m(q_\rho + mn_\rho)}{(q \cdot n + m)} (\bar{f}_V \Delta_\sigma \Sigma_{V\nu} - f_V \Delta_\sigma \bar{\Sigma}_{V\nu}) + 2(\Sigma_{V\nu} \bar{f}_V - \bar{\Sigma}_{V\nu} f_V) \right. \\ & \quad \left. \times \left(q \cdot \Delta S_{m(n)}^{\mu\nu} - F_\lambda^\mu S_{m(n)}^{\lambda\nu} \right) \right\} - 2S_{m(n)}^{\mu\nu} q^\lambda F_{\lambda\nu} \delta'(q^2 - m^2) (C_V[f_V] + mC_S[f_V]) - \delta(q^2 - m^2) \\ & \quad \times \left[(\Delta_\nu S_{m(n)}^{\mu\nu}) (C_V[f_V] + mC_S[f_V]) + mS_{m(n)}^{\mu\nu} \Delta_\nu C_S[f_V] \right] \\ & \quad + \tilde{F}^{\mu\nu} q_\nu \delta'(q^2 - m^2) (C_V[f_V] + mC_S[f_V]) + \frac{\delta(q^2 - m^2)}{2m} \left[q^\mu (f_V q \cdot \Delta \bar{\Sigma}_P - \bar{f}_V q \cdot \Delta \Sigma_P) \right. \\ & \quad \left. - m^2 ((\partial^\mu \bar{\Sigma}_P) f_V - (\partial^\mu \Sigma_P) \bar{f}_V) \right]. \end{aligned} \quad (C.4)$$

Next, applying the Schouton identity again, it is found

$$\begin{aligned}
 & \delta(q^2 - m^2) q \cdot \Delta S_{m(n)}^{\mu\nu} \\
 &= \delta(q^2 - m^2) \left[q^\mu \Delta_\alpha S_{m(n)}^{\alpha\nu} + q^\nu \Delta_\alpha S_{m(n)}^{\mu\alpha} + q^\rho \epsilon^{\mu\nu\alpha\sigma} \Delta_\alpha \frac{q_\rho n_\sigma}{2(q \cdot n + m)} + q^\sigma \epsilon^{\mu\nu\rho\alpha} \Delta_\alpha \frac{q_\rho n_\sigma}{2(q \cdot n + m)} \right] \\
 &= \delta(q^2 - m^2) \left[q^\mu \Delta_\alpha S_{m(n)}^{\alpha\nu} + q^\nu \Delta_\alpha S_{m(n)}^{\mu\alpha} + m^2 \Delta_\alpha \frac{\epsilon^{\mu\nu\alpha\sigma} n_\sigma}{2(q \cdot n + m)} + \frac{q^\rho F_{\rho\alpha} \epsilon^{\mu\nu\alpha\sigma} n_\sigma}{2(q \cdot n + m)} + \tilde{F}^{\mu\nu} \right. \\
 & \quad \left. - m \Delta_\alpha \frac{\epsilon^{\mu\nu\rho\alpha} q_\rho}{2(q \cdot n + m)} - \frac{\epsilon^{\mu\nu\rho\alpha} F_{\sigma\alpha} n^\sigma q_\rho}{2(q \cdot n + m)} \right] \tag{C.5}
 \end{aligned}$$

and

$$-S_{m(n)}^{\lambda\nu} F^\mu{}_\lambda = -\frac{q_\rho n_\sigma}{2(q \cdot n + m)} (\epsilon^{\lambda\mu\rho\sigma} F^\nu{}_\lambda + \epsilon^{\lambda\nu\mu\sigma} F^\rho{}_\lambda + \epsilon^{\lambda\nu\rho\mu} F^\sigma{}_\lambda), \tag{C.6}$$

which lead to

$$\begin{aligned}
 & \delta(q^2 - m^2) (q \cdot \Delta S_{m(n)}^{\mu\nu} - S_{m(n)}^{\lambda\nu} F^\mu{}_\lambda) \\
 &= \delta(q^2 - m^2) \left[q^\mu \Delta_\alpha S_{m(n)}^{\alpha\nu} + \tilde{F}^{\mu\nu} + q^\nu \Delta_\alpha S_{m(n)}^{\mu\alpha} - m \Delta_\alpha \frac{\epsilon^{\mu\nu\rho\alpha} (q_\rho + m n_\rho)}{2(q \cdot n + m)} + S_{m(n)}^{\mu\rho} F^\nu{}_\rho \right. \\
 & \quad \left. + \frac{q^\rho F_{\rho\alpha} \epsilon^{\mu\nu\alpha\sigma} n_\sigma}{(q \cdot n + m)} \right]. \tag{C.7}
 \end{aligned}$$

We hence obtain

$$\begin{aligned}
 & \hat{C}_2^{(n)\mu} \\
 &= \frac{\delta(q^2 - m^2)}{2} \left\{ 2q^\mu S_{m(n)}^{\rho\nu} (\bar{f}_V \Delta_\rho \Sigma_{V\nu} - f_V \Delta_\rho \bar{\Sigma}_{V\nu}) + 2S_{m(n)}^{\mu\nu} m (\Sigma_S \Delta_\nu \bar{f}_V - \bar{\Sigma}_S \Delta_\nu f_V) \right. \\
 & \quad \left. - \frac{\epsilon^{\mu\nu\rho\sigma} m (q_\rho + m n_\rho)}{(q \cdot n + m)} (\bar{f}_V \Delta_\sigma \Sigma_{V\nu} - f_V \Delta_\sigma \bar{\Sigma}_{V\nu}) + 2(\bar{f}_V \Sigma_{V\nu} - f_V \bar{\Sigma}_{V\nu}) \left[q^\mu \Delta_\alpha S_{m(n)}^{\alpha\nu} + \tilde{F}^{\mu\nu} \right. \right. \\
 & \quad \left. \left. + \frac{q^\rho F_{\rho\alpha} \epsilon^{\mu\nu\alpha\sigma} n_\sigma}{(q \cdot n + m)} - m \Delta_\alpha \frac{\epsilon^{\mu\nu\rho\alpha} (q_\rho + m n_\rho)}{2(q \cdot n + m)} \right] \right\} - 2S_{m(n)}^{\mu\nu} q^\lambda F_{\lambda\nu} \delta'(q^2 - m^2) C_V[f_V] + \delta'(q^2 - m^2) \\
 & \quad \times \tilde{F}^{\mu\nu} q_\nu (C_V[f_V] + m C_S[f_V]) \\
 & \quad - m \left(2S_{m(n)}^{\mu\nu} q^\lambda F_{\lambda\nu} \delta'(q^2 - m^2) + \delta(q^2 - m^2) ((\Delta_\nu S_{m(n)}^{\mu\nu}) + S_{m(n)}^{\mu\nu} \Delta_\nu) \right) C_S[f_V] \\
 & \quad + \frac{\delta(q^2 - m^2)}{2m} \left[q^\mu (f_V q \cdot \Delta \bar{\Sigma}_P - \bar{f}_V q \cdot \Delta \Sigma_P) - m^2 ((\partial^\mu \bar{\Sigma}_P) f_V - (\partial^\mu \Sigma_P) \bar{f}_V) \right]. \tag{C.8}
 \end{aligned}$$

On the other hand, one finds

$$\begin{aligned}
 & -2S_{m(n)}^{\mu\nu} \delta'(q^2 - m^2) q^\lambda F_{\lambda\nu} = -2\delta'(q^2 - m^2) (q^\mu S_{m(n)}^{\rho\nu} + q^\nu S_{m(n)}^{\mu\rho}) F_{\rho\nu} - 2\tilde{F}^{\mu\rho} q_\rho \delta'(q^2 - m^2) \\
 & \quad + \frac{2\delta'(q^2 - m^2) \tilde{F}^{\mu\rho}}{(q \cdot n + m)} m (q_\rho + m n_\rho) - \frac{2\delta(q^2 - m^2)}{q \cdot n + m} \tilde{F}^{\mu\rho} n_\rho, \tag{C.9}
 \end{aligned}$$

which yields

$$-2S_{m(n)}^{\mu\nu}\delta'(q^2-m^2)q^\lambda F_{\lambda\nu} = -\delta'(q^2-m^2)\left(q^\mu S_{m(n)}^{\rho\nu}F_{\rho\nu} + \tilde{F}^{\mu\rho}q_\rho - \frac{m(q_\rho + mn_\rho)\tilde{F}^{\mu\rho}}{(q\cdot n + m)}\right) - \frac{\delta(q^2-m^2)}{q\cdot n + m}\tilde{F}^{\mu\rho}n_\rho. \quad (\text{C.10})$$

In addition, one can show

$$\frac{q^\rho F_{\rho\alpha}\epsilon^{\mu\nu\alpha\sigma}n_\sigma}{(q\cdot n + m)} = \frac{1}{q\cdot n + m}(-q^\mu\tilde{F}^{\nu\sigma}n_\sigma + \tilde{F}^{\mu\sigma}n_\sigma q^\nu + m\tilde{F}^{\mu\nu}) - \tilde{F}^{\mu\nu}. \quad (\text{C.11})$$

Accordingly, $\hat{C}_2^{(n)}$ takes the form

$$\begin{aligned} \hat{C}_2^{(n)\mu} = & \frac{\delta(q^2-m^2)}{2}\left\{2q^\mu S_{m(n)}^{\rho\nu}(\bar{f}_V\Delta_\rho\Sigma_{V\nu} - f_V\Delta_\rho\bar{\Sigma}_{V\nu}) + 2S_{m(n)}^{\mu\nu}m(\Sigma_S\Delta_\nu\bar{f}_V - \bar{\Sigma}_S\Delta_\nu f_V)\right. \\ & - \frac{\epsilon^{\mu\nu\rho\sigma}m(q_\rho + mn_\rho)}{(q\cdot n + m)}(\bar{f}_V\Delta_\sigma\Sigma_{V\nu} - f_V\Delta_\sigma\bar{\Sigma}_{V\nu}) + 2(\bar{f}_V\Sigma_{V\nu} - f_V\bar{\Sigma}_{V\nu})\left[q^\mu\Delta_\alpha S_{m(n)}^{\alpha\nu}\right. \\ & \left. - \frac{q^\mu\tilde{F}^{\nu\sigma}n_\sigma}{(q\cdot n + m)} + \frac{m\tilde{F}^{\mu\nu}}{(q\cdot n + m)} - m\Delta_\alpha\frac{\epsilon^{\mu\nu\rho\alpha}(q_\rho + mn_\rho)}{2(q\cdot n + m)}\right]\left\} \right. \\ & + \frac{\delta(q^2-m^2)}{2m}\left[q^\mu(f_Vq\cdot\Delta\bar{\Sigma}_P - \bar{f}_Vq\cdot\Delta\Sigma_P) - m^2((\partial^\mu\bar{\Sigma}_P)f_V - (\partial^\mu\Sigma_P)\bar{f}_V)\right] \\ & - \delta'(q^2-m^2)\left(q^\mu S_{m(n)}^{\rho\nu}F_{\rho\nu} - \frac{m(q_\rho + mn_\rho)\tilde{F}^{\mu\rho}}{(q\cdot n + m)}\right)C_V[f_V] - m\left((2S_{m(n)}^{\mu\nu}q^\lambda F_{\lambda\nu} - \tilde{F}^{\mu\nu}q_\nu)\right. \\ & \left.\times\delta'(q^2-m^2) + \delta(q^2-m^2)(\Delta_\nu S_{m(n)}^{\mu\nu}) + \delta(q^2-m^2)S_{m(n)}^{\mu\nu}\Delta_\nu\right)C_S[f_V], \quad (\text{C.12}) \end{aligned}$$

which can be further rearranged as

$$\begin{aligned} & \hat{C}_2^{(n)\mu} \\ = & q^\mu\left\{\delta(q^2-m^2)\left[S_{m(n)}^{\rho\nu}(\bar{f}_V\Delta_\rho\Sigma_{V\nu} - f_V\Delta_\rho\bar{\Sigma}_{V\nu}) + (\bar{f}_V\Sigma_{V\nu} - f_V\bar{\Sigma}_{V\nu})\left(\partial_\alpha S_{m(n)}^{\alpha\nu} + \frac{S_{m(n)}^{\rho\nu}F_{\rho\sigma}n^\sigma}{q\cdot n + m}\right)\right.\right. \\ & \left. + \frac{1}{2m}(f_Vq\cdot\Delta\bar{\Sigma}_P - \bar{f}_Vq\cdot\Delta\Sigma_P)\right] - \delta'(q^2-m^2)S_{m(n)}^{\rho\nu}F_{\rho\nu}C_V[f_V]\left\} \right. \\ & + m\left\{\delta(q^2-m^2)\left[S_{m(n)}^{\mu\nu}(\Sigma_S\Delta_\nu\bar{f}_V - \bar{\Sigma}_S\Delta_\nu f_V) - \Delta_\nu(S_{m(n)}^{\mu\nu}C_S[f_V]) - \frac{\epsilon^{\mu\nu\rho\sigma}(q_\rho + mn_\rho)}{2(q\cdot n + m)}\right.\right. \\ & \left.\times(\bar{f}_V\Delta_\sigma\Sigma_{V\nu} - f_V\Delta_\sigma\bar{\Sigma}_{V\nu}) + (\bar{f}_V\Sigma_{V\nu} - f_V\bar{\Sigma}_{V\nu})\left(\frac{\tilde{F}^{\mu\nu}}{(q\cdot n + m)} - \epsilon^{\mu\nu\rho\alpha}\Delta_\alpha\frac{(q_\rho + mn_\rho)}{2(q\cdot n + m)}\right)\right. \\ & \left. - \frac{1}{2}((\partial^\mu\bar{\Sigma}_P)f_V - (\partial^\mu\Sigma_P)\bar{f}_V)\right] \\ & \left. + \delta'(q^2-m^2)\left[\frac{(q_\rho + mn_\rho)\tilde{F}^{\mu\rho}}{(q\cdot n + m)}C_V[f_V] - (2S_{m(n)}^{\mu\nu}q^\lambda F_{\lambda\nu} - \tilde{F}^{\mu\nu}q_\nu)C_S[f_V]\right]\right\}, \quad (\text{C.13}) \end{aligned}$$

where we utilize

$$\Delta_\alpha S_{m(n)}^{\alpha\nu} - \frac{\tilde{F}^{\nu\sigma} n_\sigma}{(q \cdot n + m)} = \partial_\alpha S_{m(n)}^{\alpha\nu} + \frac{S_{m(n)}^{\rho\nu} F_{\rho\sigma} n^\sigma}{q \cdot n + m}. \quad (\text{C.14})$$

We can now write $\hat{\mathcal{C}}_2^{(n)\mu}$ in a more succinct form as

$$\hat{\mathcal{C}}_2^{(n)\mu} = q^\mu \hat{\mathcal{C}}_{q^2}^{(n)} + m \hat{\mathcal{C}}_{m^2}^{(n)}, \quad (\text{C.15})$$

where

$$\begin{aligned} \hat{\mathcal{C}}_{q^2}^{(n)} = & \delta(q^2 - m^2) \left[-S_{m(n)}^{\rho\nu} (\Delta_\rho \widehat{\Sigma_{V\nu}}) f_V - \widehat{\Sigma_{V\nu}} f_V \left(\partial_\alpha S_{m(n)}^{\alpha\nu} + \frac{S_{m(n)}^{\rho\nu} F_{\rho\sigma} n^\sigma}{q \cdot n + m} \right) \right. \\ & \left. + \frac{1}{2m} (q \cdot \Delta \widehat{\Sigma_P}) f_V \right] - \delta'(q^2 - m^2) S_{m(n)}^{\rho\nu} F_{\rho\nu} C_V[f_V], \end{aligned} \quad (\text{C.16})$$

and

$$\begin{aligned} \hat{\mathcal{C}}_{m^2}^{(n)\mu} = & \delta(q^2 - m^2) \left[(\Delta_\nu S_{m(n)}^{\nu\mu}) C_S[f_V] - S_{m(n)}^{\nu\mu} (\Delta_\nu \widehat{\Sigma_S}) f_V + \frac{\epsilon^{\mu\nu\rho\sigma} (q_\rho + mn_\rho)}{2(q \cdot n + m)} (\Delta_\sigma \widehat{\Sigma_{V\nu}}) f_V \right. \\ & \left. - \frac{1}{2} (\partial^\mu \widehat{\Sigma_P}) f_V - \frac{\epsilon^{\mu\nu\alpha\beta} \widehat{\Sigma_{V\nu}} f_V}{2(q \cdot n + m)} \left(m \partial_\alpha n_\beta - (q_\beta + mn_\beta) \frac{(\partial_\alpha q \cdot n + n^\rho F_{\rho\alpha})}{q \cdot n + m} \right) \right] \\ & + \delta'(q^2 - m^2) \left[\frac{(q_\rho + mn_\rho) \tilde{F}^{\mu\rho}}{(q \cdot n + m)} C_V[f_V] - 2S_{m(n)}^{\mu\nu} q^\lambda F_{\lambda\nu} C_S[f_V] + \tilde{F}^{\mu\nu} q_\nu C_S[f_V] \right]. \end{aligned} \quad (\text{C.17})$$

Here, $C_V[f_V] \equiv -q_\mu \widehat{\Sigma_V^\mu} f_V$ and $C_S[f_V] \equiv -\widehat{\Sigma_S} f_V$. It is clear to see that $\hat{\mathcal{C}}_{q^2}^{(n)}$ reduces to the $\mathcal{O}(\hbar)$ correction in the collision term of the CKT found in refs. [30, 31] by taking $m = 0$ ⁴. Combining with $\hat{\mathcal{C}}_1^{(n)\mu}$ and collisionless part, one finds that the CKT with collisions is smoothly reproduced by AKE in the massless limit when $m = 0$ and $a^\mu = q^\mu$.

D Derivation of the spin diffusion for SKE

From eqs. (4.9) and (4.16), we find

$$\begin{aligned} Q_1(p) = & 2\pi \delta(q'^2 - m^2) \left(2q'^\mu G_{\mu\nu}^< q^\nu - p \cdot q' G_\mu^< \right) \\ = & 2\pi \delta(q'^2 - m^2) g_{0p} \frac{p_0}{T} \left(2\hat{\rho}_T (|\mathbf{q}|^2 (1 - z^2) - p_0 q_0 + |\mathbf{q}| |\mathbf{p}| z) + \hat{\rho}_L (2q_0^2 - p_0 q_0 - |\mathbf{q}| |\mathbf{p}| z) \right), \end{aligned} \quad (\text{D.1})$$

where we defined $\hat{\rho}_{T/L} \equiv \rho_{T/L} T / p_0$ and $z \equiv \hat{\mathbf{q}} \cdot \hat{\mathbf{p}}$. We also used the transversality $p_\mu P_T^{\mu\nu} = 0$ and the on-shell conditions $m^2 = q^2 = p^2 + q'^2 + 2p \cdot q'$ and $m^2 = q'^2 = p^2 + q^2 - 2p \cdot q$ that lead to relations $2p \cdot q = p^2 = -2p \cdot q'$.

⁴In refs. [30, 31], Σ_P and $\bar{\Sigma}_P$ are set to zero. As argued previously, such terms are actually expected to be at higher order and vanishing in the massless limit.

The on-shell condition can be also arranged as

$$q'^2 - m^2 = p^2 - 2q \cdot p = (p^0 - q^0)^2 + (2|\mathbf{q}||\mathbf{p}|z - \mathbf{p}^2 - q_0^2). \quad (\text{D.2})$$

Therefore, the delta function is nonzero when the momenta satisfy the conditions

$$p^0 = q^0 \pm \sqrt{\mathbf{p}^2 - 2|\mathbf{q}||\mathbf{p}|z + q_0^2} \sim q^0 \pm \left(q^0 - \frac{|\mathbf{q}||\mathbf{p}|z}{q_0} + \frac{\mathbf{p}^2(q_0^2 - \mathbf{q}^2 z^2)}{2q_0^3} \right), \quad (\text{D.3})$$

where the square root is expanded for a small momentum transfer $|\mathbf{p}|$. We take the positive-energy solution $q'^0 = q^0 - p^0 > 0$ for particles from the lower sign. Defining $\bar{p} \equiv |\mathbf{q}||\mathbf{p}|z/q_0 - \mathbf{p}^2(q_0^2 - \mathbf{q}^2 z^2)/(2q_0^3)$, we have⁵

$$\delta(q'^2 - m^2) \sim \frac{1}{2q^0} \left(1 + \frac{|\mathbf{q}||\mathbf{p}|z}{q_0^2} + \mathcal{O}(|\mathbf{p}|^2) \right) \delta(p^0 - \bar{p}). \quad (\text{D.4})$$

Based on the dispersion relation above, we can approximate the spectral function as

$$\hat{\rho}_T(p) \rightarrow \frac{\pi T m_D^2}{2|\mathbf{p}|^5} I_T(z) \left(1 - \frac{|\mathbf{q}||\mathbf{p}|z}{q_0^2} + \mathcal{O}(|\mathbf{p}|^2) \right), \quad (\text{D.5})$$

where $I_T(z) \equiv [1 - (|\mathbf{q}|z/q_0)^2]^{-1}$. Here, the presence of the delta function $\delta(p^0 - \bar{p})$ is assumed on the right-hand side of arrow. On the other hand, we have $\hat{\rho}_L(p) = \pi T m_D^2/|\mathbf{p}|^5$.

Note that the difference between the distribution functions provides positive powers in the small momentum transfer limit:

$$f_{Vq} - f_{Vq-p} \approx p^\beta \partial_{q^\beta} f_{Vq} - \frac{p^\alpha p^\beta}{2} \partial_{q^\alpha} \partial_{q^\beta} f_{Vq}, \quad (\text{D.6})$$

where $p^0 \sim \mathcal{O}(|\mathbf{p}|^1)$. Therefore, we have found the momentum dependences

$$\hat{\rho}_{L,T} \sim |\mathbf{p}|^{-5}, \quad f_{Vq} - f_{Vq-p} \sim |\mathbf{p}|, \quad d^3\mathbf{p} \sim d|\mathbf{p}||\mathbf{p}|^2. \quad (\text{D.7})$$

This order counting suggests that one should maintain the other factors up to $\mathcal{O}(|\mathbf{p}|)$ to get the leading-log result. Therefore, we expand the remaining factor as

$$\delta(q'^2 - m^2) g_{0p} \frac{p_0}{T} \approx \frac{1}{2|q_0|} \left[1 + \frac{|\mathbf{p}||\mathbf{q}|z}{q_0^2} \left(1 - \frac{q_0}{2T} \right) + \mathcal{O}(|\mathbf{p}|^2) \right] \delta(p^0 - \bar{p}), \quad (\text{D.8})$$

where we used $g_{0p} \approx T/p_0 - 1/2$. Combining all pieces together, it is found

$$\begin{aligned} \int_p Q_1(f_{Vq} - f_{Vq-p}) &\approx \frac{\pi T m_D^2}{2|q_0|} \int_p \frac{1}{|\mathbf{p}|^5} (2\pi) \delta \left(p_0 - \frac{\mathbf{p} \cdot \mathbf{q}}{q_0} \right) \\ &\times \left(1 - \frac{|\mathbf{p}||\mathbf{q}|z}{2q_0 T} \right) \left[2q_0^2 + |\mathbf{q}|^2 (1 - z^2) I_T(z) \right] (f_{Vq} - f_{Vq-p}). \end{aligned} \quad (\text{D.9})$$

⁵It turns out that the $\mathcal{O}(|\mathbf{p}|^2)$ term in the Delta function does not contribute to the leading logarithmic order.

In eq. (D.6), we apply the following decomposition of p_{\perp}^{β} in terms of the transverse and longitudinal components with respect to q_{\perp}^{μ} :

$$p_{\perp}^{\mu} = \hat{q}_{\perp}^{\mu} |\mathbf{p}| z + \hat{\Theta}_q^{\mu\nu} p_{\nu}, \quad \hat{\Theta}_q^{\alpha\beta} = \Theta^{\alpha\beta} + \hat{q}_{\perp}^{\alpha} \hat{q}_{\perp}^{\beta}, \quad (\text{D.10})$$

where $q_{\alpha} \hat{\Theta}_q^{\alpha\beta} = 0$. Moreover, The distribution function f_{Vq} in general can be a function of q_0 before implementing the on-shell condition. It is in fact more convenient to keep the off-shell form when $F^{\mu\nu} \neq 0$ due to the presence of $\delta'(q^2 - m^2)$ term in the AKE. Nonetheless, in the absence of background fields, it is more practical and convenient to write down the on-shell kinetic equations. We hence take $f_{Vq} = f_{Vq}(\mathbf{q}, X)$ as just a function of \mathbf{q} and X by using $q_0 = E_q = \sqrt{|\mathbf{q}|^2 + m^2}$ for fermions (here we neglect anti-fermions) in the Wigner functions. Accordingly, all the terms proportional to $\partial_{q_0} f_{Vq}$ can be dropped.

Applying the replacement

$$p^{\alpha} p^{\beta} \rightarrow p_0^2 u^{\alpha} u^{\beta} + p_0 |\mathbf{p}| z (u^{\alpha} \hat{q}_{\perp}^{\beta} + u^{\beta} \hat{q}_{\perp}^{\alpha}) + |\mathbf{p}|^2 \left(\hat{q}_{\perp}^{\alpha} \hat{q}_{\perp}^{\beta} z^2 - \frac{\hat{\Theta}_q^{\alpha\beta}}{2} (1 - z^2) \right) \quad (\text{D.11})$$

for the integrand and carrying out the integration, eq. (D.9) results in

$$\begin{aligned} & \delta(q^2 - m^2) \int_p Q_1 (f_{Vq} - f_{Vq-p}) \quad (\text{D.12}) \\ & \approx -\frac{\pi m_D^2 \delta(q^2 - m^2)}{2q_0} \int_{m_D}^T \frac{d|\mathbf{p}|}{(2\pi)^2 |\mathbf{p}|} \int_{-1}^1 dz \left(\frac{|\mathbf{q}|^2 (1 - z^2) I_T(z)}{2} + q_0^2 \right) \left[\frac{|\mathbf{q}| z^2}{q_0} \hat{q}_{\perp}^{\beta} + \left(\hat{q}_{\perp}^{\alpha} \hat{q}_{\perp}^{\beta} z^2 \right. \right. \\ & \quad \left. \left. - \frac{\hat{\Theta}_q^{\alpha\beta}}{2} (1 - z^2) \right) T \partial_{q^{\alpha}} \right] \partial_{q^{\beta}} f_{Vq} \\ & = -\frac{m_D^2 \delta(q^2 - m^2)}{8\pi q_0} \ln(1/g_c) \left[q_0 \left(j_1^T - \frac{q_0^2}{|\mathbf{q}|^2} j_2^T + j_1^L \right) q_{\perp}^{\beta} \partial_{q_{\perp}^{\beta}} - T \left(J_0^T - \frac{4q_0^2}{|\mathbf{q}|^2} j_1^T + \frac{3q_0^4}{|\mathbf{q}|^4} j_2^T + \frac{q_0^2}{|\mathbf{q}|^2} \right. \right. \\ & \quad \left. \left. \times (j_0^L - 3j_1^L) \right) \frac{q_{\perp}^{\beta} q_{\perp}^{\alpha}}{2} \partial_{q_{\perp}^{\alpha}} \partial_{q_{\perp}^{\beta}} - T \left(|\mathbf{q}|^2 \left(J_0^T - \frac{2q_0^2}{|\mathbf{q}|^2} j_1^T + \frac{q_0^4}{|\mathbf{q}|^4} j_2^T \right) + q_0^2 (j_0^L - j_1^L) \right) \frac{\eta^{\alpha\beta}}{2} \partial_{q^{\alpha}} \partial_{q^{\beta}} \right] f_{Vq}, \end{aligned}$$

where we take $u^{\beta} \partial_{q^{\beta}} f_{Vq} = 0$ and

$$\begin{aligned} j_0^T & \equiv \int_{-1}^1 dz \frac{1}{2 \left(1 - \frac{|\mathbf{q}|^2 z^2}{q_0^2} \right)} = \frac{q_0}{|\mathbf{q}|} \eta_q, \quad j_1^T \equiv \int_{-1}^1 dz \frac{\left(\frac{|\mathbf{q}| z}{q_0} \right)^2}{2 \left(1 - \frac{|\mathbf{q}|^2 z^2}{q_0^2} \right)} = \frac{q_0}{|\mathbf{q}|} \eta_q - 1, \\ j_2^T & \equiv \int_{-1}^1 dz \frac{\left(\frac{|\mathbf{q}| z}{q_0} \right)^4}{2 \left(1 - \frac{|\mathbf{q}|^2 z^2}{q_0^2} \right)} = \frac{q_0}{|\mathbf{q}|} \eta_q - \frac{|\mathbf{q}|^2}{3q_0^2} - 1, \end{aligned} \quad (\text{D.13})$$

where $\eta_q = 2^{-1} \ln [(q_0 + |\mathbf{q}|)/(q_0 - |\mathbf{q}|)]$, and

$$j_0^L \equiv \int_{-1}^1 dz = 2, \quad j_1^L \equiv \int_{-1}^1 z^2 dz = \frac{2}{3}. \quad (\text{D.14})$$

By inputting explicit forms of $j_i^{L/T}$ into (D.12), we eventually obtain

$$\begin{aligned} & \delta(q^2 - m^2) \int_p Q_1(f_{Vq} - f_{Vq-p}) \\ &= -\frac{m_D^2 \delta(q^2 - m^2)}{8\pi} \ln(1/g_c) \left[\frac{E_q^2}{|\mathbf{q}|^2} \left(1 - \frac{m^2 \eta_q}{q_0 |\mathbf{q}|} \right) q_\perp^\beta + \frac{m^2 T}{2|\mathbf{q}|^3} \left(\left(1 - \frac{3E_q^2}{|\mathbf{q}|^2} \right) \eta_q + \frac{3E_q}{|\mathbf{q}|} \right) q_\perp^\alpha q_\perp^\beta \partial_{q_\perp^\alpha} \right. \\ & \quad \left. - \frac{E_q T}{2} \left(\left(3 - \frac{E_q^2}{|\mathbf{q}|^2} \right) + \frac{m^4}{E_q |\mathbf{q}|^3} \right) \eta^{\alpha\beta} \partial_{q_\perp^\alpha} \right] \partial_{q_\perp^\beta} f_{Vq}, \end{aligned} \quad (\text{D.15})$$

where we take $q_0 = E_q$.

Subsequently, we should also calculate the $\tilde{Q}_1 f_{Vq}$ term in eq. (4.17). While the same counting (D.7) is applied to this case, there is not the order-one suppression from the difference $f_{Vq} - f_{Vq-p}$. Nevertheless, a factor of g_{0p}^{-1} provides the same order of suppression instead. Therefore, maintaining the linear terms from the other factors as in the computation above, we find

$$\begin{aligned} & \delta(q^2 - m^2) \int_p \tilde{Q}_1 f_{Vq} \\ & \approx -\frac{\pi m_D^2 \delta(q^2 - m^2)}{2q_0} \int_{m_D}^T \frac{d|\mathbf{p}|}{(2\pi)^2 |\mathbf{p}|} \int_{-1}^1 dz \left(\frac{|\mathbf{q}|^2 I_T(1-z^2)}{2} + q_0^2 \right) \frac{1}{q_0} \left(1 - \frac{|\mathbf{q}|^2 z^2}{q_0^2} \right) f_{Vq} \\ &= -\frac{m_D^2 \delta(q^2 - m^2)}{8\pi q_0} \ln(1/g_c) \left[q_0 (j_0^L - j_1^T + j_2^T) - \frac{|\mathbf{q}|^2}{q_0} (j_1^L - j_0^T + j_1^T) \right] f_{Vq}, \end{aligned} \quad (\text{D.16})$$

which yields

$$\begin{aligned} & \delta(q^2 - m^2) \int_p \tilde{Q}_1 f_{Vq} (1 - f_{Vq-p}) \\ & \approx -\frac{m_D^2 \delta(q^2 - m^2)}{4\pi} \ln(1/g_c) \left(f_{Vq} (1 - f_{Vq}) - \frac{q_0^2}{|\mathbf{q}|^2} \left(1 - \frac{\eta_q m^2}{q_0 |\mathbf{q}|} \right) f_{Vq} q_\perp^\beta \partial_{q_\perp^\beta} f_{Vq} \right). \end{aligned} \quad (\text{D.17})$$

Accordingly, the SKE results in the form given in eq. (4.20).

E Derivation of the spin diffusion for AKE

To evaluate the collision term in AKE, we implement the same strategy as in the SKE.

E.1 Half of the terms: $[Q_2^{\mu\nu} \tilde{a}_{q'\nu} - Q_1 \tilde{a}_q^\mu]$

For convenience, we first apply the following decompositions to two of the four terms in eq. (4.18):

$$Q_2^{\mu\nu} \tilde{a}_{q'\nu} - Q_1 \tilde{a}_q^\mu = 2\pi \delta(q'^2 - m^2) (Q_{\text{cl}(a)}^\mu + Q_{\text{cl}(b)}^\mu + Q_{\text{cl}(c)}^\mu), \quad (\text{E.1})$$

where

$$Q_{\text{cl}(a)}^\mu \equiv 2(q'^\rho G_{\rho\nu}^< q^\nu) (\tilde{a}_{q'}^\mu - \tilde{a}_q^\mu) + G_\rho^< (p \cdot q' \tilde{a}_q^\mu + p \cdot q \tilde{a}_{q'}^\mu) = g_{0p} \frac{p_0}{T} \left(\hat{Q}_{\text{cl}(a)}^{T\mu} + \hat{Q}_{\text{cl}(a)}^{L\mu} \right), \quad (\text{E.2a})$$

$$Q_{\text{cl}(b)}^\mu \equiv -(G_\rho^< p^\mu + 2q'_\rho G^{<\rho\mu}) q \cdot \tilde{a}_{q'} = g_{0p} \frac{p_0}{T} \left(\hat{Q}_{\text{cl}(b)}^{T\mu} + \hat{Q}_{\text{cl}(b)}^{L\mu} \right), \quad (\text{E.2b})$$

$$Q_{\text{cl}(c)}^\mu \equiv 2(p^\mu G_\rho^< q^\rho - q \cdot p G^{<\nu\mu}) \tilde{a}_{q'\nu} = g_{0p} \frac{p_0}{T} \left(\hat{Q}_{\text{cl}(c)}^{T\mu} + \hat{Q}_{\text{cl}(c)}^{L\mu} \right). \quad (\text{E.2c})$$

The remaining two terms will be addressed in section E.2. We utilized $q' \cdot a_{q'} = (q'^2 - m^2)f_{Aq'} = 0$ from the on-shell condition. On the rightmost side, the terms proportional to the transverse and longitudinal spectral functions are denoted as $\hat{Q}_{\text{cl}}^{T\mu}$ and $\hat{Q}_{\text{cl}}^{L\mu}$, respectively. Note that, similar to f_{Vq-p} in the SKE, the expansion of \tilde{a}_{q-p}^μ give rise to the factors of p for the small momentum transfer limit. Recalling the order counting in eq. (D.7), we thus maintain the terms in $\hat{Q}_{\text{cl}}^{T,L\mu}$ up to $\mathcal{O}(|\mathbf{p}|^2)$ in the following. For $Q_{\text{cl}(a)}^\mu$, we have

$$\begin{aligned}\hat{Q}_{\text{cl}(a)}^{T\mu} &= 2\hat{\rho}_T \left[|\mathbf{q}|^2(1-z^2)(\tilde{a}_{q-p}^\mu - \tilde{a}_q^\mu) - \left((p_0q_0 - |\mathbf{p}||\mathbf{q}|z)(\tilde{a}_{q-p}^\mu + \tilde{a}_q^\mu) - (p_0^2 - |\mathbf{p}|^2)\tilde{a}_{q-p}^\mu \right) \right] \\ &\approx 2\hat{\rho}_T \left[\left(2(|\mathbf{p}||\mathbf{q}|z - p_0q_0) + p_0^2 - |\mathbf{p}|^2 \right) \tilde{a}_q^\mu - \left(|\mathbf{q}|^2(1-z^2) - q_0p_0 + |\mathbf{p}||\mathbf{q}|z \right) p^\beta \partial_{q^\beta} \tilde{a}_q^\mu \right. \\ &\quad \left. + |\mathbf{q}|^2(1-z^2) \frac{p^\alpha p^\beta}{2} \partial_{q^\beta} \partial_{q^\alpha} \tilde{a}_q^\mu + \mathcal{O}(|\mathbf{p}|^3) \right],\end{aligned}\quad (\text{E.3a})$$

$$\begin{aligned}\hat{Q}_{\text{cl}(a)}^{L\mu} &= \hat{\rho}_L \left[2(q_0^2 - q_0p_0)(\tilde{a}_{q-p}^\mu - \tilde{a}_q^\mu) + \left((p_0q_0 - |\mathbf{p}||\mathbf{q}|z)(\tilde{a}_{q-p}^\mu + \tilde{a}_q^\mu) - (p_0^2 - |\mathbf{p}|^2)\tilde{a}_q^\mu \right) \right] \\ &\approx \hat{\rho}_L \left[\left(|\mathbf{p}|^2 - p_0^2 - 2(|\mathbf{p}||\mathbf{q}|z - p_0q_0) \right) \tilde{a}_q^\mu - \left(2q_0^2 - q_0p_0 - |\mathbf{p}||\mathbf{q}|z \right) p^\beta \partial_{q^\beta} \tilde{a}_q^\mu \right. \\ &\quad \left. + q_0^2 p^\alpha p^\beta \partial_{q^\alpha} \partial_{q^\beta} \tilde{a}_q^\mu + \mathcal{O}(|\mathbf{p}|^3) \right].\end{aligned}\quad (\text{E.3b})$$

As for $Q_{\text{cl}(b)}^\mu$, we have

$$\begin{aligned}\hat{Q}_{\text{cl}(b)}^{T\mu} &= 2\hat{\rho}_T (p^\mu + q_\perp^\mu - |\mathbf{q}|\hat{p}_\perp^\mu z) p \cdot \tilde{a}_{q-p} \\ &\approx 2\hat{\rho}_T \left[(p^\mu + q_\perp^\mu - |\mathbf{q}|\hat{p}_\perp^\mu z) p^\nu \tilde{a}_{\nu q} + (|\mathbf{q}|\hat{p}_\perp^\mu z - q_\perp^\mu) p^\nu p^\rho \partial_{q^\rho} \tilde{a}_{\nu q} + \mathcal{O}(|\mathbf{p}|^3) \right], \\ \hat{Q}_{\text{cl}(b)}^{L\mu} &= \hat{\rho}_L \left(2(p_0 - q_0)u^\mu - p^\mu \right) p \cdot \tilde{a}_{q-p} \\ &\approx \hat{\rho}_L \left[(p_0u^\mu - p_\perp^\mu - 2q_0u^\mu) p^\nu \tilde{a}_{q\nu} + 2q_0u^\mu p^\nu p^\rho \partial_{q^\rho} \tilde{a}_{q\nu} + \mathcal{O}(|\mathbf{p}|^3) \right].\end{aligned}\quad (\text{E.4a})$$

Here, we drop the terms proportional to $q \cdot a_q = (q^2 - m^2)f_{Aq}$ based on the on-shell condition. Finally, for $Q_{\text{cl}(c)}^\mu$, we have

$$\begin{aligned}\hat{Q}_{\text{cl}(c)}^{T\mu} &= 2\hat{\rho}_T \left(p^\mu (|\mathbf{q}|z\hat{p}_\perp^\nu - q_\perp^\nu) + (q_0p_0 - |\mathbf{q}||\mathbf{p}|z)(\Theta^{\mu\nu} + \hat{p}_\perp^\mu \hat{p}_\perp^\nu) \right) \tilde{a}_{q-p\nu} \\ &\approx 2\hat{\rho}_T \left((p_0u^\mu + p_\perp^\mu) (|\mathbf{q}|z\hat{p}_\perp^\nu - q_\perp^\nu) + (q_0p_0 - |\mathbf{q}||\mathbf{p}|z)(\Theta^{\mu\nu} + \hat{p}_\perp^\mu \hat{p}_\perp^\nu) \right) \\ &\quad \times \left(\tilde{a}_{q\nu} - p^\beta \partial_{q^\beta} \tilde{a}_{q\nu} + \mathcal{O}(|\mathbf{p}|^2) \right),\end{aligned}\quad (\text{E.5a})$$

$$\begin{aligned}\hat{Q}_{\text{cl}(c)}^{L\mu} &= \hat{\rho}_L \left(2q_0u^\nu p^\mu - 2(q_0p_0 - |\mathbf{q}||\mathbf{p}|z)u^\mu u^\nu \right) \tilde{a}_{q-p\nu} \\ &\approx \hat{\rho}_L \left(2q_0u^\nu p_\perp^\mu + 2|\mathbf{q}||\mathbf{p}|z u^\mu u^\nu \right) \left(\tilde{a}_{q\nu} - p^\beta \partial_{q^\beta} \tilde{a}_{q\nu} + \mathcal{O}(|\mathbf{p}|^2) \right).\end{aligned}\quad (\text{E.5b})$$

In the following, we evaluate those terms one by one. As in the case of SKE, we first retrieve the integrals for $|\mathbf{p}|$ that give rise to the logarithm in the following subsections. The angle integrals, with respect to z , will be performed in section E.3 afterwards.

E.1.1 Evaluating $Q_{\text{cl}(a)}^\mu$

Combining with eqs. (D.8) and (D.5), we find

$$\begin{aligned} & \delta(q^2 - m^2) \int_p 2\pi\delta((q-p)^2 - m^2) Q_{\text{cl}(a)}^\mu \\ & \approx \frac{\pi T m_D^2 \delta(q^2 - m^2)}{2q_0} \int_{m_D}^T \frac{d|\mathbf{p}|}{(2\pi)^2 |\mathbf{p}|} \int_{-1}^1 dz \left[\left(\frac{q_0 z^2}{T} + \frac{I_T z^2 (1-z^2) |\mathbf{q}|^2}{2q_0 T} \right) q_\perp^\beta \partial_{q_\perp^\beta} \right. \\ & \quad \left. - \left(q_0^2 + \frac{I_T |\mathbf{q}|^2 (1-z^2)}{2} \right) \left((1-z^2) \eta^{\alpha\beta} + (1-3z^2) \hat{q}_\perp^\alpha \hat{q}_\perp^\beta \right) \frac{\partial_{q_\perp^\alpha} \partial_{q_\perp^\beta}}{2} \right] \tilde{a}_q^\mu, \end{aligned} \quad (\text{E.6})$$

where we take $u^\beta \partial_{q^\beta} \tilde{a}_q^\mu = 0$ by imposing $q_0 = E_q = \sqrt{|\mathbf{q}|^2 + m^2}$ upon \tilde{a}_q^μ in advance and utilize

$$\left(\hat{q}_\perp^\alpha \hat{q}_\perp^\beta z^2 - \frac{\hat{\Theta}_q^{\alpha\beta}}{2} (1-z^2) \right) = -\frac{\Theta^{\alpha\beta}}{2} (1-z^2) - \frac{(1-3z^2)}{2} \hat{q}_\perp^\alpha \hat{q}_\perp^\beta. \quad (\text{E.7})$$

E.1.2 Evaluating $Q_{\text{cl}(b)}^\mu$

Next, we evaluate the term proportional to $Q_{\text{cl}(b)}^\mu$:

$$\delta(q^2 - m^2) \int_p 2\pi\delta((q-p)^2 - m^2) Q_{\text{cl}(b)}^\mu \approx \frac{\pi T m_D^2 \delta(q^2 - m^2)}{2q_0} \int \frac{d^3 \mathbf{p}}{(2\pi)^3 |\mathbf{p}|^5} (\tilde{\mathcal{K}}_{\text{cl}(b)}^{T\mu} + \tilde{\mathcal{K}}_{\text{cl}(b)}^{L\mu}), \quad (\text{E.8})$$

where

$$\begin{aligned} \tilde{\mathcal{K}}_{\text{cl}(b)}^{T\mu} = & I_T \left[|\mathbf{p}| (q_\perp^\mu - \hat{p}_\perp^\mu |\mathbf{q}| z) \left(\frac{|\mathbf{q}|}{q_0} u^\nu z + \hat{p}_\perp^\nu \right) + \frac{|\mathbf{p}|^2}{2q_0^3 T} \left(|\mathbf{q}|^2 z^2 (u^\nu ((T-q_0)q_\perp^\mu + 2q_0 T u^\mu) + \hat{p}_\perp^\mu \hat{p}_\perp^\nu q_0^2) \right. \right. \\ & \left. \left. + q_0^2 T (2\hat{p}_\perp^\mu \hat{p}_\perp^\nu q_0 - q_\perp^\mu u^\nu) + |\mathbf{q}| q_0^2 z (3\hat{p}_\perp^\mu T u^\nu - \hat{p}_\perp^\nu q_\perp^\mu + 2\hat{p}_\perp^\nu T u^\mu) + |\mathbf{q}|^3 \hat{p}_\perp^\mu u^\nu z^3 (q_0 - T) \right) \right. \\ & \left. - (q_\perp^\mu - \hat{p}_\perp^\mu |\mathbf{q}| z) \left(\frac{|\mathbf{q}|}{q_0} u^\nu z + \hat{p}_\perp^\nu \right) \left(\frac{|\mathbf{q}|}{q_0} u^\rho z + \hat{p}_\perp^\rho \right) |\mathbf{p}|^2 \partial_{q^\rho} + \mathcal{O}(|\mathbf{p}|^3) \right] \tilde{a}_{q\nu}, \end{aligned} \quad (\text{E.9a})$$

$$\begin{aligned} \tilde{\mathcal{K}}_{\text{cl}(b)}^{L\mu} = & -2\tilde{a}_{q\nu} u^\mu |\mathbf{p}| (\hat{p}_\perp^\nu q_0 + |\mathbf{q}| u^\nu z) + \frac{|\mathbf{p}|^2}{q_0^2 T} \left(u^\mu (u^\nu (|\mathbf{q}|^2 z^2 (q_0 - 2T) + q_0^2 T) + \hat{p}_\perp^\nu |\mathbf{q}| q_0 z (q_0 - T)) \right. \\ & \left. - \hat{p}_\perp^\mu q_0 T (|\mathbf{q}| u^\nu z + \hat{p}_\perp^\nu q_0) \right) \tilde{a}_{q\nu} + 2q_0 u^\mu \left(\frac{|\mathbf{q}|}{q_0} u^\nu z + \hat{p}_\perp^\nu \right) \left(\frac{|\mathbf{q}|}{q_0} u^\rho z + \hat{p}_\perp^\rho \right) |\mathbf{p}|^2 \partial_{q^\rho} \tilde{a}_{q\nu} + \mathcal{O}(|\mathbf{p}|^3). \end{aligned} \quad (\text{E.9b})$$

It is clear that $\mathcal{O}(|\mathbf{p}|)$ terms in $\tilde{\mathcal{K}}_{\text{cl}(b)}^{T/L\mu}$ can be dropped by symmetry of the integration. We should further employ the useful decomposition and replacements in eqs. (D.10), (D.11), and the following relation in the integral,

$$\begin{aligned} z p_\perp^\nu p_\perp^\mu p_\perp^\rho & = z |\mathbf{p}|^3 (\hat{q}_\perp^\nu z + \hat{p}_T^\nu) (\hat{q}_\perp^\mu z + \hat{p}_T^\mu) (\hat{q}_\perp^\rho z + \hat{p}_T^\rho) \\ & \rightarrow |\mathbf{p}|^3 \left(z^4 \hat{q}_\perp^\nu \hat{q}_\perp^\mu \hat{q}_\perp^\rho - \frac{z^2 (1-z^2)}{2} \left(\hat{q}_\perp^\nu \hat{\Theta}_q^{\mu\rho} + \hat{q}_\perp^\mu \hat{\Theta}_q^{\nu\rho} + \hat{q}_\perp^\rho \hat{\Theta}_q^{\mu\nu} \right) \right), \end{aligned} \quad (\text{E.10})$$

where $\hat{p}_T^\mu = \hat{\Theta}_q^{\mu\nu} \hat{p}_{\perp\nu}$ and we drop terms with odd powers of z . Dropping the vanishing terms by symmetry in eq. (E.8), we obtain

$$\begin{aligned} & \delta(q^2 - m^2) \int_p 2\pi\delta((q-p)^2 - m^2) Q_{\text{cl}(b)}^\mu \\ & \approx \frac{\pi T m_D^2 \delta(q^2 - m^2)}{2q_0} \int_{m_D}^T \frac{d|\mathbf{p}|}{(2\pi)^2 |\mathbf{p}|} \int_{-1}^1 dz \left(\mathcal{K}_{\text{cl}(b)}^{T\mu} + \mathcal{K}_{\text{cl}(b)}^{L\mu} \right), \end{aligned} \quad (\text{E.11})$$

where

$$\begin{aligned} \mathcal{K}_{\text{cl}(b)}^{T\mu} = I_T \left\{ \left[u^\mu z^2 \left(\frac{q_\perp^\nu}{q_0} + \frac{|\mathbf{q}|^2 u^\nu}{q_0^2} \right) - \frac{\hat{q}_\perp^\mu}{4q_0^3 T} \left(q_0^2 \hat{q}_\perp^\nu \left(3|\mathbf{q}|^2 z^2 (1-z^2) + 2q_0 T (1-3z^2) \right) \right. \right. \right. \\ \left. \left. \left. + 2u^\nu |\mathbf{q}| \left(z^2 (1-z^2) |\mathbf{q}|^2 (q_0 - T) + q_0^2 T (1-3z^2) \right) \right) - \frac{\Theta^{\mu\nu}}{4} (1-z^2) \left(2 + \frac{|\mathbf{q}|^2 z^2}{q_0 T} \right) \right] \tilde{a}_{q\nu} \\ + \frac{1}{2} \left[q_\perp^\mu (1-z^2) \left(\hat{q}_\perp^\nu \hat{q}_\perp^\rho (1-5z^2) + (1-z^2) \Theta^{\rho\nu} - 3 \frac{q_\perp^\rho}{q_0} u^\nu z^2 \right) \right. \\ \left. - z^2 (1-z^2) \left(\Theta^{\mu\nu} q_\perp^\rho + \Theta^{\rho\mu} \left(q_\perp^\nu + \frac{|\mathbf{q}|^2}{q_0} u^\nu \right) \right) \right] \partial_{q_\perp^\rho} \tilde{a}_{q\nu} \right\} \end{aligned} \quad (\text{E.12})$$

and

$$\begin{aligned} \mathcal{K}_{\text{cl}(b)}^{L\mu} = u^\mu \left[u \cdot \tilde{a}_q \left(1 + \frac{|\mathbf{q}|^2 z^2 (q_0 - 2T)}{q_0^2 T} \right) + \hat{q}_\perp \cdot \tilde{a}_q \frac{|\mathbf{q}| z^2 (q_0 - T)}{q_0 T} \right] + \hat{q}_\perp^\mu \left[\hat{q}_\perp \cdot \tilde{a}_q \frac{(1-3z^2)}{2} \right. \\ \left. - u \cdot \tilde{a}_q \frac{|\mathbf{q}| z^2}{q_0} \right] + \Theta^{\mu\nu} \tilde{a}_\nu \frac{(1-z^2)}{2} + u^\mu \left(2z^2 q_\perp^\rho u^\nu - \hat{q}_\perp^\rho \hat{q}_\perp^\nu q_0 (1-3z^2) - \Theta^{\rho\nu} q_0 (1-z^2) \right) \partial_{q_\perp^\rho} \tilde{a}_{q\nu}. \end{aligned} \quad (\text{E.13})$$

Since $\delta(q^2 - m^2) q \cdot \tilde{a}_q = \delta(q^2 - m^2) (E_q u \cdot \tilde{a}_q + q_\perp \cdot \tilde{a}_q) = 0$, we can use $u \cdot \tilde{a}_q = -q_\perp \cdot \tilde{a}_q / q_0$ to further simplify $\mathcal{K}_{\text{cl}(b)}^{T/L\mu}$, which results in

$$\begin{aligned} \mathcal{K}_{\text{cl}(b)}^{T\mu} = I_T \left\{ \hat{q}_\perp \cdot \tilde{a}_q \left[u^\mu \frac{m^2 |\mathbf{q}| z^2}{q_0^3} - \frac{\hat{q}_\perp^\mu}{2} \left(z^2 (1-z^2) \left(1 + \frac{q_0}{2T} + \frac{m^4}{q_0^4} \left(1 - \frac{q_0}{T} \right) \right) + \frac{m^2}{q_0^2} \left(1 - 5z^2 + 2z^4 \right. \right. \right. \right. \\ \left. \left. \left. + \frac{q_0}{2T} z^2 (1-z^2) \right) \right] - \frac{\Theta^{\mu\nu} \tilde{a}_{q\nu}}{2} (1-z^2) \left(1 + \frac{|\mathbf{q}|^2 z^2}{2q_0 T} \right) + \frac{1}{2} (1-z^2) \left[\hat{q}_\perp^\mu \left(|\mathbf{q}| \hat{q}_\perp^\rho \hat{q}_\perp^\nu (1-5z^2) \right. \right. \right. \\ \left. \left. \left. + |\mathbf{q}| \Theta^{\rho\nu} (1-z^2) - \frac{3|\mathbf{q}|^2 z^2}{q_0} \hat{q}_\perp^\rho u^\nu \right) - |\mathbf{q}| z^2 \Theta^{\mu\nu} \hat{q}_\perp^\rho - \left(|\mathbf{q}| \hat{q}_\perp^\nu + \frac{|\mathbf{q}|^2}{q_0} u^\nu \right) z^2 \Theta^{\rho\mu} \right] \partial_{q_\perp^\rho} \tilde{a}_{q\nu} \right\}, \end{aligned} \quad (\text{E.14})$$

and

$$\begin{aligned} \mathcal{K}_{\text{cl}(b)}^{L\mu} = u^\mu \left[\frac{m^2 |\mathbf{q}| z^2}{q_0^2 T} - \frac{|\mathbf{q}|}{q_0} \left(1 - \frac{|\mathbf{q}|^2 z^2}{q_0^2} + \frac{m^2 z^2}{q_0^2} \right) \right] \hat{q}_\perp \cdot \tilde{a}_q + \hat{q}_\perp^\mu \left[\frac{(1-z^2)}{2} - \frac{m^2 z^2}{q_0^2} \right] \hat{q}_\perp \cdot \tilde{a}_q \\ + \Theta^{\mu\nu} \tilde{a}_\nu \frac{(1-z^2)}{2} + u^\mu \left(2z^2 q_\perp^\rho u^\nu - \hat{q}_\perp^\rho \hat{q}_\perp^\nu q_0 (1-3z^2) - \Theta^{\rho\nu} q_0 (1-z^2) \right) \partial_{q_\perp^\rho} \tilde{a}_{q\nu}. \end{aligned} \quad (\text{E.15})$$

E.1.3 Evaluating $Q_{\text{cl}(c)}^\mu$

In the same way, we should cope with the other term

$$\delta(q^2 - m^2) \int_p 2\pi\delta((q-p)^2 - m^2) Q_{\text{cl}(c)}^\mu \approx \frac{\pi T m_D^2 \delta(q^2 - m^2)}{2q_0} \int \frac{d^3\mathbf{p}}{(2\pi)^3} \frac{1}{|\mathbf{p}|^5} \left(\tilde{\mathcal{K}}_{\text{cl}(c)}^{T\mu} + \tilde{\mathcal{K}}_{\text{cl}(c)}^{L\mu} \right), \quad (\text{E.16})$$

where

$$\begin{aligned} \tilde{\mathcal{K}}_{\text{cl}(c)}^{T\mu} = I_T \left[-|\mathbf{p}|(q_\perp^\nu - \hat{p}_\perp^\nu |\mathbf{q}|z) \left(\frac{|\mathbf{q}|}{q_0} u^\mu z + \hat{p}_\perp^\mu \right) + \frac{|\mathbf{p}|^2}{2q_0^3 T} \left(|\mathbf{q}|q_0 z (q_\perp^\nu - \hat{p}_\perp^\nu |\mathbf{q}|z) (\hat{p}_\perp^\mu q_0 + |\mathbf{q}|u^\mu z) \right. \right. \\ \left. \left. - T(q_0^2 - |\mathbf{q}|^2 z^2) (\hat{p}_\perp^\mu \hat{p}_\perp^\nu q_0 + \hat{p}_\perp^\nu |\mathbf{q}|u^\mu z + \Theta^{\mu\nu} q_0 - q_\perp^\nu u^\mu) \right) \right. \\ \left. + |\mathbf{p}|^2 (q_\perp^\nu - \hat{p}_\perp^\nu |\mathbf{q}|z) \left(\frac{|\mathbf{q}|}{q_0} u^\mu z + \hat{p}_\perp^\mu \right) \left(\frac{|\mathbf{q}|}{q_0} u^\rho z + \hat{p}_\perp^\rho \right) \partial_{q^\rho} + \mathcal{O}(|\mathbf{p}|^3) \right] \tilde{a}_{q\nu}, \quad (\text{E.17}) \end{aligned}$$

and

$$\begin{aligned} \tilde{\mathcal{K}}_{\text{cl}(c)}^{L\mu} = 2|\mathbf{p}|(\hat{p}_\perp^\mu q_0 + |\mathbf{q}|u^\mu z) u \cdot \tilde{a}_q - \frac{|\mathbf{p}|^2}{q_0^2 T} |\mathbf{q}|z (q_0 - 2T) (\hat{p}_\perp^\mu q_0 + |\mathbf{q}|u^\mu z) u \cdot \tilde{a}_q \\ - 2|\mathbf{p}|^2 (q_0 \hat{p}_\perp^\mu + |\mathbf{q}|z u^\mu) \left(\frac{|\mathbf{q}|}{q_0} u^\rho z + \hat{p}_\perp^\rho \right) \partial_{q^\rho} u \cdot \tilde{a}_q + \mathcal{O}(|\mathbf{p}|^3). \quad (\text{E.18}) \end{aligned}$$

As above, the $\mathcal{O}(|\mathbf{p}|)$ terms in $\tilde{\mathcal{K}}_{\text{cl}(c)}^{T/L\mu}$ do not contribute to the integral. Now, by employing eqs. (D.10), (D.11), and (E.10), we find

$$\delta(q^2 - m^2) \int_p 2\pi\delta((q-p)^2 - m^2) Q_{\text{cl}(c)}^\mu \approx \frac{\pi T m_D^2 \delta(q^2 - m^2)}{2q_0} \int_{m_D}^T \frac{d|\mathbf{p}|}{(2\pi)^2 |\mathbf{p}|} \int_{-1}^1 dz \left(\mathcal{K}_{\text{cl}(c)}^{T\mu} + \mathcal{K}_{\text{cl}(c)}^{L\mu} \right), \quad (\text{E.19})$$

where

$$\begin{aligned} \mathcal{K}_{\text{cl}(c)}^{T\mu} = I_T \left[\frac{u^\mu q_\perp^\nu}{2q_0} (1-z^2) \left(1 + \frac{|\mathbf{q}|^2 z^2}{q_0^2 T} (q_0 - T) \right) - \frac{\Theta^{\mu\nu}}{4} \left((1+z^2) \left(1 - \frac{|\mathbf{q}|^2 z^2}{q_0^2} \right) - \frac{|\mathbf{q}|^2}{q_0 T} z^2 (1-z^2) \right) \right. \\ \left. + \frac{\hat{q}_\perp^\mu \hat{q}_\perp^\nu}{4} \left((1-3z^2) \left(1 - \frac{|\mathbf{q}|^2 z^2}{q_0^2} \right) + \frac{3|\mathbf{q}|^2}{q_0 T} z^2 (1-z^2) \right) + \frac{u^\mu |\mathbf{q}| z^2 (1-z^2)}{2q_0} (3\hat{q}_\perp^\rho \hat{q}_\perp^\nu + |\mathbf{q}| \Theta^{\rho\nu}) \partial_{q_\perp^\rho} \right. \\ \left. + \frac{(1-z^2)}{2} \left(z^2 \Theta^{\mu\nu} q_\perp^\rho + q_\perp^\mu ((5z^2 - 1)\hat{q}_\perp^\nu \hat{q}_\perp^\rho + z^2 \Theta^{\rho\nu}) - (1-z^2) q_\perp^\nu \Theta^{\rho\mu} \right) \partial_{q_\perp^\rho} \right] \tilde{a}_{q\nu}, \quad (\text{E.20a}) \end{aligned}$$

$$\begin{aligned} \mathcal{K}_{\text{cl}(c)}^{L\mu} = -u \cdot \tilde{a}_q \frac{(q_0 - 2T) |\mathbf{q}| z^2}{q_0 T} \left(\frac{|\mathbf{q}|}{q_0} u^\mu + \hat{q}_\perp^\mu \right) + q_0 \left(\hat{q}_\perp^\mu \hat{q}_\perp^\rho (1-3z^2) + \Theta^{\mu\rho} (1-z^2) \right. \\ \left. - 2q_\perp^\rho q_0^{-1} u^\mu z^2 \right) \partial_{q_\perp^\rho} (u \cdot \tilde{a}_q). \quad (\text{E.20b}) \end{aligned}$$

E.2 Rest of the terms: $[Q_2^{\mu\nu} f_{Vq} \tilde{a}_{q'\nu} - Q_1(1 - f_{Vq'}) \tilde{a}_q^\mu]$

As we have evaluated two of the four terms in eq. (4.18), the remaining two terms are given as

$$\mathcal{C}_2 \equiv \delta(q^2 - m^2) \int_p g_{0p}^{-1} \left[Q_2^{\mu\nu} f_{Vq} \tilde{a}_{q'\nu} - Q_1(1 - f_{Vq'}) \tilde{a}_q^\mu \right]. \quad (\text{E.21})$$

While Q_1 has been given in the computation of the SKE, we now have the tensor part

$$\begin{aligned}
 Q_2^{\mu\nu} \tilde{a}_{q'\nu} = & -2\pi\delta(q^2 - m^2)g_{0p} \left\{ p^\mu \left[\rho_L (p^\nu - 2q_0 u^\nu) + 2\rho_T (q_\perp^\nu - |\mathbf{q}|z\hat{p}_\perp^\nu - p^\nu) \right] \tilde{a}_{q'\nu} \right. \\
 & - \tilde{a}_{q'}^\mu \left[\rho_L (q \cdot p + 2(q_0^2 - q_0 p_0)) + 2\rho_T (|\mathbf{q}|^2(1 - z^2) - q \cdot p) \right] \\
 & + 2u^\mu \rho_L (q \cdot p u^\nu + (q_0 - p_0)p^\nu) \tilde{a}_{q'\nu} - 2\tilde{a}_{q'\perp}^\mu q \cdot p \rho_T - 2q_\perp^\mu p \cdot \tilde{a}_{q'} \rho_T \\
 & \left. - 2\hat{p}_\perp^\mu (q \cdot p \hat{p}_\perp^\nu - |\mathbf{q}|z p^\nu) \rho_T \tilde{a}_{q'\nu} \right\}. \tag{E.22}
 \end{aligned}$$

Combining the two terms, it is found

$$C_2 \approx \frac{\pi m_D^2 \delta(q^2 - m^2)}{2q_0} \int \frac{d^3\mathbf{p}}{(2\pi)^3} \frac{1}{|\mathbf{p}|^5} \left(\tilde{\mathcal{K}}_{\text{cl}(d)}^{T\mu} + \tilde{\mathcal{K}}_{\text{cl}(d)}^{L\mu} \right), \tag{E.23}$$

where the integrand is given as

$$\begin{aligned}
 \tilde{\mathcal{K}}_{\text{cl}(d)}^{T\mu} = & -I_T \left[\frac{|\mathbf{q}|^3 |\mathbf{p}|}{q_0 T} z(1 - z^2)(1 - 2f_{Vq}) \tilde{a}_q^\mu + \frac{|\mathbf{q}|^2 |\mathbf{p}|^2}{2q_0^3 T} \left(\tilde{a}_q^\mu (z^2 - 1) (q_0^2 (1 - 2z|\mathbf{q}|\hat{p}_\perp^\rho (\partial_{q^\rho} f_{Vq}) - 2f_{Vq}) \right. \right. \\
 & + (2f_{Vq} - 1)|\mathbf{q}|^2 z^2) + 2f_{Vq} q_0 z \left(\tilde{a}_{q\nu} (\hat{p}_\perp^\mu \hat{q}_\perp^\nu q_0 + \hat{p}_\perp^\mu |\mathbf{q}|u^\nu z^2 - \hat{p}_\perp^\nu (\hat{q}_\perp^\mu q_0 + |\mathbf{q}|u^\mu z^2) \right. \\
 & \left. \left. - \hat{q}_\perp^\mu |\mathbf{q}|u^\nu z + \hat{q}_\perp^\nu |\mathbf{q}|u^\mu z) - |\mathbf{q}|\hat{p}_\perp^\rho (\partial_{q^\rho} \tilde{a}_q^\mu) q_0 (z^2 - 1) \right) \right] + \mathcal{O}(|\mathbf{p}|^3), \tag{E.24a}
 \end{aligned}$$

$$\begin{aligned}
 \tilde{\mathcal{K}}_{\text{cl}(d)}^{L\mu} = & - \left[\frac{2|\mathbf{q}||\mathbf{p}|q_0 z}{T} (1 - 2f_{Vq}) \tilde{a}_q^\mu + \frac{|\mathbf{p}|^2}{q_0^2 T} \left(\tilde{a}_q^\mu (q_0^3 (2z|\mathbf{q}|\hat{p}_\perp^\rho (\partial_{q^\rho} f_{Vq}) + 2f_{Vq} - 1) + (1 - 2f_{Vq})|\mathbf{q}|^2 q_0 z^2) \right. \right. \\
 & \left. \left. + 2f_{Vq} |\mathbf{q}|q_0^2 z \left(\hat{p}_\perp^\nu \tilde{a}_{q\nu} u^\mu + q_0 \hat{p}_\perp^\rho (\partial_{q^\rho} \tilde{a}_q^\mu) - u^\nu \tilde{a}_{q\nu} \hat{p}_\perp^\mu \right) \right) \right] + \mathcal{O}(|\mathbf{p}|^3). \tag{E.24b}
 \end{aligned}$$

Here, we utilize $u \cdot \tilde{a}_q = -q_\perp \cdot \tilde{a}_q / q_0$ in computations as above. Carrying out the integration, we obtain

$$\begin{aligned}
 C_2 \approx & \frac{m_D^2 T \delta(q^2 - m^2)}{8\pi E_q} \ln(1/g_c) \left(\tilde{a}_q^\mu \tilde{\mathcal{Q}}_{\text{cl}}^{(1)} + u^\mu \tilde{\mathcal{Q}}_{\text{cl}}^{(2)} + \hat{q}_\perp^\mu \tilde{\mathcal{Q}}_{\text{cl}}^{(3)} + \tilde{\mathcal{Q}}_{\text{cl}}^{(4)} \hat{q}_\perp^\nu \partial_{q_\perp^\mu} \tilde{a}_{q\nu} \right. \\
 & \left. + \tilde{\mathcal{Q}}_{\text{cl}}^{(5)} \hat{q}_\perp^\nu \partial_{q_\perp^\nu} \tilde{a}_q^\mu + \tilde{\mathcal{Q}}_{\text{cl}}^{(6)} \eta^{\nu\rho} \partial_{q_\perp^\nu} \partial_{q_\perp^\rho} \tilde{a}_q^\mu + \tilde{\mathcal{Q}}_{\text{cl}}^{(7)} \hat{q}_\perp^\nu \hat{q}_\perp^\rho \partial_{q_\perp^\nu} \partial_{q_\perp^\rho} \tilde{a}_q^\mu \right), \tag{E.25}
 \end{aligned}$$

where

$$\begin{aligned}
 \tilde{\mathcal{Q}}_{\text{cl}}^{(1)} = & \frac{1}{|\mathbf{q}|q_0 T} \left[|\mathbf{q}|q_0^2 \{ (1 - 2f_{Vq})(j_0^L - j_1^T + j_2^T) - 2|\mathbf{q}|(\hat{q}_\perp^\nu \partial_{q_\perp^\nu} f_{Vq})(j_1^L + j_1^T) \} \right. \\
 & \left. - (2f_{Vq} - 1)|\mathbf{q}|^3 (j_0^T - j_1^L - j_1^T) + 2(\hat{q}_\perp^\nu \partial_{q_\perp^\nu} f_{Vq}) j_2^T q_0^4 \right], \tag{E.26a}
 \end{aligned}$$

$$\tilde{\mathcal{Q}}_{\text{cl}}^{(2)} = -2 \left((j_1^L + j_1^T) |\mathbf{q}|^2 - j_2^T q_0^2 \right) \frac{f_{Vq}}{|\mathbf{q}|T} \hat{q}_\perp \cdot \tilde{a}_q, \tag{E.26b}$$

$$\tilde{\mathcal{Q}}_{\text{cl}}^{(3)} = -2 \left((j_1^L + j_1^T) |\mathbf{q}|^2 - j_2^T q_0^2 \right) \frac{f_{Vq}}{q_0 T} \hat{q}_\perp \cdot \tilde{a}_q, \tag{E.26c}$$

$$\tilde{\mathcal{Q}}_{\text{cl}}^{(5)} = 2 \left(\frac{q_0^2}{|\mathbf{q}|} j_2^T - |\mathbf{q}|(j_1^L + j_1^T) \right) \frac{q_0 f_{Vq}}{T}, \tag{E.26d}$$

$$\tilde{\mathcal{Q}}_{\text{cl}}^{(4)} = \tilde{\mathcal{Q}}_{\text{cl}}^{(6)} = \tilde{\mathcal{Q}}_{\text{cl}}^{(7)} = 0. \tag{E.26e}$$

E.3 Assembling all the pieces

Combining eqs. (E.6), (E.11), and (E.20a) and carrying out the integration, we acquire

$$\begin{aligned} \mathcal{C}_1 &\equiv \delta(q^2 - m^2) \int_p (Q_2^{\mu\nu} \tilde{a}_{q'\nu} - Q_1 \tilde{a}_q^\mu) \\ &\approx \frac{m_D^2 T \delta(q^2 - m^2)}{8\pi E_q} \ln(1/g_c) \left(\tilde{a}_q^\mu \mathcal{Q}_{\text{cl}}^{(1)} + u^\mu \mathcal{Q}_{\text{cl}}^{(2)} + \hat{q}_\perp^\mu \mathcal{Q}_{\text{cl}}^{(3)} + \mathcal{Q}_{\text{cl}}^{(4)} \hat{q}_\perp^\nu \partial_{q_\perp \mu} \tilde{a}_{q\nu} \right. \\ &\quad \left. + \mathcal{Q}_{\text{cl}}^{(5)} \hat{q}_\perp^\nu \partial_{q_\perp^\nu} \tilde{a}_q^\mu + \mathcal{Q}_{\text{cl}}^{(6)} \eta^{\nu\rho} \partial_{q_\perp^\nu} \partial_{q_\perp^\rho} \tilde{a}_q^\mu + \mathcal{Q}_{\text{cl}}^{(7)} \hat{q}_\perp^\nu \hat{q}_\perp^\rho \partial_{q_\perp^\nu} \partial_{q_\perp^\rho} \tilde{a}_q^\mu \right), \end{aligned} \quad (\text{E.27})$$

where

$$\mathcal{Q}_{\text{cl}}^{(1)} = \frac{1}{2} \left(j_1^L + 3j_1^T - j_0^L - 3j_0^T + \frac{q_0^2}{|\mathbf{q}|^2} (j_1^T - j_2^T) \right), \quad (\text{E.28a})$$

$$\begin{aligned} \mathcal{Q}_{\text{cl}}^{(2)} &= -\frac{1}{2} \left[2q_0 (\partial_{q_\perp^\nu} \tilde{a}_q^\nu) \left(j_0^L - (j_1^L + j_1^T) + j_2^T \frac{q_0^2}{|\mathbf{q}|^2} \right) + 2q_0 \hat{q}_\perp^\nu \hat{q}_\perp^\rho (\partial_{q_\perp^\rho} \tilde{a}_{q\nu}) \left(j_0^L - 3(j_1^L + j_1^T) + 3j_2^T \frac{q_0^2}{|\mathbf{q}|^2} \right) \right. \\ &\quad \left. - \hat{q}_\perp^\nu \tilde{a}_{q\nu} \frac{|\mathbf{q}|}{q_0} \left(j_1^L \frac{(2q_0 - T)}{T} - 3j_0^L - j_0^T - 2j_1^T \frac{q_0(m^2 - |\mathbf{q}|^2)}{|\mathbf{q}|^2 T} + j_1^T \frac{m^2(2q_0 + 3T)}{|\mathbf{q}|^2 T} \right. \right. \\ &\quad \left. \left. + j_2^T \frac{q_0^2(T - 2q_0)}{|\mathbf{q}|^2 T} \right) \right], \end{aligned} \quad (\text{E.28b})$$

$$\begin{aligned} \mathcal{Q}_{\text{cl}}^{(3)} &= -\frac{1}{2} \left[\frac{2}{|\mathbf{q}|} \left((q_0^2 j_1^T - |\mathbf{q}|^2 j_0^T) (\partial_{q_\perp^\nu} \tilde{a}_q^\nu) + \hat{q}_\perp^\nu \hat{q}_\perp^\rho (\partial_{q_\perp^\rho} \tilde{a}_{q\nu}) \left(j_0^L |\mathbf{q}|^2 + 3j_2^T q_0^2 - 3|\mathbf{q}|^2 (j_1^L + j_1^T) \right) \right) \right. \\ &\quad \left. - \hat{q}_\perp^\nu \tilde{a}_{q\nu} \left(\frac{|\mathbf{q}|^2}{q_0^2} (j_0^T - j_0^L) - \frac{m^2}{q_0^2} (j_0^L + j_0^T) + 3 \frac{m^2}{q_0^2} j_1^L + j_1^L \frac{|\mathbf{q}|^2}{q_0^2 T} (2q_0 - 3T) - 2j_1^T \frac{m^4}{q_0^2 |\mathbf{q}|^2} \right. \right. \\ &\quad \left. \left. + 5j_1^T \frac{m^2}{|\mathbf{q}|^2} + 2j_1^T \frac{|\mathbf{q}|^2}{q_0^2 T} (q_0 - 2T) + 2j_2^T \frac{m^2 q_0}{|\mathbf{q}|^2 T} - 6j_2^T \frac{m^2}{|\mathbf{q}|^2} - 2j_2^T \frac{q_0^3}{|\mathbf{q}|^2 T} + 3j_2^T \frac{q_0^2}{|\mathbf{q}|^2} \right) \right], \end{aligned} \quad (\text{E.28c})$$

$$\mathcal{Q}_{\text{cl}}^{(4)} = |\mathbf{q}| (j_1^L + j_1^T - j_0^L - j_0^T) + \frac{q_0^2}{|\mathbf{q}|} (j_1^T - j_2^T), \quad (\text{E.28d})$$

$$\mathcal{Q}_{\text{cl}}^{(5)} = \frac{q_0}{|\mathbf{q}| T} \left((j_1^L + j_1^T) |\mathbf{q}|^2 - j_2^T q_0^2 \right), \quad (\text{E.28e})$$

$$\mathcal{Q}_{\text{cl}}^{(6)} = -\frac{1}{2} \left(j_0^T |\mathbf{q}|^2 + (j_0^L - j_1^L - 2j_1^T) q_0^2 + j_2^T \frac{q_0^4}{|\mathbf{q}|^2} \right), \quad (\text{E.28f})$$

$$\mathcal{Q}_{\text{cl}}^{(7)} = -\frac{1}{2} \left(j_0^T |\mathbf{q}|^2 + (j_0^L - 3j_1^L - 4j_1^T) q_0^2 + 3j_2^T \frac{q_0^4}{|\mathbf{q}|^2} \right). \quad (\text{E.28g})$$

Note that we have implemented the following relations in computations,

$$\begin{aligned} \Theta^{\mu\nu} \tilde{a}_{q\nu} &= \tilde{a}_q^\mu - (\tilde{a}_q \cdot u) u^\mu = \tilde{a}_q^\mu + u^\mu \frac{|\mathbf{q}|}{E_q} \hat{q}_\perp \cdot \tilde{a}_q, \\ \partial_{q_\perp \rho} (\tilde{a}_q \cdot u) &= -\partial_{q_\perp \rho} \left(\frac{q_\perp \cdot \tilde{a}_q}{E_q} \right) = -\frac{1}{E_q} \left(\Theta^{\rho\nu} \tilde{a}_{q\nu} + \frac{|\mathbf{q}|^2}{E_q^2} \hat{q}_\perp^\rho \hat{q}_\perp \cdot \tilde{a}_q + q_\perp^\nu \partial_{q_\perp \rho} \tilde{a}_{q\nu} \right), \\ \Theta^{\mu\nu} q_\perp^\beta \partial_{q_\perp^\beta} \tilde{a}_{q\nu} &= q_\perp^\beta \partial_{q_\perp^\beta} \tilde{a}_q^\mu - u^\mu q_\perp^\beta \partial_{q_\perp^\beta} (\tilde{a}_q \cdot u) = q_\perp^\beta \partial_{q_\perp^\beta} \tilde{a}_q^\mu + \frac{u^\mu}{E_q} \left(\frac{m^2}{E_q^2} q_\perp^\beta + q_\perp^\alpha q_\perp^\beta \partial_{q_\perp^\alpha} \right) \tilde{a}_{q\beta}. \end{aligned}$$

By inputting exact expressions of $j_i^{T/L}$ from eqs. (D.13) and (D.14), it is found

$$\mathcal{Q}_{\text{cl}}^{(1)} = -2, \quad (\text{E.29a})$$

$$\begin{aligned} \mathcal{Q}_{\text{cl}}^{(2)} = & \left(\frac{q_0^2}{|\mathbf{q}|T} - \frac{2|\mathbf{q}|}{q_0} \left(2 + \frac{m^2}{|\mathbf{q}|^2} \right) - \frac{\eta_q m^2}{|\mathbf{q}|^2 T} (q_0 - 2T) \right) \hat{q}_\perp \cdot \tilde{a}_q \\ & + \left(\frac{q_0(m^2 - |\mathbf{q}|^2)}{|\mathbf{q}|^2} - \frac{\eta_q m^2 q_0^2}{|\mathbf{q}|^3} \right) \partial_{q_\perp^\nu} \tilde{a}_q^\nu + \left(\frac{q_0(3m^2 + |\mathbf{q}|^2)}{|\mathbf{q}|^2} - \frac{3\eta_q m^2 q_0^2}{|\mathbf{q}|^3} \right) \hat{q}_\perp^\nu \hat{q}_\perp^\rho \partial_{q_\perp^\rho} \tilde{a}_q^\nu, \end{aligned} \quad (\text{E.29b})$$

$$\begin{aligned} \mathcal{Q}_{\text{cl}}^{(3)} = & \left(\frac{q_0}{T} - 2 - \frac{\eta_q m^2}{|\mathbf{q}|q_0 T} (q_0 - 2T) \right) \hat{q}_\perp \cdot \tilde{a}_q + \left(\frac{q_0^2}{|\mathbf{q}|} - \frac{\eta_q m^2 q_0}{|\mathbf{q}|^2} \right) \partial_{q_\perp^\nu} \tilde{a}_q^\nu \\ & + \left(\frac{(3m^2 + |\mathbf{q}|^2)}{|\mathbf{q}|} - \frac{3\eta_q m^2 q_0}{|\mathbf{q}|^2} \right) \hat{q}_\perp^\nu \hat{q}_\perp^\rho \partial_{q_\perp^\rho} \tilde{a}_q^\nu, \end{aligned} \quad (\text{E.29c})$$

$$\mathcal{Q}_{\text{cl}}^{(4)} = -2|\mathbf{q}|, \quad (\text{E.29d})$$

$$\mathcal{Q}_{\text{cl}}^{(5)} = \frac{q_0^3}{|\mathbf{q}|T} \left(1 - \frac{\eta_q m^2}{q_0 |\mathbf{q}|} \right), \quad (\text{E.29e})$$

$$\mathcal{Q}_{\text{cl}}^{(6)} = -\frac{q_0^2}{2} \left(3 - \frac{q_0^2}{|\mathbf{q}|^2} + \frac{\eta_q m^4}{q_0 |\mathbf{q}|^3} \right), \quad (\text{E.29f})$$

$$\mathcal{Q}_{\text{cl}}^{(7)} = \frac{m^2 q_0}{2|\mathbf{q}|^3} (3|\mathbf{q}|q_0 + \eta_q (|\mathbf{q}|^2 - 3q_0^2)), \quad (\text{E.29g})$$

and

$$\tilde{\mathcal{Q}}_{\text{cl}}^{(1)} = \frac{2q_0}{|\mathbf{q}|T} \left[|\mathbf{q}|(1 - 2f_{Vq}) + (\hat{q}_\perp^\nu \partial_{q_\perp^\nu} f_{Vq}) \left(\frac{m^2 q_0}{|\mathbf{q}|} \eta_q - q_0^2 \right) \right], \quad (\text{E.30a})$$

$$\tilde{\mathcal{Q}}_{\text{cl}}^{(2)} = 2 \left(\frac{m^2 q_0}{|\mathbf{q}|} \eta_q - q_0^2 \right) \frac{f_{Vq}}{|\mathbf{q}|T} \hat{q}_\perp \cdot \tilde{a}_q, \quad (\text{E.30b})$$

$$\tilde{\mathcal{Q}}_{\text{cl}}^{(3)} = 2 \left(\frac{m^2 q_0}{|\mathbf{q}|} \eta_q - q_0^2 \right) \frac{f_{Vq}}{q_0 T} \hat{q}_\perp \cdot \tilde{a}_q, \quad (\text{E.30c})$$

$$\tilde{\mathcal{Q}}_{\text{cl}}^{(5)} = 2 \left(\frac{m^2 q_0}{|\mathbf{q}|} \eta_q - q_0^2 \right) \frac{q_0 f_{Vq}}{|\mathbf{q}|T}, \quad (\text{E.30d})$$

$$\tilde{\mathcal{Q}}_{\text{cl}}^{(4)} = \tilde{\mathcal{Q}}_{\text{cl}}^{(6)} = \tilde{\mathcal{Q}}_{\text{cl}}^{(7)} = 0. \quad (\text{E.30e})$$

Finally, the sum of all the pieces $\mathcal{C}_1 + \mathcal{C}_2$ from eqs. (E.25) and (E.27) provide the collision term of the AKE:

$$\begin{aligned} \lambda_c^{-1} \delta(q^2 - m^2) \hat{\mathcal{C}}_{\text{cl}}^\mu \approx & \frac{m_D^2 \delta(q^2 - m^2) T}{8\pi E_q} \ln(1/g_c) \left(\tilde{a}_q^\mu \dot{\mathcal{Q}}_{\text{cl}}^{(1)} + u^\mu \dot{\mathcal{Q}}_{\text{cl}}^{(2)} + \hat{q}_\perp^\mu \dot{\mathcal{Q}}_{\text{cl}}^{(3)} + \dot{\mathcal{Q}}_{\text{cl}}^{(4)} \hat{q}_\perp^\nu \partial_{q_\perp^\nu} \tilde{a}_{q\nu} \right. \\ & \left. + \dot{\mathcal{Q}}_{\text{cl}}^{(5)} \hat{q}_\perp^\nu \partial_{q_\perp^\nu} \tilde{a}_q^\mu + \dot{\mathcal{Q}}_{\text{cl}}^{(6)} \eta^{\nu\rho} \partial_{q_\perp^\nu} \partial_{q_\perp^\rho} \tilde{a}_q^\mu + \dot{\mathcal{Q}}_{\text{cl}}^{(7)} \hat{q}_\perp^\nu \hat{q}_\perp^\rho \partial_{q_\perp^\nu} \partial_{q_\perp^\rho} \tilde{a}_q^\mu \right). \end{aligned} \quad (\text{E.31})$$

The coefficients $\dot{\mathcal{Q}}_{\text{cl}}^{(i)} \equiv \tilde{\mathcal{Q}}_{\text{cl}}^{(i)} + \mathcal{Q}_{\text{cl}}^{(i)}$ ($i = 1, \dots, 7$) are shown in eqs. (4.23a)–(4.23g).

Open Access. This article is distributed under the terms of the Creative Commons Attribution License (CC-BY 4.0), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] STAR collaboration, *Global Λ hyperon polarization in nuclear collisions: evidence for the most vortical fluid*, *Nature* **548** (2017) 62 [[arXiv:1701.06657](#)] [[INSPIRE](#)].
- [2] STAR collaboration, *Global polarization of Λ hyperons in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV*, *Phys. Rev. C* **98** (2018) 014910 [[arXiv:1805.04400](#)] [[INSPIRE](#)].
- [3] Z.-T. Liang and X.-N. Wang, *Globally polarized quark-gluon plasma in non-central A+A collisions*, *Phys. Rev. Lett.* **94** (2005) 102301 [*Erratum ibid.* **96** (2006) 039901] [[nucl-th/0410079](#)] [[INSPIRE](#)].
- [4] F. Becattini, L. Csernai and D.J. Wang, *Λ polarization in peripheral heavy ion collisions*, *Phys. Rev. C* **88** (2013) 034905 [*Erratum ibid.* **93** (2016) 069901] [[arXiv:1304.4427](#)] [[INSPIRE](#)].
- [5] F. Becattini, V. Chandra, L. Del Zanna and E. Grossi, *Relativistic distribution function for particles with spin at local thermodynamical equilibrium*, *Annals Phys.* **338** (2013) 32 [[arXiv:1303.3431](#)] [[INSPIRE](#)].
- [6] R.-h. Fang, L.-g. Pang, Q. Wang and X.-n. Wang, *Polarization of massive fermions in a vortical fluid*, *Phys. Rev. C* **94** (2016) 024904 [[arXiv:1604.04036](#)] [[INSPIRE](#)].
- [7] I. Karpenko and F. Becattini, *Study of Λ polarization in relativistic nuclear collisions at $\sqrt{s_{NN}} = 7.7$ –200 GeV*, *Eur. Phys. J. C* **77** (2017) 213 [[arXiv:1610.04717](#)] [[INSPIRE](#)].
- [8] H. Li, L.-G. Pang, Q. Wang and X.-L. Xia, *Global Λ polarization in heavy-ion collisions from a transport model*, *Phys. Rev. C* **96** (2017) 054908 [[arXiv:1704.01507](#)] [[INSPIRE](#)].
- [9] F. Becattini and I. Karpenko, *Collective Longitudinal Polarization in Relativistic Heavy-Ion Collisions at Very High Energy*, *Phys. Rev. Lett.* **120** (2018) 012302 [[arXiv:1707.07984](#)] [[INSPIRE](#)].
- [10] D.-X. Wei, W.-T. Deng and X.-G. Huang, *Thermal vorticity and spin polarization in heavy-ion collisions*, *Phys. Rev. C* **99** (2019) 014905 [[arXiv:1810.00151](#)] [[INSPIRE](#)].
- [11] H.-Z. Wu, L.-G. Pang, X.-G. Huang and Q. Wang, *Local spin polarization in high energy heavy ion collisions*, *Phys. Rev. Research.* **1** (2019) 033058 [[arXiv:1906.09385](#)] [[INSPIRE](#)].
- [12] W. Florkowski, A. Kumar, R. Ryblewski and A. Mazeliauskas, *Longitudinal spin polarization in a thermal model*, *Phys. Rev. C* **100** (2019) 054907 [[arXiv:1904.00002](#)] [[INSPIRE](#)].
- [13] F. Becattini, G. Cao and E. Speranza, *Polarization transfer in hyperon decays and its effect in relativistic nuclear collisions*, *Eur. Phys. J. C* **79** (2019) 741 [[arXiv:1905.03123](#)] [[INSPIRE](#)].
- [14] X.-L. Xia, H. Li, X.-G. Huang and H.Z. Huang, *Feed-down effect on Λ spin polarization*, *Phys. Rev. C* **100** (2019) 014913 [[arXiv:1905.03120](#)] [[INSPIRE](#)].
- [15] W. Florkowski, B. Friman, A. Jaiswal and E. Speranza, *Relativistic fluid dynamics with spin*, *Phys. Rev. C* **97** (2018) 041901 [[arXiv:1705.00587](#)] [[INSPIRE](#)].
- [16] W. Florkowski, R. Ryblewski and A. Kumar, *Relativistic hydrodynamics for spin-polarized fluids*, *Prog. Part. Nucl. Phys.* **108** (2019) 103709 [[arXiv:1811.04409](#)] [[INSPIRE](#)].

- [17] K. Hattori, M. Hongo, X.-G. Huang, M. Matsuo and H. Taya, *Fate of spin polarization in a relativistic fluid: An entropy-current analysis*, *Phys. Lett. B* **795** (2019) 100 [[arXiv:1901.06615](#)] [[INSPIRE](#)].
- [18] D.T. Son and N. Yamamoto, *Berry Curvature, Triangle Anomalies and the Chiral Magnetic Effect in Fermi Liquids*, *Phys. Rev. Lett.* **109** (2012) 181602 [[arXiv:1203.2697](#)] [[INSPIRE](#)].
- [19] M.A. Stephanov and Y. Yin, *Chiral Kinetic Theory*, *Phys. Rev. Lett.* **109** (2012) 162001 [[arXiv:1207.0747](#)] [[INSPIRE](#)].
- [20] J.-W. Chen, S. Pu, Q. Wang and X.-N. Wang, *Berry Curvature and Four-Dimensional Monopoles in the Relativistic Chiral Kinetic Equation*, *Phys. Rev. Lett.* **110** (2013) 262301 [[arXiv:1210.8312](#)] [[INSPIRE](#)].
- [21] D.T. Son and N. Yamamoto, *Kinetic theory with Berry curvature from quantum field theories*, *Phys. Rev. D* **87** (2013) 085016 [[arXiv:1210.8158](#)] [[INSPIRE](#)].
- [22] A. Vilenkin, *Macroscopic parity violating effects: neutrino fluxes from rotating black holes and in rotating thermal radiation*, *Phys. Rev. D* **20** (1979) 1807 [[INSPIRE](#)].
- [23] D.E. Kharzeev, L.D. McLerran and H.J. Warringa, *The Effects of topological charge change in heavy ion collisions: ‘Event by event P and CP-violation’*, *Nucl. Phys. A* **803** (2008) 227 [[arXiv:0711.0950](#)] [[INSPIRE](#)].
- [24] K. Fukushima, D.E. Kharzeev and H.J. Warringa, *The Chiral Magnetic Effect*, *Phys. Rev. D* **78** (2008) 074033 [[arXiv:0808.3382](#)] [[INSPIRE](#)].
- [25] C. Manuel and J.M. Torres-Rincon, *Kinetic theory of chiral relativistic plasmas and energy density of their gauge collective excitations*, *Phys. Rev. D* **89** (2014) 096002 [[arXiv:1312.1158](#)] [[INSPIRE](#)].
- [26] J.-Y. Chen, D.T. Son, M.A. Stephanov, H.-U. Yee and Y. Yin, *Lorentz Invariance in Chiral Kinetic Theory*, *Phys. Rev. Lett.* **113** (2014) 182302 [[arXiv:1404.5963](#)] [[INSPIRE](#)].
- [27] J.-Y. Chen, D.T. Son and M.A. Stephanov, *Collisions in Chiral Kinetic Theory*, *Phys. Rev. Lett.* **115** (2015) 021601 [[arXiv:1502.06966](#)] [[INSPIRE](#)].
- [28] E.V. Gorbar, V.A. Miransky, I.A. Shovkovy and P.O. Sukhachov, *Consistent Chiral Kinetic Theory in Weyl Materials: Chiral Magnetic Plasmons*, *Phys. Rev. Lett.* **118** (2017) 127601 [[arXiv:1610.07625](#)] [[INSPIRE](#)].
- [29] D.E. Kharzeev, M.A. Stephanov and H.-U. Yee, *Anatomy of chiral magnetic effect in and out of equilibrium*, *Phys. Rev. D* **95** (2017) 051901 [[arXiv:1612.01674](#)] [[INSPIRE](#)].
- [30] Y. Hidaka, S. Pu and D.-L. Yang, *Relativistic Chiral Kinetic Theory from Quantum Field Theories*, *Phys. Rev. D* **95** (2017) 091901 [[arXiv:1612.04630](#)] [[INSPIRE](#)].
- [31] Y. Hidaka, S. Pu and D.-L. Yang, *Nonlinear Responses of Chiral Fluids from Kinetic Theory*, *Phys. Rev. D* **97** (2018) 016004 [[arXiv:1710.00278](#)] [[INSPIRE](#)].
- [32] Y. Hidaka and D.-L. Yang, *Nonequilibrium chiral magnetic/vortical effects in viscous fluids*, *Phys. Rev. D* **98** (2018) 016012 [[arXiv:1801.08253](#)] [[INSPIRE](#)].
- [33] N. Mueller and R. Venugopalan, *Worldline construction of a covariant chiral kinetic theory*, *Phys. Rev. D* **96** (2017) 016023 [[arXiv:1702.01233](#)] [[INSPIRE](#)].
- [34] N. Mueller and R. Venugopalan, *The chiral anomaly, Berry’s phase and chiral kinetic theory, from world-lines in quantum field theory*, *Phys. Rev. D* **97** (2018) 051901 [[arXiv:1701.03331](#)] [[INSPIRE](#)].

- [35] A. Huang, Y. Jiang, S. Shi, J. Liao and P. Zhuang, *Out-of-Equilibrium Chiral Magnetic Effect from Chiral Kinetic Theory*, *Phys. Lett. B* **777** (2018) 177 [[arXiv:1703.08856](#)] [[INSPIRE](#)].
- [36] A. Huang, S. Shi, Y. Jiang, J. Liao and P. Zhuang, *Complete and Consistent Chiral Transport from Wigner Function Formalism*, *Phys. Rev. D* **98** (2018) 036010 [[arXiv:1801.03640](#)] [[INSPIRE](#)].
- [37] S. Carignano, C. Manuel and J.M. Torres-Rincon, *Consistent relativistic chiral kinetic theory: A derivation from on-shell effective field theory*, *Phys. Rev. D* **98** (2018) 076005 [[arXiv:1806.01684](#)] [[INSPIRE](#)].
- [38] Ö.F. Dayi and E. Kilinçarslan, *Quantum Kinetic Equation in the Rotating Frame and Chiral Kinetic Theory*, *Phys. Rev. D* **98** (2018) 081701 [[arXiv:1807.05912](#)] [[INSPIRE](#)].
- [39] Y.-C. Liu, L.-L. Gao, K. Mameda and X.-G. Huang, *Chiral kinetic theory in curved spacetime*, *Phys. Rev. D* **99** (2019) 085014 [[arXiv:1812.10127](#)] [[INSPIRE](#)].
- [40] Y. Sun and C.M. Ko, *Chiral kinetic approach to the chiral magnetic effect in isobaric collisions*, *Phys. Rev. C* **98** (2018) 014911 [[arXiv:1803.06043](#)] [[INSPIRE](#)].
- [41] D.O. Rybalka, E.V. Gorbar and I.A. Shovkovy, *Hydrodynamic modes in a magnetized chiral plasma with vorticity*, *Phys. Rev. D* **99** (2019) 016017 [[arXiv:1807.07608](#)] [[INSPIRE](#)].
- [42] S. Lin and A. Shukla, *Chiral Kinetic Theory from Effective Field Theory Revisited*, *JHEP* **06** (2019) 060 [[arXiv:1901.01528](#)] [[INSPIRE](#)].
- [43] N. Yamamoto and D.-L. Yang, *Chiral Radiation Transport Theory of Neutrinos*, *Astrophys. J.* **895** (2020) 1 [[arXiv:2002.11348](#)] [[INSPIRE](#)].
- [44] D.-L. Yang, *Side-Jump Induced Spin-Orbit Interaction of Chiral Fluids from Kinetic Theory*, *Phys. Rev. D* **98** (2018) 076019 [[arXiv:1807.02395](#)] [[INSPIRE](#)].
- [45] K. Fukushima, S. Pu and Z. Qiu, *Eddy magnetization from the chiral Barnett effect*, *Phys. Rev. A* **99** (2019) 032105 [[arXiv:1808.08016](#)] [[INSPIRE](#)].
- [46] Y. Sun and C.M. Ko, *Λ hyperon polarization in relativistic heavy ion collisions from a chiral kinetic approach*, *Phys. Rev. C* **96** (2017) 024906 [[arXiv:1706.09467](#)] [[INSPIRE](#)].
- [47] S.Y.F. Liu, Y. Sun and C.M. Ko, *Spin polarizations in a covariant angular momentum conserved chiral transport model*, [arXiv:1910.06774](#) [[INSPIRE](#)].
- [48] N. Mueller and R. Venugopalan, *Constructing phase space distributions with internal symmetries*, *Phys. Rev. D* **99** (2019) 056003 [[arXiv:1901.10492](#)] [[INSPIRE](#)].
- [49] N. Weickgenannt, X.-L. Sheng, E. Speranza, Q. Wang and D.H. Rischke, *Kinetic theory for massive spin-1/2 particles from the Wigner-function formalism*, *Phys. Rev. D* **100** (2019) 056018 [[arXiv:1902.06513](#)] [[INSPIRE](#)].
- [50] J.-H. Gao and Z.-T. Liang, *Relativistic Quantum Kinetic Theory for Massive Fermions and Spin Effects*, *Phys. Rev. D* **100** (2019) 056021 [[arXiv:1902.06510](#)] [[INSPIRE](#)].
- [51] K. Hattori, Y. Hidaka and D.-L. Yang, *Axial Kinetic Theory and Spin Transport for Fermions with Arbitrary Mass*, *Phys. Rev. D* **100** (2019) 096011 [[arXiv:1903.01653](#)] [[INSPIRE](#)].
- [52] Z. Wang, X. Guo, S. Shi and P. Zhuang, *Mass Correction to Chiral Kinetic Equations*, *Phys. Rev. D* **100** (2019) 014015 [[arXiv:1903.03461](#)] [[INSPIRE](#)].
- [53] Y.-C. Liu and X.-G. Huang, *Anomalous chiral transports and spin polarization in heavy-ion collisions*, *Nucl. Sci. Tech.* **31** (2020) 56 [[arXiv:2003.12482](#)] [[INSPIRE](#)].

- [54] S. Li and H.-U. Yee, *Quantum Kinetic Theory of Spin Polarization of Massive Quarks in Perturbative QCD: Leading Log*, *Phys. Rev. D* **100** (2019) 056022 [[arXiv:1905.10463](#)] [[INSPIRE](#)].
- [55] J.-j. Zhang, R.-h. Fang, Q. Wang and X.-N. Wang, *A microscopic description for polarization in particle scatterings*, *Phys. Rev. C* **100** (2019) 064904 [[arXiv:1904.09152](#)] [[INSPIRE](#)].
- [56] J.I. Kapusta, E. Rrapaj and S. Rudaz, *Relaxation Time for Strange Quark Spin in Rotating Quark-Gluon Plasma*, *Phys. Rev. C* **101** (2020) 024907 [[arXiv:1907.10750](#)] [[INSPIRE](#)].
- [57] D. Vasak, M. Gyulassy and H.T. Elze, *Quantum Transport Theory for Abelian Plasmas*, *Annals Phys.* **173** (1987) 462 [[INSPIRE](#)].
- [58] H.T. Elze, M. Gyulassy and D. Vasak, *Transport Equations for the QCD Quark Wigner Operator*, *Nucl. Phys. B* **276** (1986) 706 [[INSPIRE](#)].
- [59] E.V. Gorbar, V.A. Miransky, I.A. Shovkovy and X. Wang, *Radiative corrections to chiral separation effect in QED*, *Phys. Rev. D* **88** (2013) 025025 [[arXiv:1304.4606](#)] [[INSPIRE](#)].
- [60] M. Buzzegoli, E. Grossi and F. Becattini, *General equilibrium second-order hydrodynamic coefficients for free quantum fields*, *JHEP* **10** (2017) 091 [*Erratum ibid.* **07** (2018) 119] [[arXiv:1704.02808](#)] [[INSPIRE](#)].
- [61] S. Lin and L. Yang, *Mass correction to chiral vortical effect and chiral separation effect*, *Phys. Rev. D* **98** (2018) 114022 [[arXiv:1810.02979](#)] [[INSPIRE](#)].
- [62] X.-G. Huang and J. Liao, *Axial Current Generation from Electric Field: Chiral Electric Separation Effect*, *Phys. Rev. Lett.* **110** (2013) 232302 [[arXiv:1303.7192](#)] [[INSPIRE](#)].
- [63] Y. Jiang, X.-G. Huang and J. Liao, *Chiral electric separation effect in the quark-gluon plasma*, *Phys. Rev. D* **91** (2015) 045001 [[arXiv:1409.6395](#)] [[INSPIRE](#)].
- [64] S. Pu, S.-Y. Wu and D.-L. Yang, *Holographic Chiral Electric Separation Effect*, *Phys. Rev. D* **89** (2014) 085024 [[arXiv:1401.6972](#)] [[INSPIRE](#)].
- [65] S. Pu, S.-Y. Wu and D.-L. Yang, *Chiral Hall Effect and Chiral Electric Waves*, *Phys. Rev. D* **91** (2015) 025011 [[arXiv:1407.3168](#)] [[INSPIRE](#)].
- [66] V. Bargmann, L. Michel and V.L. Telegdi, *Precession of the polarization of particles moving in a homogeneous electromagnetic field*, *Phys. Rev. Lett.* **2** (1959) 435 [[INSPIRE](#)].
- [67] W.-T. Deng and X.-G. Huang, *Event-by-event generation of electromagnetic fields in heavy-ion collisions*, *Phys. Rev. C* **85** (2012) 044907 [[arXiv:1201.5108](#)] [[INSPIRE](#)].
- [68] L. McLerran and V. Skokov, *Comments About the Electromagnetic Field in Heavy-Ion Collisions*, *Nucl. Phys. A* **929** (2014) 184 [[arXiv:1305.0774](#)] [[INSPIRE](#)].
- [69] B. Svetitsky, *Diffusion of charmed quarks in the quark-gluon plasma*, *Phys. Rev. D* **37** (1988) 2484 [[INSPIRE](#)].
- [70] M. Le Bellac, *Thermal field theory*, Cambridge University Press (2000).
- [71] N. Yamamoto, *Photonic chiral vortical effect*, *Phys. Rev. D* **96** (2017) 051902 [[arXiv:1702.08886](#)] [[INSPIRE](#)].
- [72] X.-G. Huang and A.V. Sadofyev, *Chiral Vortical Effect For An Arbitrary Spin*, *JHEP* **03** (2019) 084 [[arXiv:1805.08779](#)] [[INSPIRE](#)].
- [73] N. Yamamoto, *Chiral transport of neutrinos in supernovae: Neutrino-induced fluid helicity and helical plasma instability*, *Phys. Rev. D* **93** (2016) 065017 [[arXiv:1511.00933](#)] [[INSPIRE](#)].

- [74] N. Yamamoto, *Scaling laws in chiral hydrodynamic turbulence*, *Phys. Rev. D* **93** (2016) 125016 [[arXiv:1603.08864](#)] [[INSPIRE](#)].
- [75] Y. Masada, K. Kotake, T. Takiwaki and N. Yamamoto, *Chiral magnetohydrodynamic turbulence in core-collapse supernovae*, *Phys. Rev. D* **98** (2018) 083018 [[arXiv:1805.10419](#)] [[INSPIRE](#)].
- [76] A. Ohnishi and N. Yamamoto, *Magnetars and the Chiral Plasma Instabilities*, [arXiv:1402.4760](#) [[INSPIRE](#)].
- [77] D. Grabowska, D.B. Kaplan and S. Reddy, *Role of the electron mass in damping chiral plasma instability in Supernovae and neutron stars*, *Phys. Rev. D* **91** (2015) 085035 [[arXiv:1409.3602](#)] [[INSPIRE](#)].
- [78] D.B. Kaplan, S. Reddy and S. Sen, *Energy Conservation and the Chiral Magnetic Effect*, *Phys. Rev. D* **96** (2017) 016008 [[arXiv:1612.00032](#)] [[INSPIRE](#)].
- [79] M. Wakamatsu, *Gauge and frame-independent decomposition of nucleon spin*, *Phys. Rev. D* **83** (2011) 014012 [[arXiv:1007.5355](#)] [[INSPIRE](#)].
- [80] X.-D. Ji, *Gauge-Invariant Decomposition of Nucleon Spin*, *Phys. Rev. Lett.* **78** (1997) 610 [[hep-ph/9603249](#)] [[INSPIRE](#)].
- [81] E. Leader and C. Lorcé, *The angular momentum controversy: What's it all about and does it matter?*, *Phys. Rept.* **541** (2014) 163 [[arXiv:1309.4235](#)] [[INSPIRE](#)].
- [82] K. Fukushima and S. Pu, *Relativistic decomposition of the orbital and the spin angular momentum in chiral physics and Feynman's angular momentum paradox*, [arXiv:2001.00359](#) [[INSPIRE](#)].