

**UCC Library and UCC researchers have made this item openly available.  
Please [let us know](#) how this has helped you. Thanks!**

<b>Title</b>	Iatrogenic hypoglycaemia following glucose-insulin infusions for the treatment of hyperkalaemia
<b>Author(s)</b>	Tee, Su Ann; Devine, Kerri; Potts, Adam; Javaid, Usman; Razvi, Salman; Quinton, Richard; Roberts, Graham; Leech, Nicola J.
<b>Publication date</b>	2020-09-26
<b>Original citation</b>	Tee, S. A., Devine, K., Potts, A., Javaid, U., Razvi, S., Quinton, R., Roberts, G. and Leech, N. J. (2020) 'Iatrogenic hypoglycaemia following glucose-insulin infusions for the treatment of hyperkalaemia', <i>Clinical Endocrinology</i> . doi: 10.1111/cen.14343
<b>Type of publication</b>	Article (peer-reviewed)
<b>Link to publisher's version</b>	<a href="http://dx.doi.org/10.1111/cen.14343">http://dx.doi.org/10.1111/cen.14343</a> Access to the full text of the published version may require a subscription.
<b>Rights</b>	© 2020, John Wiley & Sons, Ltd. This is the peer reviewed version of the following article Tee, S. A., Devine, K., Potts, A., Javaid, U., Razvi, S., Quinton, R., Roberts, G. and Leech, N. J. (2020) 'Iatrogenic hypoglycaemia following glucose-insulin infusions for the treatment of hyperkalaemia', <i>Clinical Endocrinology</i> , doi: 10.1111/cen.14343, which has been published in final form at <a href="https://doi.org/10.1111/cen.14343">https://doi.org/10.1111/cen.14343</a> . This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.
<b>Embargo information</b>	Access to this article is restricted until 12 months after publication by request of the publisher.
<b>Embargo lift date</b>	2021-09-26
<b>Item downloaded from</b>	<a href="http://hdl.handle.net/10468/10618">http://hdl.handle.net/10468/10618</a>

Downloaded on 2021-11-27T11:18:34Z



DR SU ANN TEE (Orcid ID : 0000-0001-7893-7352)

DR KERRI DEVINE (Orcid ID : 0000-0003-0129-4373)

DR SALMAN RAZVI (Orcid ID : 0000-0002-9047-1556)

Article type : Original Article

**Title: Iatrogenic hypoglycaemia following glucose-insulin infusions for the treatment of hyperkalaemia**

**Short Title: *Gwl Hypo Evaluation***

Su Ann Tee\*<sup>1</sup>, Kerri Devine\*<sup>1</sup>, Adam Potts<sup>1</sup>, Usman Javaid<sup>2</sup>, Salman Razvi<sup>2,3</sup>, Richard Quinton<sup>1,3</sup>, Graham Roberts<sup>4,5</sup>, Nicola J Leech<sup>1</sup>

<sup>1</sup>Department of Diabetes & Endocrinology, Newcastle-upon-Tyne Hospitals NHS Foundation Trust, Newcastle, UK

<sup>2</sup>Department of Diabetes & Endocrinology, Queen Elizabeth Hospital, Gateshead Health NHS Foundation Trust, Gateshead, UK

<sup>3</sup>Translational & Clinical Research Institute, University of Newcastle-upon-Tyne, UK

<sup>4</sup>Diabetes Research Group, Swansea University, Swansea, UK

<sup>5</sup>Clinical Research Facility - Cork, University College Cork, Cork, Ireland

*\*contributed equally to this work*

**Corresponding author, reprint requests addressed to:**

Dr Nicola Leech

Royal Victoria Infirmary

Newcastle upon Tyne

United Kingdom

NE1 4LP

Email: Nicola.leech3@nhs.net

**Conflict of interest:** Nothing to disclose.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/CEN.14343](https://doi.org/10.1111/CEN.14343)

This article is protected by copyright. All rights reserved

**Funding:** None.

**Word Count:** 2500

**Data Availability Statement:** The data that support the findings of this study are available from the corresponding author upon reasonable request.

## Abstract

**Objectives:** To study the incidence of, and risk factors for, iatrogenic hypoglycaemia following Gwl infusion in our institution.

**Context:** Hyperkalaemia is a life-threatening biochemical abnormality. Glucose-with-insulin (Gwl) infusions form standard management, but risk iatrogenic hypoglycaemia (glucose  $\leq$  3.9mmol/L). Recently updated UK guidelines include an additional glucose infusion in patients with pre-treatment capillary blood glucose (CBG)  $<$ 7.0 mmol/L.

**Design:** Retrospective analysis of outcomes for Gwl infusions prescribed for hyperkalaemia from 1<sup>st</sup> January-28<sup>th</sup> February 2019, extracted from the Newcastle-upon-Tyne Hospitals NHS Foundation Trust electronic platform (eRecord).

**Participants:** 132 patients received 228 Gwl infusions for hyperkalaemia.

**Main outcome measures:** Incidence, severity and time-to-onset of hypoglycaemia.

**Results:** Hypoglycaemia incidence was 11.8%. At least 1 hypoglycaemic episode occurred in 18.2% of patients with 6.8% having at least 1 episode of severe hypoglycaemia ( $<$ 3.0 mmol/L). Most episodes (77.8%) occurred within 3 hours of treatment.

Lower pre-treatment CBG(5.9 mmol/L [4.1 mmol/L - 11.2 mmol/L];; versus 7.6 mmol/L [3.7 mmol/L - 31.3 mmol/L],  $p = 0.000$ ) was associated with hypoglycaemia risk. A diagnosis of type 2 diabetes and treatment for hyperkalaemia within the previous 24 hours were negatively associated.

**Conclusions:** Within our inpatient population, around 1 in 8 Gwl infusions delivered as treatment for hyperkalaemia resulted in iatrogenic hypoglycaemia. Higher pre-treatment CBG and a diagnosis of type 2 diabetes were protective, irrespective of renal function. Our findings support the immediate change to current management, either with additional glucose infusions, or by using glucose-only infusions in patients without diabetes. These approaches should be compared via a prospective randomised study.

**Keywords:** Hypoglycaemia, hyperkalaemia, dextrose, glucose-with-insulin

## Introduction

Hyperkalaemia is a commonly encountered but potentially life-threatening biochemical abnormality, with a reported incidence of between 1-10% in hospital inpatients, particularly among those with renal impairment<sup>1,2</sup>. Due to the effects of potassium on cardiac myocyte resting membrane potential, hyperkalaemia is associated with a significant risk of developing arrhythmias and cardiac arrest, and potassium ( $K^+$ ) levels  $\geq 6.5$  mmol/L (normal range 3.5-5.3 mmol/L) warrant urgent treatment<sup>3</sup>. There is surprisingly limited evidence behind the current guidelines for treating hyperkalaemia in hospital inpatients, and there is wide variation in clinical practice both within and between individual centres.

Glucose-with-insulin (Gwl) infusions form part of the current UK national guidelines from the Renal Association for treatment of hyperkalaemia<sup>4</sup>, based on the physiological observation that insulin stimulates the activity of the  $Na^+-K^+$  ATP pump, which leads to an influx of potassium into cells and a corresponding reduction in extracellular fluid potassium.

Previous publications have highlighted the risk of iatrogenic hypoglycaemia (capillary blood glucose  $\leq 3.9$ mmol/L) following hyperkalaemia treatment with Gwl infusions; the reported incidence ranging from 6% to 75%<sup>5-9</sup>. Patients with hyperkalaemia may have additional risk factors for hypoglycaemia, such as end-stage renal disease, or diabetes treated with insulin or insulin secretagogue therapy. Hypoglycaemia has potentially life-threatening consequences, including precipitation of acute cardiovascular events and cardiac arrhythmia, and may cause life-changing generalised brain injury in the long term. Hypoglycaemia among hospitalised inpatients has also been linked to increased morbidity, mortality and length-of-stay<sup>10</sup>.

Consideration of the above evidence has led to some changes in the updated UK Renal Association (UKRA) guidelines (July 2020<sup>11</sup>), which now recommend:

1. Treat all patients requiring Gwl with 10 units of soluble insulin and 25g dextrose
2. In addition, give 10% dextrose at a rate of 50ml/hr for 5 hrs in patients with pre treatment glucose  $< 7$  mmol/L
3. Monitor blood glucose for 12 hrs after Gwl treatment

Our aim was to determine the incidence of iatrogenic hypoglycaemia following Gwl infusions in our inpatient population, as well as to determine any predictive factors for development of hypoglycaemia.

## Methods

We performed a retrospective audit within the Newcastle-upon-Tyne Hospitals NHS Foundation Trust to determine the incidence of iatrogenic hypoglycaemia in adult patients receiving Gwl infusions for the treatment of hyperkalaemia. Caldicott approval was granted for data access and the project was registered with the trust Audit registry (Clinical Governance & Audit Registration N°9889). Data on all patients prescribed Gwl infusions as treatment for hyperkalaemia from 1<sup>st</sup> January to 28<sup>th</sup> February 2019 was extracted from the trust electronic prescribing system “eRecord”.

Our trust guidelines recommend treatment with Gwl infusion for all patients with potassium level  $\geq 6.5$  mmol/L, or potassium 6-6.4 mmol/L with ECG changes, or where there are other reasons for concern. Gwl as per trust protocol contains 10 units of Actrapid® (short acting human insulin), infused over 15 minutes, in either 50 ml of 50% dextrose (*i.e.* 25g glucose), or 100ml of 20% dextrose (20g glucose) in patients where there are concerns about the reliability of IV access. Capillary blood glucose (CBG) is recorded at baseline, and at 15 minutes, 30 minutes, 60 minutes, 90 minutes, 2 hours, 3 hours, 4 hours and 6 hours from the onset of the infusion. All patients who developed hypoglycaemia (defined as CBG  $\leq 3.9$  mmol/L) up to six hours following Gwl infusions were included in the “hypoglycaemia” cohort, and were compared against those who did not develop hypoglycaemia.

The following data were extracted from eRecord: age and sex of patient, date and time of infusion, baseline capillary glucose, pre- and post-treatment potassium level and renal function at the time of treatment. Body mass index (BMI) was not available for the majority of patients, as patient weight and height is not reliably recorded during acute admissions within our trust. Previous diagnoses of diabetes (including treatment modality), chronic kidney disease (CKD; defined as a persistent estimated glomerular filtration rate – eGFR of  $<60$  ml/min/1.73m<sup>2</sup> in the 3 to 6 months preceding admission) and chronic liver disease were also recorded. The reduction in potassium after treatment was determined from the nadir potassium level within six hours of infusion. Hypoglycaemia was defined as capillary blood glucose  $<4.0$  mmol/L, and severe hypoglycaemia as capillary blood glucose of  $<3.0$  mmol/L, based on the National Diabetes

Inpatient Audit definitions<sup>12</sup>. The data that support the findings of this study are available from the corresponding author upon reasonable request.

Results were analysed using IBM® SPSS Statistics® (Version 26, Chicago, IL). In univariate analysis for the key outcome of development of hypoglycaemia, continuous variables of interest were compared with independent-samples t-test or Mann-Whitney *U* test depending on their distribution. The categorical variables were analysed with Chi-Square statistic or Fisher's exact test. Analysis of time to hypoglycaemia against renal function was by Kruskal-Wallis test for categorical data (existing CKD diagnosis) or by Pearson correlation for continuous data (serum creatinine). Results are presented using median and range for continuous variables unless stated otherwise, and with odds ratios with 95% confidence intervals (CI) for categorical variables.

## **Results**

### ***Baseline Demographics***

Data was extracted for 132 patients (86 males and 46 females). The median age of participants was 67 years (range: 19-95 years). There was no significant difference in age between males (median: 66 years [range: 23-95 years]) and females (median: 72 years [range: 19-95 years], (Mann-Whitney *U*,  $p = 0.481$ ). A total of 261 Gwl infusions were prescribed for these patients between 01/01/2019 and 28/02/2019 for treatment of hyperkalaemia. Thirty three episodes of Gwl infusions were excluded from analysis due to lack of recorded baseline blood glucose, post-treatment potassium or non-standard time points, leaving 228 infusions (prescribed to 132 patients).

A prior diagnosis of diabetes mellitus was recorded in 37.9% (95% CI: 30.1-46.4%;  $n = 50/132$ ). As expected from a cohort experiencing hyperkalaemia, 54.6% (95% CI: 46.1-62.8%;  $n = 72/132$ ) had an established diagnosis of CKD G3 or above (on basis of prior eGFR).

### ***Incidence and severity of hypoglycaemia***

At least one episode of hypoglycaemia occurred in 18.2% of patients (95% CI: 12.5%-25.6%;  $n = 24/132$ ) with 6.82% (95% CI: 3.63%-12.45%;  $n = 9/132$ ) having at least one episode of severe hypoglycaemia ( $<3.0$ mmol/L). There were 27 (11.8%, 95% CI: 8.3-16.7%) episodes of hypoglycaemia within the six hours after treatment recorded among 228 infusions prescribed.

### ***Factors influencing the risk of developing hypoglycaemia***

Patient Factors

In univariate analysis, the age (OR 0.987/year, 95% CI: 0.960-1.014,  $p = 0.345$ ) and sex (OR 2.325, males versus females; 95% CI: 0.806-6.705,  $p = 0.118$ ) of the patient were not significant predictors of having at least one episode of hypoglycaemia following Gwtreatment.

In wider univariate analysis of patient-level factors, a prior diagnosis of type 2 diabetes mellitus appeared to be protective against hypoglycaemia (OR 0.079, 95% CI: 0.010-0.616,  $p = 0.015$ ). A diagnosis of type 1 diabetes, treatment with insulin or a sulfonylurea and degree of chronic kidney disease were not significantly associated with hypoglycaemia (Table 1). No patients (0/19) in the hypoglycaemia group had a recorded diagnosis of chronic liver disease, therefore we did not pursue further analysis.

#### Episode-Level Factors

With the data available, we expressed the following epidemiological data by discrete hyperkalaemia episode, rather than per patient (Table 2).

The median pre-treatment glucose was significantly lower in those that developed iatrogenic hypoglycaemia (5.9 mmol/L [range: 4.1 mmol/L - 11.2 mmol/L],  $n = 27$ ; versus 7.6 mmol/L [range: 3.7 mmol/L - 31.3 mmol/L],  $n = 201$ ; Mann-Whitney  $U$ ,  $p = 0.000$ ). There was a strong association between higher pre-treatment glucose levels and reduction in hypoglycaemia events (OR 0.669/mmol, 95% CI: 0.520-0.860,  $p = 0.002$ ). Indeed, the 2020 UKRA guidelines recommend the use of a  $<7$  mmol/L threshold for pre-infusion glucose as being at increased risk of hypoglycaemia<sup>11</sup>. We support the use of this threshold, identifying a strong relationship (OR 4.146; 95% CI: 1.676-10.255,  $p = 0.002$ ) between hypoglycaemia and pre-infusion glucose  $<7$  mmol/L (Figure 2).

The median number of treatments per patient was 1 (range 1-12), but 34.1% ( $n = 45/132$ ) of patients had more than one treatment episode. The number of treatments across the patient's admission was inversely associated with episodes of hypoglycaemia (OR 0.347, 95% CI: 0.123-0.975,  $p = 0.045$ ) following Gwl infusion. This may be a marker of insulin resistance in this group.

Whilst there was a slight statistically significant difference between the median pre-treatment potassium it was not a significant predictor in univariate odds analysis of hypoglycaemia (OR 0.402, 95% CI: 0.138 – 1.175,  $p = 0.096$ ).



Of the infusions prescribed, only 4.4% (95% CI: 2.4%-7.9%;  $n = 10/228$ ) utilised 20% dextrose, whilst the majority (95.6%) used 50% dextrose. Due to the small number in the first group, and indeed the small difference in administered glucose (20g versus 25g), we have not compared these variables separately.

There were no significant differences found in age, current renal function, or previous diagnosis of CKD between those developing hypoglycaemia and those who did not. Similarly, there was no difference in potassium response to treatment (*Table 2*).

### **Time to developing hypoglycaemia**

Figure 1 demonstrates that most episodes of hypoglycaemia occurred within the first 3 hours of receiving Gwl infusions, with 51.9% of hypoglycaemia episodes occurring within 2 hours, and 77.8% within 3 hours of the infusion. The median time to development of hypoglycaemia was 110 minutes (range 35-221 minutes). There were no episodes of hypoglycaemia recorded between 4 and 6 hours after infusion. There was no effect on time to hypoglycaemia demonstrated in patients with existing CKD ( $H(3) = 0.669$ ;  $p = 0.880$ ), or depending on serum creatinine at the time of treatment ( $r = -0.09$ ;  $R^2 = 0.008$ ;  $p = 0.677$ ).

### **Discussion**

The incidence of iatrogenic hypoglycaemia following Gwl infusions for hyperkalaemia within our study population was 11.8%, which is similar to the rates (6.1% to 17.5%) reported in other studies (*Table 3*)<sup>6-8,13-16</sup>. The rate of hypoglycaemia we observed was slightly lower than in another recent UK study by Boughton et al<sup>13</sup>; however, this is most probably because their protocol involved administering 10 units of insulin in a lower (20g) glucose load, whereas 95.6% of the infusions in our cohort contained a 25g total glucose load (as recommended in the 2014 Renal Association guidelines). Taking into account the number of patients having recurrent infusions, 18.2% of patients (95% CI: 12.5%-25.6%;  $n = 24/132$ ) developed hypoglycaemia after receiving Gwl infusions for the treatment of hyperkalaemia which, in our view, represents an unacceptable level of risk to patients.

Lower pre-treatment capillary blood glucose was identified as the main risk factor for development of hypoglycaemia in our analysis. A prior recorded diagnosis of type 2 diabetes appeared protective, reflecting insulin resistance in this group. Only one previous study<sup>6</sup> has mentioned this as a potential risk factor, but the study involved patients exclusively with end-

stage renal disease, and therefore generalisability is limited. Our findings show importantly and uniquely, that a diagnosis of type 2 diabetes appears protective regardless of pre-existing renal function.

Several suggestions have been made to reduce risk of hypoglycaemia from Gwl infusions<sup>9,17</sup>, including administering a lower dose of insulin, pre-loading with additional glucose or administering a higher overall glucose load.

It is evident that graded release of endogenous insulin occurs as part of the normal physiological response to a glucose load<sup>18</sup>. Hence for hyperkalaemic patients without a diagnosis of diabetes or other disorders of insulin regulation, administering intravenous glucose alone is potentially an effective treatment, and eliminates the risk of iatrogenic hypoglycaemia.

A small crossover study by Chothia *et al.* in non-diabetic patients with hyperkalaemia examined this principle. After infusion of 50g dextrose, with- or without- 10 units of insulin, they did find a mean K<sup>+</sup> drop of 0.5 (+/- 0.31) mmol/L with dextrose alone, however this was slightly less than the 0.83 (+/- 0.53) mmol/L reduction in the group treated with conventional Gwl ( $p=0.01$ )<sup>17</sup>. Although this approach is currently not recommended in the latest UKRA guideline iteration, we would strongly support further investigation of this approach, with a view to future guidelines being multidisciplinary, rather than monospecialty-derived.

We believe there needs to be an immediate change to practice in order to reduce the present unacceptable risk of developing hypoglycaemia with Gwl infusions for hyperkalaemia, particularly in patients with a lower pre-treatment glucose level and without type 2 diabetes. However, the question is whether to adopt the proposal in recently updated UKRA guidelines to administer an additional glucose load of 25g, or more simply, to administer glucose-only infusions for this patient group. We propose that the safety and efficacy of these two approaches be subject to prospective study. In the interim, we propose the following changes (Figure 3) to the hyperkalaemia management protocol within our trust and suggest that this can be adopted by other institutions, due to concerns regarding patient safety with existing protocols as highlighted by our audit.

There were limitations to our analysis. Due to insufficient data, we were unable to perform rigorous within-person risk factor analysis for those who underwent multiple Gwl treatments. We did however distinguish factors on a per-patient (e.g. sex) versus per-episode (e.g. pre-treatment

glucose) basis. Examination of whether repeated Gwl infusions increased hypoglycaemia risk over a 24-hour time period in fact found the opposite association. This may be another marker of insulin resistance in such patients. Similarly, for patients with diabetes, medication, including insulin injections, given on the same day could theoretically have contributed to greater risk of hypoglycaemia, but we found these patients to have a lower incidence of hypoglycaemia. We did not have sufficient data available to assess whether body mass index affected risk of hypoglycaemia, as patient weight and height are currently not systematically or reliably recorded as part of acute inpatient admissions to our trust. We also did not look at the effect of potential concurrent medication (other than insulin or oral hypoglycaemic medication) with hypoglycaemia listed as a possible side-effect; however there are likely only small numbers of patients on such medication.

The July 2020 UKRA guidelines recommend actively monitoring for hypoglycaemia for twelve hours post infusion, which is not our current trust policy. We did not attempt to collect data beyond six hours due to potential for introducing bias (e.g. those with a diagnosis of diabetes would have had additional blood glucose monitoring). Reassuringly, we did not identify any episodes of hypoglycaemia occurring between four and six hours after treatment.

### **Conclusions**

Consistent with previous publications, we determined that around 1 in 8 Gwl infusions (11.8%) delivered as treatment for hyperkalaemia resulted in iatrogenic hypoglycaemia within our inpatient population, with 18.2% patients (almost 1 in 5) experiencing this complication.

Having a lower pre-treatment capillary blood glucose level was the main factor associated with hypoglycaemia risk, and a prior diagnosis of type 2 diabetes may be protective. Our findings support UKRA recommendations for giving additional glucose loading for patients with low pre-treatment glucose level (<7.0mmol/L), but would equally support a robust exploration of using glucose-only infusions. These two approaches should ideally be compared via a prospective randomised study, but for the interim, we recommend the immediate revision of current approaches to hyperkalaemia treatment for overwhelming reasons of patient safety.

### **References**

1. Khanagavi J, Gupta T, Aronow WS, et al. Hyperkalemia among hospitalized patients and association between duration of hyperkalemia and outcomes. *Arch Med Sci.* 2014;10(2):251-257.

2. Stevens MS, Dunlay RW. Hyperkalemia in hospitalized patients. *Int Urol Nephrol*. 2000;32(2):177-180.
3. Soar J, Perkins GD, Abbas G, et al. European Resuscitation Council Guidelines for Resuscitation 2010 Section 8. Cardiac arrest in special circumstances: Electrolyte abnormalities, poisoning, drowning, accidental hypothermia, hyperthermia, asthma, anaphylaxis, cardiac surgery, trauma, pregnancy, electrocution. *Resuscitation*. 2010;81(10):1400-1433.
4. UK Renal Association. Clinical Practice Guidelines: Treatment of acute hyperkalaemia in adults. . <https://renal.org/wp-content/uploads/2017/06/hyperkalaemia-guideline-1.pdf>. Published 2014. Accessed 11/05/2020.
5. Allon M, Copkney C. Albuterol and insulin for treatment of hyperkalemia in hemodialysis patients. *Kidney Int*. 1990;38(5):869-872.
6. Apel J, Reutrakul S, Baldwin D. Hypoglycemia in the treatment of hyperkalemia with insulin in patients with end-stage renal disease. *Clin Kidney J*. 2014;7(3):248-250.
7. Coca A, Valencia AL, Bustamante J, Mendiluce A, Floege J. Hypoglycemia following intravenous insulin plus glucose for hyperkalemia in patients with impaired renal function. *PLoS One*. 2017;12(2):e0172961.
8. Schafers S, Naunheim R, Vijayan A, Tobin G. Incidence of hypoglycemia following insulin-based acute stabilization of hyperkalemia treatment. *J Hosp Med*. 2012;7(3):239-242.
9. Al-Sharefi A, Quinton R, & Roberts G. In reference to: "Preventing hypoglycemia following treatment of hyperkalemia in hospitalized patients." *J Hosp Med* 2019;14(6). <https://doi.org/10.12788/jhm.3209>.
10. Nirantharakumar K, Marshall T, Kennedy A, Narendran P, Hemming K, Coleman JJ. Hypoglycaemia is associated with increased length of stay and mortality in people with diabetes who are hospitalized. *Diabet Med*. 2012;29(12):e445-448.
11. UK Renal Association. Clinical Practice Guidelines: Treatment of acute hyperkalaemia in adults. <https://renal.org/wp-content/uploads/2020/07/RENAL-ASSOCIATION-HYPERKALAEMIA-GUIDELINE-2020.pdf>. Published 2020. Updated 21st July 2020. Accessed 17th August 2020.
12. National Diabetes Inpatient Audit. National Diabetes Inpatient Audit:are hospitals providing good quality care? A summary report of the National Diabetes Inpatient Audit for England and Wales 2017. [https://www.diabetes.org.uk/resources-s3/2018-06/NaDIA\\_%20Summary\\_v6.pdf?\\_ga=2.204443347.250004555.1529573739-1251873676.1529573739](https://www.diabetes.org.uk/resources-s3/2018-06/NaDIA_%20Summary_v6.pdf?_ga=2.204443347.250004555.1529573739-1251873676.1529573739). Published 2017. Accessed 27/06/2020.

- Accepted Article
13. Boughton CK, Dixon D, Goble E, et al. Preventing Hypoglycemia Following Treatment of Hyperkalemia in Hospitalized Patients. *J Hosp Med*. 2019;14(5):284-287.
  14. Estep P, Efird L. Evaluation of Hypoglycemia Incidence and Risk Factors in Patients Treated with IV Insulin Aspart for Hyperkalemia. *Endocrinol Diabetes Res*. 2015;1(1).
  15. Scott NL, Klein LR, Cales E, Driver BE. Hypoglycemia as a complication of intravenous insulin to treat hyperkalemia in the emergency department. *Am J Emerg Med*. 2019;37(2):209-213.
  16. Jacob BC, Peasah SK, Chan HL, Niculas D, Shogbon Nwaesei A. Hypoglycemia Associated With Insulin Use During Treatment of Hyperkalemia Among Emergency Department Patients. *Hosp Pharm*. 2019;54(3):197-202.
  17. Chothia MY, Halperin ML, Rensburg MA, Hassan MS, Davids MR. Bolus administration of intravenous glucose in the treatment of hyperkalemia: a randomized controlled trial. *Nephron Physiol*. 2014;126(1):1-8.
  18. Groen J, Willebrands AF, Kamminga CE, Van Schothorst HK, Godfried EG. Effects of glucose administration on the potassium and inorganic phosphate content of the blood serum and the electrocardiogram in normal individuals and in non-diabetic patients. *Acta Med Scand*. 1952;141(5):352-366.

## Tables and Figures

Factor	Hypoglycaemia group (n=24)	No hypoglycaemia group (n=108)	Univariate OR	95% CI	p-value
Age in years [median (range)] <sup>a</sup>	63.0 (19-95)	67.5 (19-95)	0.987/year	0.960-1.014	0.345
Sex (M versus F) <sup>b</sup>	79.2% male (n = 19/24)	62.0% male (n = 67/108)	2.325	0.806-6.705	0.118
Recorded diagnosis of diabetes <sup>b</sup>	20.8% (n=4/24)	42.6% (n=46/108)	0.355	0.123-1.020	0.054
Prescribed diabetes treatment <sup>b</sup>	16.7% (n=4/24)	35.2% (n=38/108)	0.368	0.117-1.156	0.087
Prescribed insulin for diabetes <sup>b</sup>	16.7 % (n=4/24)	25.0% (n=27/108)	0.600	0.188-1.911	0.387
Prescribed sulphonylurea for diabetes <sup>NB</sup>	0 % (n=0/24)	10.2 % (n=11/108)	-	-	-
CKD Diagnosis (Compared with CKD Stages 1 and 2) <sup>b</sup>					
Stage 3	29.2% (n = 7/24)	30.6% (n = 33/108)	1.594	0.572-4.438	0.372
Stage 4	12.5% (n = 3/27)	15.7% (n = 17/108)	1.511	0.407-5.606	0.537
Stage 5	12.5% (n = 3/27)	8.3% (n = 9/108)	1.133	0.212-6.053	0.884

**Table 1. Predictors of post-treatment hypoglycaemia – Patient Factors**

*OR –odds ratio; CI – confidence interval; CKD- chronic kidney disease*

*Statistical test applied: <sup>a</sup> Mann Whitney U Test, <sup>b</sup>Chi-Square test*

*<sup>NB</sup> – Chi Square test could not be applied to this variable due to no responses in one group*

	Hypoglycaemia (n=27)	No hypoglycaemia (n=201)	<i>p</i> -value
Median pre-treatment K <sup>+</sup> level <i>mmol/L</i> <sup>a</sup> (Range)	6.1 (5.2-8)	6.3 (5.4-8.2)	<b>0.024</b>
Median pre-treatment CBG level <i>mmol/L</i> <sup>a</sup> (Range)	5.9 (4.1-11.2)	7.6 (3.7-31.3)	<b>0.000</b>
Median creatinine <i>mmol/L</i> <sup>a</sup> (Range)	246 (53-684)	170 (50-2114)	0.669
Mean maximal reduction in K <sup>+</sup> <i>mmol/L</i> <sup>b</sup> (+/- SEM)	0.77 +/- 0.09	0.74 +/- 0.05	p = 0.779

**Table 2. Predictors of post-treatment hypoglycaemia – Episode-level Factors**

*CBG – capillary blood glucose; SEM – standard error of the mean*

*Statistical test applied: <sup>a</sup> Mann Whitney U Test, <sup>b</sup> independent samples t-test*







Figure 1. Time to development of hypoglycaemia following Gwl infusion

**Incidence of Post Treatment Hypoglycaemia Depending on Pre Treatment Glucose**




**Figure 2. Incidence of Post Treatment Hypoglycaemia with Glucose Above and Below 7 mmol/l**

Authors	Publication date	Population	Treatment protocol	% developing hypoglycaemia	Identified Risk Factors	Limitations
Apel et al <sup>6</sup>	2014	Inpatients on haemodialysis (USA) (n = 221 episodes)	10 units insulin + 25g dextrose (94%)	13.1% (29/221 episodes)	Absence of DM  Not taking medication for DM  Lower pre-treatment glucose	Hypoglycaemia defined as CBG <3.3 mmol/L  Did not distinguish type of DM  Haemodialysis patients only  Analysed "per episode", not "per patient"
Boughton et al <sup>13</sup>	2019	Adult (non critical care) inpatients (UK) (n = 662 episodes)	10 units insulin + 20g dextrose	17.5% (116/662 episodes)	Older age  Low body weight  Lower pre-treatment glucose	Did not distinguish type of DM  Analysed "per episode", not "per patient"
Coca et al <sup>7</sup>	2017	Adult (non haemodialysis) inpatients (Spain) (n = 164 episodes)	10 units insulin + 50g dextrose + 40mg furosemide	6.7% (11/164 episodes)	Lower pre-treatment glucose	No factors reached significance  Did not distinguish type of DM  Analysed "per episode", not "per patient"
Estep et al <sup>14</sup>	2015	All adult inpatients (USA) (n= 86)	Variable dose of both insulin and dextrose	17.4% (15/86 patients)	Lower pre-treatment glucose	No standardised treatment protocol  Limited data collection
Jacob et al <sup>16</sup>	2019	Adult admissions via emergency department (USA) (n = 172)	10 units of insulin with 25g dextrose	19.8% (34/172 patients)	Lower pre-treatment glucose	Did not distinguish type of DM

					Higher pre-treatment potassium	
Schafer et al <sup>8</sup>	2012	Adult inpatients (USA) (n = 219)	Variable dose of both insulin and glucose	8.7% (19/219 patients)	Low body weight	No standardised treatment protocol  No/minimal statistical analysis  Did not distinguish type of DM
Scott et al <sup>15</sup>	2019	Adult admissions via emergency department (USA) (n = 409)	Variable dose of both insulin and dextrose	16.6% (68/409 patients)	Lower pre-treatment glucose  Higher insulin dose  Lower dose of dextrose	No standardised treatment protocol  Did not distinguish type of DM

**Table 3. Previous studies investigating iatrogenic hypoglycaemia following Gwl infusions**



Give 10 units soluble insulin in 25g glucose **PLUS**  
Give 25g glucose load (10% glucose @50ml/hr) over 5 hours

**OR**

*Give 75-100g\* glucose-only infusion; if moderate hyperkalaemia and clinically stable  
(in approved pilot study setting)*

Give 10 units soluble insulin in 25g glucose

**Figure 3. Proposed changes to glucose-insulin infusion treatment section in UK Renal Association Hyperkalaemia guideline**

*\* optimum dose to be determined by further study*