



Tikhonov-Fenichel reduction for parameterized critical manifolds with applications to chemical reaction networks

Feliu, Elisenda; Kruff, Niclas; Walcher, Sebastian

Published in:
Journal of Nonlinear Science

DOI:
[10.1007/s00332-020-09610-3](https://doi.org/10.1007/s00332-020-09610-3)

Publication date:
2020

Document version
Publisher's PDF, also known as Version of record

Document license:
[CC BY](#)

Citation for published version (APA):
Feliu, E., Kruff, N., & Walcher, S. (2020). Tikhonov-Fenichel reduction for parameterized critical manifolds with applications to chemical reaction networks. *Journal of Nonlinear Science*, 30(4), 1355-1380.
<https://doi.org/10.1007/s00332-020-09610-3>



Tikhonov–Fenichel Reduction for Parameterized Critical Manifolds with Applications to Chemical Reaction Networks

Elisenda Feliu¹ · Niclas Kruff² · Sebastian Walcher²

Received: 23 May 2019 / Accepted: 14 January 2020
© The Author(s) 2020

Abstract

We derive a reduction formula for singularly perturbed ordinary differential equations (in the sense of Tikhonov and Fenichel) with a known parameterization of the critical manifold. No a priori assumptions concerning separation of slow and fast variables are made, or necessary. We apply the theoretical results to chemical reaction networks with mass action kinetics admitting slow and fast reactions. For some relevant classes of such systems, there exist canonical parameterizations of the variety of stationary points; hence, the theory is applicable in a natural manner. In particular, we obtain a closed form expression for the reduced system when the fast subsystem admits complex-balanced steady states.

Keywords Singular perturbation · Critical manifold · Chemical reaction network · Complex balancing

Mathematics Subject Classification 92C45 · 34E15 · 80A30 · 13P10

Communicated by Dr. Alain Goriely.

✉ Sebastian Walcher
walcher@matha.rwth-aachen.de

Elisenda Feliu
efeliu@math.ku.dk

Niclas Kruff
niclas.kruff@matha.rwth-aachen.de

¹ Department of Mathematical Sciences, University of Copenhagen, Universitetsparken 5, 2100 Copenhagen, Denmark

² Lehrstuhl A für Mathematik, RWTH Aachen, 52056 Aachen, Germany

1 Introduction

A fundamental result on singular perturbation reductions, due to Tikhonov and Fenichel, allows to reduce the dimension of an ordinary differential equation with a small positive parameter in the asymptotic limit when the parameter approaches zero. This theorem has numerous applications in the sciences, in particular to chemical and biochemical reaction networks, especially quasi-steady state (QSS) for certain chemical species, and partial equilibrium approximation (PEA) for slow and fast reactions. However, the application of Tikhonov's and Fenichel's theory to reaction networks may pose some computational problems, and the purpose of the present paper is to address and resolve one of these problems.

Throughout the present work, we assume that a suitable small parameter (in a system possibly depending on several parameters) has been identified; hence, we deal with a singularly perturbed ordinary differential equation. However, we do not assume the equation to be given in separated fast and slow variables, so the usual version of the reduction theorem is not directly applicable. In applications, fast–slow variable separation is frequently not satisfied a priori and worse, there may be no explicit way to rewrite the system in fast–slow form. Generally one may circumvent (and to some extent resolve) this problem by resorting to an “implicit” version of the reduction, which admits the critical submanifold of phase space as an invariant set, but this approach may also encounter computational feasibility problems.

Given this background, we derive in the present paper an explicit singular perturbation reduction that is applicable whenever a parameterization of the critical manifold is known. This reduction formalism seems particularly useful for reaction networks when the partial equilibrium approximation is applicable, since in many instances varieties of stationary points admit a canonical parameterization. We study this scenario in detail, and our results include a general closed form reduction formula, based on stoichiometry alone, for fast subsystems that are complex balanced, as well as a discussion of linear attractivity properties of slow manifolds.

The paper is organized as follows. After some preliminary work (mostly recalling notions and results from the literature) we derive in Sect. 2 a general formula for Tikhonov–Fenichel reduction when a (possibly local) parameterization of the critical manifold is given, and also consider some special cases. We illustrate the procedure by some small examples, briefly indicating that the range of applications is not restricted to chemical reaction networks. The setting of reaction networks is studied in Sect. 3. To finish the paper we discuss some special examples.

2 A Reduction Formula

We consider the singular perturbation reduction of ordinary differential equations, with no a priori assumption on separated (slow and fast) variables. Thus let $U \subseteq \mathbb{R}^n$ be open, $\varepsilon_0 > 0$, and let h be a smooth function in some neighborhood of $U \times [0, \varepsilon_0)$, with values in \mathbb{R}^n . This defines a parameter-dependent system of ordinary differential equations, viz.

$$\dot{x} = h^{(0)}(x) + \varepsilon h^{(1)}(x) + \varepsilon^2 \dots, \quad x \in U, \quad \varepsilon \geq 0, \tag{1}$$

and rewritten in slow time scale $\tau = \varepsilon t$ we have a singularly perturbed system

$$x' = \frac{1}{\varepsilon} h^{(0)}(x) + h^{(1)}(x) + \varepsilon \dots, \quad x \in U, \quad \varepsilon \geq 0. \tag{2}$$

Here $h^{(0)}$ is called the fast part and $h^{(1)}$ the slow part of either system. We focus on the behavior of (1), (2) as $\varepsilon \rightarrow 0$, and we will restrict attention to scenarios for which, modulo a coordinate transformation, the classical singular perturbation theorems of Tikhonov (1952) and Fenichel (1979) are applicable. (To be specific, we refer to the version of Tikhonov’s theorem as given in Verhulst 2005, Theorem 8.1 ff.; see also Goeke and Walcher 2014, Sect. 2.1.)

The focus of the present paper is on singularly perturbed systems which are not in “Tikhonov standard form” with separated slow and fast variables, but admit a transformation to such a form. Finding reductions explicitly is relevant for various applications. In Sect. 3 we will discuss in detail the application to chemical reaction networks.

The earliest result about reduction in a coordinate-free setting is due to Fenichel (see Fenichel 1979, Sect. 5, in particular Lemma 5.4), who discussed the case when the critical manifold is given explicitly as the graph of some function. Generally there exist intrinsic conditions for the existence of a coordinate transformation to Tikhonov standard form (see Fenichel 1979, Sect. 5 and Noethen and Walcher 2011, Proposition 2.1), but such a transformation frequently cannot be obtained in explicit form. A general implicit reduction procedure was developed in Goeke and Walcher (2014); Noethen and Walcher (2011). Building on this, in the present paper we derive a version of the reduced system that can be computed explicitly when a parameterization of the critical manifold is known.

2.1 Review: Tikhonov–Fenichel Reduction

We recall some basic results on coordinate-independent Tikhonov–Fenichel reduction from Goeke and Walcher (2014); Noethen and Walcher (2011) which are applicable whenever the critical manifold is given implicitly. In particular we refer to Goeke and Walcher (2014), Theorem 1 and the subsequent remarks.

Proposition 1 *Let system (1) be given, and denote by $\mathcal{V}(h^{(0)})$ the zero set of $h^{(0)}$. Moreover, let $0 < r < n$ and set $s := n - r > 0$.*

(a) *Assume that $a \in \mathcal{V}(h^{(0)})$ has the following properties.*

- *There exists a neighborhood \tilde{U} of a such that $\text{rank } Dh^{(0)}(x) = r$ for all $x \in Z := \mathcal{V}(h^{(0)}) \cap \tilde{U}$; in particular Z is an s -dimensional submanifold of \mathbb{R}^n .*
- *For all $x \in Z$ there is a direct sum decomposition*

$$\mathbb{R}^n = \text{Ker } Dh^{(0)}(x) \oplus \text{Im } Dh^{(0)}(x).$$

- For all $x \in Z$ the nonzero eigenvalues of $Dh^{(0)}(x)$ have real part < 0 .

Then in some neighborhood of a there exists an invertible coordinate transformation from (1) to Tikhonov standard form

$$\begin{aligned} \dot{y}_1 &= \varepsilon f_1(y_1, y_2) + O(\varepsilon^2) \\ \dot{y}_2 &= f_2(y_1, y_2) + O(\varepsilon) \end{aligned}$$

with separated slow and fast variables; moreover, the fast system satisfies a linear stability condition.

- (b) Conversely, the conditions in part (a) are necessary for the existence of a local coordinate transformation to Tikhonov standard form.
- (c) One may choose \tilde{U} such that there exists a product decomposition with functions $\mu(x)$ taking values in $\mathbb{R}^{r \times 1}$, $P(x)$ taking values in $\mathbb{R}^{n \times r}$, such that

$$h^{(0)}(x) = P(x)\mu(x), \quad \text{for all } x \in Z; \tag{3}$$

moreover $\text{rank } P(a) = r$, $\text{rank } D\mu(a) = r$ and

$$Z = \mathcal{V}(\mu) \cap \tilde{U}.$$

Here the entries of μ may be taken as any r entries of $h^{(0)}$ that are functionally independent at a .

- (d) The following system (in slow time) is defined on \tilde{U} , and admits Z as an invariant set:

$$x' = \left(I_n - P(x)A(x)^{-1}D\mu(x) \right) h^{(1)}(x), \tag{4}$$

with

$$A(x) := D\mu(x)P(x).$$

The restriction of this system to Z corresponds to the reduced equation in Tikhonov's theorem.

We refer to Z as the local critical manifold (or local asymptotic slow manifold) of system (1). The decomposition (3) for smooth vector fields is a consequence of the implicit function theorem, and thus in general an explicit determination may not be possible. But the decomposition can be determined algorithmically whenever the right-hand side of (1) is polynomial or rational; see Goeke and Walcher (2014).

- Remark 1**
- (a) Note that $Dh^{(0)}(x) = P(x)D\mu(x)$ on Z , due to $\mu(x) = 0$. Since $P(x)$ has full rank on Z , $Dh^{(0)}(x)$ and $P(x)$ have the same column space.
 - (b) The eigenvalues of $A(x)$, $x \in Z$, are the nonzero eigenvalues of $Dh^{(0)}(x)$ whenever the latter has rank r ; see Goeke and Walcher (2014), Remark 3.

(c) We call

$$Q(x) := I_n - P(x)A(x)^{-1}D\mu(x) \tag{5}$$

the *projection operator* of the reduction. For each x this is a linear projection of rank $s = n - r$ which sends every element of \mathbb{R}^n to its kernel component from the kernel-image decomposition with respect to $Dh^{(0)}(x)$. (In the special case when the critical manifold is the graph of some function, a projection matrix was determined in Fenichel 1979, Lemma 5.4.)

(d) Formally system (4) is defined whenever $A(x)$ is invertible, and by Fenichel’s results it corresponds to a reduced system as $\varepsilon \rightarrow 0$ whenever all eigenvalues of $A(x)$ have nonzero real part (normal hyperbolicity).

Remark 2 The reduced system may just have the form $x' = 0$; in particular this occurs in the following scenario: $h^{(0)}$ always admits $n - s$ independent first integrals near any point of Z , and locally every point of Z is uniquely determined as an intersection of Z with suitable level sets of these first integrals; see Goeke and Walcher (2014), Sect. 2.3. Now, in the special case when $h^{(1)}$ admits the same first integrals, then Z as well as every intersection of Z with level sets is invariant for the reduced equation, meaning that every point of Z is invariant, thus stationary. However, the only information to be gained from $x' = 0$ for small $\varepsilon > 0$ is that system (2) restricted to the invariant manifold has right-hand side of order ε or higher. (Generally the reduced system (4) in slow time represents only the $O(1)$ term in ε .)

While Proposition 1 provides a general coordinate-free approach to singular perturbation reduction, the critical manifold Z is given only implicitly via the zeros of $h^{(0)}$, and one cannot generally expect an explicit reduction to a system in \mathbb{R}^s . Moreover, there may be a problem with the feasibility of the computations, in particular with the computation of the projection matrix Q . Therefore, it is natural to search for simplified reduction procedures in special circumstances. One notable scenario appears when a parameterization for the critical manifold is explicitly known, and we will next discuss reduction in this case.

2.2 Parameterized Critical Manifolds

We keep the assumptions and notation from Proposition 1, in particular the decomposition (3), the s -dimensional local critical manifold Z (being the zero set of μ , as well as of $h^{(0)}$), and the reduced system

$$x' = Q(x)h^{(1)}(x) \quad \text{on } Z. \tag{6}$$

Now assume that there is an open set $W \subseteq \mathbb{R}^s$ and a smooth parameterization

$$\Phi: W \rightarrow Z, \quad \text{rank } D\Phi(v) = s \text{ for all } v \in W. \tag{7}$$

Then every solution $x(t)$ of (6) with initial value in $\Phi(W)$ can be written in the form

$$x(t) = \Phi(v(t)),$$

for t in some neighborhood of 0, and differentiation yields

$$D\Phi(v(t)) v'(t) = x'(t) = Q(\Phi(v(t))) \cdot h^{(1)}(\Phi(v(t))). \quad (8)$$

The remaining task is to simplify this expression.

Theorem 1 (a) For every $v \in W$ there exists a unique $R(v) \in \mathbb{R}^{s \times n}$ such that

$$Q(\Phi(v)) = D\Phi(v) \cdot R(v).$$

(b) The reduced system, in parameterized version (8), is given by

$$v' = R(v) \cdot h^{(1)}(\Phi(v)). \quad (9)$$

(c) The matrix $R(v)$ is uniquely determined by the conditions

$$R(v) \cdot P(\Phi(v)) = 0 \quad \text{and} \quad R(v) \cdot D\Phi(v) = I_s,$$

and therefore can be obtained from the matrix equation

$$R(v) \cdot (D\Phi(v) \mid P(\Phi(v))) = (I_s \mid 0)$$

with $(D\Phi(v) \mid P(\Phi(v)))$ invertible. In particular, $v \mapsto R(v)$ is smooth.

(d) For every $x \in Z$ let $L(x) \in \mathbb{R}^{s \times n}$ be of full rank s and such that $L(x) Dh^{(0)}(x) = 0$; equivalently $L(x)P(x) = 0$. Moreover, define $L^*(v) := L(\Phi(v))$. Then

$$R(v) = (L^*(v) D\Phi(v))^{-1} L^*(v),$$

and the reduced system, in parameterized form, is given by

$$v' = (L^*(v) D\Phi(v))^{-1} L^*(v) h^{(1)}(\Phi(v)). \quad (10)$$

Proof For every $v \in W$ one has $h^{(0)}(\Phi(v)) = 0$, and by differentiation

$$Dh^{(0)}(\Phi(v))D\Phi(v) = 0.$$

Thus the image of $D\Phi(v)$ is contained in the kernel of $Dh^{(0)}(\Phi(v))$, and these two vector spaces have dimension s ; hence, they are equal. In turn, for $x \in Z$ the kernel of $Dh^{(0)}(x)$ is by construction equal to the image of $Q(x)$. Thus, for every v the matrices $Q(\Phi(v))$ and $D\Phi(v)$ have the same column space, and the latter has full rank. Therefore, every column of $Q(\Phi(v))$ is a unique linear combination of the

columns of $D\Phi(v)$. Rewritten in matrix language, this is the assertion of part (a). Part (b) is now obvious from Eq. (8) and injectivity of $D\Phi(v)$.

The first condition given in part (c) is a consequence of part (a), the identity $Q(x) \cdot P(x) = 0$ for all $x \in Z$ (which is readily verified from the defining Eq. (5)), and the fact that $D\Phi(v)$ is an injective linear map. The second condition follows from the fact that $D\Phi(v) \cdot R(v) = Q(\Phi(v))$ is a projection of rank s , by using Lemma 1 in ‘‘Appendix’’. Invertibility of the matrix $(D\Phi(v) \mid P(\Phi(v)))$ follows from the direct kernel–image decomposition with respect to $Dh^{(0)}(\Phi(v))$, since the columns of $D\Phi(v)$ span the kernel and the columns of $P(\Phi(v))$ span the image (see Goeke and Walcher 2014 for more details).

To prove (d), first recall from Remark 1 that $Dh^{(0)}(x)$ and $P(x)$ have the same column space, therefore $L(x)P(x) = 0$ on Z . This and $R(v)P(\Phi(v)) = 0$ from part (c) imply that $R(v) = \Lambda(v)L(\Phi(v))$ for all $v \in W$, with $\Lambda(v) \in \mathbb{R}^{s \times s}$ uniquely determined. Using now the second condition in part (c), we have

$$\Lambda(v)L(\Phi(v))D\Phi(v) = I_s,$$

hence $L(\Phi(v))D\Phi(v)$ is invertible and

$$\Lambda(v) = (L(\Phi(v))D\Phi(v))^{-1},$$

which leads to the asserted expression. □

Remark 3 (a) The second equation in part (c) of the theorem shows that $R(v)$ is (according to definition) a left inverse of $D\Phi(v)$. This observation allows to obtain (9) directly from (8). As a consequence, one obtains the identities

$$R(v) D\Phi(v) R(v) = R(v) \text{ and } D\Phi(v) R(v) D\Phi(v) = D\Phi(v);$$

but we note that in general $R(v)$ will not be the Moore–Penrose inverse of $D\Phi(v)$.

- (b) The characterization in part (d) of the theorem shows that $R(v)$ can be computed by standard linear algebra: One only needs to determine the left kernel of $Dh^{(0)}(x)$ to find $L(x)$, and then compute products and inverses of certain matrices. There is no need for explicit knowledge of the projection matrix Q , or of the matrix P from the decomposition, as part (c) might suggest. However, the column space of $Dh^{(0)}(x)$, $x \in Z$, is a crucial ingredient.
- (c) On the other hand, knowledge of P and μ seems indispensable for the computation of $A(x) = D\mu(x)P(x)$, and of $A(\Phi(v))$. Note that the eigenvalues of the latter provide direct information on the stability of the critical manifold; see Remark 1 (b).

We consider some special cases in more detail.

Corollary 1 (a) *Assume that*

$$\Phi(v) = \begin{pmatrix} \Phi_1(v) \\ \Phi_2(v) \end{pmatrix}, \quad \text{with } \Phi_1(v) \in \mathbb{R}^s \text{ and } D\Phi_1(v) \text{ invertible, for all } v \in W,$$

and partition

$$P(x) = \begin{pmatrix} P_1(x) \\ P_2(x) \end{pmatrix} \quad \text{with } P_1(x) \in \mathbb{R}^{s \times r}.$$

Then

$$R(v) = (R_1(v) \mid R_2(v))$$

with

$$\begin{aligned} R_1(v) &= I_s - P_1 \left(D\Phi_2 D\Phi_1^{-1} P_1 - P_2 \right)^{-1} D\Phi_2 D\Phi_1^{-1} \\ R_2(v) &= P_1 \left(D\Phi_2 D\Phi_1^{-1} P_1 - P_2 \right)^{-1}. \end{aligned}$$

In these expressions the argument of $D\Phi_1$ and $D\Phi_2$ is v and the argument of P_1 and P_2 is $\Phi(v)$.

- (b) (Fenichel 1979, Lemma 5.4:) In the special case when $\Phi_1(v) = v$, thus the critical manifold is the graph of Φ_2 , we get

$$\begin{aligned} R_1(v) &= I_s - P_1 (D\Phi_2 P_1 - P_2)^{-1} D\Phi_2 \\ R_2(v) &= P_1 (D\Phi_2 P_1 - P_2)^{-1}. \end{aligned}$$

Proof With R_i given as above one verifies

$$R_1 D\Phi_1 + R_2 D\Phi_2 = I_s \text{ and } R_1 P_1 + R_2 P_2 = 0$$

by direct computation. Rewriting, one obtains

$$(R_1 \mid R_2) \begin{pmatrix} D\Phi_1 P_1 \\ D\Phi_2 P_2 \end{pmatrix} = (I_s \mid 0),$$

which is the defining property of R in Theorem 1. □

Remark 4 (a) Up to a relabeling of variables in \mathbb{R}^n , a partitioning for $\Phi(v)$ as required in Corollary 1 always exists locally, due to the rank condition on the derivative. Moreover, by local invertibility of Φ_1 there exists a parameterization

$$\Psi : w \mapsto \begin{pmatrix} w \\ \Phi_2 \circ \Phi_1^{-1}(w) \end{pmatrix}$$

as in part (b) of the corollary. But in general an explicit determination of the inverse of Φ_1 is not possible.

- (b) In the special case $\Phi_1(v) = v$ the reduction formula was also derived by Stiefenhofer (1998), Eq. (2.13), with a different proof.

(c) In the yet more special case that $\Phi_1(v) = v$ and $\Phi_2(v) = 0$ the procedure yields the familiar quasi-steady-state reduction. Indeed, in this case one has $\mu(x) = x_2$ for $x = (x_1, x_2)^{\text{tr}}$ and $x_1 \in \mathbb{R}^s$, thus $R_1 = I_s$, $R_2 = -P_1 P_2^{-1}$ and the reduced equation is

$$v' = \left(I_s \mid -P_1((v, 0)^{\text{tr}})P_2^{-1}((v, 0)^{\text{tr}}) \right) \begin{pmatrix} h_1^{(1)}((v, 0)^{\text{tr}}) \\ h_2^{(1)}((v, 0)^{\text{tr}}) \end{pmatrix}.$$

Ignoring higher order terms in ε (which are irrelevant for Tikhonov–Fenichel reduction) and renaming variables, one obtains the same system by setting the second part of

$$\begin{pmatrix} \dot{x}_1 \\ \dot{x}_2 \end{pmatrix} = \begin{pmatrix} P_1 \\ P_2 \end{pmatrix} \cdot x_2 + \varepsilon \begin{pmatrix} h_1^{(1)} \\ h_2^{(1)} \end{pmatrix} + \dots$$

equal to zero, solving for x_2 , substituting into the first part and passing to slow time. This is another proof of the fact that singular perturbation reduction and QSS reduction agree when the critical manifold is a coordinate subspace. (The first proof was given in Goeke et al. 2017, Proposition 5.)

Finally, with a view on chemical reaction networks, we address conservation laws.

Proposition 2 *Let the smooth real-valued function ψ be defined on some open subset of U which has non-empty intersection with $\Phi(W)$, and assume that ψ is a first integral of system (1) for every ε . Then $\tilde{\psi} := \psi \circ \Phi$ is constant or a first integral of the parameterized reduced system (9).*

Proof By Lax and Walcher (2018), Proposition 8 the restriction of ψ to the critical manifold Z is also a first integral of the reduced system (4), thus $D\psi(x)Q(x)h^{(1)}(x) = 0$ for all $x \in Z$. Therefore,


$$\begin{aligned} D\tilde{\psi}(v)R(v)h^{(1)}(\Phi(v)) &= D\psi(\Phi(v))D\Phi(v)R(v)h^{(1)}(\Phi(v)) \\ &= D\psi(\Phi(v))Q(\Phi(v))h^{(1)}(\Phi(v)) \\ &= 0, \end{aligned}$$

which is the characterizing property for first integrals of system (9). □

2.3 Illustrative Examples

The following small examples have the primary function to illustrate the arguments and reduction procedures from the previous subsection.

Example 1 We consider a (hypothetical) slow–fast system, with fast reaction



and slow reaction

$$X_1 + X_3 \rightleftharpoons 2X_2,$$

with associated differential equation (according to the procedure from Sect. 3.1)

$$\begin{aligned}\dot{x}_1 &= -k_1x_1x_2 + k_{-1}x_3 - \varepsilon k_2x_1x_3 + \varepsilon k_{-2}x_2^2 \\ \dot{x}_2 &= -k_1x_1x_2 + k_{-1}x_3 + 2\varepsilon k_2x_1x_3 - 2\varepsilon k_{-2}x_2^2 \\ \dot{x}_3 &= k_1x_1x_2 - k_{-1}x_3 - \varepsilon k_2x_1x_3 + \varepsilon k_{-2}x_2^2.\end{aligned}$$

The critical manifold Z is determined by $Kx_1x_2 = x_3$, with $K = k_1/k_{-1}$, and we have $P = (1, 1, -1)^t$, $\mu(x) = (-k_1x_1x_2 + k_{-1}x_3)$. A parameterization of Z is given by

$$\Phi: \mathbb{R}_{\geq 0}^2 \rightarrow \mathbb{R}^3, \quad \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} \mapsto \begin{pmatrix} v_1 \\ v_2 \\ K v_1 v_2 \end{pmatrix},$$

hence to determine $R(v)$ via Theorem 1(c) we have to solve

$$R(v) \cdot \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ K v_2 & K v_1 & -1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}.$$

By straightforward calculations, one obtains

$$R(v) = \frac{1}{1 + K(v_1 + v_2)} \begin{pmatrix} 1 + K v_1 & -K v_1 & 1 \\ -K v_2 & 1 + K v_2 & 1 \end{pmatrix}.$$

With

$$h^{(1)}(\Phi(v)) = (-k_2 K v_1^2 v_2 + k_{-2} v_2^2) \cdot \begin{pmatrix} 1 \\ -2 \\ 1 \end{pmatrix}$$

the reduced system becomes


$$\begin{pmatrix} v_1' \\ v_2' \end{pmatrix} = \frac{k_2 K v_1^2 v_2 - k_{-2} v_2^2}{1 + K(v_1 + v_2)} \begin{pmatrix} -2 - 3K v_1 \\ 1 + 3K v_2 \end{pmatrix}.$$

This is a case where the critical manifold is the graph of the rational function $(x_1, x_2) \mapsto Kx_1x_2$; thus Corollary 1 would also be applicable. Moreover, we get

$$A(\Phi(v)) = D\mu(\Phi(v))P(\Phi(v)) = -(k_1(v_1 + v_2) + k_{-1}) < 0 \text{ on } \mathbb{R}_{\geq 0}^2,$$

hence linear stability of the critical manifold follows by Remark 3(b).

Example 2 As a non-hypothetical variant, we discuss the system with the same fast reaction as in Example 1, but with slow reaction



and associated differential equation

$$\begin{aligned} \dot{x}_1 &= -k_1 x_1 x_2 + k_{-1} x_3 + \varepsilon k_2 x_3 \\ \dot{x}_2 &= -k_1 x_1 x_2 + k_{-1} x_3 \\ \dot{x}_3 &= k_1 x_1 x_2 - k_{-1} x_3 - \varepsilon k_2 x_3 \end{aligned}$$

after discarding the equation for x_4 . This is Michaelis–Menten with slow degradation of complex to enzyme and product. Here $R(v)$ is the same as in the previous example, and


$$h^{(1)}(\Phi(v)) = k_2 K v_1 v_2 \cdot \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}.$$

The reduced system becomes

$$\begin{pmatrix} v'_1 \\ v'_2 \end{pmatrix} = \frac{k_2 K v_1 v_2}{1 + K(v_1 + v_2)} \begin{pmatrix} K v_1 \\ -(1 + K v_2) \end{pmatrix},$$

and we note a further built-in reduction: The differential equation for the reaction network admits the first integral $\psi = x_1 + x_3$ from stoichiometry; hence, by Proposition 2 the reduced equation inherits the first integral $\tilde{\psi} = v_1 + K v_1 v_2$. Thus one ends up with a one-dimensional reduced equation, as it should be.

Example 3 For contrast, consider the hypothetical slow–fast system with fast reaction



and the same slow reaction as in Example 1. The differential equation now becomes

$$\begin{aligned} \dot{x}_1 &= -2k_1 x_1^2 x_2^2 + 2k_{-1} x_3^3 - \varepsilon k_2 x_1 x_3 + \varepsilon k_{-2} x_2^2 \\ \dot{x}_2 &= -2k_1 x_1^2 x_2^2 + 2k_{-1} x_3^3 + 2\varepsilon k_2 x_1 x_3 - 2\varepsilon k_{-2} x_2^2 \\ \dot{x}_3 &= 3k_1 x_1^2 x_2^2 - 3k_{-1} x_3^3 - \varepsilon k_2 x_1 x_3 + \varepsilon k_{-2} x_2^2, \end{aligned}$$

and the critical manifold Z is given by $K x_1^2 x_2^2 = x_3^3$, with $K = k_1/k_{-1}$, and $P = (2, 2, -3)^{\text{tr}}$. A parameterization of Z is given by

$$\Phi: \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} \mapsto \begin{pmatrix} v_1^3 \\ v_2^3 \\ K v_1^2 v_2^2 \end{pmatrix}.$$

It is obvious that

$$L = \begin{pmatrix} 1 & -1 & 0 \\ 3 & 0 & 2 \end{pmatrix}$$

is of rank two and satisfies $L \cdot P = 0$. This yields, by Theorem 1(d),

$$L \cdot D\Phi(v) = L \cdot \begin{pmatrix} 3v_1^2 & 0 \\ 0 & 3v_2^2 \\ 2Kv_1v_2^2 & 2Kv_1^2v_2 \end{pmatrix} = \begin{pmatrix} 3v_1^2 & -3v_2^2 \\ 9v_1^2 + 4Kv_1v_2^2 & 4Kv_1^2v_2 \end{pmatrix}.$$

Finally the reduced system is given by

$$\begin{aligned} v' &= \frac{1}{3v_1v_2(4K(v_1^3 + v_2^3) + 9v_1v_2)} \begin{pmatrix} 4Kv_1^2v_2 & 3v_2^2 \\ -9v_1^2 - 4Kv_1v_2^2 & 3v_1^2 \end{pmatrix} \cdot L \cdot h^{(1)} \\ &= \frac{1}{3v_1v_2(4K(v_1^3 + v_2^3) + 9v_1v_2)} \cdot \\ &\quad \begin{pmatrix} -12K^2v_1^7v_2^3k_2 + 12Kv_1^2v_2^7k_{-2} - 15Kv_1^2v_1^3v_2^4k_2 + 15v_2^8k_{-2} \\ 12K^2v_1^6v_2^4k_2 + 27Kv_1^7v_2^2k_2 - 12Kv_1v_2^8k_{-2} - 15Kv_1^2v_1^5v_2^2k_2 - 12v_1^2v_2^6k_{-2} \end{pmatrix}. \end{aligned}$$

One can further reduce the dimension to one by utilizing the first integral $\psi = 4x_1 + 5x_2 + 6x_3$ from stoichiometry.

In this example we could have used Remark 4(a) and chosen a different parameterization

$$\Psi : \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} \mapsto \begin{pmatrix} v_1 \\ v_2 \\ Kv_1^{2/3}v_2^{2/3} \end{pmatrix},$$

which directly represents Z as the graph of a function. But it seems more natural and convenient to work with a reduced system that has rational right-hand side. Moreover, the second parameterization may obscure the fact that Z is an algebraic variety.

Example 4 Finally we sketch an example that is motivated by mechanics, to indicate that the range of applications does not only include reaction networks. (See Arnold and Anosov 1988 for background, and also Walcher 1991.) Specifically we look at a pair of coupled nonlinear oscillators

$$\begin{pmatrix} \dot{y}_1 \\ \dot{y}_2 \\ \dot{y}_3 \\ \dot{y}_4 \end{pmatrix} = \begin{pmatrix} y_2 \\ -y_1 \\ \omega y_4 \\ -\omega y_3 \end{pmatrix} + f_3(y) + \text{t.h.o.}$$

with irrational $\omega > 0$; thus, we are in a non-resonant scenario, f_3 a homogeneous polynomial vector field of degree three, and “t.h.o.” denoting terms of higher order. We assume that the system is in normal form up to degree three, with the special form

$$f_3(y) = \frac{1}{2} \left(a \cdot (y_1^2 + y_2^2) + b \cdot (y_3^2 + y_4^2) \right) \begin{pmatrix} y_1 \\ y_2 \\ cy_3 \\ cy_4 \end{pmatrix} + \begin{pmatrix} -p_1 y_2 \\ p_1 y_1 \\ -p_2 y_4 \\ p_2 y_3 \end{pmatrix}$$

where $a < 0$, $b > 0$, $c < 0$ are real constants and p_1 , p_2 are linear combinations of $y_1^2 + y_2^2$ and $y_3^2 + y_4^2$. Reduction of the degree three Taylor polynomial by the invariants $y_1^2 + y_2^2$, $y_3^2 + y_4^2$ yields a two-dimensional system, given by

$$\begin{aligned} \dot{x}_1 &= x_1 (ax_1 + bx_2) \\ \dot{x}_2 &= cx_2 (ax_1 + bx_2). \end{aligned} \quad (11)$$

The reduced system is degenerate in our case, admitting a line of stationary points given by $\mu := ax_1 + bx_2 = 0$. The choice of signs ensures that this line lies in the positive quadrant (which is positively invariant), and also that solutions on the invariant lines $x_1 = 0$ resp. $x_2 = 0$ converge to 0 as $t \rightarrow \infty$.

We consider (11) as fast part $h^{(0)}$ of a singularly perturbed system, thus we have

$$P(x) = \begin{pmatrix} x_1 \\ cx_2 \end{pmatrix} \text{ and choose } \Phi(v) = \begin{pmatrix} bv \\ -av \end{pmatrix}, \quad v > 0.$$

A straightforward calculation shows that $A(\Phi(v)) = ab(1-c)v < 0$ whenever $v > 0$, so the line $\mu = 0$ is attracting in the positive quadrant. Moreover, one may choose $L(x) = (cx_2, -x_1)$ and thus obtains the reduced equation

$$v' = \frac{-1}{ab(1-c)} \cdot (ac \ b) h^{(1)}(\Phi(v)) \quad (12)$$

for arbitrary small perturbation $h^{(1)}$. The choice

$$h^{(1)}(x) = \begin{pmatrix} x_1^3 \\ -x_2^4 \end{pmatrix}, \quad h^{(1)}(\Phi(v)) = \begin{pmatrix} b^3 v^3 \\ -a^4 v^4 \end{pmatrix}$$

is compatible with the mechanical context under consideration here and yields a positive stationary point for (12) as well as (11). For the original system, this yields the existence of an invariant torus.

3 Applications to Reaction Networks

While the range of applications of Theorem 1 is not restricted to chemical reaction networks, it is natural to discuss reduction of these reaction networks, for the following reason: The consideration of slow and fast reactions leads to critical manifolds that consist of stationary points of a subnetwork. Extensive previous work on parametrizations of the positive part of the variety of stationary points in chemical reaction networks

(see, e.g., Horn 1972; Craciun et al. 2009; Müller et al. 2016) shows that parametrizations exist and can be explicitly found for several relevant and familiar classes of networks. Thus the results of the previous section can often be applied and reduced systems can be explicitly obtained. In particular, in the case where the fast subnetwork admits a complex-balanced steady state (see below), a closed formula for the reduced system can be readily expressed in terms of the stoichiometry of the network alone (see Proposition 4).

We first recall some general facts about reaction networks and then discuss two special classes. The results will be illustrated by examples.

3.1 Reaction Networks

We briefly recall here the mathematical description of reaction networks according to Feinberg (1995), Horn and Jackson (1972) (see also the recent monograph by Feinberg (2019)), and then outline the general setup for networks with fast and slow reactions, as already suggested by some illustrative examples in the previous section.

A *reaction network* on a set of species $\{X_1, \dots, X_n\}$ is a digraph whose nodes are finite linear combinations of species with nonnegative integer coefficients; each edge is called a reaction. Thus a node is of the form $y = \sum_{i=1}^n a_i X_i$ and is identified with the vector $y = (a_1, \dots, a_n) \in \mathbb{R}^n$. We let x_i denote the concentration of X_i and $x = (x_1, \dots, x_n)$. For each reaction (denoted by $y \rightarrow y'$) we assume given a rate function $w_{y \rightarrow y'}(x) \in \mathbb{R}_{\geq 0}$ for $x \in \mathbb{R}_{\geq 0}^n$. This leads to a system of differential equations describing the evolution of the concentrations in time:

$$\dot{x} = \sum_{\text{reactions } y \rightarrow y'} w_{y \rightarrow y'}(x)(y' - y), \quad x \in \mathbb{R}_{\geq 0}^n. \quad (13)$$

Note that $y' - y \in \mathbb{R}^n$ encodes the net production of each species by the occurrence of the reaction $y \rightarrow y'$. The vector subspace spanned by all the $y' - y$ is called the *stoichiometric subspace* of the reaction network.

It is convenient to write the system in matrix–vector form by introducing the *matrix* N whose columns are the vectors $y' - y$ (after fixing an order of the set of reactions). Then, with $w(x)$ denoting the vector of rate functions in the same order, (13) can be rewritten as

$$\dot{x} = Nw(x), \quad x \in \mathbb{R}_{\geq 0}^n. \quad (14)$$

A frequent choice of rate function is the one from *mass action kinetics*, with

$$w_{y \rightarrow y'}(x) = k_{y \rightarrow y'} \prod_{i=1}^n x_i^{y_i},$$

with $k_{y \rightarrow y'} > 0$ called reaction rate constants and using the convention $0^0 = 1$. For the following we recall some definitions:

Definition 1 Let $x, y \in \mathbb{R}^n$ and $M \in \mathbb{R}^{n \times m}$, with columns $M_1, \dots, M_m \in \mathbb{R}^n$.

(a) For $x \in \mathbb{R}_{>0}^n$ we define

$$x^y := \prod_{i=1}^n x_i^{y_i},$$

noting that the definition may be extended to all $x \in \mathbb{R}^n$ when all y_i are nonnegative integers.

(b) For $x \in \mathbb{R}_{>0}^n$ we define

$$x^M := \begin{pmatrix} x^{M_1} \\ \vdots \\ x^{M_m} \end{pmatrix} \in \mathbb{R}^m,$$

noting that the definition may be extended to all $x \in \mathbb{R}^n$ when all entries of M are nonnegative integers.

(c) The *Hadamard product* $x \circ y$ is defined as the component-wise multiplication of the two vectors x, y , i.e.,

$$x \circ y = \begin{pmatrix} x_1 \cdot y_1 \\ \vdots \\ x_n \cdot y_n \end{pmatrix}.$$

In view of the last definition we can rewrite the reaction network for mass action kinetics in the form

$$\dot{x} = N \cdot w(x) = N \cdot (K \circ x^Y), \tag{15}$$

where $K \in \mathbb{R}_{>0}^m$ (m is the number of reactions) is a vector containing the reaction rate constants and $Y \in \mathbb{R}^{n \times m}$ is the matrix whose columns are the reactant vectors of each reaction, called the kinetic order matrix.

It follows directly from (14) that any vector in the left kernel of N defines a linear first integral, regardless of the form of $w(x)$. These linear first integrals are commonly referred to as conservation laws, and their common level sets are called stoichiometric compatibility classes. If each connected component of the reaction network has exactly one terminal strongly connected component, then all linear first integrals of (14) arise in this way; see Feinberg and Horn (1977). Finally we recall the notion of *deficiency* of the reaction network, which is defined as the number of nodes minus the rank of N minus the number of connected components; see, e.g., Horn (1972) or Feinberg (1972).

We turn now to a scenario with prescribed slow and fast reactions; see also the discussions in Heinrich and Schauer (1983), Lee and Othmer (2009). The subdigraph induced by the fast reactions is itself a reaction network with the same set of species, which we call the *fast subnetwork*. We stipulate that even if some species are not part of any fast reaction, we still consider them as part of the fast subnetwork. We

have a corresponding stoichiometric matrix N_f and rate vector $w_f(x)$, such that, in the notation of Sect. 2,

$$h^{(0)}(x) = N_f w_f(x) = N_f \cdot (K_f \circ x^{Y_f}). \quad (16)$$

Analogously, we have

$$h^{(1)}(x) = N_s w_s(x) = N_s \cdot (K_s \circ x^{Y_s}) \quad (17)$$

for the slow subsystem. Keeping the notation from Sect. 2, we let Z be the zero set of $h^{(0)}$ (possibly restricted to a neighborhood \tilde{U} of some point), and let r denote the rank of $Dh^{(0)}(x)$, $x \in Z$. Then clearly $\text{rank } N_f \geq r$, but the inequality may be strict; see Heinrich and Schauer (1983) and also Sect. 3 of Goeke and Walcher (2014). In the present paper we will, however, restrict attention to the case when equality holds:

Blanket hypothesis We impose on system (16) the conditions of Proposition 1(a) and the additional condition that $\text{rank } N_f = \text{rank } Dh^{(0)}(x) = r$, $x \in Z$.

Due to nonnegativity of concentrations, the points of Z will be in $\mathbb{R}_{\geq 0}^n$. In some instances we will require that the neighborhood \tilde{U} in Proposition 1 is even a subset of $\mathbb{R}_{> 0}^n$, and likewise we will occasionally require that the domain W of the parameterization Φ is a subset of $\mathbb{R}_{> 0}^s$. By our assumption the zero set Z of $h^{(0)}$ has dimension $s = n - r$. Assume now that there exists a smooth parameterization

$$\Phi: W \rightarrow Z$$

with $\text{rank } D\Phi(v) = s$ on W .

Proposition 3 Let system (16) be given, with a parameterization Φ of the critical manifold as in (7), and assume the blanket hypothesis holds on $\Phi(W)$. Let $L_f \in \mathbb{R}^{s \times n}$ be a matrix whose rows form a basis of the left kernel of N_f . Then the matrix $R(v)$ in Theorem 1 is given as

$$R(v) = (L_f D\Phi(v))^{-1} L_f,$$

and the reduced system is

$$v' = (L_f D\Phi(v))^{-1} L_f h^{(1)}(\Phi(v)), \quad v \in W.$$

Furthermore, the column space of the matrix $P(x)$ in any decomposition of $h^{(0)}(x)$ as in Proposition 1(c) equals the column space of N_f .

Proof This follows from Theorem 1(d), since $L_f Dh^{(0)}(x) = L_f N_f D w_f(x) = 0$ on Z . \square

Note that by Remark 2, if $\text{rank } N = \text{rank } N_f$, then the reduced system is given by $v' = 0$.

3.2 Canonical Parameterizations for Some Classes

To find a function $\Phi(v)$ which yields a parameterization of positive steady states of the fast subnetwork, several strategies can be employed. We review here the two most common approaches. Throughout we use the blanket hypothesis, denoting the rank of the stoichiometric matrix N_f by r , the number of species of the full network by n , and let $s = n - r$. For simplicity, we consider mass action kinetics, although several results hold for more general classes of rate functions.

3.2.1 Non-interacting Sets and Rational Parameterizations

As was shown in Feliu and Wiuf (2012), non-interacting sets of species may be utilized to find rational parameterizations of the steady states, given certain conditions. Thus consider a subset of species $\mathcal{Y} = \{X_{i_1}, \dots, X_{i_r}\}$, with the following assumptions.

- (i) For every fast reaction $y \rightarrow y'$, both the sum of the coefficients of the species in \mathcal{Y} in y and the sum of the coefficients in y' are at most one. This means that no pair of species in \mathcal{Y} appear together at one side of a reaction, and further no species appears with coefficient greater than 1.
- (ii) The rank of the submatrix of N_f given by the rows i_1, \dots, i_r is equal to r .
- (iii) Consider the network induced by the fast subnetwork by setting all species not in \mathcal{Y} to zero. For each species X_{i_j} in \mathcal{Y} , there is a directed path from X_{i_j} to 0 in this induced network.

In the nomenclature of Feliu and Wiuf (2012), assumption (i) means that \mathcal{Y} is non-interacting, (ii) means that no conservation law has support in \mathcal{Y} , and (iii) means that there exists a spanning tree rooted at $*$ in the appropriate digraph (see Feliu and Wiuf 2012, Sect. 8 for details).

Let $X_{\ell_1}, \dots, X_{\ell_s}$ be the species not in \mathcal{Y} . If \mathcal{Y} satisfies (i), (ii) and (iii), then the components i_1, \dots, i_r of $h^{(0)}(x)$ form a linear system in x_{i_1}, \dots, x_{i_r} that has a unique solution in terms of $x_{\ell_1}, \dots, x_{\ell_s}$. Furthermore, the solution is a rational function in $x_{\ell_1}, \dots, x_{\ell_s}$ and in the reaction rate constants $k_{y \rightarrow y'} > 0$, with all coefficients positive (Feliu and Wiuf 2012). The solution can be found using graphical procedures, but in practice, solving the system of linear equations is the preferred approach (see Feliu and Wiuf 2012 for more on this).

By this procedure one obtains a parameterization of the zero set Z of $h^{(0)}$ in s variables $v_i = x_{\ell_i}, i = 1, \dots, s$. Further, clearly $\text{rank } D\Phi(v) = s$. In Feliu et al. (2019) some conditions are stated which guarantee that the assumptions in Proposition 1(a) are satisfied.

3.2.2 Monomial Parameterizations and Deficiency Zero Networks

We next consider another common scenario occurring, for instance, for so-called complex-balanced steady states (see Feinberg 1995; Horn and Jackson 1972) and networks with toric steady states (see Perez Millan et al. 2012; Müller et al. 2016). In this scenario the zero set Z of $h^{(0)}$ in $\mathbb{R}_{>0}^n$ agrees with the solution set of a collection of binomial equations

$$a_\ell(k)x^{u_\ell} - b_\ell(k)x^{c_\ell} = 0, \quad x \in \mathbb{R}_{>0}^n, \ell = 1, \dots, q, \quad (18)$$

where $u_\ell, c_\ell \in \mathbb{R}^n$ and $a_\ell(k), b_\ell(k)$ are polynomials in the parameters of the rate functions that only attain positive values for valid k . Here, for the sake of simplicity, we restrict attention to the case when all $x_i > 0$. Under these assumptions, the solution set Z to (18) equals the solution set of

$$x^{u_\ell - c_\ell} = \frac{b_\ell(k)}{a_\ell(k)}, \quad x \in \mathbb{R}_{>0}^n, \ell = 1, \dots, q. \quad (19)$$

The solution set of (19), if non-empty, admits a monomial parameterization of the following form. Let x^* be any fixed solution of (19) and $M \in \mathbb{R}^{n \times q}$ the matrix whose columns are $u_\ell - c_\ell$. If x is a solution to (19), then

$$x^M = (x^*)^M.$$

It is a classical result (see for example Lemma 3.7 in Müller et al. 2016) that the solution set to this equation, and hence Z , can be parameterized in the form

$$\Phi(v) = x^* \circ v^B = (x_i^* v^{b_i})_{i=1, \dots, n}, \quad v \in \mathbb{R}_{>0}^d. \quad (20)$$

Here $d = \dim \ker M^{\text{tr}}$, and b_1, \dots, b_n are the columns of a matrix $B \in \mathbb{R}^{d \times n}$ with row span equal to $\ker M^{\text{tr}}$ and $\ker B^{\text{tr}} = \{0\}$ (thus $d = \text{rank } B$). With an easy computation one verifies the well-known identity

$$D\Phi(v) = \text{diag}(x^* \circ v^B) B^{\text{tr}} \text{diag}(1/v), \quad (21)$$

where, for a vector α , $\text{diag}(\alpha)$ denotes the diagonal matrix with the entries of α in the diagonal, and $1/v$ is defined component-wise. By this identity, the rank of $D\Phi(v)$ equals d , the rank of B , and therefore we are in the setting of Sect. 2.2 provided that $d = s$. With this in place, by Proposition 3 the matrix $R(v)$ in Theorem 1 becomes

$$\begin{aligned} R(v) &= (L_f \text{diag}(x^* \circ v^B) B^{\text{tr}} \text{diag}(1/v))^{-1} L_f \\ &= \text{diag}(v) (L_f \text{diag}(x^* \circ v^B) B^{\text{tr}})^{-1} L_f. \end{aligned} \quad (22)$$

We turn now to the special case of *complex-balanced steady states* for mass action kinetics. These are steady states such that for each fixed node y of the fast subnetwork, it holds that

$$\sum_{\text{reaction } y' \rightarrow y} w_{y' \rightarrow y}(x)(y - y') = \sum_{\text{reaction } y \rightarrow y'} w_{y \rightarrow y'}(x)(y' - y). \quad (23)$$

As shown in Feinberg (1995) and Horn (1972), a necessary condition for complex-balanced steady states to exist is that each connected component of the fast subnetwork is strongly connected; this is what is known as a weakly reversible reaction network. In this case, if the parameters $k_{y \rightarrow y'}$ satisfy certain algebraic conditions, then there are

positive complex-balanced steady states and any positive steady state is complex balanced. Furthermore, if the deficiency of the fast subnetwork is zero, then any positive steady state is complex balanced, independent of the values of the parameters.

The set Z of positive complex-balanced steady states agrees with the solution set of a collection of binomial equations of the form

$$K_{ij}x^{y_i} - K_{ji}x^{y_j} = 0, \quad x \in \mathbb{R}_{>0}^n,$$

for every pair of nodes y_i, y_j in the same connected component of the reaction network, and with K_{ij} and K_{ji} positive polynomials in the parameters $k_{y \rightarrow y'}$ for the reactions in the same connected component (Craciun et al. 2009). This implies that the column span of the matrix M above agrees with the column span of N_f , and therefore $\ker M^{\text{tr}}$ has rank s . As a suitable matrix B one can choose L_f as in Proposition 3. We thus obtain an explicit expression for the reduced system.

Proposition 4 *Assume that the fast subsystem (16) has a positive complex-balanced steady state x^* . Then:*

- (a) *With the notation of Theorem 1 and Proposition 3, there is a parameterization (20) of the critical manifold with $B = L_f$, parameter space $\mathbb{R}_{>0}^s$ and*

$$R(v) = \text{diag}(v)(L_f \text{diag}(x^* \circ v^{L_f})L_f^{\text{tr}})^{-1}L_f.$$

- (b) *The reduced system is given by:*

$$\begin{aligned} v' &= R(v) \cdot h^{(1)}(\Phi(v)) = R(v) \cdot N_s \cdot (K_s \circ (x^*)^{Y_s} \circ v^{L_f Y_s}) \\ &= \text{diag}(v)(L_f \text{diag}(x^* \circ v^{L_f})L_f^{\text{tr}})^{-1}L_f \cdot N_s \cdot (K_s \circ (x^*)^{Y_s} \circ v^{L_f Y_s}). \end{aligned} \tag{24}$$

Proof Part (a) is clear, while part (b) follows immediately with part (a) and

$$\begin{aligned} h^{(1)}(\Phi(v)) &= h^{(1)}(x^* \circ v^{L_f}) = N_s \cdot (K_s \circ (x^* \circ v^{L_f})^{Y_s}) \\ &= N_s \cdot (K_s \circ (x^*)^{Y_s} \circ (v^{L_f})^{Y_s}) \\ &= N_s \cdot (K_s \circ (x^*)^{Y_s} \circ v^{L_f Y_s}). \end{aligned}$$

□

3.2.3 Attractivity of the Critical Manifold


In the discussion so far we restricted attention to computing a reduced system on the critical manifold and did not address the question whether the attractivity condition from Proposition 1(a) is satisfied. Of course, Remarks 1 and 3 are available, but due to our consideration of slow and fast reaction networks one also may resort to known properties of certain classes of reaction networks. There are general attractivity results available for complex-balanced *mechanisms* as introduced by Horn (1972), which are

of primary interest in our setting, since we only require positivity of reaction constants in our considerations. By Horn (1972), Theorem 4A a mechanism is complex balanced for all choices of reaction rate constants if and only if it is weakly reversible and has deficiency zero. For these systems Feinberg (1995) proved in Remark C.2 that every steady state is linearly attractive (within its stoichiometric compatibility class). We therefore have:

Proposition 5 *If the fast subsystem (16) is weakly reversible and of deficiency zero, then all nonzero eigenvalues of the Jacobian have negative real part, hence all the conditions of Proposition 1(a) hold.*


3.2.4 Examples

Example 5 We consider the following reaction network with mass action kinetics, where the numbers $k_{y \rightarrow y'}$ are written as labels of the reactions:



This network is a two-component system where X_1, X_2 are the unphosphorylated and phosphorylated forms of the histidine kinase and X_3, X_4 are the unphosphorylated and phosphorylated forms of the response regulator. Further we have a dead-end complex between the unphosphorylated forms of both proteins.

We now look at the slow-fast scenario where the fast reactions are those with labels k_1, \dots, k_6 , such that the fast subnetwork is



and the slow reactions have labels k_7, k_8, k_9 . With this choice of slow-fast reactions, we have

$$N_f = \begin{pmatrix} -1 & 1 & 0 & 0 & 1 & -1 \\ 1 & -1 & -1 & 1 & 0 & 0 \\ 0 & 0 & -1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 1 & -1 & -1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \quad N_s = \begin{pmatrix} 0 & -1 & 1 \\ 0 & 0 & 0 \\ 1 & -1 & 1 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & -1 \end{pmatrix}$$

and

$$Y_5 = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad K_5 = \begin{pmatrix} k_7 \\ k_8 \\ k_9 \end{pmatrix}.$$

The fast reaction network in (26) has 6 nodes and three connected components, and the rank of N_f is $r = 3$ (hence also $s = 3$). Therefore, the deficiency is zero and any positive steady state, that is, any element of Z , is complex balanced and a solution to a set of binomial equations. Under mass action, the steady states of this fast subnetwork are the solutions to

$$\begin{aligned} -k_3x_2x_3 + k_1x_1 - k_2x_2 + k_4x_5 &= 0, \\ -k_6x_1x_4 + k_5x_5 &= 0, \\ k_3x_2x_3 + k_6x_4x_1 - k_4x_5 - k_5x_5 &= 0, \end{aligned} \tag{27}$$

and we easily verify that

$$x^* = \left(1, \frac{k_1}{k_2}, \frac{k_2k_4k_6}{k_1k_3k_5}, 1, \frac{k_6}{k_5}, 1\right)^{\text{tr}}$$

is a positive steady state of the fast subnetwork. With Proposition 4(b) the reduced system can be computed. We choose

$$L_f = \begin{pmatrix} 1 & 1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix} \tag{28}$$

and obtain the following parameterization of Z :

$$\Phi(v) = x^* \circ \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix}^{L_f} = x^* \circ \begin{pmatrix} v_1 \\ v_1 \\ v_2 \\ v_2 \\ v_1v_2 \\ v_3 \end{pmatrix} = \begin{pmatrix} v_1 \\ \frac{k_1}{k_2}v_1 \\ v_2 \\ \frac{k_2k_4k_6}{k_1k_3k_5}v_2 \\ v_2 \\ \frac{k_6}{k_5}v_1v_2 \\ v_3 \end{pmatrix}.$$

Using Eq. (24), the reduced system is found:

$$v' = \frac{1}{\xi} \cdot \begin{pmatrix} \frac{1}{k_1k_3}k_2(k_1k_3k_5 + k_2k_4k_6)(-k_2k_4k_6k_8v_1v_2 + k_1k_3k_5k_9v_3) \\ k_5(k_1 + k_2)(-k_2k_4k_6k_8v_1v_2 + k_1k_3k_5k_9v_3) \\ \frac{\xi}{k_1k_3k_5} \cdot (k_2k_4k_6k_8v_1v_2 - k_1k_3k_5k_9v_3) \end{pmatrix},$$

where ξ is given by

$$\xi = k_1(k_1k_3k_5 + k_2k_3k_5)k_6v_1 + k_2(k_1k_3k_5 + k_2k_4k_6)k_6v_2 + k_5(k_1 + k_2)(k_1k_3k_5 + k_2k_4k_6).$$

In addition, we conclude by Proposition 5 that all nonzero eigenvalues of $Dh^{(0)}$ have negative real part on Z .

Observe that the parameterization $\Phi(v)$ is not unique. For example, choosing another starting steady state

$$x^* = \left(1, \frac{k_1}{k_2}, 1, \frac{k_1k_3k_5}{k_2k_4k_6}, \frac{k_1k_3}{k_2k_4}, 1\right)^{\text{tr}},$$

we obtain the parameterization

$$\Phi(v) = \left(v_1, \frac{k_1v_1}{k_2}, v_2, \frac{k_1k_3k_5v_2}{k_2k_4k_6}, \frac{k_1k_3v_1v_2}{k_2k_4}, v_3\right)^{\text{tr}},$$

and the reduced system

$$\begin{aligned} v_1' &= -\frac{k_2k_4}{q(v)}(k_1k_3k_5 + k_2k_4k_6)(k_8v_1v_2 - k_9v_3) \\ v_2' &= -\frac{k_2k_4}{q(v)}k_4k_6(k_1 + k_2)(k_8v_1v_2 - k_9v_3) \\ v_3' &= k_8v_1v_2 - k_9v_3, \end{aligned}$$


with

$$\begin{aligned} q(v) &= k_1k_3(k_1k_3k_5 + k_2k_4k_6)v_2 + k_1k_3k_4k_6(k_1 + k_2)v_1 \\ &\quad + k_4(k_1 + k_2)(k_1k_3k_5 + k_2k_4k_6). \end{aligned}$$

The same parameterization is obtained by eliminating x_2, x_4, x_5 after realizing that the set of species $\{X_2, X_4, X_5\}$ satisfies (i), (ii) and (iii) in Sect. 3.2.1.

Example 6 If we remove the reactions with label k_8, k_9 from the network (25), then the stoichiometric matrices of both the fast subnetwork and the full network have rank 3. Hence, if the reduction with a parameterized critical manifold is possible, the reduced system is $v' = 0$.

Example 7 We analyze the reaction network



This system can be interpreted as a dual phosphorylation cycle with X_1 the kinase catalyzing the phosphorylation of a substrate S with two phosphorylation sites. Then

X_2, X_5, X_6 correspond to the phosphoforms with no, one, two phosphate groups, respectively, and X_3 and X_4 are intermediate enzyme-substrate forms. Dephosphorylation proceeds without a phosphatase. We let the fast system be all reactions involved in the conversion $X_2 \leftrightarrow X_5$, namely those with label k_1, \dots, k_6 . Hence the reactions with label k_7, k_8 are slow. With this choice, we have


$$N_f = \begin{pmatrix} -1 & 1 & 0 & 1 & -1 & 0 \\ -1 & 1 & 0 & 0 & 0 & 1 \\ 1 & -1 & -1 & 0 & 0 & 0 \\ 0 & 0 & 1 & -1 & 1 & 0 \\ 0 & 0 & 0 & 1 & -1 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \quad L_f = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 1 \\ -1 & 1 & 0 & 0 & 1 & 0 \\ 1 & 0 & 1 & 1 & 0 & 0 \end{pmatrix},$$

and

$$h^{(0)}(x) = \left(-k_1x_1x_2 - k_5x_1x_5 + k_2x_3 + k_4x_4, -k_1x_1x_2 + k_2x_3 + k_6x_5, \right. \\ \left. k_1x_1x_2 - k_2x_3 - k_3x_3, k_5x_1x_5 + k_3x_3 - k_4x_4, \right. \\ \left. -k_5x_1x_5 + k_4x_4 - k_6x_5, 0 \right)^{tr}$$

$$h^{(1)}(x) = \left(0, 0, 0, 0, -k_7x_1x_5 + k_8x_6, k_7x_1x_5 - k_8x_6 \right)^{tr}.$$

The fast network has deficiency 1, since the rank of N_f is 3, and the network has 6 nodes and two connected components. Thus, the steady states are not complex balanced for all k . Instead, we observe that the set $\{X_3, X_4, X_5\}$ satisfies (i)–(iii) in Sect. 3.2.1. Indeed, (i) and (ii) are easy to check. For (iii), the induced network obtained after setting the species not in this set to zero is



and clearly there is directed path to 0 from every species. This implies that x_3, x_4, x_5 can be solved from the system $h^{(0)}(x)_{3,4,5} = 0$ to obtain the following parameterization (where $v_1 = x_1, v_2 = x_2, v_3 = x_6$)

$$\Phi: (v_1, v_2, v_3) \mapsto \begin{pmatrix} v_1 \\ v_2 \\ \frac{k_1}{k_2+k_3} v_1 v_2 \\ \frac{k_1 k_3}{k_4 k_6 (k_2+k_3)} (k_5 v_1 + k_6) v_1 v_2 \\ \frac{k_1 k_3}{k_6 (k_2+k_3)} v_1 v_2 \\ v_3 \end{pmatrix},$$

which has an entry that is not monomial. Next we compute the matrix $R(v)$ and the reduced system using Proposition 3. We have

$$D\Phi(v) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ \frac{k_1 v_2}{k_2+k_3} & \frac{k_1 v_1}{k_2+k_3} & 0 \\ \frac{k_1 k_3 (2k_5 v_1 + k_6) v_2}{k_4 k_6 (k_2+k_3)} & \frac{(k_5 v_1 + k_6) k_1 k_3 v_1}{k_4 k_6 (k_2+k_3)} & 0 \\ \frac{k_1 k_3 v_2}{k_6 (k_2+k_3)} & \frac{k_1 k_3 v_1}{k_6 (k_2+k_3)} & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

and using $R(v_1, v_2, v_3) = (L_f D\Phi(v))^{-1} L_f$ the reduced system is given by

$$\begin{pmatrix} v_1' \\ v_2' \\ v_3' \end{pmatrix} = \begin{pmatrix} \xi_1 \\ \xi_2 \\ \frac{\xi_2}{k_6(k_3+k_2)} \end{pmatrix} \begin{pmatrix} k_1(k_3 k_5 v_1 + k_3 k_6 + k_4 k_6) v_1 \\ -(2k_1 k_3 k_5 v_1 v_2 + k_1 k_3 k_6 v_2 + k_1 k_4 k_6 v_2 + k_2 k_4 k_6 + k_3 k_4 k_6) \\ \xi_2 \end{pmatrix}$$

where

$$\begin{aligned} \xi_1 &= k_1 k_3 k_7 v_1^2 v_2 - (k_2 + k_3) k_6 k_8 v_3 \\ \xi_2 &= k_1^2 k_3^2 k_5 v_1^2 v_2 + (k_2 + k_3) \left(k_1 k_3 k_5 k_6 v_1^2 + 2k_1 k_3 k_5 k_6 v_1 v_2 \right. \\ &\quad \left. + k_1 k_6 (k_3 k_4 + k_3 k_6 + k_4 k_6) v_1 + k_1 (k_3 + k_4) k_6^2 v_2 + k_4 (k_2 + k_3) k_6^2 \right). \end{aligned}$$

We next verify that the eigenvalue condition in Proposition 1(a) is satisfied. To this end we check the eigenvalues of the matrix $A(x) = D\mu(x) \cdot P(x)$ with

$$P = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -1 & 0 & -1 \\ 0 & 0 & 1 \\ 1 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \mu(x) = \begin{pmatrix} -k_1 x_1 x_2 - k_5 x_1 x_5 + k_2 x_3 + k_4 x_4 \\ -k_1 x_1 x_2 + k_2 x_3 + k_6 x_5 \\ k_5 x_1 x_5 + k_3 x_3 - k_4 x_4 \end{pmatrix}.$$

Using the Routh–Hurwitz conditions (see Gantmacher 2005, Ch. V, §6) for its characteristic polynomial $\chi_A(\lambda) = \lambda^3 + \sigma_1 \lambda^2 + \sigma_2 \lambda + \sigma_3$ of degree 3, we obtain:

$$\begin{aligned} \sigma_1 &= k_1 x_1 + k_1 x_2 + k_5 x_1 + k_5 x_5 + k_2 + k_3 + k_4 + k_6 \\ \sigma_2 &= k_1 k_5 x_1 (x_1 + x_2 + x_5) + (k_1 k_3 + k_1 k_4 + k_1 k_6 + k_2 k_5 + k_3 k_5) x_1 \\ &\quad + k_1 (k_3 + k_4 + k_6) x_2 + k_5 (k_2 + k_3 + k_6) x_5 + (k_2 + k_3) (k_4 + k_6) + k_4 k_6 \\ \sigma_3 &= k_1 k_3 k_5 x_1 (x_1 + x_2 + x_5) + k_1 (k_3 k_4 + k_3 k_6 + k_4 k_6) x_1 \\ &\quad + k_1 k_6 (k_3 + k_4) x_2 + k_6 (k_2 + k_3) (k_5 x_5 + k_4). \end{aligned}$$

In order to verify that all eigenvalues have negative real part, we use the Hurwitz conditions for polynomials of degree three:

$$\sigma_1 > 0, \quad \sigma_3 > 0, \quad \sigma_1 \cdot \sigma_2 - \sigma_3 > 0.$$

The three expressions on the left side of each inequality are polynomials in the parameters and x with all coefficients positive, and hence are positive when evaluated at positive values of k and x .

Acknowledgements Open Access funding provided by Projekt DEAL. The work of NK and SW has been supported by the bilateral Project ANR-17-CE40-0036 and DFG-391322026 SYMBIONT. EF has been supported by the Independent Research Fund of Denmark. SW thanks the Department of Mathematical Sciences of the University of Copenhagen for its hospitality during a research visit when essential parts of the present manuscript were devised. Likewise, EF thanks the hospitality of RWTH Aachen where this project was initiated.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Appendix

Lemma 1 *Let $0 < s < n$ and $A \in \mathbb{R}^{n \times s}$, $B \in \mathbb{R}^{s \times n}$ such that $\text{rank } AB = s$ and $(AB)^2 = AB$. Then $BA = I_s$.*

Proof Recall that \mathbb{R}^n is the direct sum of the eigenspaces of AB for eigenvalues 1 and 0. Let z_1, \dots, z_s be a basis of the former. Then Bz_1, \dots, Bz_s are linearly independent due to

$$\sum \lambda_i Bz_i = 0 \Rightarrow 0 = \sum \lambda_i ABz_i = \sum \lambda_i z_i,$$

hence they form a basis of \mathbb{R}^s . Finally, $ABz_i = z_i$ implies $(BA)Bz_i = Bz_i$ for $1 \leq i \leq s$. \square

References

- Arnold, V.I., Anosov, D.V. (eds.): Dynamical Systems I. Springer, Berlin (1988)
- Craciun, G., Dickenstein, A., Shiu, A., Sturmfels, B.: Toric dynamical systems. *J. Symb. Comput.* **44**(11), 1551–1565 (2009)
- Feinberg, M.: Complex balancing in general kinetic systems. *Arch. Ration. Mech. Anal.* **49**, 187–194 (1972)
- Feinberg, M., Horn, F.J.M.: Chemical mechanism structure and the coincidence of the stoichiometric and kinetic subspaces. *Arch. Ration. Mech. Anal.* **66**(1), 83–97 (1977)
- Feinberg, M.: The existence and uniqueness of steady states for a class of chemical reaction networks. *Arch. Ration. Mech. Anal.* **132**, 311–370 (1995)

- Feinberg, M.: Foundations of Chemical Reaction Network Theory. Springer, Cham (2019)
- Feliu, E., Wiuf, C.: Variable elimination in chemical reaction networks with mass action kinetics. *SIAM J. Appl. Math.* **72**, 959–981 (2012)
- Feliu, E., Lax, C., Walcher, S., Wiuf, C.: Non-interacting species and reduction of reaction networks. Preprint, [arXiv:1908.11270](https://arxiv.org/abs/1908.11270) (2019)
- Fenichel, N.: Geometric singular perturbation theory for ordinary differential equations. *J. Differ. Equ.* **31**(1), 53–98 (1979)
- Gantmacher, F.R.: Applications of the Theory of Matrices. Dover, Mineola (2005)
- Goeke, A., Walcher, S.: A constructive approach to quasi-steady state reduction. *J. Math. Chem.* **52**, 2596–2626 (2014)
- Goeke, A., Walcher, S., Zerz, E.: Classical quasi-steady state reduction—a mathematical characterization. *Physica D* **345**, 11–26 (2017)
- Heinrich, R., Schauer, M.: Quasi-steady-state approximation in the mathematical modeling of biochemical networks. *Math. Biosci.* **65**, 155–170 (1983)
- Horn, F.: Necessary and sufficient conditions for complex balancing in chemical kinetics. *Arch. Ration. Mech. Anal.* **49**, 172–186 (1972)
- Horn, F., Jackson, R.: General mass action kinetics. *Arch. Ration. Mech. Anal.* **47**, 81–116 (1972)
- Lax, C., Walcher, S.: Singular perturbations and scaling. *Discrete Contin. Dyn. Syst. Ser. B.* **25**, 1–29 (2020)
- Lee, C.H., Othmer, H.G.: A multi-time-scale analysis of chemical reaction networks: I. Deterministic systems. *J. Math. Biol.* **60**, 387–450 (2009)
- Müller, S., Feliu, E., Regensburger, G., Conradi, C., Shiu, A., Dickenstein, A.: Sign conditions for injectivity of generalized polynomial maps with applications to chemical reaction networks and real algebraic geometry. *Found. Comput. Math.* **16**(1), 69–97 (2016)
- Noethen, L., Walcher, S.: Tikhonov’s theorem and quasi-steady state. *Discrete Contin. Dyn. Syst. Ser. B* **16**(3), 945–961 (2011)
- Perez Millan, M., Dickenstein, A., Shiu, A., Conradi, C.: Chemical reaction systems with toric steady states. *Bull. Math. Biol.* **74**(5), 1027–1065 (2012)
- Stiefenhofer, M.: Quasi-steady-state approximation for chemical reaction networks. *J. Math. Biol.* **36**, 593–609 (1998)
- Tikhonov, A.N.: Systems of differential equations containing a small parameter multiplying the derivative. *Math. Sb.* **31**, 575–586 (1952). (in Russian)
- Verhulst, F.: Methods and Applications of Singular Perturbations. Boundary Layers and Multiple Timescale Dynamics. Springer, New York (2005)
- Walcher, S.: On differential equations in normal form. *Math. Ann.* **291**(2), 293–314 (1991)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.