

1 LRH: Groundwater food webs

D. C. Weitowitz et al.

2 RRH: Volume 38

September 2019

3

4 **Obligate groundwater crustaceans mediate biofilm interactions in a subsurface food**
5 **web**

6

7 **Damiano C. Weitowitz^{1,3}, Anne L. Robertson^{1,4}, John P. Bloomfield^{2,5}, Louise Maurice^{2,6}**
8 **and Julia Reiss^{1,7}**

9

10

11 ¹Department of Life Sciences, University of Roehampton, Holybourne Avenue, London
12 SW15 4JD, United Kingdom

13

14 ²British Geological Survey, Maclean Building, Crowmarsh Gifford, Wallingford,
15 Oxfordshire, OX10 8BB, United Kingdom

16

17 E-mail addresses: ³weitowid@roehampton.ac.uk; ⁴a.robertson@roehampton.ac.uk;

18 ⁵jpb@bgs.ac.uk; ⁶loma@bgs.ac.uk; ⁷julia.reiss@roehampton.ac.uk

19

20

21 Received 10 January 2018; Accepted 23 October 2018; Published online XX Month 2019.

22

23

24

25

26

27 **Abstract:** Food webs in groundwater ecosystems are dominated by only a few top-level
28 consumers, mainly crustaceans. These obligate groundwater dwellers—or stygobites—clearly
29 interact with groundwater biofilm, but it is uncertain whether they affect the abundance and
30 structure of biofilm assemblages. We hypothesized that crustacean stygobites would reduce
31 bacteria and protozoan abundance and alter biofilm assemblage structure. We also
32 hypothesized that high densities of stygobites would remove more bacteria and protozoa than
33 would low densities, and that this difference would become more pronounced over time.
34 First, we established that the amphipods *Niphargus fontanus* and *Niphargus kochianus* both
35 ingest biofilm by examining their gut contents. We then conducted two microcosm
36 experiments. The first experiment showed that both *N. fontanus* and the isopod *Proasellus*
37 *cavaticus* increased protozoan abundance but that bacterial abundance was only slightly
38 reduced in the presence of *P. cavaticus*. In the second experiment, we determined how zero,
39 low, and high densities of *N. kochianus* affected the biofilm. The high-density treatment of *N.*
40 *kochianus* had significantly higher protozoan abundance than the control and the low-density
41 treatment, and high densities of *N. kochianus* significantly increased the relative proportions
42 of small and medium-sized bacteria over time compared with controls. Our controlled
43 microcosm experiments demonstrate that macroinvertebrate stygobites can influence
44 groundwater biofilm assemblages, although the exact mechanisms are not clear. These results
45 support the hypothesis that stygobites influence essential ecosystem services supplied by
46 groundwater ecosystems.

47

48 **Keywords:** protozoa, microcosms, bacteria, biofilm, flow cytometer, stygobite, *Niphargus*,
49 *Proasellus*.

50

51 Groundwater is a critical resource for the ~2 billion people worldwide who depend on it for
52 drinking water (Morris et al. 2003). Moreover, many terrestrial and aquatic ecosystems rely
53 wholly or partially on access to groundwater (Boulton 2005). Biotic communities within
54 groundwater contribute to the maintenance of groundwater quality via the breakdown of
55 organic matter, nutrients, and contaminants (e.g. Kota et al. 1999, Gibert and Deharveng
56 2002, Tomlinson & Boulton 2008) providing vital ecosystem services (Griebler and Avramov
57 2015). Many of the resident animals (called stygobites) are groundwater obligates (Gibert et
58 al. 1994), and they uniquely contribute to global biodiversity. Stygobite species often have
59 restricted distributions (Gibert et al. 2009), which make them especially vulnerable to
60 anthropogenic pressures such as pollution (Boulton et al. 2003).

61 Food webs in groundwater ecosystems are also unique in that they are truncated and
62 far less complex than their surface water counterparts. Their simplicity is associated with the
63 negligible primary production in most groundwater ecosystems, which are largely dependent
64 on scarce allochthonous energy sources to fuel community biomass and production (Gibert et
65 al. 1994, Gibert and Deharveng 2002). Organic matter is the basal component of these food
66 webs; prokaryotes, single-celled eukaryotes (protozoans); and microscopic metazoans are
67 primary consumers; and macroinvertebrates (principally crustaceans) or cavefish are top-
68 level consumers. In comparison with their surface water counterparts, stygobites have a
69 reduced metabolism and low growth and reproduction rates – adaptations to the limited
70 energy and constant temperature in the groundwater environment (Spicer 1998). Other
71 stygobite adaptations include lack of eyes and pigmentation and resistance to hypoxia and
72 starvation (Hervant et al. 1995, Hervant et al. 1999).

73 Groundwater food web interactions, especially those between micro- and
74 macroorganisms, are poorly understood (Griebler and Avramov 2015, but see Boulton et al.
75 2008). Few experimental studies with appropriate replication have been conducted.

76 Conflicting evidence exists for whether or not stygobitic crustaceans cause top-down control
77 in groundwater food webs. Cooney and Simon (2009) found that *Gammarus minus*, a cave
78 amphipod, reduces bacterial activity, whereas other studies demonstrated that bacteria are
79 more abundant and active when grazed by *G. minus* or *Caecidotea tridentata*, a subterranean
80 isopod (Edler and Dodds 1996, Kinsey et al. 2007). Other studies have found no consumptive
81 effects of stygobites (Foulquier et al. 2010, 2011). Researchers mainly attributed this lack of
82 effect to low metabolic rates and low abundances of top-level consumers in energy-limited
83 environments (Foulquier et al. 2010, 2011). It seems likely that grazer density and feeding
84 time are important predictors to consider when investigating the effects of stygobites on
85 groundwater assemblages. Similarly, there is contradictory evidence for bottom-up control of
86 groundwater food webs. Foulquier et al. (2010, 2011) found that bacterial assemblages were
87 more abundant and active at higher levels of dissolved organic carbon (DOC). However,
88 Weitowitz (2017) found that higher nutrient concentrations did not result in higher bacterial
89 abundances.

90 Trophic relationships in surface water ecosystems have received considerable
91 attention in recent decades (e.g. Sih et al. 1985, Billen and Servais 1990, Muylaert et al.
92 2002, Shurin et al. 2012). These studies clearly show that both bottom-up and top-down
93 forces are important in structuring biological communities (McQueen et al. 1989, Menge
94 2000). Macrofaunal isopod and amphipod crustaceans such as *Gammarus* spp. and *Asellus*
95 spp. are known to play a critical role in surface waters both as food for higher trophic levels
96 and as decomposers of organic material (Graca et al. 1994a, 1994b). These taxa can also
97 affect biofilm groups such as small metazoans (Rosemond et al. 2001) and algae (Duffy and
98 Hay 2000, Bruno et al. 2008), but they are not known to purposefully predate on protozoans.
99 However, surface water protozoans can strongly influence bacterial populations in both
100 positive and negative ways (e.g. Wey et al. 2012, Huws et al. 2005, Humphreys 2009). Given

101 the importance and strength of consumer-mediated interactions in surface waters it is likely
102 that such interactions also occur in groundwater ecosystems.

103 In addition to feeding interactions, aquatic invertebrates can have indirect effects on
104 the microbial food web and ecosystem functioning. For example, macrofauna are known to
105 both bioturbate sediments and compact fine sediments into fecal pellets (Boulton et al.,
106 2008). Furthermore, interstitial bacterial activity can be stimulated by invertebrate
107 bioturbation in sediments (Mermillod-Blondin et al. 2000), and microbial activity can be
108 enhanced through nutrients provided by hyporheic invertebrates in the form of fecal pellets
109 (Boulton 2000, Marshall and Hall 2004).

110 Macrofaunal invertebrate stygobites are the top consumers in many groundwater
111 ecosystems. However, amphipods and isopods move and appear to acquire food differently.
112 The amphipods *N. fontanus* (Bate 1859) and *N. kochianus* (Schellenberg 1932) preferentially
113 use their gnathopods to pick up, manipulate and ingest pieces of sediment. The isopod *P.*
114 *cavaticus* (Leydig 1871), however, is a bottom crawler, directly grazing on sediment surfaces
115 (personal observation). Previous authors showed that sedimentary biofilm provides up to 83%
116 of the diet for *P. cavaticus* (e.g. Francois et al. 2016). However, the evidence is less clear for
117 *Niphargus* spp., which have been described as being both polyphagous (Fiser et al. 2008,
118 Arnscheidt et al. 2012) and predatory (Knight and Johns 2015).

119

120 In this study, we tested two hypotheses: (1) The presence of stygobites will
121 significantly reduce bacterial and protozoan abundances and alter biofilm assemblage
122 structure. *Proasellus cavaticus* will exert a stronger effect than *N. fontanus* because of its
123 scraping ‘lawn mower’ feeding strategy, which has also been observed in some surface
124 isopods (Naylor 1955, Jones 1972). (2) High stygobite densities will remove more bacteria

125 and protozoans than low densities, and this effect will become more pronounced over time as
126 fewer and fewer reproductive bacteria and protozoa remain in the system.

127

128 **METHODS**

129 To test our hypotheses, we first quantified the diets of the 3 target species. We then
130 conducted 2 manipulative experiments.

131 **Study species**

132 All 3 target species are 8 – 11 mm long and commonly occur in the UK. *Niphargus*
133 *kochianus* (Fig. 1A) is the most abundant and widespread amphipod species in UK chalk
134 aquifers (Maurice et al. 2016). The isopod, *P. cavaticus* (Fig. 1B), occurs mainly in carbonate
135 aquifers (Johns et al. 2015). *N. fontanus* (Fig. 1C) is found in a wide range of groundwater
136 habitats in the UK (Johns et al. 2015).

137

138 **Gut content study**

139 We conducted a preliminary study to confirm that the *Niphargus* species used in our study
140 feed on and ingest sedimentary biofilm. We collected 45 individuals of *N. kochianus* and 2 *N.*
141 *fontanus* from a chalk borehole (Berkshire, UK) and then starved the animals in ultrapure
142 water for 14 days to promote gut clearing. We then incubated individuals (one per
143 microcosm) with a biofilm-coated stone tile (Fiji, B&Q, dimension - 3.1 x 1.4 x 0.8 cm) at 11
144 °C in the dark for 96 h. These tiles were previously exposed to groundwater for 4 weeks to
145 allow the natural colonization of biofilm. Tiles were placed in the same chalk borehole used
146 to source the stygobite amphipods. Individuals were then stored in > 98 % ethanol. Those
147 that had expelled their guts on preservation were discarded. We followed the approach of
148 Navarro-Barranco et al. (2013) to better observe gut contents. Specimens were placed in
149 vials of Hertwig's liquid (270 g of chloral hydrate, 19 mL of 1N chloric acid, 60 mL of

150 glycerine, and 150 mL of distilled water) in an oven at 65 °C for 4 hours. Individuals were
151 then mounted on a slide and the contents of the foregut (we were only interested in food
152 intake over the last 96 hours) studied under an Olympus BX53 microscope and photographed
153 at x400 magnification.

154

155 **Experiment 1. Testing the hypothesis that stygobite presence will reduce bacterial and**
156 **protozoan abundances and alter biofilm assemblage structure.**

157 *Experimental setup and design* Nine *N. fontanus* and 9 *Proasellus cavaticus* were
158 collected over 2 days in November 2013 from a cave system in Wales (Elm Hole; latitude
159 51.81, longitude -3.14) and kept in the dark in containers of cave water at 11 °C.

160 We exposed stone tiles in a borehole (chalk, Berkshire, UK) to obtain natural
161 groundwater biofilms. Stone tiles of equal size (Fiji, B&Q, dimension - 3.1 x 1.4 x 0.8 cm)
162 were autoclaved and washed in ultrapure water, placed in mesh nets with a mesh diameter of
163 500 µm, and suspended in the borehole for 3 weeks to colonize. Griebler et al. (2002) showed
164 that numbers of attached bacteria on sediment in similarly clean groundwater near Salzburg,
165 Austria reached $500 * 10^5$ cells per cm³ within 4 weeks of exposure. On retrieval, tiles were
166 transported to the laboratory in a cool box and stored in unfiltered groundwater in the dark at
167 11 °C (the same temperature as water in the borehole) for four weeks until the start of the
168 experiment, which allowed for further growth of the biofilm.

169 For this experiment, we used 3 treatments (consumer *N. fontanus*, consumer *P.*
170 *cavaticus*, and a control) each with 27 replicates ($3*27=81$ microcosms). We used a block
171 design running 6 replicates on days 0 to 4 (Run 1), another 6 replicates on days 8 to 12 (Run
172 2), and another 6 on days 16 to 20 (Run 3). For the last block, we ran nine replicates on day
173 24 (Run 4) (see Table S1). We employed this design because we had to ‘re-use’ individuals
174 to obtain a high replication. This temporal block design enabled us to statistically account for

175 any differences in starting conditions such as the condition of the biofilm tiles (Bailey and
176 Reiss 2014). One individual represented one replicate in each of the 4 runs (e.g. *N. fontanus*
177 individuals 1 to 6 and *P. cavaticus* individuals 1 to 6 were used for day 0-4 (see Table S1).
178 All individuals were used 3 different times—twice in the 4-day trials (runs 1-3) and once in
179 run 4).

180 Prior to each experimental run, the crustaceans were starved in filtered groundwater
181 for 4 days to allow them to empty most of their intestines. Only animals with empty foreguts
182 were used in the experiments. Microcosms were set up in 50 mL glass beakers containing 20
183 mL of filtered and autoclaved borehole water and were kept at 11 °C in darkness to mirror
184 groundwater conditions. One tile was placed in each microcosm to provide a food source for
185 the stygobites, and 1 individual of each species was introduced into the respective treatments.
186 Stygobites were checked for mortality every 24 h (two died during the experiment and were
187 replaced with an individual of equal size on discovery).

188 Each run was terminated after 96 h. We then retrieved crustaceans from the
189 microcosms, measured the abundance of bacteria and protozoa on the tiles, and assessed the
190 structure of each biofilm community.

191

192 ***Response variables*** We used a toothbrush to brush the biofilm on each tile into 10 mL of
193 0.25 µm filtered, autoclaved water, a widely used method to detach biofilm from various
194 substrates (see Wipfli et al. 1998, Cardinale et al. 2002, Bouletreau et al. 2006, Vercraene-
195 Eairmal et al. 2010). We used 10 standardized downstrokes on each side of the tiles. We then
196 homogenized the samples with a magnetic stirrer before further processing.

197 To assess the protozoa, we fixed two 500-µl subsamples of the homogenate for
198 microscopic analysis in 2% glutaraldehyde. We used a gridded Sedgwick Rafter cell to count
199 and measure protozoa in each sample under an Olympus CX 21 microscope at x400

200 magnification. We followed Adl et al. (2006) to assign all protozoan cells to 10 morphotype
201 categories, including different types of ciliates, flagellates, and testate amoebae. We used
202 Foissner and Berger (1996) to aid in protozoan identification and morphotype assignments.

203 For the bacterial analysis, we poured a 1 mL subsample of the initial homogenate
204 through a 40- μ m filter. We used a C6 flow cytometer (BD Technologies, North Carolina) to
205 analyze 495- μ l of this filtrate. Preliminary trials in which samples were both sonicated and
206 homogenized resulted in significantly higher counts of non-bacterial debris but did not
207 significantly increase bacterial counts (Weitowitz 2017). We therefore chose not to use
208 sonification to further separate clumps of bacterial cells. Preliminary trials (Weitowitz 2017)
209 also helped us determine the best possible threshold level to identify bacteria and exclude
210 noise. The primary threshold was set at SSC-H (side scatter) 4000 and a secondary threshold
211 at FSC-H (forward scatter) 8000. A dual threshold applies more stringent conditions before
212 counting a particle and excludes more potential noise (BD Biosciences, 2011, p. 5).

213 We used SYTO-9 (Molecular Probes, Life Technologies; Massachusetts) to stain
214 bacteria and distinguish them from soil particles (Lebaron et al. 1998, Gasol and Del Giorgio
215 2000). After preliminary staining trials (Weitowitz 2017), we selected a final SYTO-9
216 concentration of 5 μ M (see also Lebaron et al. 1998). We mixed 495 μ l of microcosm
217 homogenate with 5 μ l of SYTO-9 stock solution resulting in a total volume of 500 μ l for flow
218 cytometric analysis. After adding stain, we incubated the samples in the dark at room
219 temperature for 15 minutes to allow the stain to bind to the DNA.

220 Before counting bacteria, we gated out noise caused by the applied electrical voltage
221 and the running of filtered water using FSC-H vs FL-1 (green fluorescence) dot plots
222 (Troussellier et al. 1999). We kept these bacterial gates constant throughout the experiment.
223 Different bacterial size groups were identified according to their clustering along the FL-1
224 fluorescence axis, allowing for a discrimination of different bacterial populations (see

225 Troussellier et al. 1999). We then ran each 500 μ l sample for 1 minute at slow flow to
226 minimize doublet counts.

227

228 **Experiment 2. Testing the hypothesis that high densities will remove more bacteria and**
229 **protozoans than low densities and that this effect will become more pronounced over**
230 **time.**

231 *Experimental setup and design* For the second experiment, we collected 250 individuals
232 of *N. kochianus* from two boreholes in the Berkshire Chalk aquifer. Collected animals were
233 transported to the laboratory in a cool box filled with groundwater that was maintained at 11
234 °C. In one of the boreholes we suspended 2 tile sizes (Fiji, B&Q, large = 3.1 x 1.4 x 0.8 cm,
235 small = 1.5 x 1.5 x 1 cm) in mesh bags to allow groundwater biofilm to colonize over a
236 period of 5 wk. Next the tiles were stored for 4 wk in the dark at 11 °C until the start of the
237 experiment. This storage period allowed additional growth of the biofilm.

238 This experiment featured 3 treatments: ungrazed biofilm as a control, ‘low *Niphargus*
239 density’ and ‘high *Niphargus* density’. We used nine *N. kochianus* for the low-density
240 treatment and 18 individuals for the high-density treatment. The densities were based on
241 invertebrate sampling (standardized net hauls) conducted in the same chalk aquifer
242 (Weitowitz 2017). Each treatment had 10 replicates, resulting in 30 microcosms (Table S2).

243 To create the microcosms, we filled 250-mL glass beakers with 100 mL of filtered
244 and autoclaved groundwater. We placed 2 large rectangular tiles for bacterial analysis and six
245 small tiles for protozoan analysis in PARAFILM-sealed microcosms. A single control tile in
246 a mesh bag (mesh size 0.1 mm²) was suspended in all microcosms of the treatments and
247 control, which the crustaceans were not able to access. This tile was used to assess biofilm
248 dynamics in the absence of grazing. We then added the stygobites. Over the course of 32
249 days, we sampled protozoans from 5 random replicates of each treatment on six occasions

250 (days 2, 5, 11, 16, 23, 32 for a total of 90 samples). Bacteria were sampled in all replicates on
251 9 occasions (days 2, 3, 5, 9, 11, 16, 18, 23, 27, 32 for a total of 270 samples).

252

253 **Response variables** We obtained samples for protozoan analysis by sacrificing one small
254 tile on each sampling occasion. We carefully brushed the biofilm on each protozoan tile into
255 10 mL of autoclaved water by applying 10 standardized downstrokes with a toothbrush. We
256 then fixed samples with glutaraldehyde and counted protozoa under a microscope as in
257 experiment 1.

258 We sampled bacteria from two large tiles each marked by a grid of 15 evenly sized
259 (0.6 x 0.4 cm) sections (Weitowitz 2017). On each sampling occasion, we pooled three 200-
260 μ l samples directly pipetted from randomly selected sections in each microcosm, and we
261 ensured that no section was sampled more than once. After pipetting, clear patches became
262 visible on the tiles suggesting that biofilm was present and was sampled effectively. The
263 bacterial samples were then thoroughly homogenized in Eppendorff tubes, before being
264 processed in the flow cytometer as in experiment 1. We assigned each counted bacterium to
265 one of 3 body size categories: small, medium and large.

266

267 **Statistical analyses**

268 We performed all statistical analyses in the open source statistical environment R (R
269 Development Core Team 2013). Initially all response variables were checked for normality
270 and homogeneity of variance with the Shapiro-Wilk normality and Levene variance tests. If a
271 response variable violated parametric assumptions, we used the Box-Cox transformation
272 method of package 'MASS' (Venables and Ripley 2002) to identify the best form of power-
273 transformation for the dependent variable.

274 For the first experiment, we assessed if differences in protozoan and bacterial
275 abundance occurred between the 3 treatments (see Table 1). Because we reused individuals in
276 this experiment and because the experiment was run in blocks (see Table S1), we analyzed
277 the data with linear mixed effects models (LMMs) in the R package ‘lme4’ (Bates et al.,
278 2015). LMMs are commonly used to analyze ecological data when multiple measurements
279 (e.g. on a single individual) constitute pseudoreplicates (Perkins et al. 2012, Zuur et al.2009).
280 Because we ‘reused’ individual stygobites three times, we fitted individual ID as a random
281 effect to account for differences in which particular individuals affect the biofilm. We also
282 fitted Block as a predictor in the models, because some of the replicates were run on different
283 days. The R-code for the LMM was: `lmer(log10(Response) ~ Consumer + Block + (1|`
284 `Individual)`. We used a Tukey post-hoc test to find out which treatments were different from
285 each other.

286 In the second experiment we also used LMMs in the R package ‘lme4’ (because this
287 approach is identical to repeated measures ANOVA) to test for the effect of different
288 stygobite densities, time, and their interaction on bacterial and protozoan response variables.
289 The R-code was: `lmer(Response ~ Treatment*Day + (1 | Unique ID)`, where Treatment is
290 high, low, or zero (control) *Niphargus* density, and Unique ID is the microcosm that was
291 measured repeatedly over time, represented by the variable Day. We used a Tukey test for
292 post-hoc comparisons.

293

294 **RESULTS**

295 **Gut content analyses**

296 The gut content analyses established that both *Niphargus* species ingested biofilm.
297 We found a homogeneous mass of recently ingested organic material (e.g. bacteria,
298 protozoans) and sediment particles in the foreguts of the *Niphargus* individuals (Fig 1C-F).

299 All individuals initially had empty foreguts, indicating that the material found came from
300 biofilm associated with the tiles. Overall, we detected organic material in 32 of the 45 *N.*
301 *kochianus* individuals and both of the *N. fontanus* individuals.

302

303 **Experiment 1. Hypothesis: Stygobite presence will reduce bacterial and protozoan**
304 **abundances and alter biofilm assemblage structure.**

305 The presence of both *N. fontanus* and *Proasellus cavaticus* had a significant positive
306 effect on protozoan abundances found on tiles (Table 1, Fig. 2) compared with the control
307 without stygobites. In *N. fontanus* and *P. cavaticus* microcosms, the number of protozoans
308 was double that of the control (Fig. 2). The post-hoc test for protozoan abundance showed
309 that the effects of both species were different from the control (Tukey-test; *Niphargus* vs
310 control, $P < 0.01$ and *Proasellus* vs control, $P < 0.05$). In the LMM, the random effect explained
311 only 1% of protozoan abundance, i.e. the identity of the individual stygobite used was not a
312 significant predictor of the response.

313 The effect of stygobites on bacterial abundances was less marked (Table 1). Although
314 *P. cavaticus* seemed to reduce the number of bacteria (Fig. 3), this effect was not significant
315 and variation in bacteria abundance was much greater between blocks (Table 1). Neither
316 stygobite species changed the bacterial assemblage structure in terms of altering the relative
317 proportion of small, medium, and large bacteria (data not shown). Block had a highly
318 significant effect on bacterial abundance (Table 1). For example, bacterial abundance was
319 significantly lower in block 4 than in block 1, indicating that bacterial abundance changed
320 significantly with time. Thus, it was important to fit Block as a predictor in the LMMs.

321

322 **Experiment 2. Hypothesis: High densities will remove more bacteria and protozoans**
323 **than low densities and this effect will become more pronounced over time.**

324 As in the first experiment, protozoan abundances were significantly affected by the
325 density of *N. kochianus*, by time, and by the interaction between density and time (Table 2).
326 As with the other two stygobites, the presence of *N. kochianus* at high densities resulted in
327 more protozoans than in the control treatment (Fig. 4). In fact, when averaged across all time
328 points and density treatments, protozoan abundances were twice as high when *Niphargus* was
329 present (Fig. 4). However, these differences did not occur during the first part of the
330 experiment. Abundances remained at comparably low levels in all treatments from day 2 to
331 day 16 (Fig. 4). However, from day 23 on, protozoan abundance increased in the high density
332 *N. kochianus* treatment relative to the control (Fig. 4).

333 No significant differences in the number of protozoan morphotypes occurred across
334 treatments, but the number of protozoan morphotypes in all treatments increased significantly
335 over time (Table 2, Fig. 4).

336 The density treatments did not significantly affect bacterial abundance (Table 2).
337 However, bacterial assemblage structure was significantly affected by *N. kochianus* density,
338 by time, and by the interaction of the two predictors. ‘*Niphargus* Density’ was a significant
339 predictor of the proportion of small and medium bacteria, but not of large bacteria (Fig. 5,
340 Table 2). On day two of the experiment, small bacteria tended to make up a larger proportion
341 of the total bacterial population in the high-density treatment relative to either the low-density
342 or control treatment (Fig. 5). Conversely, the initial relative proportions of medium and large
343 bacteria tended to be higher in the low-density and control treatments (Fig. 5). Throughout
344 the course of the experiment the relative proportions of small, medium, and large bacteria
345 continuously changed. The percentage of small bacteria decreased in the high-density
346 treatment, while the proportion of medium and large bacteria tended to increase (Fig. 5). In
347 the low-density and control treatments, the proportion of medium and large bacterial size

348 classes tended to slightly decline over time. By day 32, the proportion of bacterial size classes
349 was very similar between treatments (Fig. 5).

350 On the mesh tiles excluded from stygobite access, bacterial abundance ($F_{2,243} = 0.5$, P
351 > 0.05) and the proportion of small ($F_{2,243} = 0.1$, $P > 0.05$), medium ($F_{2,243} = 0.02$, $P > 0.05$)
352 and large bacteria ($F_{2,243} = 0.06$, $P > 0.05$) did not differ between treatments.

353

354 **DISCUSSION**

355 Our experiments showed that the *Niphargus* species can ingest biofilm and that the
356 presence of each of the three species altered the biofilm. The strength and nature of this effect
357 depended on stygobite density and the duration of exposure to the biofilm.

358 Our microcosm experiments offer a unique glimpse of macroinvertebrate stygobite
359 behavior and their influence on primary resources within experimental microcosms.
360 However, our experimental design did not enable us to determine whether these are direct
361 food web effects, facilitation via increased nutrient recycling, or a combination of processes.
362 The role of stygobites in groundwater food webs has been intensely debated in recent years
363 (e.g. Boulton et al. 2008). Despite their widespread prevalence and the absence of other top-
364 level consumers, most studies have attributed little importance to obligate groundwater
365 animals, because of the temporal stability of groundwater ecosystems and the low metabolic
366 rates and perceived low abundance of stygobites (Gibert et al. 1994, Boulton et al. 2003,
367 Wilhelm et al. 2006, Sorensen et al. 2013). However, controlled experiments investigating
368 groundwater food webs are scarce (but see Edler and Dodds 1996, Cooney and Simon 2009,
369 Foulquier et al. 2010).

370

371 **Effects on Protozoa**

372 Both single individuals of *N. fontanus* and *P. cavaticus*, as well as *N. kochianus* at
373 high densities, significantly increased protozoan abundance in our experimental microcosms.
374 As there is currently little information on the role of stygobites in groundwater food webs, the
375 consistency of this effect across all experimental species is noteworthy. It remains to be
376 determined whether the stimulatory link to protozoans is mediated directly by feeding activity
377 or indirectly via excretion or bioturbation.

378 Previous studies have shown that microscopically small surface-water crustaceans
379 such as copepods and cladocerans selectively feed on specific protozoan species (Sanders and
380 Wickham 1993, Reiss and Schmid-Araya 2010) and size classes (Stoecker and Capuzzo
381 1990, Sommer et al. 2001), thus demonstrating that these crustaceans can actively target
382 protozoans. Stygobites are also thought to obtain their nutrients from biofilm coating
383 sediments and rocks, including associated protozoans (Baerlocher and Murdoch 1989,
384 Fenwick et al. 2004, Boulton et al. 2008). However, for our study species, predation on
385 protozoans does not appear to be substantial, given that predators tend to reduce prey
386 abundances (Sih et al. 1985, Mamilov et al. 2000) and protozoan abundance did not decline.
387 It is possible that rapid turnover and recruitment of Protozoa completely compensated for
388 losses due to predation. Another possibility is that stygobites may either bioturbate or graze
389 the biofilm, causing tightly bound biofilm fragments to be dislodged from the substratum
390 (e.g. Gibert et al. 1994). These activities would provide a greater surface area for grazing by
391 bacterivorous protozoans, allowing them to reproduce faster and attain higher abundances.

392 Stygobite presence increased morphotype diversity in experiment 2. Protozoans such
393 as flagellates and ciliates are omnipresent in groundwater (Novarino et al. 1997), so the
394 resting spores (Finlay 2002) of many protozoan species would have been present on the
395 biofilm tiles. However, it seems that when stygobites were absent, the spores remained

396 dormant. It is possible that the proliferation of protozoans (caused by stygobites) increased
397 the likelihood that rarer protozoan morphotypes would be detected in our subsamples.

398

399 **Effects on bacteria**

400 Only *P. cavaticus* reduced bacterial abundances in experiment 1, and the effect was
401 not strong compared with changes detected for protozoans. The gut contents of both
402 *Niphargus* species show they clearly ingest tile-associated biofilm. This result indicates that
403 bacteria in biofilm are likely to provide at least some of the diet for stygobites (Boulton et al.
404 2008). However, previous studies have found both strong positive and negative correlations
405 between bacterial responses and stygobite grazing (Griebler et al. 2002, Cook et al. 2007,
406 Foulquier et al. 2010, 2011).

407 In experiment 2 we measured respiration rates from 5 replicates of one small tile in all
408 treatments (reported in Weitowitz 2017). We measured respiration both halfway through and
409 at the end of the experiment. The bacterial activity rates were higher in the presence of
410 stygobites, perhaps because either their grazing or bioturbation removed senescent bacteria
411 and enhanced solute uptake by active bacteria. Such effects may explain the relatively small
412 difference in bacterial abundances between stygobite treatments (Weitowitz 2017). The
413 relatively small amount of bacterial biomass removed by invertebrate and protozoan grazing
414 might be offset by the increase in bacterial growth. Other studies in surface waters and
415 terrestrial ecosystems have also shown an effect of higher-order animals on bacterial activity
416 rates across a range of taxa, including collembolans (Hanlon and Anderson 1979), nematodes
417 (Traunspurger et al. 1997), and protozoans (Hahn and Hoefle 2001).

418 We also observed time-dependent effects on bacterial assemblage structure. These
419 effects might be a direct result of stygobite grazing, an indirect effect associated with
420 increased protozoan grazing, or both given that protozoans were more abundant in the

421 presence of stygobites. In other aquatic systems, protozoan grazing is size-selective
422 (Chrzanowski et al. 1990, Gonzalez et al. 1990, Simek and Chrzanowski 1992) and has been
423 shown to affect bacterial assemblage structure (Hahn and Hoefle 1999, 2001). For example,
424 the uptake efficiency of bacteria by flagellates and ciliates, the dominant protozoans in our
425 biofilm, decreases with prey cell size. No lower uptake limit exists (Hahn and Hoefle 2001).
426 Stygobites consume microbes (Simon et al. 2003, Hallam et al. 2008), so they may also
427 directly affect bacterial assemblage structure. In the presence of stygobites, small and
428 medium-sized bacteria were initially present at lower frequencies than large-sized bacteria,
429 but this pattern quickly disappeared. One explanation for this observation is that the smaller
430 sizes of bacteria responded by increasing their activity and rate of cell division. Such
431 compensatory reactions in response to predation have been observed previously and were
432 attributed to rapid bacterial generation rates (Hanlon and Anderson 1979, Traunspurger et al.
433 1997).

434 Both *N. fontanus* and *P. cavaticus* increased protozoan abundance in the biofilm, but
435 only *P. cavaticus* reduced bacterial abundance and only slightly. These responses may have
436 been at least partly caused by the different feeding strategies of the species, with *P. cavaticus*
437 harvesting the bacterial ‘carpet’ more efficiently than *N. fontanus*. However, both species
438 appear to increase the nutrient availability to protozoans, but through different behaviors.
439 *Proasellus cavaticus* may dislodge biofilm by browsing over sediment and scraping off
440 bacteria, whereas *N. fontanus*, an active swimmer, may dislodge biofilm via bioturbation as it
441 passes over and disturbs the sediments.

442 The relationships between components in groundwater food webs are not limited to
443 organismal interactions. Stygobites also provide food directly to microbes and protozoa by
444 excreting feces or producing pellets of fine interstitial materials (Boulton et al. 2008). We
445 observed these activities in our experimental microcosms. Bacteria are known to process

446 fecal pellets in aquatic habitats (e.g. Yoon et al. 1996, Wotton and Malmqvist 2001), and this
447 activity may partly explain how the bacteria overcame increased grazing pressure. In the
448 control microcosms, however, the nutrient-poor conditions in combination with the reduced
449 nutrient cycling likely provided unfavorable conditions for bacterial reproduction.

450 Aquifers and their associated organisms, particularly protozoa and bacteria, support
451 important ecosystem services such as nutrient (e.g. denitrification, nitrification) and
452 contaminant transformation (e.g. biodegradation) (Mattison et al. 2002, 2005, Tomlinson and
453 Boulton 2008). They also maintain carbon flux through food webs. The effect of stygobites
454 on groundwater biofilm demonstrated here could have important implications for these
455 services, and stygobites may also play a significant role in maintaining clean drinking water.
456 Future studies should address these important issues.

457

458 **CONCLUSIONS**

459 Our experiments suggest that stygobites can increase abundances of protozoa and
460 alter the structure of both protozoa and bacteria assemblages. As for species from surface
461 ecosystems, their impact is likely to depend on their abundance in the systems. To date,
462 however, estimates of stygobite abundance in aquifers are rare (Maurice and Bloomfield
463 2012, Sorensen et al. 2013).

464 Maintaining groundwater ecosystem functionality and stability is becoming
465 increasingly important in the face of environmental pollution and global climate change.
466 Groundwater biota, and particularly stygobites, are adapted for the constant temperature and
467 low-nutrient conditions in groundwater. A change in groundwater temperatures or nutrient
468 levels could therefore lead to the disappearance of whole functional groups of organisms in
469 these simple systems, leading to ecosystem destabilization (Avramov et al. 2013). Further
470 experiments are needed to identify the mechanisms by which stygobites affect groundwater

471 biofilms and influence ecosystem services, and thus build a foundation for an informed
472 approach to the conservation of these systems.

473

474 **ACKNOWLEDGMENTS**

475 Author contributions. DW, JR and ALR conceived the study and designed the
476 experiments. DW carried out the experiments and processed the laboratory samples. DW and
477 JR carried out the statistical analysis. DW wrote the manuscript with significant contributions
478 from JR, ALR, LM and JB.

479 DW was supported by a joint studentship from the National Environment Research
480 Council (NERC) and the University of Roehampton, London. LM and JB publish with the
481 permission of the Executive Director of the British Geological Survey (UKRI). We are
482 indebted to Lee Knight for organizing caving trips and assisting in the collection of stygobite
483 individuals. We would also like to thank Tim Johns from the Environment Agency for
484 organizing access to the chalk boreholes. Additional thanks goes to Professor Rosemary
485 Bailey for statistical help, the University of Roehampton technical staff for providing
486 assistance during the lab work, and to Dr. Robert Busch for the provision of theoretical and
487 practical training on the BD C6 flow cytometer. Szymon Szary, lab technician at the
488 University of Roehampton, contributed valuable ideas in numerous fruitful discussions. We
489 thank the four anonymous reviewers whose comments improved the manuscript.

490

491 **LITERATURE CITED**

492 Adl, S. M., D. C. Coleman and F. Read. 2006. Slow recovery of soil biodiversity in sandy
493 loam soils of Georgia after 25 years of no-tillage management. *Agriculture, Ecosystems*
494 *& Environment* 114: 323-334.

- 495 Arnscheidt, J., J. Dooley, K. Eriksson, C. Hack, H. J. Hahn, T. Higgins, T. K. McCarthy, C.
496 McInerney, P. Wood. 2012. Biogeography and ecology of Irish groundwater fauna:
497 Assessment of the distribution, structure and functioning of subterranean fauna within
498 Irish groundwater systems. (2007-W-MS-1-S1). STRIVE Report of the Environmental
499 Protection Agency Report Series No. 95.
500 https://www.epa.ie/pubs/reports/research/water/STRIVE_95_web.pdf
- 501 Avramov, M., S. I. Schmidt and C. Griebler. 2013. A new bioassay for the ecotoxicological
502 testing of VOCs on groundwater invertebrates and the effects of toluene on *Niphargus*
503 *inopinatus*. *Aquatic Toxicology* 130:1-8.
- 504 BD Biosciences. 2011. Threshold and analysis of small particles on the BD accuri C6 flow
505 cytometer. Technical Bulletin, London, United Kingdom.
- 506 Bailey, R.A. and Reiss, J., 2014. Design and analysis of experiments testing for biodiversity
507 effects in ecology. *Journal of Statistical Planning and Inference*, 144:69-80.
- 508 Billen, G., P. Servais and S. Becquevort. 1990. Dynamics of bacterioplankton in oligotrophic
509 and eutrophic aquatic environments: Bottom-up or top-down control? *Hydrobiologia*
510 207:37-42.
- 511 Bouletreau, S., F. Garabetian, S. Sauvage, J-M. Sanchez-Perez. 2006. Assessing the
512 importance of a self-generated detachment process in river biofilm models. *Freshwater*
513 *Biology* 51:901-912.
- 514 Boulton, A.J. 2000. The subsurface macrofauna. Pages 37 -361 in J. Jones and P. Mulholland
515 (editors) *Streams and Ground Waters*. Academic Press, New York, USA.
- 516 Boulton, A.J. 2005. Chances and challenges in the conservation of groundwater and their
517 dependent ecosystems. *Aquatic Conservation: Marine and Freshwater Ecosystems*. 15:
518 319-323.

- 519 Boulton, A.J., W.F. Humphreys and S.M. Eberhard. 2003. Imperiled subsurface waters in
520 Australia: Biodiversity, threatening processes and conservation. *Aquatic Ecosystem*
521 *Health & Management* 6:41-54.
- 522 Boulton, A. J., G. D. Fenwick, P. J. Hancock and M. S. Harvey. 2008. Biodiversity,
523 functional roles and ecosystem services of groundwater invertebrates. *Invertebrate*
524 *Systematics* 22:103-116.
- 525 Bruno, J. F., K. E. Boyer, J. E. Duffy and S. C. Lee. 2008. Relative and interactive effects of
526 plant and grazer richness in a benthic marine community. *Ecology* 89:2518-2528.
- 527 Cardinale B. J., M. A. Palmer, C. M. Swan, S. Brooks, N. LR. Poff. 2002. The influence of
528 substrate heterogeneity on biofilm metabolism in a stream ecosystem. *Ecology* 83:412-
529 422.
- 530 Chrzanowski, T. H. and K. Simek. 1990. Prey-size selection by freshwater flagellated
531 protozoa. *Limnology & Oceanography* 35:1429-1436.
- 532 Cook, P., B. Veuger, S. Böer and J. J. Middelburg. 2007. Effect of nutrient availability on
533 carbon and nitrogen incorporation and flows through benthic algae and bacteria in near-
534 shore sandy sediment. *Aquatic Microbial Ecology* 49: 65-180.
- 535 Cooney, T. J., and K. S. Simon. 2009. Influence of dissolved organic matter and invertebrates
536 on the function of microbial films in groundwater. *Microbial Ecology* 58:599-610.
- 537 Crowder, L. B., D. D. Squires and J. A. Rice. 1997. Nonadditive effects of terrestrial and
538 aquatic predators on juvenile estuarine fish. *Ecology* 78:1796-1804.
- 539 Duffy, J. E. and M. E. Hay. 2000. Strong impacts of grazing amphipods on the organization
540 of a benthic community. *Ecological Monographs* 70:237-263.
- 541 Edler, C. and W. Dodds. 1996. The ecology of a subterranean isopod, *Caecidotea tridentata*.
542 *Freshwater Biology* 35:249-259.

- 543 Field, A. 2013. Discovering statistics using IBM SPSS statistics. Sage Publications, London,
544 United Kingdom.
- 545 Finlay, B.J. 2002. Global dispersal of free-living microbial eukaryote species. *Science*
546 296:1061-1063.
- 547 Fiser, C., B. Sket, P. Trontelj. 2008. A phylogenetic perspective on 160 years of troubled
548 taxonomy of *Niphargus* (Crustacea: Amphipoda). *Zoologica Scripta* 37:665-680.
- 549 Foissner, W., and H. Berger. 1996. A user- friendly guide to the ciliates (Protozoa,
550 Ciliophora) commonly used by hydrobiologists as bioindicators in rivers, lakes, and
551 waste waters, with notes on their ecology. *Freshwater Biology* 35:375-482.
- 552 Foulquier, A., F. Mermillod-Blondin, F. Malard and J. Gibert. 2011. Response of sediment
553 biofilm to increased dissolved organic carbon supply in groundwater artificially
554 recharged with stormwater. *Journal of Soils and Sediments* 11:382-393.
- 555 Foulquier, A., L. Simon, F. Gilbert, F. Fourel, F. Malard and F. Mermillod- Blondin. 2010.
556 Relative influences of DOC flux and subterranean fauna on microbial abundance and
557 activity in aquifer sediments: New insights from ¹³C- tracer experiments. *Freshwater*
558 *Biology* 55:1560-1576.
- 559 Francois, C. M., F. Mermillod-Blondin, F. Malard, F. Fourel, C. Lecuyer, C. J. Douady, L.
560 Simon. 2016. Trophic ecology of groundwater species reveals specialization in a low-
561 productivity environment. *Functional Ecology* 30:262-273.
- 562 Gasol, J. M. and P. A. Del Giorgio. 2000. Using flow cytometry for counting natural
563 planktonic bacteria and understanding the structure of planktonic bacterial
564 communities. *Scientia Marina* 64:197-224.
- 565 Gibert, J., D. C. Culver, M. Dole- Olivier, F. Malard, M. C. Christman and L. Deharveng.
566 2009. Assessing and conserving groundwater biodiversity: Synthesis and perspectives.
567 *Freshwater Biology* 54:930-941.

- 568 Gibert, J., D. L. Danielopol and J. A. Stanford. 1994. Groundwater ecology. Academic Press,
569 San Diego, California.
- 570 Gibert, J. and L. Deharveng. 2002. Subterranean ecosystems: A truncated functional
571 biodiversity. *BioScience* 52:473-481.
- 572 Gonzalez, J. M., E. B. Sherr, B. F. Sherr. 1990. Size-selective grazing on bacteria by natural
573 assemblages of estuarine flagellates and ciliates. *Applied and Environmental*
574 *Microbiology* 56:583-589.
- 575 Graca, M., L. Maltby and P. Calow. 1994a. Comparative ecology of *Gammarus pulex* (L.)
576 and *Asellus aquaticus* (L.) I: Population dynamics and microdistribution. *Hydrobiologia*
577 281:155-162.
- 578 Graça, M., L. Maltby and P. Calow. 1994b. Comparative ecology of *Gammarus pulex* (L.)
579 and *Asellus aquaticus* (L.) II: Fungal preferences. *Hydrobiologia* 281:163-170.
- 580 Griebler, C. and M. Avramov. 2015. Groundwater ecosystem services: A review. *Freshwater*
581 *Science* 34:355-367.
- 582 Griebler, C., , B. Mindl, D. Slezak, M. Geiger-Kaiser. 2002. Distribution patterns of attached
583 and suspended bacteria in pristine and contaminated shallow aquifers studied with an
584 *in-situ* sediment exposure microcosm. *Aquatic Microbial Ecology* 28:117-129.
- 585 Hahn, M. W. and M. G. Hofle. 1999. Flagellate predation on a bacterial model community:
586 Interplay of size-selective grazing, specific bacterial cell size, and bacterial community
587 composition. *Applied and Environmental Microbiology* 65:4863-4872.
- 588 Hahn, M. W. and M. G. Hofle. 2001. Grazing of Protozoa and its effect on populations of
589 aquatic bacteria. *FEMS Microbiology Ecology* 35:113-121.
- 590 Hallam, F., M. Yacoubi-Khebiza, K. Oufdou, M. Boulanouar. 2008. Groundwater quality in
591 an arid area of Morocco: Impact of pollution on the biodiversity and relationships

- 592 between crustaceans and bacteria of health interest. *Environmental Technology*
593 29:1179-1189.
- 594 Hanlon, R. and J. Anderson. 1979. The effects of Collembola grazing on microbial activity in
595 decomposing leaf litter. *Oecologia* 38:93-99.
- 596 Hervant, F., J. Mathieu, D. Garin and A. Fréminet. 1995. Behavioral, ventilatory, and
597 metabolic responses to severe hypoxia and subsequent recovery of the hypogean
598 *Niphargus rhenorhodanensis* and the epigeal *Gammarus fossarum* (Crustacea:
599 Amphipoda). *Physiological Zoology* 68:223-244.
- 600 Hervant, F., J. Mathieu and H. Barre. 1999. Comparative study on the metabolic responses of
601 subterranean and surface-dwelling amphipods to long-term starvation and subsequent
602 refeeding. *The Journal of Experimental Biology* 202:3587-3595.
- 603 Humphreys, W. 2009. Hydrogeology and groundwater ecology: Does each inform the other?
604 *Hydrogeology Journal* 17:5-21.
- 605 Huws, S., A. McBain and P. Gilbert. 2005. Protozoan grazing and its impact upon population
606 dynamics in biofilm communities. *Journal of Applied Microbiology* 98:238-244.
- 607 Johns, T., J. I. Jones, L. Knight, L. Maurice, P. Wood and A. Robertson. 2015. Regional-
608 scale drivers of groundwater faunal distributions. *Freshwater Science* 34:316-328.
- 609 Jones, M. 1972. The mouthparts of the members of the *Jaera albifrons* group of species
610 (Crustacea: Isopoda). *Marine Biology* 14:264-270.
- 611 Kinsey, J., T. J. Cooney and K. S. Simon. 2007. A comparison of the leaf shredding ability
612 and influence on microbial films of surface and cave forms of *Gammarus minus* Say.
613 *Hydrobiologia* 589:199-205.
- 614 Knight, L. R. F. D., T. Johns. 2015. Auto-ecological studies on *Niphargus aquilex*
615 (Schioedte, 1855) and *Niphargus glenniei* (Spooner, 1952) (Crustacea: Amphipoda:
616 Niphargidae). *Cave and Karst Science* 42:63-77.

- 617 Kota, S., R. C. Borden, and M. A. Barlaz. 1999. Influence of protozoan grazing on
618 contaminant biodegradation. *FEMS Microbiology Ecology* 29:179-189
- 619 Lawrence, M.A. 2015. ez: Easy Analysis and Visualization of Factorial Experiments. R
620 package version 4.3. <https://CRAN.R-project.org/package=ez>.
- 621 Lebaron, P., N. Parthuisot and P. Catala. 1998. Comparison of blue nucleic acid dyes for flow
622 cytometric enumeration of bacteria in aquatic systems. *Applied and Environmental*
623 *Microbiology* 64:1725-1730.
- 624 Mamilov, A. S., B. Byzov, A. Pokarzhevskii and D. Zvyagintsev. 2000. Regulation of the
625 biomass and activity of soil microorganisms by microfauna. *Microbiology* 69:612-621.
- 626 Marshall, M. C. and R. O. Hall. 2004. Hyporheic invertebrates affect N cycling and
627 respiration in stream sediment microcosms. *Journal of the North American*
628 *Benthological Society* 23:416-428.
- 629 Mattison, R., H. Taki and S. Harayama. 2005. The soil flagellate *Heteromita globosa*
630 accelerates bacterial degradation of alkylbenzenes through grazing and acetate excretion
631 in batch culture. *Microbial Ecology* 49:142-150.
- 632 Mattison, R. G., H. Taki and S. Harayama. 2002. The bacterivorous soil flagellate *Heteromita*
633 *globosa* reduces bacterial clogging under denitrifying conditions in sand-filled aquifer
634 columns. *Applied and Environmental Microbiology* 68:4539-4545.
- 635 Maurice, L. and J. P. Bloomfield. 2012. Stygobitic invertebrates in groundwater – A review
636 from a hydrogeological perspective. *Freshwater Reviews* 5:51-71.
- 637 Maurice, L., A. Robertson, D. White, L. Knight, T. Johns, F. Edwards, M. Arietti, J. P. R.
638 Sorensen, D. Weitowitz, B. P. Marchant and J. P. Bloomfield. 2015. The invertebrate
639 ecology of the Chalk aquifer in England (UK). *Hydrogeology Journal* 24:1-16.

- 640 McQueen, D. J., M. R. Johannes, J. R. Post, T. J. Stewart and D. R. Lean. 1989. Bottom-up
641 and top-down impacts on freshwater pelagic community structure. *Ecological*
642 *Monographs* 59:289-309.
- 643 Menge, B. A. 2000. Top-down and bottom-up community regulation in marine rocky
644 intertidal habitats. *Journal of Experimental Marine Biology and Ecology* 250:257-289.
- 645 Mermillod-Blondin, F., M. Creuze des Chatelliers, M. Gerino, J. P. Gaudet. 2000. Testing the
646 effect of *Limnodrilus* sp. (Oligochaeta, Tubificidae) on organic matter and nutrient
647 processing in the hyporheic zone: A microcosm method. *Fundamental and Applied*
648 *Limnology* 149:467-487.
- 649 Morris, B. L., A. R. Lawrence, P. J. Chilton, B. Adams, R. C. Caylow and B. A. Klinck.
650 2003. Groundwater and its susceptibility to degradation: a global assessment of the
651 problems and options for management. UNEP Early Warning & Assessment Report
652 Series RS. 03-3, Nairobi, Kenya 126pp.
- 653 Muylaert, K., K. Van Der Gucht, N. Vloemans, L. D. Meester, M. Gillis and W. Vyverman.
654 2002. Relationship between bacterial community composition and bottom-up versus
655 top-down variables in four eutrophic shallow lakes. *Applied and Environmental*
656 *Microbiology* 68:4740-4750.
- 657 Navarro-Barranco, C., J. M. Tierno-de-Figueroa, J. M. Guerra-Garcia, L. Sanchez-Tocino, J.
658 C. Garcia-Gomez. 2013. Feeding habits of amphipods (Crustacea: Malacostraca) from
659 shallow soft bottom communities: Comparison between marine caves and open habitats.
660 *Journal of Sea Research* 78: 1-7.
- 661 Naylor, E. 1955. The diet and feeding mechanism of *Idotea*. *Journal of the Marine Biological*
662 *Association of the United Kingdom* 34:347-355.

- 663 Novarino, G. A., H. Warren, G. Butler, A. Lambourne, J. Boxshall, N.E. Bateman, R.W.
664 Kinner, R.A. Harvey and B. Teltsch. 1997. Protistan communities in aquifers: A review.
665 FEMS Microbiology Reviews. 20:261-275.
- 666 Nyström, P. and J. Strand. 1996. Grazing by a native and an exotic crayfish on aquatic
667 macrophytes. *Freshwater Biology* 36:673-682.
- 668 Perkins, D.M., G. Yvon- Durocher, B.O. Demars, Reiss, J., D.E. Pichler, N. Friberg, M.
669 Trimmer and G. Woodward. 2012. Consistent temperature dependence of respiration
670 across ecosystems contrasting in thermal history. *Global Change Biology* 18:1300-
671 1311.
- 672 R Development Core Team. 2013. R: A language and environment for statistical computing.
673 R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL
674 <http://www.R-project.org/>.
- 675 Reiss, J. and J. M. Schmid-Araya. 2010. Life history allometries and production of small
676 fauna. *Ecology* 91: 497-507.
- 677 Rosemond, A. D., C. M. Pringle, A. Ramírez and M. J. Paul. 2001. A test of top-down and
678 bottom-up control in a detritus-based food web. *Ecology* 82: 2279-2293.
- 679 Sanders, R. W. and S. A. Wickham. 1993. Planktonic Protozoa and metazoa: Predation, food
680 quality and population control. *Aquatic Microbial Ecology* 7: 197-223.
- 681 Shurin, J. B., J. L. Clasen, H. S. Greig, P. Kratina and P. L. Thompson. 2012 Warming shifts
682 top-down and bottom-up control of pond food web structure and function. *Philosophical*
683 *Transactions of the Royal Society of London: Series B, Biological Sciences* 367: 3008-
684 3017.
- 685 Sih, A., P. Crowley, M. McPeck, J. Petranka and K. Strohmeier. 1985. Predation,
686 competition, and prey communities: A review of field experiments. *Annual Review of*
687 *Ecology and Systematics* 16: 269-311.

- 688 Simek, K. and T. H. Chrzanowski. 1992. Direct and indirect evidence of size-selective
689 grazing on pelagic bacteria by freshwater nanoflagellates. *Applied and Environmental*
690 *Microbiology* 58: 3715-3720.
- 691 Simon, K.S., E. F. Benfield, S. A. Macko. 2003. Food web structure and the role of epilithic
692 biofilms in cave streams. *Ecological Society of America* 84: 2395-2406.
- 693 Sommer, U., F. Sommer, B. Santer, C. Jamieson, M. Boersma, C. Becker and T. Hansen.
694 2001. Complementary impact of copepods and cladocerans on phytoplankton. *Ecology*
695 *Letters* 4: 545-550.
- 696 Sorensen, J.P.R., L. Maurice, F. K. Edwards, D. J. Lapworth, D. S. Read, D. Allen, A. S.
697 Butcher, L. K. Newbold, B. R. Townsend, P. J. Williams. 2013. Using boreholes as
698 windows into groundwater ecosystems. *PLOS One* 8: 1-13.
- 699 Spicer, J. I. 1998. Is the reduced metabolism of hypogean amphipods solely a result of food
700 limitation? *Hydrobiologia* 377: 201-204.
- 701 Stoecker, D. K. and J. M. Capuzzo. 1990. Predation on protozoa: Its importance to
702 zooplankton. *Journal of Plankton Research* 12: 891-908.
- 703 Tomlinson, M. and A. Boulton. 2008. Subsurface groundwater dependent ecosystems: A
704 review of their biodiversity, ecological processes and ecosystem services. *Waterlines*
705 *Occasional Paper* 8: pp89.
- 706 Traunspurger, W., M. Bergtold and W. Goedkoop. 1997. The effects of nematodes on
707 bacterial activity and abundance in a freshwater sediment. *Oecologia* 112: 118-122.
- 708 Troussellier, M., C. Courties, P. Lebaron and P. Servais. 1999. Flow cytometric
709 discrimination of bacterial populations in seawater based on SYTO 13 staining of
710 nucleic acids. *FEMS Microbiology Ecology* 29: 319-330.
- 711 Venables, W. N. and B. D. Ripley. 2002. *Modern Applied Statistics with S*. Fourth Edition.
712 Springer, New York. ISBN 0-387-95457-0.

- 713 Vercaene-Eairmal, M., B. Lauga, S. Saint Laurent, N. Mazzella, S. Boutry, M. Simon, S.
714 Karama, F. Delmas, R. Duran. 2010. Diuron biotransformation and its effects on
715 biofilm bacterial community structure. *Chemosphere* 81: 837-843.
- 716 Weitowitz, D.C. 2017. An investigation into the distribution of obligate groundwater animals
717 (stylobites) in England and Wales. (Unpublished doctoral dissertation). University of
718 Roehampton, London, United Kingdom.
- 719 Wey, J. K., K. Jurgens and M. Weitere. 2012. Seasonal and successional influences on
720 bacterial community composition exceed that of protozoan grazing in river biofilms.
721 *Applied and Environmental Microbiology* 78: 2013-2024.
- 722 Wilhelm, F. M., S. J. Taylor and G. L. Adams. 2006. Comparison of routine metabolic rates
723 of the stygobite, *Gammarus acherondytes* (Amphipoda: Gammaridae) and the
724 stygophile, *Gammarus troglophilus*. *Freshwater Biology* 51: 1162-1174.
- 725 Wipfli, M. S., J. Hudson and J. Caouette. 1998. Influence of salmon carcasses on stream
726 productivity: Response of biofilm and benthic macroinvertebrates in southeastern
727 Alaska, U.S.A. *Canadian Journal of Fisheries and Aquatic Science* 55: 1503-1511.
- 728 Wotton, R. S. and B. Malmqvist. 2001. Feces in aquatic ecosystems: Feeding animals
729 transform organic matter into fecal pellets, which sink or are transported horizontally by
730 currents; these fluxes relocate organic matter in aquatic ecosystems. *BioScience* 51:
731 537-544.
- 732 Yoon, W. D., J.-C. Marty, D. Sylvain, P. Nival. 1996. Degradation of faecal pellets in *Pegea*
733 *confoederata* (Salpidae, Thaliacea) and its implication in the vertical flux of organic
734 matter. *Journal of Experimental Marine Biology and Ecology* 203: 147-177.
- 735 Zuur, A. F., Ieno, E. N., Walker, N. J., Savaliev, A. A. & Smith, G. M. 2009. *Mixed Effects*
736 *Models and Extensions in Ecology with R*. Springer, New York.

737 **FIGURE CAPTIONS**

738 Fig. 1. Photos of the stygobite grazer species (A) *Niphargus fontanus*, (B) *Proasellus*
 739 *cavaticus* and (C) *N. kochianus*. Panels C–F show the gut content of *N. kochianus*.
 740 These contents are shown at x400 magnification in (E) and (F) and reveal a
 741 homogeneous mass of organic material and sediment particles.

742 Fig. 2. The effect of control, *N. fontanus*, and *P. cavaticus* treatments (one individual per
 743 replicate) on protozoan abundances (individuals / mL H₂O) in feeding microcosms
 744 (experiment 1). The box and whisker plots summarize replicates from four different
 745 experimental time blocks, with individual data points superimposed to visualize the
 746 distribution of the data. The horizontal line within the box indicates the median and
 747 the boundaries of the box indicate the 25th and 75th percentiles.

748 Fig. 3. The effect of control, *N. fontanus*, and *P. cavaticus* treatments (one individual per
 749 replicate) on bacterial abundances (individuals / μL H₂O) in feeding microcosms
 750 (experiment 1). The box and whicker plots summarize replicates from four different
 751 experimental time blocks, with individual data points superimposed to visualize the
 752 distribution of the data. The horizontal line within the box indicates the median and
 753 the boundaries of the box indicate the 25th and 75th percentiles.

754 Fig. 4. The effect of control and different density treatments (low and high) of *N. kochianus*
 755 on protozoan abundance (individuals / mL⁻¹ H₂O) and number of protozoan
 756 morphotypes (number / mL) over time in experiment 2. Different density treatments
 757 are symbolized by dotted (control), dashed (low density), and solid (high density)
 758 lines. Protozoan responses were sampled on six occasions (days 2, 5, 11, 16, 23 and
 759 32).

760 Fig. 5. The effect of control and different density treatments (low and high) of *N. kochianus*
 761 on the relative proportion of small, medium, and large bacterial size classes (as % of

762 total bacteria) over time in experiment 2. Different density treatments are symbolized
763 by dotted (control), dashed (low density), and solid (high density) lines. Bacteria
764 responses were sampled on nine occasions (days 2, 3, 5, 9, 11, 16, 18, 23, 27, 32).