



**Manchester  
Metropolitan  
University**

---

Cirocco, Robert M and Watling, Jennifer R and Facelli, José M (2020) The combined effects of water and nitrogen on the relationship between a native hemiparasite and its invasive host. *New Phytologist*. ISSN 0028-646X

---

**Downloaded from:** <https://e-space.mmu.ac.uk/626585/>

**Version:** Accepted Version

**Publisher:** Wiley

**DOI:** <https://doi.org/10.1111/nph.16944>

Please cite the published version

<https://e-space.mmu.ac.uk>



DR ROBERT MICHAEL CIROCCO (Orcid ID : 0000-0002-1609-2531)

Article type : Regular Manuscript

**The combined effects of water and nitrogen on the relationship between a native hemiparasite and its invasive host.**

**Robert M. Cirocco<sup>1\*</sup>, Jennifer R. Watling<sup>1,2</sup> and José M. Facelli<sup>1</sup>**

<sup>1</sup>School of Biological Sciences, The University of Adelaide, Adelaide, SA 5005, Australia;

<sup>2</sup>Ecology and Environment Research Centre, Manchester Metropolitan University, Manchester, M15 6BH, UK

Robert Cirocco ORCID ID: 0000-0002-1609-2531

Jennifer Watling ORCID ID: 0000-0001-6305-9905

José Facelli ORCID ID: 0000-0002-0408-0082

Author for correspondence:

*Robert M. Cirocco*

*Tel: +61 8313 5281*

*Email: robert.cirocco@adelaide.edu.au*

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/nph.16944](https://doi.org/10.1111/nph.16944)

This article is protected by copyright. All rights reserved

Received: 12 May 2020

Accepted: 5 September 2020

## Summary

- Stem hemiparasites are dependent on their hosts for water and nitrogen. Most studies, however, assess the influence of one factor on parasite:host associations, thus limiting our mechanistic understanding of their performance in nature.
- We investigated the combined effects of water and nitrogen (N) availability on both host (*Ulex europaeus*) and parasite (*Cassitha pubescens*).
- Parasite infection significantly decreased host shoot biomass and shoot:root ratio more severely in high water than low water, irrespective of N supply. Parasite stem [N] was significantly higher in high water than low water treatments, regardless of N supply, but parasite biomass didn't vary among treatments. Irrespective of water and N supply, infected plants had significantly lower total, root and nodule biomass, predawn and midday quantum yields, maximum electron transport rates, water potentials and nitrogen concentration [N]. Parasite  $\delta^{13}\text{C}$  was significantly higher than that of the host.
- Our results suggest that stem hemiparasites can better extract resources from hosts when water availability is high, resulting in greater impact on the host in these conditions. Where hemiparasitic plants are being investigated as biocontrol for invasive weeds, they may be more effective in wetter habitats than in dry ones.

**Key words:** biomass, carbon isotope, chlorophyll fluorescence, hemiparasite, legume, nitrogen, *Ulex europaeus*, water potential.

## Introduction

Parasitic plants comprise *c.* 1% of all angiosperms and occur on all continents apart from Antarctica (Heide-Jørgensen, 2013). They are an important group because of their direct and indirect impacts on ecosystems, communities and individual species across their global distribution (Press & Phoenix, 2005). For example, parasitic plants can increase nutrient cycling or affect plant community structure and diversity via their impacts at the host population level

(Bardgett *et al.*, 2006; Queded, 2008; Hartley *et al.*, 2015; Těšitel *et al.*, 2020). Numerous studies have investigated parasitic plant effects on their hosts (Press & Graves, 1995; Watling & Press, 2001; Press & Phoenix, 2005; Bell & Adams, 2011), but few have studied the impact of multiple environmental factors on these associations (e.g. Sui *et al.*, 2019). For instance, it is well known that abiotic factors such as nitrogen and water availability alter the impacts of parasitic plants on host performance (e.g. Cechin & Press, 1993; Le *et al.*, 2015), but they are rarely studied in combination, and where they have, they have largely focused on annual parasites of agricultural crops.

There have been several papers on the influence of nitrogen (N) on parasitic plant effects on their hosts, with mixed results. Some studies found that N had no influence on the association between *Striga hermonthica* and some cereal hosts (Gurney *et al.*, 1995; Aflakpui *et al.*, 1998, 2002, 2005; Sinebo & Drennan, 2001). By contrast, high N was found to reduce the negative effect of other annual root hemiparasites such as *Rhinanthus minor*, *Phtheirospermum japonicum* and *S. hermonthica* on hosts (Gibson & Watkinson, 1991; Cechin & Press, 1993, 1994; Irving *et al.*, 2019). High N was also found to reduce the negative effect of the annual stem holoparasite *Cuscuta* on its hosts (Jeschke & Hilpert, 1997; Shen *et al.*, 2013; but see Jeschke *et al.*, 1997). Cirocco *et al.* (2017) reported that in high N, the negative effect of the perennial stem hemiparasite, *Cassytha pubescens*, on root biomass of the native and invasive hosts, *Acacia paradoxa* and *Ulex europaeus*, was ameliorated and enhanced, respectively. In addition, this stem hemiparasitic vine strongly decreased nodulation of both these leguminous hosts, regardless of N supply (Cirocco *et al.*, 2017).

Surprisingly few papers have explored the influence of water on host:parasite associations, and most have investigated stem holo- or hemiparasitic vines. For instance, although the holoparasitic vine *Cuscuta australis* negatively affected photosynthesis of *Mikania micrantha*, irrespective of water supply, its negative impact on host stomatal conductance and transpiration was enhanced in low water conditions (Le *et al.*, 2015). By contrast, Evans and Borowicz (2013, 2015) found that *Cuscuta gronovii* performed better and had a greater impact on host biomass in high water conditions. Similarly, the stem hemiparasite *Cassytha pubescens* also grew more vigorously and had a more severe impact on total biomass of the invasive host, *Ulex europaeus* in high water conditions (Cirocco *et al.*, 2016a). Cirocco *et al.* (2018) also found in a field study that the effect of *C. pubescens* on predawn quantum yield ( $F_v/F_m$ ) of *U. europaeus* was strongest at the site with

highest water availability. Miller *et al.* (2003) investigated the impact of the mistletoe *Amyema miquelii* on the tree host *Eucalyptus largiflorens* at sites varying in soil salinity. Hosts at the sites with lowest salinity were more likely to be infected than those at high salinity sites. The impact of the mistletoe on host predawn water potential and  $\delta^{13}\text{C}$ , however, did not vary significantly across sites, or with parasite load (Miller *et al.*, 2003).

Manipulating more than one biotic or abiotic factor is desired to better mimic complex field conditions. However, these studies are rare likely due to their large size and associated logistical difficulties and, to our knowledge, none have investigated stem hemiparasites (Matthies & Egli, 1999; Gao *et al.*, 2019; Jokien & Irving, 2019; Sui *et al.*, 2019). Two studies investigated the influence of water and nitrogen on host:parasite associations involving the annual root hemiparasite *Rhinanthus alectorolophus*, one with perennial hosts (Korell *et al.*, 2019), and another with annual grasses (Těšitel *et al.*, 2015). Korell *et al.* (2019) found that water and nitrogen had no interactive effect on the association. In contrast, water and nitrogen did have an interactive influence on parasite effects on the annual grasses (Těšitel *et al.*, 2015). These contrasting results using the same parasite species may be related to differences in experimental design and/or host species. For example, annual hosts might be more sensitive to infection than perennial hosts because they have less capacity for resource storage to buffer parasite resource removal. With so few studies, however, it is difficult to generalise.

Here, we investigated the effect of water and nitrogen in combination on performance of a perennial, stem hemiparasite, *Cassytha pubescens*, and its impact on the perennial leguminous shrub, *Ulex europaeus*. We hypothesised that the parasite would negatively impact this major invasive shrub across water  $\times$  nitrogen treatments, based on our previous field study which demonstrated a significant effect of the parasite regardless of environmental conditions (Cirocco *et al.*, 2018). We also hypothesised that the parasite would perform better and have a more pronounced effect on the host in high water treatments, regardless of N supply. This is because we previously found that high water availability enhanced the impact of the parasite on total biomass of *U. europaeus* (Cirocco *et al.*, 2016a), but nitrogen had no effect (Cirocco *et al.*, 2017). To gauge parasite impacts on health of *U. europaeus*, we measured host photosynthetic performance, water and nutrient status,  $\delta^{13}\text{C}$  and biomass. These same measures were quantified for *C. pubescens* to assess its performance on the host in the various treatments.

## Materials and Methods

### *Study species*

*Ulex europaeus* L. (Fabaceae) is a spiny shrub that can grow to 1–4 m tall and has a life span of 30 years (Hornoy *et al.*, 2011). Being a legume it can grow well on nutrient poor soils by obtaining nitrogen in reduced form via engagement with nitrogen-fixing bacteria, namely strains of *Bradyrhizobium* (Weir *et al.*, 2004; Rodríguez-Echeverría, 2010). It produces vast numbers of long-lived seed, has vigorous growth and can rapidly invade disturbed areas (Parsons & Cuthbertson, 2001). *Ulex europaeus* is native to the Iberian Peninsula, but over time has been introduced to all continents (apart from Antarctica), and has become so problematic that it is on the world's 100 worst invasive alien species list (Lowe *et al.*, 2000; see Hornoy *et al.*, 2013).

*Cassytha pubescens* R. Br. (Lauraceae) is an Australian native, perennial, hemiparasitic vine (*c.* 0.5–1.5 mm in diameter) that has indeterminate growth and thus, can infect multiple hosts at any one time (Weber, 1981; Kokubugata *et al.*, 2012). It is an obligate parasite without roots that coils around and attaches to host stems not greater than *c.* 2.5 cm in diameter (McLuckie, 1924; Weber, 1981). *C. pubescens* is a generalist parasite and commonly infects *U. europaeus* in temperate southern Australia.

### *Experimental design*

*U. europaeus* plants (*c.* 35 cm in height) were acquired from the field as described in Cirocco *et al.* (2017) in early September 2017. They were transplanted into 1.65-l pots containing sand with pH 4.75, which is similar to the pH of soil in the field where these parasite:host associations occur (Cirocco *et al.*, 2018). These plants were then randomly allocated into treatments (although treatments were not imposed until later; see below): high water or low water, and with high or low nitrogen availability. Each treatment combination had 10 uninfected and 10 *Cassytha* infected *U. europaeus*. Plants were also randomly allocated into 8 blocks with each block containing all factorial combinations.

*U. europaeus* were infected with the parasite in early November, 2017, according to Shen *et al.* (2010), and prior to water and nitrogen treatments being imposed. Briefly, host plants already infected with *C. pubescens* were positioned adjacent plants to be infected. The parasite was allowed to establish separate, independent connections with the experimental plants. Once

haustoria of the parasite were attached and appeared fully developed, the connection was severed from the donor plant. This process took 3 months (Supporting Information Fig. S1). During this time, plants were supplied monthly with 100 ml of liquid fertiliser (Nitrosol, Rural Research Ltd, Auckland, New Zealand; NPK 8 : 3 : 6), as per manufacturer's recommended dosage. Plants were grown in an evaporatively cooled glasshouse at The University of Adelaide.

All plants were subsequently transplanted into 5-l pots, containing the same soil medium, in late February 2018. Water  $\times$  nitrogen treatments began at this time and plants were re-randomized within blocks fortnightly to negate any small light differences within the glasshouse (Supporting Information Fig. S1). Plants in the high and low water treatments were kept at 100% and 60% field capacity, respectively. Field capacity of the sand was determined using a modification of the filter-paper technique (Bouyoucos, 1929; see Cirocco *et al.*, 2016a for details). Plants in the high and low nitrogen treatments were supplied with standard or modified Hoagland's solution, respectively (refer to Cirocco *et al.*, 2017 for details). As *U. europaeus* is a legume low nitrogen plants were not supplied with an N source ( $\text{Ca}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$  and  $\text{KNO}_3$  were substituted with  $\text{CaCl}_2$  and  $\text{KCL}$ , respectively). All plants were supplied with 100 ml of the relevant Hoagland's solution weekly for the first 6 weeks, and then fortnightly following that. Treatments ran for *c.* 3 months and the experiment ended in late May-early June 2018. As the latter is the transition between autumn and winter in the southern hemisphere, plants were supplied with a supplemental light source for the final 2 weeks of the experiment when chlorophyll fluorescence and water potential measurements were made.

#### *Chlorophyll fluorescence and water potential ( $\Psi$ )*

Predawn quantum yield ( $F_v/F_m$ ) of both host and parasite ( $n = 7-8$ ) was measured 81 days after treatments were imposed (DAT) with a portable chlorophyll fluorometer (MINI-PAM, Walz, Effeltrich, Germany) equipped with a leaf-clip (2030-B, Walz, Effeltrich, Germany). The MINI-PAM and leaf clip were also used to generate rapid light response curves (RLCs) (see Cirocco *et al.*, 2017 for details). The RLC measurements for both host and parasite were made between 10:00 and 13:30 h over 2 days (83 and 84 DAT). Efficiency of PSII in light ( $\Phi_{\text{PSII}}$ ) of host and parasite ( $n = 7-8$ ) was recorded at the sixth light step of the RLCs ( $\text{PPFD} = 981 \pm 5 \mu\text{mol m}^{-2} \text{s}^{-1}$ ,  $n = 94$ ). Maximum electron transport rate ( $\text{ETR}_{\text{max}}$ ) for both host and parasite ( $n = 7-8$ ) was calculated using the RLCs by the Win-Control 3 software (ver 3.25, Walz).

Midday water potentials ( $\Psi$ ) were determined on freshly cut shoots of *U. europaeus* and tendrils of *C. pubescens* using a Scholander-type pressure chamber with a digital gauge (PMS Instrument Company, Albany, OR). Midday water potential measurements were made between 12:00 and 14:45 h over 3 days ( $n = 7-8$ ).

#### *Biomass, $\delta^{13}\text{C}$ and nitrogen concentration*

A destructive harvest was conducted using the six healthiest blocks (including parasite) at the end of the experiment (93–112 DAT). Above and below ground biomass as well as nodules of *U. europaeus* and parasite stems ( $n = 6$ ) were oven-dried at 60°C for seven days. Carbon isotope composition ( $\delta^{13}\text{C}$ ) and N concentration of *U. europaeus* spines and tendrils of *C. pubescens* were determined by mass spectrometry (GV Instruments, Manchester, UK) and elemental analysis (CUBE Elemental Analyser, Elementar Analysensysteme, Hanau, Germany) at Flinders Analytical, Adelaide.

#### *Statistical analysis*

Variances were homogeneous for both host and parasite data, unless otherwise stated. Three-way ANOVAs were used to analyse host parameters. If no three-way interactions (infection  $\times$  water  $\times$  nitrogen) were detected for a particular host parameter, then two-way interactions were considered, if these were also not detected then significant independent effects were treated as valid. For example a valid independent effect of infection would signify that the parasite had an effect on the host parameter regardless of the water or nitrogen conditions. Parasite data were analysed with two-way ANOVAs testing for interactions between water  $\times$  nitrogen. Again, if no two-way interaction was detected for a particular parasite parameter, then independent effects of water or nitrogen were considered. All analyses were conducted with JMP software v.4.0.3 (SAS institute Inc., Cary, NC, USA) and  $\alpha = 0.05$ .

## **Results**

### *Host and parasite growth*

There were no significant treatment interactions for total biomass of *U. europaeus*, but it was independently affected by infection (Table 1; Fig. 1a). On average, infection suppressed total biomass by 71% (Fig. 1b). Water also independently affected this parameter: total biomass (gram



dry weight) of low water plants ( $53.3 \pm 6.7$ ) was 16% lower than that of high water plants ( $63.7 \pm 7.8$ ) (Table 1). There was an infection  $\times$  water interaction for shoot biomass (Table 1; Fig. 1d), with infected plants being around 75% smaller than uninfected plants in high water compared with 68% smaller in low water (Fig. 1e). No treatment interactions were found for root biomass, but it was independently affected by infection which resulted in root biomass being 67% lower, on average across all treatments (Table 1; Fig. 1f,g).

Similar to shoot biomass, there was an infection  $\times$  water interaction for shoot:root ratio of *U. europaeus* (Tables 1, 2). Shoot:root ratio (S:R) of high water infected plants was 15% lower than that of high water uninfected plants, whereas S:R of low water infected plants was not significantly different from that of low water uninfected plants (Table 2). There were no treatment interactions detected for nodule biomass or nodule biomass  $g^{-1}$  root biomass, but these parameters were independently affected by infection and nitrogen (Tables 1, 2). Infection suppressed nodule biomass by 39%, but increased nodule biomass  $g^{-1}$  root biomass by 47% (Table 2). Nodule biomass (including on a  $g^{-1}$  root biomass basis) of low nitrogen plants was on average 20% higher than that of high nitrogen plants (Table 2). Parasite biomass (including on a  $g^{-1}$  host total biomass basis) was unaffected by water or nitrogen treatments (Table 3; Fig. 2a,b).

#### *Photosynthetic performance and water status of host and parasite*

There were no treatment interactions detected for  $F_v/F_m$ ,  $\Phi_{PSII}$  or  $ETR_{max}$  of *U. europaeus*, but these parameters were significantly affected by infection (Table 4; Fig. 3a,c,e). Infection suppressed host  $F_v/F_m$ ,  $\Phi_{PSII}$  and  $ETR_{max}$  by 3, 26 and 34%, respectively (Fig. 3b,d,f). Treatments had no significant impact on  $F_v/F_m$ ,  $\Phi_{PSII}$  or  $ETR_{max}$  of *C. pubescens* (Table 3; Fig. 4a,b,c). There were also no interactions detected for midday  $\Psi$  of *U. europaeus*, however, there were significant independent effects of infection, water and nitrogen for this variable (Table 4; Fig. 5a). On average, midday  $\Psi$  of infected plants was 15% lower than uninfected plants (Fig. 5b). In addition, low water and low nitrogen plants both had an 11% more negative midday  $\Psi$  relative to high water and high nitrogen plants, respectively (Fig. 5c,d). Midday  $\Psi$  of *C. pubescens* was significantly lower (by 30%) than that of its host ( $F_{1,47} = 84.5$ ;  $P = <0.0001$ ) (Fig. 5e). There was no interactive effect of water  $\times$  nitrogen on midday  $\Psi$  of the parasite, but it was significantly affected by water supply (Table 3; Fig. 5f). As found for the host, midday  $\Psi$  of *C. pubescens* in the low water treatment was 10% more negative than the parasite growing in high water conditions (Fig. 5g).

### Carbon isotope and nitrogen status of host and parasite

There were no treatment interactions detected for  $\delta^{13}\text{C}$  of *U. europaeus*, however, infection, water and nitrogen all had independent effects on this host parameter (Table 4). The  $\delta^{13}\text{C}$  of uninfected plants was significantly lower than that of infected plants (Table 2). Moreover,  $\delta^{13}\text{C}$  of *U. europaeus* in either high water or high nitrogen was significantly lower relative to those in low water or low nitrogen conditions, respectively (Table 2). Also,  $\delta^{13}\text{C}$  differed significantly between infected *U. europaeus* and *C. pubescens* ( $F_{1,35} = 123$ ;  $P = <0.0001$ ).  $\delta^{13}\text{C}$  of *C. pubescens* ( $-28.2 \pm 0.17\text{‰}$ ) was significantly higher than that of infected *U. europaeus* ( $-30.2 \pm 0.17\text{‰}$ ) ( $n = 24$ ).

There was no interaction between water and nitrogen on  $\delta^{13}\text{C}$  of *C. pubescens*, however, similar to the host, both these factors independently affected  $\delta^{13}\text{C}$  of the parasite (Table 3; Fig. 4d).  $\delta^{13}\text{C}$  of *C. pubescens* in high water ( $-28.4 \pm 0.17\text{‰}$ ) was significantly lower than in low water conditions ( $-27.9 \pm 0.29\text{‰}$ ) ( $n = 12$ ). Parasite  $\delta^{13}\text{C}$  in high nitrogen ( $-28.6 \pm 0.15\text{‰}$ ) was significantly lower than that of *C. pubescens* in the low nitrogen treatment ( $-27.8 \pm 0.25\text{‰}$ ) ( $n = 12$ ).

There were no treatment interactions found for foliar (i.e. spine) N concentration of *U. europaeus*, but it was independently affected by infection (Table 4; Fig. 6a). Foliar nitrogen concentration of infected plants was 10% lower than that of uninfected ones (Fig. 6b). No water  $\times$  nitrogen interaction was detected for parasite stem N concentration, however, it was independently affected by water (Table 3; Fig. 6c). Nitrogen concentration of *C. pubescens* in low water was 18% lower compared with that in high water (Fig. 6d).

### Discussion

Our hypothesis that water and nitrogen would not have an interactive influence on the impact of *C. pubescens* on *U. europaeus* was supported by the majority of the data. Irrespective of water and nitrogen treatments, infection with the native parasitic vine strongly decreased total biomass of the invasive shrub by 70%. Similarly, Korell *et al.* (2019) found that *Rhinanthus alectorolophus* negatively affected growth of three perennial hosts, regardless of water  $\times$  nitrogen treatments. We also observed that infection had a greater impact on shoot biomass and shoot:root ratio of hosts in high water than low water, irrespective of N supply. In contrast, Těšitel *et al.* (2015) found that *R. alectorolophus* grew better and had a stronger impact on aboveground biomass of maize and wheat in low water conditions (when N supply was high). This disparity between findings may be related to *R. alectorolophus* being able to maintain sufficient stomatal conductance and resource

removal under water deficits (Těšitel *et al.*, 2015). Conversely, *C. pubescens* appears to have lower stomatal conductance (parasite  $\delta^{13}\text{C}$  higher in low water) and resource removal (less parasite N in low water, Fig. 6d) and thus, less impact in low water conditions. Relative to invasive hosts, some native hosts may tolerate infection due to their physiology (e.g. less profligate resource use and supply to the parasite) while others may display defence at the haustorial interface (Cameron & Seel, 2007; Facelli *et al.*, 2020). Thus, although native hemiparasites may still provide a potential means of controlling invasive weeds, their impact will be greater when water is not limited, and may also decline as drought becomes more likely in some regions as a consequence of climate change (Sala *et al.*, 2000; Těšitel *et al.*, 2017).

The impact of *C. pubescens* on growth of *U. europaeus* may be partly due to its effect on host photosynthesis (Fig. 3b,d,f). In the current study, host  $F_v/F_m$  was 3% lower for infected *U. europaeus*, irrespective of water or nitrogen treatment. In contrast, Těšitel *et al.* (2015) found that *R. alectorolophus* grew more and had a greater negative effect on  $F_v/F_m$  of maize when water was low and nitrogen was high. Thus, more parasite growth may translate to greater impact on this host parameter, and as *C. pubescens* did not grow better in any of the treatments this may explain why findings differed between studies. In two glasshouse experiments, *C. pubescens* also had a significant negative impact on  $F_v/F_m$  of *U. europaeus*, irrespective of water treatment or host size (Cirocco *et al.*, 2016a, 2020). In addition, despite environmental variation across field sites  $F_v/F_m$  of infected *U. europaeus* was significantly lower (by 5–10%) relative to uninfected plants (Cirocco *et al.*, 2018).  $F_v/F_m$  of another invasive host, *Cytisus scoparius*, was also significantly lower when infected with *C. pubescens* in a glasshouse experiment (Shen *et al.*, 2010), but not when measured in the field (Prider *et al.*, 2009). In both field and glasshouse conditions, the native parasite had no significant impact on  $F_v/F_m$  of *Leptospermum myrsinoides* likely due to this native host's ability (possibly through co-evolution) to maintain photoprotective capacity and xanthophyll engagement in response to infection (Prider *et al.*, 2009; Girocco *et al.*, 2015).

Whether invasive hosts such as *U. europaeus* have insufficient photoprotective capacity to cope with infection, increasing susceptibility to chronic photoinhibition requires investigation.

*Ulex europaeus*  $\text{ETR}_{\text{max}}$  was also significantly lower (by 34%) when infected with *C. pubescens*, regardless of water or nitrogen treatment. We found a similar impact of infection on  $\text{ETR}_{\text{max}}$  of this host (36%), irrespective of environmental variation among three field sites (Cirocco *et al.*, 2018). In three glasshouse experiments,  $\text{ETR}_{\text{max}}$  and midday ETR of infected *U. europaeus* were

also both significantly lower regardless of variations in host size, nitrogen or light (Cirocco *et al.*, 2016b, 2017, 2020). Shen *et al.* (2010) also found that  $ETR_{max}$  was significantly lower for *Cytisus scoparius* when infected with *C. pubescens*. By contrast, *C. pubescens* had no significant impact on  $ETR_{max}$  of *A. paradoxa* regardless of nitrogen (Cirocco *et al.*, 2017). Although midday ETR of infected *L. myrsinoides* was 36% lower when grown in high light, it was unaffected by *C. pubescens* in low light grown hosts (Cirocco *et al.*, 2016b). Shen *et al.* (2013) found that photosynthesis of the invasive *Mikania micrantha* significantly declined in response to infection with the stem holoparasite *Cuscuta campestris*, irrespective of nitrogen. Jokinen & Irving (2019) found that for the *Orobancha minor:Trifolium pratense* native association, photosynthesis of infected plants was significantly lower than uninfected plants, regardless of manipulations of light and nitrogen. Negative effects of parasitic plants on host photosynthesis are typically attributed to parasite acquisition of resources, resulting in lower host nitrogen (Cirocco *et al.*, 2018; Jokinen & Irving, 2019), stomatal conductance (Shen *et al.*, 2010; Girocco *et al.*, 2017), or both (Shen *et al.*, 2013). Infected plants in our study did show signs of water and nitrogen depletion (Figs 5b, 6b) and possibly lower stomatal conductance as suggested by infected plants having significantly higher  $\delta^{13}C$  than uninfected plants (Table 2).

In our study we found that *U. europaeus* foliar (spine) N concentration was 10% lower in infected plants, irrespective of water or nitrogen treatment. Similar impacts of infection on *U. europaeus* were observed both in the field and in earlier glasshouse studies (Cirocco *et al.*, 2016a, 2018). In contrast, we have also found that *C. pubescens* had no significant impact on foliar N concentration of *U. europaeus*, irrespective of host size, light or nitrogen availability (Cirocco *et al.*, 2016b, 2017, 2020). There is no clear reason for this disparity among results, but the parasite can clearly adversely affect foliar N of this invasive host. In contrast, foliar N of native hosts, such as *A. paradoxa* and *L. myrsinoides*, does not seem to be impacted by *C. pubescens* (Cirocco *et al.*, 2016b, 2017). Irving *et al.* (2019) also found that for the *Phtheirospermum japonicum:Medicago sativa* native association, host N was unaffected by infection with this root hemiparasite, irrespective of nitrogen supply. On the other hand, Jokinen & Irving (2019) found that shoot N concentration of *Trifolium pratense* was significantly lower in plants infected with *O. minor*, regardless of light and nitrogen treatments. In our experiment, parasite removal of N may partly explain the lower foliar concentrations of this nutrient for infected *U. europaeus*, but it is not clear why this is not always observed for this association.

Based on previous studies (Cirocco *et al.*, 2016a, 2018), our hypothesis that parasite impact would be greatest in high water treatments was supported. Parasite impact on shoot biomass of *U. europaeus* was 7% greater in high water relative to low water conditions (Fig. 1e). *C. pubescens* (2016a) similarly found that the significant negative effect of *C. pubescens* on shoot and total biomass of *U. europaeus* was 25% stronger in high water relative to low water treatments. The difference in the magnitude of the high water effect between the two studies may be due to the low water treatments not being low enough in the current study, compared with low water treatments in *C. pubescens* (2016a). The reason why *C. pubescens* has a greater effect on *U. europaeus* in high water conditions may be due to it being more difficult for the parasite to remove host resources in low water conditions. The parasite did have a significantly lower water potential in low water relative to high water, presumably to assist in resource uptake (Fig. 5g). This lower  $\Psi$  may also have led to the parasite lowering its stomatal conductance over the long-term as inferred from parasite  $\delta^{13}\text{C}$  in low water being significantly higher (by 0.5‰) compared with high water (Table 3). *C. pubescens* (2016a) also found  $\delta^{13}\text{C}$  of *C. pubescens* in low water to be significantly higher (by 1.5‰) relative to high water conditions. Also, comparing  $\delta^{13}\text{C}$  of *C. pubescens* across three field sites, this parameter was significantly higher at the driest site (*C. pubescens*, 2018). Nitrogen stress can also lead to decreased stomatal conductance (Chapin, 1991), and this may have occurred for the parasite with significantly higher  $\delta^{13}\text{C}$  in low nitrogen compared with high nitrogen treatments (Table 3). Decreases in parasite stomatal conductance would help ameliorate its water status to some extent, but impede resource uptake from the host. The fact that low water treatments resulted in significantly lower N for the parasite but not for the host (Fig. 6d) implies that the parasite has difficulty in extracting N from the host in low water relative to high water. This explanation does not seem to be confounded by a diluting or concentrating N effect resulting from changes in parasite or host growth, as parasite biomass per gram host dry weight did not change among treatments (Fig. 2b). Thus, all of the above, in terms of compromised parasite water-status, stomatal conductance and N concentration in low water, help explain why its degree of impact on host growth was less severe in these conditions.

Comparing parasite and host sensitivity to water availability,  $\delta^{13}\text{C}$  of *C. pubescens* was significantly higher (by 2‰) than its host, *U. europaeus*, and similar results have been previously reported by our lab for both glasshouse (*C. pubescens*, 2016a, 2020) and field studies (*C. pubescens*, 2018). Thus, it seems that *C. pubescens* is more conservative in its water-use than infected *U.*

*europaeus*. By contrast, mistletoes typically have lower  $\delta^{13}\text{C}$  (less conservative in water-use) relative to their hosts particularly as temperature increases (Bannister & Strong, 2001; Scalon & Wright, 2015). This difference might be due to mistletoes having higher leaf tissue water capacitance than their hosts, thereby enabling higher stomatal conductances and lower water-use efficiencies than the plants they infect (Glatzel, 1983; Davidson *et al.*, 1989). Whereas, *C. pubescens* being a thin vine is likely more vulnerable to desiccation and probably does not share this feature with mistletoes, although confirmation is required. Another possibility explaining higher parasite  $\delta^{13}\text{C}$  relative to the host is that *C. pubescens* is achieving a substantial level of heterotrophy (Cernusak *et al.*, 2004), however, this also requires investigation.

### **Conclusion**

Irrespective of water and nitrogen, the native stem hemiparasite *C. pubescens* had a large and significant impact on growth of *U. europaeus*, one of the world's worst invasive species. We also found that water and nitrogen status of the parasite were higher in well-watered conditions, suggesting that it was better able to extract resources under these conditions and may explain why its impact on host shoot biomass and S:R was greater in high water. Our data continue to support the potential use of some native hemiparasites as biocontrol for major invasive shrubby weeds (Těšitel *et al.*, 2020) and suggest that parasite impact is likely to be greater in higher rainfall areas regardless of soil nitrogen conditions. However, as some regions become drier and warmer in the future (Klausmeyer & Shaw, 2009), we may expect the impact of hemiparasitic plants on host performance to decline.

### **Acknowledgements**

Special thanks to Jason Young (Flinders Analytical) for his expert analysis with the IRMS, Wendy Sullivan for providing access to grinding facilities for sample preparation and Prof. Robert J. Reid for advice on imposing experimental treatments. The Plant Accelerator, Australian Plant Phenomics Facility, is funded by the Australian National Collaborative Research Infrastructure Strategy (NCRIS). This study was part funded by the Australian Department of Agriculture, Water and the Environment (55119480), Primary Industries and Regions SA (56119496) and the Department of Environment, Water and Natural Resources (Adelaide and Mount Lofty Ranges Natural Resources Management Board) (56121016).

## Author contributions

RMC and JMF conceived and designed the experiment. RMC performed the experiment and analysed the data. RMC, JMF and JRW interpreted the analysis and wrote the manuscript.

## References

**Aflakpui GKS, Gregory PJ, Froud-Williams RJ. 1998.** Uptake and partitioning of nitrogen by maize infected with *Striga hermonthica*. *Annals of Botany* **81**: 287–294.

**Aflakpui GKS, Gregory PJ, Froud-Williams RJ. 2002.** Growth and biomass partitioning of maize during vegetative growth in response to *Striga hermonthica* infection and nitrogen supply. *Experimental Agriculture* **38**: 265–276.

**Aflakpui GKS, Gregory PJ, Froud-Williams RJ. 2005.** Carbon ( $^{13}\text{C}$ ) and nitrogen ( $^{15}\text{N}$ ) translocation in a maize-*Striga hermonthica* association. *Experimental Agriculture* **41**: 321–333.

**Bannister P, Strong GL. 2001.** Carbon and nitrogen isotope ratios, nitrogen content and heterotrophy in New Zealand mistletoes. *Oecologia* **126**: 10–20.

**Bardgett RD, Smith RS, Shiel RS, Peacock S, Simkin JM, Quirk H, Hobbs PJ. 2006.** Parasitic plants indirectly regulate below-ground properties in grassland ecosystems. *Nature* **439**: 969–972.

**Bell TL, Adams MA. 2011.** Attack on all fronts: functional relationships between aerial and root parasitic plants and their woody hosts and consequences for ecosystems. *Tree Physiology* **31**: 3–15.

**Bouyoucos GJ. 1929.** A new, simple, and rapid method for determining the moisture equivalent of soils, and the role of soil colloids on this moisture equivalent. *Soil Science* **27**: 233–242.

**Cameron DD, Seel WE. 2007.** Functional anatomy of haustoria formed by *Rhinanthus minor*: linking evidence from histology and isotope tracing. *New Phytologist* **174**: 412–419.

**Cechin I, Press MC. 1993.** Nitrogen relations of the sorghum-*Striga hermonthica* host-parasite association: growth and photosynthesis. *Plant, Cell & Environment* **16**: 237–247.

**Cechin I, Press MC. 1994.** Influence of nitrogen on growth and photosynthesis of a  $\text{C}_3$  cereal, *Oryza sativa*, infected with the root hemiparasite *Striga hermonthica*. *Journal of Experimental Botany* **45**: 925–930.

- Cernusak LA, Pate JS, Farquhar GD. 2004.** Oxygen and carbon isotope composition of parasitic plants and their hosts in southwestern Australia. *Oecologia* **139**: 199–213.
- Chapin III FS. 1991.** Integrated responses of plants to stress: a centralised system of physiological responses. *Bioscience* **41**: 29–36.
- Ciocco RM, Waterman MJ, Robinson SA, Facelli JM, Watling JR. 2015.** Native hemiparasite and light effects on photoprotection and photodamage in a native host. *Functional Plant Biology* **42**: 1168–1178.
- Ciocco RM, Facelli JM, Watling JR. 2016a.** High water availability increases the negative impact of a native hemiparasite on its non-native host. *Journal of Experimental Botany* **67**: 1567–1575.
- Ciocco RM, Facelli JM, Watling JR. 2016b.** Does light influence the relationship between a native stem hemiparasite and a native or introduced host? *Annals of Botany* **117**: 521–531.
- Ciocco RM, Facelli JM, Watling JR. 2017.** Does nitrogen affect the interaction between a native hemiparasite and its native or introduced leguminous hosts? *New Phytologist* **213**: 812–821.
- Ciocco RM, Facelli JM, Watling JR. 2018.** A native parasitic plant affects the performance of an introduced host regardless of environmental variation across field sites. *Functional Plant Biology* **45**: 1128–1137.
- Ciocco RM, Facelli JM, Watling JR. 2020.** The impact of a native hemiparasite on a major invasive shrub is affected by host size at time of infection. *Journal of Experimental Botany* **71**: 140.
- Davidson NJ, True KC, Pate JS. 1989.** Water relations of the parasite: host relationship between the mistletoe *Amyema linophyllum* (Fenzl) Tieghem and *Casuarina obesa* Miq. *Oecologia* **80**: 321–330.
- Evans B, Borowicz V. 2013.** *Verbesina alternifolia* tolerance to the holoparasite *Cuscuta gronovii* and the impact of drought. *Plants* **2**: 635–649.
- Evans BA, Borowicz VA. 2015.** The plant vigor hypothesis applies to a holoparasitic plant on a drought-stressed host. *Botany* **93**: 685–689.
- Facelli E, Wynn N, Tsang HT, Watling JR, Facelli JM. 2020.** Defense responses of native and invasive plants to the native generalist vine parasite *Cassytha pubescens* – anatomical and functional studies. *Australian Journal of Botany*. doi.org/10.1071/BT19136



- Gao F-L, Che X-X, Yu F-H, Li J-M. 2019.** Cascading effects of nitrogen, rhizobia and parasitism via a host plant. *Flora* **251**: 62–67.
- Gibson CC, Watkinson AR. 1991.** Host selectivity and the mediation of competition by the root hemiparasite *Rhinanthus minor*. *Oecologia* **86**: 81–87.
- Glatzel G. 1983.** Mineral nutrition and water relations of hemiparasitic mistletoes: a question of partitioning. Experiments with *Loranthus europaeus* on *Quercus petraea* and *Quercus robur*. *Oecologia* **56**: 193–201.
- Gurney AL, Press MC, Ransom JK. 1995.** The parasitic angiosperm *Striga hermonthica* can reduce photosynthesis of its sorghum and maize hosts in the field. *Journal of Experimental Botany* **46**: 1817–1823.
- Hartley SE, Green JP, Massey FP, Press MCP, Stewart AJA, John EA. 2015.** Hemiparasitic plant impacts animal and plant communities across four trophic levels. *Ecology* **96**: 2408–2416.
- Heide-Jørgensen HS. 2013.** Introduction: the parasitic syndrome in higher plants. In: Joel D, Gressel J, Musselman L, eds. *Parasitic Orobanchaceae: parasitic mechanisms and control strategies*. Berlin, Germany: Springer-Verlag, 1–14.
- Hornoy B, Atlan A, Roussel V, Buckley YM, Tarayre M. 2013.** Two colonisation stages generate two different patterns of genetic diversity within native and invasive ranges of *Ulex europaeus*. *Heredity* **111**: 355–363.
- Irving LJ, Kim D, Schwier N, Vaughan JKE, Ong G, Hama T. 2019.** Host nutrient supply affects the interaction between the hemiparasite *Phtheirospermum japonicum* and its host *Medicago sativa*. *Environmental and Experimental Botany* **162**: 125–132.
- Jeschke WD, Hilpert A. 1997.** Sink-stimulated photosynthesis and sink-dependent increase in nitrate uptake: nitrogen and carbon relations of the parasitic association *Cuscuta reflexa*–*Ricinus communis*. *Plant, Cell & Environment* **20**: 47–56.
- Jeschke WD, Baig A, Hilpert A. 1997.** Sink-stimulated photosynthesis, increased transpiration and increased demand-dependent stimulation of nitrate uptake: nitrogen and carbon relations in the parasitic association *Cuscuta reflexa*–*Coleus blumei*. *Journal of Experimental Botany* **48**: 915–925.
- Jokinen JI, Irving LJ. 2019.** Effects of light level and nitrogen supply on the red clover–*Orobanche minor* host-parasite interaction. *Plants* **8**: 146.

**Klausmeyer KR, Shaw MR. 2009.** Climate change, habitat loss, protected areas and the climate adaptation potential of species in Mediterranean ecosystems worldwide. *PloS One* **4**: e6392.

**Kokubugata G, Nakamura K, Forster PI, Wilson GW, Holland AE, Hirayama Y, Yokota M. 2012.** *Cassytha pubescens* and *C. glabella* (Lauraceae) are not disjunctly distributed between Australia and the Ryukyu Archipelago of Japan—evidence from morphological and molecular data. *Australian Systematic Botany* **25**: 364–373.

**Korell L, Sandner TM, Matthies D, Ludewig K. 2019.** Effects of drought and N level on the interactions of the root hemiparasite *Rhinanthus alectorolophus* with a combination of three host species. *Plant Biology* **22**: 84–92.

**Le QV, Tennakoon KU, Metali F, Lim LB, Bolin JF. 2015.** Impact of *Cuscuta australis* infection on the photosynthesis of the invasive host, *Mikania micrantha*, under drought condition. *Weed Biology and Management* **15**: 138–146.

**Lowe S, Browne M, Boudjelas S, De Poorter M. 2000.** *100 of the world's worst invasive alien species: a selection from the global invasive species database*. Auckland, New Zealand: IUCN/SSC Invasive Species Specialist Group (ISSG).

**Matthies D, Egli P. 1999.** Response of a root hemiparasite to elevated CO<sub>2</sub> depends on host type and soil nutrients. *Oecologia* **120**: 156–161.

**McLuckie J. 1924.** Studies in Parasitism. I. A contribution to the physiology of the genus *Cassytha*, Part 1. *Proceedings of the Linnaen Society of New South Wales* **49**: 333–369.

**Miller AC, Watling JR, Overton IC, Sinclair R. 2003.** Does water status of *Eucalyptus largiflorens* (Myrtaceae) affect infection by the mistletoe *Amyema miquelii* (Loranthaceae)? *Functional Plant Biology* **30**: 1239–1247.

**Parsons WT, Cuthbertson EG. 2001.** *Noxious Weeds of Australia* (second edition). Victoria, Australia: CSIRO Publishing.

**Press MC, Graves JD. 1995.** *Parasitic plants*. London, UK: Chapman & Hall.

**Press MC, Phoenix GK. 2005.** Impacts of parasitic plants on natural communities. *New Phytologist* **166**: 737–751.

**Prider JN, Watling JR, Facelli JM. 2009.** Impacts of a native parasitic plant on an introduced and a native host species: implications for the control of an invasive weed. *Annals of Botany* **103**: 107–115.

**Quasted HM. 2008.** Parasitic plants—impacts on nutrient cycling. *Plant and Soil* **311**: 269–272.

- Rodríguez-Echeverría S. 2010.** Rhizobial hitchhikers from Down Under: invasional meltdown in a plant–bacteria mutualism? *Journal of Biogeography* **37**: 1611–1622.
- Sala OE, Chapin III FS, Armesto JJ, Berlow E, Bloomfield J, Dirzo R, Huber-Sanwald E, Huenneke LF, Jackson RB, Kinzig A et al. 2000.** Global biodiversity scenarios for the year 2100. *Science* **287**: 1770–1774.
- Scalon MC, Wright IJ. 2015.** A global analysis of water and nitrogen relationships between mistletoes and their hosts: broad-scale tests of old and enduring hypotheses. *Functional Ecology* **29**: 1114–1124.
- Shen H, Prider JN, Facelli JM, Watling JR. 2010.** The influence of the hemiparasitic angiosperm *Cassytha pubescens* on photosynthesis of its host *Cytisus scoparius*. *Functional Plant Biology* **37**: 14–21.
- Shen H, Xu S-J, Hong L, Wang Z-M, Ye W-H. 2013.** Growth but not photosynthesis response of a host plant to infection by a holoparasitic plant depends on nitrogen supply. *PloS One* **8**: e75555.
- Sinebo W, Drennan DSH. 2001.** Vegetative growth of sorghum and *Striga hermonthica* in response to nitrogen and the degree of host root infection. *European Journal of Plant Pathology* **107**: 849–860.
- Sui X-L, Zhang T, Tian Y-Q, Xue R-J, Li A-R. 2019.** A neglected alliance in battles against parasitic plants: arbuscular mycorrhizal and rhizobial symbioses alleviate damage to a legume host by root hemiparasitic *Pedicularis* species. *New Phytologist* **221**: 470–481.
- Těšitel J, Těšitelová T, Fisher JP, Lepš J, Cameron DD. 2015.** Integrating ecology and physiology of root-hemiparasitic interaction: interactive effects of abiotic resources shape the interplay between parasitism and autotrophy. *New Phytologist* **205**: 350–360.
- Těšitel J, Mládek J, Horník J, Těšitelová T, Adamec V, Tichy L. 2017.** Suppressing competitive dominants and community restoration with native parasitic plants using the hemiparasitic *Rhinanthus alectorolophus* and the dominant grass *Calamagrostis epigejos*. *Journal of Applied Ecology* **54**: 1487–1495.
- Těšitel J, Cirocco RM, Facelli JM, Watling JR. 2020.** Native parasitic plants: biological control for plant invasions? *Applied Vegetation Science* **23**: 464–469.
- Watling JR, Press MC. 2001.** Impacts of infection by parasitic angiosperms on host photosynthesis. *Plant Biology* **3**: 244–250.

**Weber JZ. 1981.** A taxonomic revision of *Cassytha* (Lauraceae) in Australia. *Journal of the Adelaide Botanic Garden* **3**: 187–262.

**Weir BS, Turner SJ, Silvester WB, Park D-C, Young JM. 2004.** Unexpectedly diverse *Mesorhizobium* strains and *Rhizobium leguminosarum* nodulate native legume genera of New Zealand, while introduced legume weeds are nodulated by *Bradyrhizobium* species. *Applied and Environmental Microbiology* **70**: 5980–5987.

### Supporting Information

Additional supporting information may be found in the online version of this article.

**Fig. S1** Photos of first and second phases of experiment.

**Table S1** *F* and sum of squares for host growth parameters.

**Table S2** *F* and sum of squares for all parasite parameters.

**Table S3** *F* and sum of squares for host physiological measures.

### Figure legends

**Fig. 1** (a) Total, (c) shoot and (e) root biomasses of *Ulex europaeus* either uninfected (white bars) or infected with *Cassytha pubescens* (light grey bars) growing in high (H) or low (L) water (W) and nitrogen (N) conditions. (b) Independent effect of infection on total biomass. (d) Infection × water effect on shoot biomass and (f) independent effect of infection on root biomass. Data are means ± 1SE; different letters signify significant differences; (a, c, e)  $n = 6$ , (b, f)  $n = 24$  and (d)  $n = 12$ .

**Fig. 2** (a) Parasite biomass and (b) parasite biomass  $g^{-1}$  host total biomass of *Cassytha pubescens* when infecting *Ulex europaeus* growing in high (H) or low (L) water (W) and nitrogen (N) conditions. Data are means ± 1SE and  $n = 6$ .

**Fig. 3** (a) Predawn and (c) midday quantum yield and (e) maximum electron transport rate of *Ulex europaeus* either uninfected (white bars) or infected with *Cassytha pubescens* (light grey bars)

growing in high (H) or low (L) water (W) and nitrogen (N) conditions. (b), (d) and (f) Independent effect of infection on all three host parameters. Data are means  $\pm$  1SE; different letters signify significant differences; (a, c, e)  $n = 7-8$ , (b)  $n = 30-32$  and (d, f)  $n = 31-32$ .

**Fig. 4** (a) Predawn and (b) midday quantum yield, (c) maximum electron transport rate and (d) carbon isotope composition of *Cassitha pubescens* when infecting *Ulex europaeus* growing in high (H) or low (L) water (W) and nitrogen (N) conditions. Data are means  $\pm$  1SE and (a, b, c)  $n = 7-8$  and (d)  $n = 6$ .

**Fig. 5** (a) Midday shoot water potential ( $\Psi$ ) of *Ulex europaeus* either uninfected (white bars) or infected with *Cassitha pubescens* (light grey bars) growing in high (H) or low (L) water (W) and nitrogen (N) conditions. (b), (c) and (d) Independent effect of infection, water (dotted bars) and nitrogen (hatched bars) on host  $\Psi$ , respectively. (e)  $\Psi$  of *U. europaeus* infected with *C. pubescens* relative to that of the parasite (independent species effect: i.e. water and nitrogen plants pooled for each species). (f)  $\Psi$  of parasite growing in all treatment combinations. (g) Independent effect of water on parasite  $\Psi$ . Data are means  $\pm$  1SE; different letters signify significant differences; (a)  $n = 7-8$ , (b, c, d)  $n = 31-32$ , (e)  $n = 31$ , (f)  $n = 7-8$  and (g)  $n = 15-16$ .

**Fig. 6** (a) Foliar nitrogen concentration (N) of *Ulex europaeus* either uninfected (white bars) or infected with *Cassitha pubescens* (light grey bars) growing in high (H) or low (L) water (W) and nitrogen (N) conditions. (b) Independent effect of infection on host N. (c) N of parasite growing in all treatments. (d) Independent effect of water on parasite N. Data are means  $\pm$  1SE; different letters signify significant differences; (a, c)  $n = 6$ , (b)  $n = 24$  and (d)  $n = 12$ .

**Table 1** Three-way ANOVA results (*P*-values) for the effects of infection with *Cassiope* *pubescens* (I), water (W) and nitrogen supply (N) on total, shoot and root biomass, shoot:root ratio (S:R), nodule biomass (Nod) and nodule biomass g<sup>-1</sup> root biomass (Nod g<sup>-1</sup> root) of *Ulex europaeus*.

	<b>Total</b>	<b>Shoot</b>	<b>Root</b>	<b>S:R</b>	<b>Nod</b>	<b>Nod g<sup>-1</sup> root</b>
I	<b>&lt;0.0001</b>	<b>&lt;0.0001</b>	<b>&lt;0.0001</b>	<b>0.016</b>	<b>&lt;0.0001</b>	<b>&lt;0.0001</b>
W	<b>0.022</b>	<b>0.006</b>	0.349	0.127	0.154	0.577
I × W	0.126	<b>0.037</b>	0.569	<b>0.033</b>	0.335	0.541
N	0.417	0.316	0.930	0.151	<b>0.021</b>	<b>0.006</b>
I × N	0.955	0.927	0.746	0.190	0.510	0.996
W × N	0.936	0.911	0.922	0.843	0.507	0.721
I × W × N	0.859	0.740	0.746	0.292	0.407	0.585
Block	0.743	0.776	0.834	0.096	0.932	0.118

Significant effects are in bold; *F* and sum of square values are presented in Supporting Information Table S1 and root biomass and Nod g<sup>-1</sup> root data log transformed to achieve homoscedasticity.

**Table 2** Shoot:root ratio (S:R), nodule biomass (Nod; g dry weight), nodule biomass g<sup>-1</sup> root biomass (Nod g<sup>-1</sup> root) and carbon isotope composition ( $\delta^{13}\text{C}$  ; ‰) of *Ulex europaeus* either uninfected (minus) or infected (plus) with *Cassiope pubescens* growing in high (HW) or low water (LW) conditions and supplied (HN) or not supplied (LN) with nitrogen.

<b>Treatment</b>	<b>S:R</b>	<b>Nod</b>	<b>Nod g<sup>-1</sup> root</b>	<b><math>\delta^{13}\text{C}</math></b>
HW/HN-	3.63 ± 0.26	0.763 ± 0.076	0.036 ± 0.005	-31.4 ± 0.23
HW/HN+	2.79 ± 0.27	0.573 ± 0.089	0.074 ± 0.011	-30.8 ± 0.33
HW/LN-	3.44 ± 0.31	1.05 ± 0.10	0.049 ± 0.005	-31.2 ± 0.18
HW/LN+	2.50 ± 0.26	0.674 ± 0.085	0.091 ± 0.009	-29.9 ± 0.36
LW/HN-	2.76 ± 0.10	0.828 ± 0.080	0.038 ± 0.003	-31.3 ± 0.23

LW/HN+	3.15 ± 0.26	0.408 ± 0.058	0.066 ± 0.009	-30.1 ± 0.35
LW/LN-	2.89 ± 0.33	0.928 ± 0.125	0.045 ± 0.004	-30.1 ± 0.58
LW/LN+	2.39 ± 0.38	0.531 ± 0.050	0.081 ± 0.011	-29.9 ± 0.23
Infection × Water				
HW-	3.53 ± 0.19 <b>a</b>	N/A	N/A	N/A
HW+	2.65 ± 0.19 <b>b</b>	N/A	N/A	N/A
LW-	2.83 ± 0.16 <b>ab</b>	N/A	N/A	N/A
LW+	2.77 ± 0.25 <b>b</b>	N/A	N/A	N/A
Infection effect				
Uninfected	3.18 ± 0.15	0.893 ± 0.051 <b>a</b>	0.042 ± 0.002 <b>a</b>	-31.0 ± 0.19 <b>a</b>
Infected	2.71 ± 0.15	0.547 ± 0.039 <b>b</b>	0.079 ± 0.005 <b>b</b>	-30.2 ± 0.17 <b>b</b>
Water effect				
HW	3.09 ± 0.16	0.766 ± 0.055	0.062 ± 0.006	-30.8 ± 0.18 <b>a</b>
LW	2.80 ± 0.15	0.674 ± 0.059	0.058 ± 0.005	-30.4 ± 0.21 <b>b</b>
Nitrogen effect				
HN	3.08 ± 0.13	0.643 ± 0.050 <b>a</b>	0.053 ± 0.005 <b>a</b>	-30.9 ± 0.17 <b>a</b>
LN	2.81 ± 0.17	0.797 ± 0.061 <b>b</b>	0.067 ± 0.006 <b>b</b>	-30.3 ± 0.20 <b>b</b>

No infection × water × nitrogen interaction for all parameters; significant infection × water effect only for S:R; significant independent effects of either infection or nitrogen on Nod, Nod g<sup>-1</sup> root and δ<sup>13</sup>C; significant independent effect of water on δ<sup>13</sup>C. Different letters in bold signify significant differences (vertically), data are means ± 1SE (N/A = data are not applicable); for each treatment: *n* = 6, infection × water effect: *n* = 12 and for all independent effects: *n* = 24.

**Table 3** Two-way ANOVA results (*P*-values) for effects of water (W) and nitrogen (N) supply on parasite biomass, parasite biomass g<sup>-1</sup> host total biomass, predawn and midday quantum yield ( $F_v/F_m$  and  $\Phi_{PSII}$ ), maximum electron transport rates ( $ETR_{max}$ ), midday water potential (MD  $\Psi$ ), carbon isotope composition (δ<sup>13</sup>C) and stem nitrogen concentration [N] of *Cassitha pubescens* infecting *Ulex europaeus*.

Parasite biomass	Parasite biomass g <sup>-1</sup> host	$F_v/F_m$	$\Phi_{PSII}$	$ETR_{max}$	MD $\Psi$	δ <sup>13</sup> C	[N]
------------------	---------------------------------------	-----------	---------------	-------------	-----------	-------------------	-----

W	0.411	0.516	0.971	0.742	0.619	<b>0.019</b>	<b>0.050</b>	<b>0.027</b>
N	0.172	0.748	0.452	0.691	0.846	0.330	<b>0.002</b>	0.658
W × N	0.419	0.398	0.113	0.657	0.581	0.518	0.154	0.681
Block	0.386	0.989	<b>0.032</b>	0.235	0.298	<b>0.008</b>	<b>0.037</b>	0.553

Significant effects are in bold;  $F$  and sum of square values are presented in Supporting Information Table S2.

**Table 4** Three-way ANOVA results ( $P$ -values) for the effects of infection with *Cassytha pubescens* (I), water (W) and nitrogen supply (N) on predawn and midday quantum yield ( $F_v/F_m$  and  $\Phi_{PSII}$ ), maximum electron transport rates ( $ETR_{max}$ ), midday water potential (MD  $\Psi$ ), carbon isotope composition  $\delta^{13}C$  and foliar nitrogen concentration [N] of *Ulex europaeus*.

	$F_v/F_m$	$\Phi_{PSII}$	$ETR_{max}$	MD $\Psi$	$\delta^{13}C$	[N]
I	<b>&lt;0.0001</b>	<b>&lt;0.0001</b>	<b>&lt;0.0001</b>	<b>0.0001</b>	<b>0.0009</b>	<b>0.002</b>
W	0.388	0.222	0.139	<b>0.003</b>	<b>0.046</b>	0.837



I × W	0.407	0.991	0.661	0.537	0.581	0.238
N	0.649	0.495	0.903	<b>0.003</b>	<b>0.011</b>	0.918
I × N	0.446	0.479	0.274	0.308	0.683	0.702
W × N	0.978	0.838	0.750	0.471	0.871	0.372
I × W × N	0.737	0.402	0.211	0.627	0.098	0.380
Block	0.537	0.361	0.332	0.203	0.135	0.599

---

Significant effects are in bold; *F* and sum of square values are presented in Supporting Information Table S3.











