



Manchester Metropolitan University

Leal, Cecília G and Lennox, Gareth D and Ferraz, Silvio FB and Ferreira, Joice and Gardner, Toby A and Thomson, James R and Berenguer, Erika and Lees, Alexander and Hughes, Robert M and Mac Nally, Ralph and Aragão, Luiz EOC and Brito, Janaina G de and Castello, Leandro and Garrett, Rachael D and Hamada, Neusa and Juen, Leandro and Leitão, Rafael P and Louzada, Julio and Morello, Thiago F and Moura, Nárgila G and Nessimian, Jorge L and Oliveira-Junior, José Max B and Oliveira, Victor Hugo F and Oliveira, Vívian C de and Parry, Luke and Pompeu, Paulo S and Solar, Ricardo RC and Zuanon, Jansen and Barlow, Jos (2020) Integrated terrestrial-freshwater planning doubles conservation of tropical aquatic species. *Science*, 370 (6512). pp. 117-121. ISSN 0036-8075

Downloaded from: <http://e-space.mmu.ac.uk/626622/>

Version: Accepted Version

Publisher: AAS

DOI: <https://doi.org/10.1126/science.aba7580>

Please cite the published version

<https://e-space.mmu.ac.uk>

Title: Integrated terrestrial-freshwater planning doubles conservation of tropical aquatic species

Authors: Cecília G. Leal^{1,2*§}, Gareth D. Lennox^{3*§}, Silvio F. B. Ferraz¹, Joice Ferreira⁴, Toby A. Gardner⁵, James R. Thomson⁶, Erika Berenguer^{3,7}, Alexander C. Lees^{8,9}, Robert M. Hughes^{10,11}, Ralph Mac Nally¹², Luiz E. O. C. Aragão^{13,14}, Janaina G. de Brito¹⁵, Leandro Castello¹⁶, Rachael D. Garrett¹⁷, Neusa Hamada¹⁸, Leandro Juen¹⁹, Rafael P. Leitão²⁰, Julio Louzada², Thiago F. Morello²¹, Nárgila G. Moura²², Jorge L. Nessimian²³, José Max B. Oliveira-Junior²⁴, Victor H. F. de Oliveira², Vivian C. de Oliveira¹⁸, Luke Parry³, Paulo S. Pompeu², Ricardo R. C. Solar²⁰, Jansen Zuanon¹⁸, Jos Barlow^{2,3}

Affiliations:

1. Luiz de Queiroz College of Agriculture, University of São Paulo, CEP 13418-900, Piracicaba, SP, Brazil
2. Departamento de Ecologia e Conservação, Universidade Federal de Lavras, CEP 37200-900, Lavras, MG, Brazil
3. Lancaster Environment Centre, Lancaster University, Lancaster, UK
4. EMBRAPA Amazônia Oriental, CEP 66095-100, Belém, Pará, Brazil
5. Stockholm Environment Institute, Linegatan 87D, 11523, Stockholm, Sweden
6. Department of Environment, Land, Water and Planning, Arthur Rylah Institute for Environmental Research, Heidelberg, Vic, Australia
7. Environmental Change Institute, University of Oxford, Oxford, UK
8. Department of Natural Sciences, Manchester Metropolitan University, Manchester, M1 5GD, UK
9. Cornell Lab of Ornithology, Cornell University, Ithaca, New York, USA
10. Amnis Opes Institute, Corvallis, Oregon, USA
11. Department of Fisheries & Wildlife, Oregon State University, Corvallis, Oregon, USA
12. School of BioSciences, The University of Melbourne, Parkville 3052, VIC, Australia
13. Tropical Ecosystems and Environmental Sciences Group (TREES), Remote Sensing Division, National Institute for Space Research - INPE, Avenida dos Astronautas, São José dos Campos, SP, Brazil
14. College of Life and Environmental Sciences, University of Exeter, Exeter, UK
15. Escola Estadual Maria Miranda Araújo, Secretaria de Educação do Estado de Mato Grosso, Av. Aeroporto, s/n, CEP 78336-000, Colniza, MT, Brazil.
16. Department of Fish and Wildlife Conservation, Virginia Polytechnic Institute and State University, Blacksburg, VA, USA
17. Environmental Policy Group, Departments of Environmental System Science and Humanities, Social, and Political Science, ETH Zürich, 8092 Zürich, Switzerland

18. Coordenação de Biodiversidade, Instituto Nacional de Pesquisas da Amazônia, Avenida André Araújo, 2.936, Petrópolis, CEP 69067-375, Manaus, AM, Brazil
19. Laboratório de Ecologia e Conservação, Instituto de Ciências Biológicas, Universidade Federal do Pará, Rua Augusto Correia, No. 1, Bairro Guamá, CEP 66075-110, Belém, PA, Brazil
20. Departamento de Genética, Ecologia e Evolução, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais, Avenida Antônio Carlos 6627, CP 486, CEP 31270-901, Belo Horizonte, MG, Brazil
21. Universidade Federal do ABC, São Bernardo do Campo, SP, Brazil
22. Museu Paraense Emílio Goeldi, Belém, PA, Brazil
23. Departamento de Zoologia, Instituto de Biologia, Universidade Federal do Rio de Janeiro, Av. Carlos Chagas Filho 373, CEP 21941-590, Rio de Janeiro, RJ, Brazil
24. Instituto de Ciências e Tecnologia das Águas, Universidade Federal do Oeste do Pará, Rua Vera Paz, s/n (Unidade Tapajós), Bairro Salé, CEP 68040-255, Santarém, PA, Brazil

* These authors contributed equally

§ Corresponding authors. Email: c.gontijoleal@gmail.com

1 **Abstract:** Conservation initiatives overwhelmingly focus on terrestrial biodiversity and little is
2 known about the freshwater co-benefits of terrestrial conservation actions. We sampled >1,500
3 terrestrial and freshwater species in the Amazon and simulated conservation for species from both
4 realms. Prioritizations based on terrestrial species yielded on average just 22% of the freshwater
5 benefits achieved through freshwater-focused conservation. However, using integrated cross-realm
6 planning, freshwater benefits could be increased by up to 600% for a 1% reduction in terrestrial
7 benefits. Where freshwater biodiversity data are unavailable but aquatic connectivity is accounted for,
8 freshwater benefits could still be doubled for negligible losses of terrestrial coverage. Conservation
9 actions are urgently needed to improve the status of freshwater species globally. Our results suggest
10 such gains can be achieved without compromising terrestrial conservation goals.

11 **One Sentence Summary:** Integrated conservation planning increases freshwater species protection
12 by up to 600% without compromising terrestrial conservation.

13 Main Text

14 Freshwater ecosystems occupy less than 1% of the Earth's surface, make up only 0.01% of all water,
15 yet host c. 10% of all known species, including a third of all vertebrates (1). They also deliver vital
16 ecosystem services, such as climate regulation and the provision of food, fuel and fiber (2).

17 Nevertheless, freshwater ecosystems are far more imperilled than their terrestrial or marine
18 counterparts; since 1970, for example, populations of freshwater vertebrates have declined by 83%
19 compared to a c. 40% decline of terrestrial and marine vertebrates (3,4). A range of threats have long
20 been linked to this collapse in freshwater biodiversity, including habitat loss and degradation,
21 overexploitation, eutrophication, flow modification, and the introduction of non-native species (5).
22 These are now amplified by emerging stressors, including climate change and contamination from
23 microplastics and biochemicals (3).

24 Despite the freshwater biodiversity crisis (6), freshwater species are rarely considered in broad-scale
25 conservation strategies (7-9). Although distributions of terrestrial and freshwater vertebrates display a
26 degree of spatial congruence (10), there are three key reasons why freshwater conservation based on
27 terrestrial priorities cannot be taken for granted. First, studies that reveal terrestrial-freshwater
28 congruence rely on coarse-grained data, and such congruence might not occur at local scales where
29 conservation decisions are implemented. Second, assessments of the distribution of freshwater biota
30 are often restricted to small scales or specific taxonomic groups (11). Third, and most importantly,
31 terrestrial prioritizations do not account for aquatic connectivity, which strongly affects the
32 distribution of freshwater species, facilitates nutrient flows and mediates the cumulative effects of
33 stressors along watercourses (12-15). Given these limitations, there is an urgent need to understand the
34 extent to which freshwater biodiversity can benefit from terrestrial conservation actions, and whether
35 freshwater protection can be increased through integrated planning for both realms. This is
36 particularly critical in tropical regions, which harbor >80% of the world's freshwater fish and are
37 undergoing the most rapid land-use changes on Earth (16).

38 Here, we addressed these knowledge gaps using data from extensive terrestrial and freshwater
39 biodiversity surveys in two biogeographically distinct regions of Brazilian Amazonia: Paragominas
40 and Santarém (Fig. S1; 17). With >40% of their forests having been converted to agricultural land-
41 uses, these regions typify the agricultural-forest frontier in the Amazon (18). In terrestrial sites ($n =$
42 377; Fig. S2), we sampled plants ($n = 812$ species), birds ($n = 327$ species), and dung beetles ($n = 141$
43 species). In freshwater sites ($n = 99$ streams; Fig. S3), we sampled fish ($n = 143$ species); Odonata
44 (i.e. dragonflies and damselflies; $n = 134$ species); and Ephemeroptera (mayflies), Plecoptera
45 (stoneflies) and Trichoptera (caddisflies; hereafter, "EPT"), which are frequently used as a measure of

46 freshwater ecosystem health (19). We could identify EPT individuals only to genus level ($n = 59$
47 genera; 17). All taxa are referred to as “species” hereafter.

48 Using these data, we first investigated the extent to which one species group (e.g. fish) is protected
49 under conservation strategies directed at another species group (e.g. plants), which we refer to as
50 “incidental conservation”. To do so, we built regional species distribution maps with an array of
51 biophysical predictors (Table S1; 17). We then used the distribution maps and the Zonation
52 conservation planning framework (20) to simulate terrestrial and freshwater conservation at the
53 catchment scale, a natural landscape unit that integrates hydrological processes. Zonation selects
54 catchments that maximize the weighted average proportion of species distributions under conservation
55 while accounting for species complementarity, and we use this as our conservation benefit function
56 (17). For the freshwater analyses, we used the *directed-connectivity* algorithm, which produces
57 aquatically connected conservation networks appropriate for freshwater species (21). To focus on
58 biodiversity (i.e. without socio-economic considerations), we first ran the optimization analyses
59 constrained by the proportion of the landscape that could be conserved. We then tested the robustness
60 of these findings to budget-constrained analyses by incorporating two region-specific estimates of
61 agricultural opportunity costs (Fig. S4; 17). Finally, we undertook sensitivity analyses by varying
62 available conservation resources. We report results for the area-constrained analysis in which 20% of
63 landscape could be conserved, which aligns with the Aichi target to conserve at least 17% of
64 terrestrial and inland water areas (4). For an overview of all analyses, see Fig. S1.

65 Terrestrially focused conservation planning provided limited incidental conservation benefits for
66 freshwater species (Fig. 1). Among taxa and regions, on average just 22% (range: 14-29%) of the
67 freshwater benefits achieved through freshwater conservation were secured through terrestrial
68 conservation. In contrast, freshwater species prioritisations achieved on average 84% (range: 70-96%)
69 of the terrestrial benefits achieved through terrestrial prioritisations. Within both freshwater and
70 terrestrial realms, prioritizing for any one taxonomic group provided >92% of the maximum
71 achievable benefits to other groups in the same realm. These results were similar whether the
72 optimisations were constrained by area or financial budgets (Fig. 1A-C).

73 Differences in the incidental conservation outcomes can be explained by (i) the correlations in
74 catchment priority rankings among species groups (Figs. S5 & S6) and (ii) the spatial distribution of
75 conservation priorities (Fig. 2, S7 & S8). Terrestrial and freshwater groups act as good surrogates for,
76 respectively, other terrestrial and freshwater groups because of the strong correlation in catchment
77 priority rankings: a catchment with high marginal conservation value for one terrestrial group is likely
78 to be of high marginal conservation value for other terrestrial groups, and the same holds for
79 freshwater taxa. Catchment priority ranking correlations were somewhat weaker between terrestrial

80 and freshwater groups, leading to smaller but nonetheless high incidental terrestrial benefits when
81 focused on freshwater species. However, the failure to incorporate aquatic connectivity into terrestrial
82 planning produced conservation network designs that were inadequate for freshwater species (Figs. 2,
83 S7 & S8), resulting in poor freshwater outcomes from terrestrial planning.

84 Next, we considered the extent to which freshwater benefits could be increased through conservation
85 planning mechanisms targeted at both terrestrial and freshwater species. To do so, we developed two
86 integrated planning techniques (17). Our first approach utilised both terrestrial and freshwater
87 biodiversity data to determine a prioritisation optimized for species from both realms (hereafter, “joint
88 planning”). Given the general paucity of freshwater biodiversity data, our second approach
89 incorporated aquatic connectivity into the terrestrial optimizations to account for freshwater species
90 habitat requirements (hereafter, “terrestrial-plus-connectivity”). Using these approaches, we
91 undertook two trade-off analyses. We first determined the increase in freshwater benefits that could be
92 achieved for a given reduction in terrestrial benefits from their optimum. We focus on this trade-off
93 analysis in the main text. We also considered the increase in freshwater benefits for a given resource
94 increase (e.g. increase in landscape covered or financial budgets) while maintaining terrestrial benefits
95 at their optimum. As above, we focused on area-constrained optimizations in which 20% of a
96 landscape could be conserved.

97 Using the joint planning approach, freshwater benefits could be increased by on average 62% and
98 345% in Paragominas and Santarém, respectively, for a negligible 1% reduction in terrestrial benefits
99 relative to their optimum (Fig. 3). A 5% reduction in terrestrial benefits, on the other hand, resulted in
100 an average increase in freshwater benefits of 184% in Paragominas and 365% in Santarém. The
101 terrestrial-plus-connectivity approach generally produced lower freshwater conservation gains.
102 Nonetheless, a 1% and 5% reduction in terrestrial benefits increased freshwater benefits by 75-100%
103 and 130-175% in both Paragominas and Santarém. Alternatively, the freshwater gains we document
104 for a 1% and 5% reduction in terrestrial benefits could be achieved without any terrestrial losses for,
105 respectively, a <1% and <5% increase in conservation resources (Fig. S9). Trade-offs were
106 qualitatively similar with the incorporation of opportunity costs (Fig. 3) and more and less
107 pronounced for, respectively, lower and higher conservation resource levels (Fig. S10).

108 While the freshwater gains we found for negligible reductions in terrestrial protection were substantial
109 in both Paragominas and Santarém, there were large regional differences when using the joint
110 planning approach that incorporates both terrestrial and freshwater biodiversity data (Fig. 3). These
111 differences arise from variation in the spatial overlap of conservation priorities between regions. In
112 Santarém, many of the highest priority catchments for terrestrial and freshwater groups were in the
113 south-west (where the Tapajós National Forest is located; Fig. 2). In Paragominas, the same spatial

114 overlap in priorities was not apparent (Fig. 2). Thus, in Paragominas, substantial deviation from the
115 optimal catchment prioritization for terrestrial species was required to achieve the largest increases in
116 freshwater benefits. In Santarém, by contrast, large freshwater gains were possible simply by selecting
117 catchments in the region of high conservation value for both realms that produced the requisite
118 aquatic connectivity. Therefore, the realized magnitude of the freshwater gains possible from
119 integrated planning will depend on the underlying spatial covariance in species distributions, which
120 determines the spatial overlap in conservation priorities.

121 These results provide compelling evidence that the protection of freshwater species can be vastly
122 improved without undermining terrestrial conservation goals. However, there are factors for which we
123 did not account that could lead to significantly different terrestrial-freshwater trade-offs than we
124 found. First, we did not incorporate the many additional socio-ecological benefits of freshwater
125 conservation, meaning our results are likely to be conservative. For example, in addition to the direct
126 provisioning, supporting, regulating and cultural services freshwater ecosystems provide (2), by
127 enhancing landscape connectivity, freshwater conservation can also promote movement of terrestrial
128 species, recolonization of defaunated areas, and seed dispersal and pollination services (22). Second
129 and conversely, where freshwater conservation imposes external opportunity costs beyond a loss of
130 agricultural profits, by, for example, precluding the development of hydropower or imposing water-
131 use restrictions in the surrounding landscape, the overall scope for conservation investment may be
132 reduced, leading to fewer net benefits from integrated planning. The manifestation of these additional
133 socio-ecological trade-offs that emerge when protecting freshwater ecosystems is likely to be highly
134 dependent on local circumstances, but their consideration will be essential for designing effective and
135 sustainable conservation projects. Finally, our optimization analyses were static. As freshwater
136 biodiversity data were collected in different years in Paragominas (2011) and Santarém (2010), and as
137 the regions experienced significantly different climatic conditions during this time (17), some of the
138 observed regional differences in trade-offs could result from temporal variation. Understanding and
139 incorporating environmentally mediated changes in species distributions will be important for
140 estimating the long-term benefits of integrated terrestrial-freshwater planning.

141 Identifying promising new approaches for biodiversity conservation is only the first step towards
142 improving conservation outcomes. Given that evidence is lacking for the translation of systematic
143 conservation planning exercises into tangible benefits (23), how best to turn our findings into
144 meaningful action? First, while previous global conservation agendas – such as the UN’s Sustainable
145 Development Goals and the Convention on Biological Diversity – have recognized the need to
146 conserve both terrestrial and freshwater ecosystems (SDG 15, Aichi target 11), recognition of their
147 interdependence remains largely absent from conservation planning. As the world prepares to
148 consider new, post-2020 conservation targets (9,24), we show that a truly integrated approach to

149 conservation on land, which accounts for trade-offs and harnesses synergies among ecosystems and
150 realms, can provide a cost-effective means to significantly improve outcomes. Understanding where
151 such gains are highest and lowest should be a focus of future research efforts. Crucially, our findings
152 from two biogeographically distinct regions with different biophysical drivers of species distributions
153 (Fig. S11) suggest substantial freshwater gains ought to be attainable across the biodiverse
154 agricultural frontier regions of the forested tropics. Second, conservation remains hampered by a
155 severe lack of biodiversity data, especially in tropical regions (11,25). Resolving these data shortfalls
156 will be necessary to unlock the benefits we document, and this will require more investment in large-
157 scale ecological surveys and taxonomy (16,26). Third, to be effective and feasible, integrated
158 terrestrial-freshwater strategies need to be aligned with or incorporated into current environmental
159 policies and laws. In particular, freshwater-orientated planning should not come at the expense of
160 existing protected areas, which often hold the last populations of endangered species and are coming
161 under increasing pressure globally (27) and in the Amazon (28). Overcoming these challenges will be
162 difficult, but the task is small compared to the enormous gains that can be made for the world's
163 diverse and highly threatened freshwater biota.

164 **References**

- 165 1. E. V. Balian, H. Segers, C. Lévêque, K. Martens, K., The Freshwater Animal Diversity
166 Assessment: an overview of the results. *Hydrobiologia* **595**, 627-637 (2008).
- 167 2. Millennium Ecosystem Assessment. Ecosystems and human well-being: wetlands and water—
168 Synthesis (World Resources Institute, Washington, D. C., 2005).
- 169 3. A. J. Reid *et al.*, Emerging threats and persistent conservation challenges for freshwater
170 biodiversity. *Biol. Rev.* **94**, 849-873 (2018).
- 171 4. UNEP-WCMC and IUCN. Protected Planet Report 2016 (UNEP-WCMC and IUCN, Cambridge,
172 UK and Gland, Switzerland, 2016).
- 173 5. D. Dudgeon *et al.*, Freshwater biodiversity: importance, threats, status and conservation
174 challenges. *Biol. Rev.* **81**, 163-182 (2006).
- 175 6. J. S. Albert *et al.*, Scientists' warning to humanity on the freshwater biodiversity crisis. *Ambio*
176 <https://doi.org/10.1007/s13280-020-01318-8> (2020).
- 177 7. T. M. Brooks *et al.*, Global biodiversity conservation priorities. *Science* **313**, 58-61 (2006).
- 178 8. J. E. M. Watson, N. Dudley, D. B. Segan, M. Hockings, The performance and potential of
179 protected areas. *Nature* **515**, 67-73 (2014).
- 180 9. D. Tickner *et al.*, Bending the curve of global freshwater biodiversity loss: an emergency
181 recovery plan. *Bioscience* **4**, 330-342 (2020).
- 182 10. R. Abell *et al.*, Concordance of freshwater and terrestrial biodiversity. *Conserv. Lett.* **4**, 127-136
183 (2011).

- 184 11. A. S. L. Rodrigues, T. M. Brooks, Shortcuts for biodiversity conservation planning: effectiveness
185 of surrogates. *Annu. Rev. Ecol. Evol. S.* **38**, 713-737 (2007).
- 186 12. L. Castello *et al.*, The vulnerability of Amazon freshwater ecosystems. *Conserv. Lett.* **6**, 217-229
187 (2013).
- 188 13. R. Abell, J. Allan, B. Lehner, Unlocking the potential of protected areas for freshwaters. *Biol.*
189 *Conserv.* **134**, 48-63 (2007).
- 190 14. S. Linke, E. Turak, J. Nel, Freshwater conservation planning: the case for systematic approaches,
191 *Freshwater Biol.* **56**, 6-20 (2011).
- 192 15. J. Carvajal-Quintero *et al.*, Drainage network position and historical connectivity explain global
193 patterns in freshwater fishes' range size. *Proc. Natl Acad. Sci. USA*, **116**, 134-13439 (2019).
- 194 16. J. Barlow *et al.*, The future of hyperdiverse tropical ecosystems. *Nature*, **559**, 517-526 (2018).
- 195 17. See supplementary materials.
- 196 18. T. A. Gardner *et al.*, A social and ecological assessment of tropical land uses at multiple scales:
197 The Sustainable Amazon Network. *Phil. T. R. Soc. B.* **368**, 20120166 (2013).
- 198 19. J. L. Nessimian *et al.*, Land use, habitat integrity, and aquatic insect assemblages in Central
199 Amazonian streams. *Hydrobiologia*, **614**, 117-131 (2008).
- 200 20. A. Moilanen, Landscape zonation, benefit functions and target-based planning: Unifying reserve
201 selection strategies. *Biol. Conserv.* **134**, 571-579 (2007).
- 202 21. A. Moilanen *et al.*, A method for freshwater conservation prioritization. *Freshwater Biol.* **53**,
203 577-592 (2008).
- 204 22. J. J. Tewksbury *et al.*, Corridors affect plants, animals, and their interactions in fragmented
205 landscapes. *Proc. Natl Acad. Sci. USA* **99**, 12923-12929 (2002).
- 206 23. E. J. McIntosh *et al.*, Absence of evidence for the conservation outcomes of systematic
207 conservation planning around the globe: a systematic map. *Environ. Evid.* **7**, 22 (2018).
- 208 24. G. M. Mace *et al.*, Aiming higher to bend the curve of biodiversity loss. *Nat. Sustainability*, **1**,
209 448-451 (2018).
- 210 25. W. R. T. Darwall *et al.*, Implications of bias in conservation research and investment for
211 freshwater species. *Conserv. Lett.* **4**, 474-482 (2011).
- 212 26. L. W. Drew, Are We Losing the Science of Taxonomy? *BioScience* **61**, 942-946 (2011).
- 213 27. M. B. Mascia *et al.*, Protected area downgrading, downsizing, and degazettement (PADDD) in
214 Africa, Asia, and Latin America and the Caribbean, 1900-2010. *Biol. Conserv.* **169**, 355-361
215 (2014).
- 216 28. S. M. Pack *et al.*, Protected area downgrading, downsizing, and degazettement (PADDD) in the
217 Amazon. *Biol. Conserv.*, **197**, 32-39 (2016).
- 218 29. J. M. C. da Silva, A. B. Rylands, G. A. B. Da Fonseca, The fate of the Amazonian areas of
219 endemism. *Conserv. Biol.* **19**, 689-694 (2005).

- 220 30. Instituto Nacional de Pesquisas Espaciais. TerraClass Data 2010. [http://www.inpe.br/cra/](http://www.inpe.br/cra/projetos_pesquisas/terraclass2010)
221 projetos_pesquisas/terraclass2010 (2013).
- 222 31. J. Barlow *et al.*, Anthropogenic disturbance in tropical forests can double biodiversity loss from
223 deforestation. *Nature* **535**, 144-147 (2016).
- 224 32. M. Winchell, R. Srinivasan, M. Di Luzio, J. G. Arnold, Arcswat Interface for SWAT2012: User's
225 Guide (AgriLife Research, Texas, 2013).
- 226 33. A. C. Lees, *et al.*, One hundred and thirty-five years of avifaunal surveys around Santarem,
227 central Brazilian Amazon. *Rev. Bras. Ornitol.* **21**, 16-57 (2013).
- 228 34. A. C. Lees *et al.*, Paragominas: a quantitative baseline inventory of an eastern Amazonian
229 avifauna. *Rev. Bras. Ornitol.* **20**, 93-118 (2012).
- 230 35. J. M. B. Oliveira-Junior *et al.*, Neotropical dragonflies (Insecta: Odonata) as indicators of
231 ecological condition of small streams in the eastern Amazon. *Austral Ecol.* **40**, 733-744 (2015).
- 232 36. J. G. Brito *et al.*, Biological indicators of diversity in tropical streams: Congruence in the
233 similarity of invertebrate assemblages. *Ecol. Indic.* **85**, 85-92 (2018).
- 234 37. C. G. Leal *et al.*, Is environmental legislation conserving tropical stream faunas? A large-scale
235 assessment of local, riparian and catchment-scales influences on Amazonian fish. *J. Appl. Ecol.*
236 **55**, 1312-1326 (2018).
- 237 38. N. Ellis *et al.*, Gradient forests: Calculating importance gradients on physical predictors. *Ecology*
238 **93**, 156-168 (2012).
- 239 39. J. Pearce, S. Ferrier, Evaluating the predictive performance of habitat models developed using
240 logistic regression. *Ecol. Model.* **133**, 225-245 (2000).
- 241 40. F. E. Putz, K. H. Redford, The importance of defining "forest": Tropical forest degradation,
242 deforestation, long-term phase shifts, and further transitions. *Biotropica* **42**, 10-20 (2010).
- 243 41. G. D. Lennox *et al.*, Second rate or a second chance? Assessing biomass and biodiversity
244 recovery in regenerating Amazonian forests. *Glob. Change Biol.* **24**, 5680-5694 (2018).
- 245 42. N. R. Bond, J. R. Thomson, P. Reich, Incorporating climate change in conservation planning for
246 freshwater fishes. *Divers. Distrib.* **20**, 931-942 (2014).
- 247 43. R. D. Garrett *et al.*, Explaining the persistence of low income and environmentally degrading
248 land uses in the Brazilian Amazon. *Ecol. Soc.* **22**, 27 (2017).
- 249 44. J. A. Robalino, A. Pfaff, Contagious development: Neighbor interactions in deforestation. *J. Dev.*
250 *Econ.* **97**, 427-436 (2012).
- 251 45. M. D. C. Vera-Diaz, R. K. Kaufmann, D. Nepstad, P. Schlesinger, An interdisciplinary model of
252 soybean yield in the Amazon Basin: The climatic, edaphic, and economic determinants. *Ecol.*
253 *Econ.* **65**, 420-431 (2008).
- 254 46. A. Gusso, J. R. Ducati, Algorithm for Soybean Classification Using Medium Resolution Satellite
255 Images. *Remote Sens.* **4**, 3127-3142 (2012).

- 256 47. A. Gusso, D. Arvor, J. R. Ducati, M. R. Veronez, L. G. da Silveira Junior, Assessing the MODIS
257 Crop Detection Algorithm for Soybean Crop Area Mapping and Expansion in the Mato Grosso
258 State, Brazil. *Sci. World J.* **2014**, 863141 (2014).
- 259 48. S. M. Vincente-Serrano, S. Beguería, J. I. López-Moreno, A Multiscalar Drought Index Sensitive
260 to Global Warming: The Standardized Precipitation Evapotranspiration Index. *J. Clim.* **23**, 1696-
261 1718 (2010).
- 262 49. J. T. Abatzoglou *et al.*, TerraClimate, a high-resolution global dataset of monthly climate and
263 climatic water balance from 1958–2015. *Sci. Data* **5**, 170191 (2018).
- 264 50. S. Begueria. S. M. Vincente-Serrano, SPEI: Calculation of the Standardised Precipitation-
265 Evapotranspiration Index <https://CRAN.R-project.org/package=SPEI> (2017).

266 **Acknowledgements:** We thank Ima Vieira for coordinating the INCT Biodiversidade e Uso da Terra
267 na Amazônia and the Large Scale Biosphere-Atmosphere Program (LBA) for logistical and
268 infrastructure support during field measurements. **Funding:** This work was supported by grants from
269 Brazil (CNPq 574008/2008-0, 458022/2013-6, and 400640/2012-0; Embrapa SEG:02.08.06.005.00;
270 The Nature Conservancy – Brasil; CAPES scholarships) the UK (Darwin Initiative 17-023;
271 NE/F01614X/1; NE/G000816/1; NE/F015356/2; NE/I018123/1; NE/K016431/1), Formas 2013-1571,
272 and Australian Research Council grant DP120100797. Individual support included FAPESP funding
273 (2017/25383-0) to CGL, H2020-MSCA-RISE (691053-ODYSSEA) funding to EB and JB, Fulbright
274 Brasil funding to RMH, NSF-Belmont Forum award (1852113) to LC, and CNPq funding
275 (144452/2010-3; 307597/2016-4; 436007/2018-5 and 313183/2014-7 respectively) to VCO, LJ, RPL
276 and JZ. Institutional support was provided by the Herbário IAN in Belém, LBA in Santarém and
277 FAPEMAT. This is paper no. 76 in the Sustainable Amazon Network publication series. **Author**
278 **contributions:** TAG, JF, JB, LP, CGL, GDL and SFBF designed the research with additional taxon-
279 specific input from EB, ACL, RMH, LEOCA, JGB, NH, LJ, RPL, JL, NGM, JLN, JMBO, VHFO,
280 VCO, PSP, RRCS and JZ. CGL, EB, ACL, RMH, JGB, NH, LJ, RPL, JL, NGM, JLN, JMBO,
281 VHFO, VCO, LP, RRCS and JZ collected the field data or analyzed biological samples. SFBF and
282 TAG processed the remote sensing data. GDL and CGL undertook the analyses with input from JB,
283 JRT, RMN, RDG, TFM and SFBF. CGL, GDL and JB led writing with all authors contributing to
284 reviewing and editing. **Competing interests:** Authors declare no competing interests. **Data and**
285 **materials availability:** The data and computer code used in these analyses are available at:
286 <https://doi.org/10.6084/m9.figshare.12423881>. The Zonation conservation planning software is
287 available at: <https://github.com/cbig/zonation-core>.

288 **Ethics statement:** All biodiversity sampling was undertaken in compliance with Brazilian
289 environmental regulations under the following licenses: (i) Sisbio license #24164 for collecting plants,
290 issued by the Chico Mendes Institute for Biodiversity Conservation (ICMBio); (ii) Sisbio license

291 #10061-1 for collecting dung beetles, issued by the Institute of Environment and Renewable Natural
292 Resources (IBAMA); (iii) Sisbio licenses #10199-2 and #24355-2 for collecting fish, both issued by
293 ICMBio (iv) Sisbio license #10873-1 for collecting insects, issued by IBAMA; (v) Sisbio license
294 #19102-4 for collecting Odonata, Ephemeroptera, Plecoptera and Trichoptera, issued by ICMBio. No
295 license was required for bird sampling because the methods were observational and did not involve
296 collecting or handling of specimens. Socio-economic data was collected following the UK Research
297 Integrity Office Principles for Research involving human participants, human material, and personal
298 data and was collected with informed consent. Further approval for opportunity cost data collection
299 was obtained from the Brazilian Agricultural Research Corporation (Embrapa) under CAAE
300 29054920.4.0000.5173 and Stanford University under IRB Protocol 19044.

301 **Supplementary Materials:**

302 Materials and Methods

303 Figures S1-S13

304 Table S1

305 References 29-50


306 Figure legends

307 **Fig. 1. Incidental conservation.** The incidental conservation benefits achieved for one species group
308 when focused on another. The *x*-axis ticks are labelled with the focal group first. For example, *T-F*
309 shows the incidental conservation benefits achieved for a freshwater group when prioritizing for a
310 terrestrial group. Points show results for each taxonomic pair. Boxplots show the interquartile range.
311 The center line shows the median. Results are shown for the area-constrained analysis (**A**) with the
312 constraint that 10%, 20% or 30% of landscape can be conserved, and the budget-constrained analyses
313 with two opportunity cost estimates (**B-C**) and with budget levels such that approximately 10%, 20%
314 and 30% of the landscape can be conserved (17). Letters next to the boxplots show results of pairwise
315 comparisons of group means within resource levels (17). Variables not sharing a letter have
316 statistically different means.

317 **Fig. 2. Catchment prioritizations for terrestrial and freshwater biodiversity.** Catchment
318 conservation priority rankings in Paragominas (**A-F**) and Santarém (**G-L**) for terrestrial (**A-C, G-I**)
319 and freshwater (**D-F, J-L**) taxa. Rankings are based on catchment marginal conservation value, with 1
320 indicating the catchment with the highest marginal conservation value and 0 that with the lowest
321 marginal conservation value. Results are shown for the area-constrained analysis.


322 **Fig. 3. Terrestrial-freshwater trade-offs.** The decrease in terrestrial benefits from their optimum
323 required to achieve an increase in freshwater benefits through the joint-planning and the terrestrial-
324 plus-connectivity approaches in Paragominas (**A, C, E**) and Santarém (**B, D, F**). The thin lines show
325 the results for each terrestrial-freshwater taxonomic pair. The thick lines show one s.e.m., where the
326 mean was estimated using Holling type-II curves, for each integrated planning approach. Results are
327 shown for the area-constrained analysis (**A-B**) with the constraint that 20% of landscape could be
328 conserved, and the budget-constrained analyzes with two opportunity cost estimates (**C-F**) and with
329 budget levels such that approximately 20% of the landscape could be conserved (17).

330
331




332

333



334

335



336

337