

Journal of Experimental Botany

doi:10.1093/jxb/eru264

This paper is available online free of all access charges (see http://jxb.oxfordjournals.org/open_access.html for further details)


Journal of
Experimental
Botany
www.jxb.oxfordjournals.org

RESEARCH PAPER

Overexpression of the kiwifruit *SVP3* gene affects reproductive development and suppresses anthocyanin biosynthesis in petals, but has no effect on vegetative growth, dormancy, or flowering time

Rongmei Wu¹, Tianchi Wang¹, Tony McGie², Charlotte Voogd¹, Andrew C. Allan^{1,3}, Roger P. Hellens¹ and Erika Varkonyi-Gasic^{1,*}

¹ The New Zealand Institute for Plant & Food Research Limited (Plant & Food Research) Mt Albert, Private Bag 92169, Auckland 1142, New Zealand

² The New Zealand Institute for Plant & Food Research Limited (Plant & Food Research) Palmerston North, Private Bag 11600, Palmerston North 4442, New Zealand

³ School of Biological Sciences, University of Auckland, Private Bag 92019, Auckland, New Zealand

* To whom correspondence should be addressed. E-mail: erika.varkonyi-gasic@plantandfood.co.nz

Received 13 March 2014; Revised 15 May 2014; Accepted 19 May 2014

Abstract

SVP-like MADS domain transcription factors have been shown to regulate flowering time and both inflorescence and flower development in annual plants, while having effects on growth cessation and terminal bud formation in perennial species. Previously, four SVP genes were described in woody perennial vine kiwifruit (*Actinidia* spp.), with possible distinct roles in bud dormancy and flowering. Kiwifruit *SVP3* transcript was confined to vegetative tissues and acted as a repressor of flowering as it was able to rescue the *Arabidopsis svp41* mutant. To characterize kiwifruit *SVP3* further, ectopic expression in kiwifruit species was performed. Ectopic expression of *SVP3* in *A. deliciosa* did not affect general plant growth or the duration of endodormancy. Ectopic expression of *SVP3* in *A. eriantha* also resulted in plants with normal vegetative growth, bud break, and flowering time. However, significantly prolonged and abnormal flower, fruit, and seed development were observed, arising from *SVP3* interactions with kiwifruit floral homeotic MADS-domain proteins. Petal pigmentation was reduced as a result of *SVP3*-mediated interference with transcription of the kiwifruit flower tissue-specific *R2R3 MYB* regulator, *MYB110a*, and the gene encoding the key anthocyanin biosynthetic step, *F3GT1*. Constitutive expression of *SVP3* had a similar impact on reproductive development in transgenic tobacco. The flowering time was not affected in day-neutral and photoperiod-responsive *Nicotiana tabacum* cultivars, but anthesis and seed germination were significantly delayed. The accumulation of anthocyanin in petals was reduced and the same underlying mechanism of *R2R3 MYB NtAN2* transcript reduction was demonstrated.

Key words: *Actinidia*, bud break, dormancy, flowering, kiwifruit, MYB, *Nicotiana*, petal colour, SVP.

Introduction

Plant MADS-domain transcription factors act as key regulators of many developmental processes (Smaczniak *et al.*, 2012). The STMADS subfamily of MADS-box genes are predominantly expressed in vegetative tissues and play roles in vegetative development and transition to flowering

in diverse plant species (Carmona *et al.*, 1998; Hartmann *et al.*, 2000; Mao *et al.*, 2000; Yu *et al.*, 2002; Masiero *et al.*, 2004; Aswath and Kim, 2005; Liu *et al.*, 2007; Trevaskis *et al.*, 2007; He *et al.*, 2010; Cohen *et al.*, 2012; Cooke *et al.*, 2012).

Relatively small changes in sequence can have a major effect on STMADS gene function. In *Arabidopsis thaliana*, the two STMADS genes *SVP* and *AGL24* act in an opposite manner despite their close homology. *SVP* plays a key role as a repressor of the transition to flowering; a single amino acid substitution causes the loss of function resulting in an early flowering phenotype (Hartmann *et al.*, 2000; Méndez-Vigo *et al.*, 2013). In contrast, *AGL24* acts as a promoter of flowering because its inactivation results in late flowering (Yu *et al.*, 2002). Both proteins have further roles during early stages of flower development to repress expression of class B and C floral homeotic genes and maintain floral meristem identity (Gregis *et al.*, 2009). Protein interactions and formation of complexes with other MADS-domain proteins are essential for STMADS function (Liu *et al.*, 2009; Gregis *et al.*, 2013). Interaction with *APETALA1* (*API*) in particular resulted in floral reversion phenotypes observed in transgenic *Arabidopsis* upon ectopic expression of *SVP*-like genes (Liu *et al.*, 2007; Gregis *et al.*, 2008; Fornara *et al.*, 2008; Wu *et al.*, 2012; Jaudal *et al.*, 2013). For these and perhaps other reasons, the function of STMADS observed upon ectopic expression in *Arabidopsis* might be different from their function in the species of origin (Trevaskis *et al.*, 2007; Jaudal *et al.*, 2013).

In woody perennial plants, the STMADS subfamily genes have been reported to act as growth inhibitors, which correlated with maintenance of plant dormancy (Mazzitelli *et al.*, 2007; Bielenberg *et al.*, 2008; Diaz-Riquelme *et al.*, 2009; Horvath *et al.*, 2010; Li *et al.*, 2010; Sasaki *et al.*, 2011; Yamane *et al.*, 2011). Deletion of the six *SVP*-like *DORMANCY-ASSOCIATED MADS-BOX* (*DAM*) genes in peach (*Prunus persica*) resulted in failure to enter dormancy under cold or short-day induction, and the expression of a subset of these genes was elevated during endodormancy (Bielenberg *et al.*, 2008; Li *et al.*, 2009; Yamane *et al.*, 2011). Similarly, a negative correlation of expression with endodormancy release was observed for six tandemly arrayed *DAM* genes predicted to act as transcriptional repressors in Japanese apricot (*Prunus mume*) (Yamane *et al.*, 2008; Sasaki *et al.*, 2011). Ectopic expression of one of these genes in transgenic poplar resulted in premature growth cessation and terminal bud set, demonstrating the role in growth inhibition (Sasaki *et al.*, 2011). However, neither of these genes was functionally characterized by ectopic expression in the plant species of origin.

In the kiwifruit species *Actinidia chinensis* and *A. deliciosa*, four *SVP* genes (*SVPI–SVP4*) have been identified, with potentially distinct roles in bud dormancy and flowering (Wu *et al.*, 2012). Their expression was mostly confined to vegetative tissues, with *SVP3* showing the highest relative expression. *SVPI*, *SVP2*, and *SVP4* were elevated in buds over the winter dormancy period. In contrast, *SVP3* accumulation in buds did not demonstrate seasonal changes, but delayed flowering in transgenic *Arabidopsis* and was able to rescue the *Arabidopsis svp41* mutant. Ectopic expression resulted in floral reversion phenotypes, caused by the strong capacity for heterodimerization with *Arabidopsis* MADS box proteins responsible for flower development (Wu *et al.*, 2012).

In the current study, functional analysis of *SVP3* was performed using ectopic transgenic analysis in kiwifruit species *A. deliciosa* and *A. eriantha*, which have different chilling requirements and bloom time, and a model plant species tobacco (*Nicotiana tabacum*). Normal vegetative growth and floral transition, but abnormal flower development, reduced petal pigmentation, and abnormal fruit and seed development were observed. The underlying mechanism of reduced anthocyanin accumulation in petals was examined. Based on the results obtained, the biological function of *SVP3* is discussed.

Materials and methods

Gene isolation and vector construction

The *A. chinensis SVP3* coding sequence was cloned under the control of the *Cauliflower mosaic virus* (CaMV) 35S promoter (*35S:SVP3*) into the pART277 binary vector (Gleave, 1992) as previously described (Wu *et al.*, 2012). The resulting plasmid was transformed into *Agrobacterium tumefaciens* strain EHA105 for transformation into *A. deliciosa* and *A. eriantha*, and into GV3101 for transformation into *N. tabacum*. A construct with a reporter gene *uidA* (*GUS*; β -glucuronidase) under the control of the CaMV 35S promoter (*35S:GUS*) in appropriate *Agrobacterium* strains was used to transform control plants (Wang *et al.*, 2006). *Actinidia eriantha SVP3* (GenBank accession no. KJ123703) was amplified from the bud cDNA using gene-specific oligonucleotide primers, 5'-ATGGCGAGAGAGAAGATCAAGA-3' (forward) and 5'-TGGTGTGACATTTCAAGTTCG-3' (reverse). The predicted amino acid sequence alignment with *A. deliciosa* and *A. chinensis SVP3* homologues was performed using Vector NTI version 9.0. (Invitrogen) Clustal W (opening 15, extension penalty 0.3).

Plant transformation and growth conditions

The transformation procedures for *A. deliciosa* and *A. eriantha* were as previously described (Wang *et al.*, 2006, 2007). Once roots were established, transgenic plants were transferred to soil and grown in the containment glasshouse at the Plant & Food Research, Mt Albert, Auckland, New Zealand for 18 months. Flowering time and floral phenotype were assessed in the following spring season. *Nicotiana tabacum* 'Maryland Mammoth' and 'Samsun' transformations were carried out on young leaf discs excised from *in vitro* grown shoots (Horsch *et al.*, 1985). Transgenic tobacco plants were grown in a glasshouse at 20 °C under long-day (16/8 h light/dark; for 'Samsun') or short-day conditions (8/16 h light/dark; for 'Maryland Mammoth'). The seeds from these transgenic plants were collected and germinated on half-strength Murashige and Skoog (MS) agar medium (Murashige and Skoog, 1962) supplemented with 50 μ g ml⁻¹ kanamycin. Following the segregation tests, two homozygous lines of both cultivars at T₂ generations were used for flowering time and floral phenotype assessment.

RNA extraction and expression studies

Actinidia eriantha root, stem, leaf, axillary bud, shoot apex, flower, and fruit tissue collection was carried out on vines growing in the glasshouse at the Plant & Food Research, Mt Albert, Auckland, New Zealand, during the spring and summer season of 2011–2012. *Actinidia chinensis* root, stem, leaf, flower, fruit, and axillary and apical bud collection was carried out on vines growing at the Plant & Food Research orchard near Kerikeri, New Zealand, during the spring and summer season of 2005–2006.

Total RNA was extracted from kiwifruit tissue as previously described (Chang *et al.*, 1993). Total RNA was isolated from

tobacco flowers using the Trizol reagent (Invitrogen). A 5 µg aliquot of total RNA was treated with DNase I (Ambion) and reverse-transcribed at 37 °C using the BluePrint[®] Reagent kit for reverse transcription–PCR (RT–PCR; TaKaRa) according to the manufacturer's instructions. Amplification and quantification were carried out using the LightCycler[®] 480 System and LightCycler[®] 480 SYBR Green I Master Mix (Roche Diagnostics). A minimum of three technical replicate reactions were performed and a non-template control was included in each run. Thermal cycling conditions were 95 °C for 5 min, followed by 50 cycles of 95 °C for 10 s, 60 °C for 10 s, and 72 °C for 20 s, and a melting temperature cycle, with constant fluorescence data acquisition from 65 °C to 95 °C. The data were analysed using the Target/Reference ratio calculated with the LightCycler[®] 480 software 1.5 (Roche Diagnostics). The expression was normalized to previously characterized reference genes, kiwifruit *actin* (Wu *et al.*, 2012) and tobacco *Ntα-Tub1* (Pattanaik *et al.*, 2010).

Oligonucleotide primers for kiwifruit *SVP3*, *actin*, anthocyanin pathway genes, and regulatory genes *MYB110* and *MYB10* were as previously described (Montefiori *et al.*, 2011; Wu *et al.*, 2012; Fraser *et al.*, 2013). Oligonucleotide primers for tobacco *NtAn1a*, *NtAN1b*, *NtAn2*, and *Ntα-Tub1* were as described by Bai *et al.* (2011) and Pattanaik *et al.* (2010).

Biochemical analyses

For chlorophyll quantification, petals collected at visually similar developmental stages were frozen and ground in liquid nitrogen. Ground tissues were suspended in 80% acetone with 2.5M phosphate buffer pH 7.5 and incubated for 1 h in the dark, followed by centrifugation at 16 000 *g* for 10 min. The absorbance of 200 µl of the supernatant was measured spectrophotometrically at 663 nm and 646 nm (SpectraMax 384, Molecular Devices, Sunnyvale, CA, USA). The total chlorophyll (C_{a+b}) concentration was calculated using the equation $C_{a+b} (\mu\text{g ml}^{-1}) = 7.15A_{663} + 18.71A_{646}$ (Lichtenthaler and Rinderle, 1988).

Anthocyanins and flavonoids were extracted from lyophilized samples and analysed by high-performance liquid chromatography (HPLC) and liquid chromatography–mass spectrometry (LC-MS) as described previously (McGhie *et al.*, 2011; Dare *et al.*, 2013). Quantification of cyanidin 3-*O*-galactoside was achieved by calibration at 520 nm by HPLC analysis. The peaks were identified by a comparison of retention times with authentic standards of cyanidin 3-*O*-galactoside (Polyphenols Laboratories, Sandnes, Norway).

Microscopic analyses

Tobacco petal epidermis was obtained by careful peeling and observed under a light microscope (Olympus Vanox AHT3, Olympus Optical Co. Ltd, Tokyo, Japan). Photographs were taken using a CoolSNAP Colour camera system (Roper Scientific Ltd, Tucson, AZ, USA).

Yeast two-hybrid assay

The full-length open reading frame of kiwifruit *SVP* genes and kiwifruit *AGAMOUS* (*AG*), *FRUITFULL* class (*FUL* and *FUL*-like), *SEPALLATA* class (*SEPI*, *SEP3*, and *SEP4*), *APETALA3* (*AP3*), *PISTILLATA* (*PI*), and *CENTRORADIALIS* (*CEN*) were amplified and cloned as described previously (Varkonyi-Gasic *et al.*, 2011, 2013; Wu *et al.*, 2012). The bait vector and prey vector were mated and selected on synthetic complete (SC) plates lacking leucine and tryptophan. The affinity of two protein interactions was determined on selective SC medium lacking leucine, tryptophan, and histidine, and in the presence of 1, 3, and 5 mM 3-aminotriazole (3AT) at 30 °C. All assayed proteins were tested for autoactivation, homodimerization, and heterodimerization capacity. The strength of interactions was compared with control plasmids of pEXPTM32/Krev1 with pEXPTM22/RalGDS-wt, pEXPTM22/RalGDS-m1, and pEXPTM22/RalGDS-m2 (Invitrogen).

Results

Overexpression of SVP3 has no significant effect on growth or dormancy in A. deliciosa

To investigate the role of kiwifruit *SVP3*, transgenic *A. deliciosa* was generated using the *SVP3* cDNA driven by the CaMV 35S promoter (Wu *et al.*, 2012). Ten independent transgenic lines with high *SVP3* transgene expression levels were generated and compared with six control lines (Fig. 1A). Monitoring during callus formation, plantlet growth in tissue culture, and upon transfer to soil revealed no obvious difference between 35S:*SVP3* lines and control lines. Eight 35S:*SVP3* and three control lines were further monitored for spring bud break. The number of breaking buds or developing shoots was recorded weekly and is presented as a percentage of total buds. Large variation in the time of the first visible bud break, positions of breaking buds, number of developing branches, and rate of shoot outgrowth was observed for all lines; however, the first visible bud break time of the 35S:*SVP3* lines was in the range recorded for control lines and was followed by normal growth of multiple branches (Fig. 1B, C). Therefore, it was concluded that *SVP3* does not have a significant effect on growth and dormancy in *A. deliciosa* under glasshouse conditions.

SVP3 affects reproductive development in transgenic A. eriantha

Actinidia deliciosa has a high chilling requirement (Wall *et al.*, 2008) and late bud break, and fails to flower under glasshouse conditions. Therefore, to evaluate further the role of *SVP3*, the non-cultivated kiwifruit species *A. eriantha* was chosen for transformation, as it provided advantages including low chilling requirement and early bud break, fast maturity, and prolific flowering in glasshouse conditions (Wang *et al.*, 2006). *Actinidia eriantha SVP3* shares almost identical sequence and expression pattern with the previously studied cultivated kiwifruit species *A. deliciosa* and *A. chinensis* (Supplementary Fig. S1A, B available at JXB online; Wu *et al.*, 2012). Six transgenic lines with moderate to high levels of *SVP3* transgene expression and three control lines were generated (Fig. 2A). None of the lines showed any obvious difference during callus formation, growth in tissue culture, and establishment in the soil; similar growth rate and architecture were observed (Supplementary Fig. S1 available at JXB online), and in all lines the time of bud break and flowering time (appearance of visible floral buds) were comparable (Table 1). The number of flowers was highly variable and not correlated with transgene expression levels, but the duration of flower development and appearance of flowers were altered significantly in transgenic lines, particularly in the highest expressing Line 1 (Table 1).

For further detailed analysis, Line 1 and Line 2 were chosen as representatives of high and intermediate *SVP3* transgene expression, respectively. Flower buds in Line 1 and to a lesser extent in Line 2 were not fully enclosed with sepals (Fig. 2B); the number and order of developing floral organs appeared normal (Fig. 2C), but flowers in Line 1 remained




Fig. 1. Overexpression of *SVP3* in *Actinidia deliciosa*. (A) Relative expression of *SVP3* in 35S:*SVP3* and control plants. The expression was normalized against kiwifruit *actin*. Error bars represent the standard error (SE) of four replicate reactions. (B) Bud break time measured as a percentage of total buds with visible shoot outgrowth. (C) A subset of 35S:*SVP3* and control *A. deliciosa* plants, photographed on the same day in spring. Arrowheads indicate positions of visible bud break and shoot outgrowth. Scale bar=10cm.

smaller (Fig. 2D). Fruit in Line 1 was misshapen and smaller compared with control or Line 2 fruit (Fig. 2E, F) and the transgenic seed was smaller, lighter (Fig. 2G–I), and failed to germinate.

Floral reversion phenotypes observed upon expression of *STMADS* genes in *Arabidopsis* result from interaction with *Arabidopsis* floral homeotic MADS box proteins, particularly AP1 (Fornara *et al.*, 2008; Gregis *et al.*, 2009; Wu *et al.*, 2012). To test if phenotypes observed upon ectopic expression of *SVP3* in *A. eriantha* arose from similar MADS box protein interactions, yeast two-hybrid analysis was performed using previously characterized kiwifruit floral MADS box genes (Varkonyi-Gasic *et al.*, 2011). *SVP3* interacted with kiwifruit AG and SEP proteins SEP1, SEP3, and SEP4 with similar

strong interaction intensities; a weaker interaction was identified with FUL and FUL-like, but no interaction was detected with AP3, PI, and the negative control CEN (Fig. 2J).

SVP3 affects petal colour in transgenic *A. eriantha* by transcriptional repression of key genes in the anthocyanin pathway

In addition to the described phenotypes, the colour of the petals of 35S:*SVP3* lines was dramatically reduced (Fig. 2D, Table 1) and the chlorophyll content was higher (Fig. 3A).

The presence and intensity of the peak corresponding to the major kiwifruit anthocyanin, cyanidin 3-*O*-galactoside, correlated with the visual petal phenotype in flower buds and open




Fig. 2. Overexpression of *SVP3* in *Actinidia eriantha*. (A) Relative expression of *SVP3* in *35S:SVP3* and control plants. The expression was normalized against kiwifruit *actin*. Error bars represent the standard error (SE) of four replicate reactions. (B–D) Morphology of flower buds (B, C) and fully open flowers (D). (E, F) Fruit morphology. (G–I) Seed morphology (G, H) and weight (I). Error bars represent the standard error (SE), $n=3$. Scale bars=5mm (B, C), 10mm (D–F), and 0.5mm (G, H). (J) Protein interactions detected by yeast two-hybrid assays. ++, very strong interaction; +, strong interaction; –, no interaction.

flowers (Fig. 3B, C), and further quantification confirmed differential accumulation of cyanidin 3-*O*-galactoside (Fig. 3D) and several minor flavonoid compounds (Supplementary Fig. S2 available at *JXB* online).

Analysis of the relative expression of the genes encoding enzymes of the anthocyanin biosynthetic pathway (Montefiori *et al.*, 2011) revealed similar expression profiles in representative *35S:SVP3* and control lines, with some variation between flower bud and open flower samples. The exception was the transcript of *F3GT1*, which was absent or barely detectable in flower buds of *35S:SVP3* lines, in contrast to relatively high accumulation observed in the control flower bud (Fig. 4A). Analysis of regulatory genes of the anthocyanin biosynthetic pathway (Fraser *et al.*, 2013) revealed high relative expression of *MYB110a* and low relative expression of *MYB10* in flower buds of the control line. In *35S:SVP3* lines these transcripts were absent or barely detectable, and in the control line both transcripts showed a decrease in open flowers (Fig. 4B).

SVP3 does not alter flowering time but affects reproductive development and petal colour in transgenic tobacco

To investigate further whether kiwifruit *SVP3* affects reproductive development and anthocyanin accumulation in a conserved manner, transgenic tobacco lines were generated using the *35S:SVP3* construct. A day-neutral tobacco ‘Samsun’ and a short-day variety ‘Maryland Mammoth’ were used. This approach provided the opportunity to establish if *SVP3* affects flowering time in an annual representative of asterids, the clade of eudicots to which the perennial kiwifruit belongs. Eight independent transgenic lines of tobacco ‘Samsun’ and three of ‘Maryland Mammoth’ were generated. Two homozygous T₂ lines of each cultivar with high transgene expression and two corresponding wild-type control lines were chosen for detailed analysis. No significant differences were observed in vegetative growth and flowering time, but the duration of flower development was extended, flower fertility was diminished, and seed germination was delayed (Table 2). Spike-shaped sepals in transgenic plants failed to enclose flower buds fully, the corolla remained green, and the distal part of the petal was altered in both colour and shape (Fig. 5A). Additional phenotypes were identified, including occasional homeotic conversion of stamen to petal and change in the ovary shape and its position relative to attachment of other floral parts (Fig. 5B). Changes in total chlorophyll, anthocyanin, and flavonoid content and presence of stomata in green petal segments were also noted (Supplementary Fig. S3 available at *JXB* online). To establish if reduced pigment accumulation resulted from transcriptional repression of key regulators of anthocyanin biosynthesis, quantitative PCR (qPCR) analysis of basic helix–loop–helix (bHLH) genes, *NtAn1a* and *NtAn1b*, and the R2R3 MYB gene, *NtAn2* (Pattanaik *et al.*, 2010; Bai *et al.*, 2011), was performed. *NtAn2* was significantly and consistently down-regulated in transgenic lines, while *NtAn1a* and *NtAn1b* expression showed only minor variations (Fig. 5C).

Table 1. Phenotypic analysis of six lines of 35S:SVP3 transgenic *A. eriantha* compared with three control lines

Bud break time represents the number of days from 100% leaf drop to the first visible bud break. Flowering time represents the number of days from the first visible bud break to the first visible flower bud. Anthesis time represents the number of days from visible flower buds to fully open flowers.

<i>A. eriantha</i> transgenic lines	Bud break time (d)	Flowering time (d)	No. of flowers	Anthesis time (d)	Petal colour
Line 1	38	25	12	55.1 ± 10.3	White/green
Line 2	32	23	2	49.7 ± 4.1	Light pink
Line 3	38	24	1	43	Light pink
Line 4	38	26	2	47.6 ± 6.8	Light pink
Line 5	32	21	10	46.2 ± 7.6	Light pink
Line 6	35	21	2	45.1 ± 9.3	Light pink
Control 1	35	28	12	27.4 ± 4.1	Pink
Control 2	40	22	35	21.4 ± 5.0	Pink
Control 3	32	24	5	27.8 ± 6.2	Pink

Data are expressed as the mean ± SE.

Discussion

Overexpression of SVP3 affects reproductive development but not floral transition in transgenic kiwifruit and tobacco

Ectopic expression of the *STMADS* gene subfamily often results in altered flowering time, severe floral abnormalities, delayed flower anthesis, and senescence (Hartmann *et al.*, 2000; Yu *et al.*, 2002; Lee *et al.*, 2007; Fornara *et al.*, 2008; He *et al.*, 2010; Li *et al.*, 2010), and affects duration of endodormancy in woody perennial species (Sasaki *et al.*, 2011). Overexpression of *SVP3* in *A. deliciosa* showed no major effect on vegetative growth and the timing of first visible bud break in glasshouse conditions, although insufficient winter chilling and large variation in the position and number of synchronously breaking buds might have affected the rate of visible shoot outgrowth. Overexpression of *SVP3* in *A. eriantha* and tobacco showed no significant effect on plant growth, bud break, and flowering time, supporting the findings in *A. deliciosa*. However, flower, fruit, and seed development were severely affected. The size and shape of sepals were abnormal and they failed to enclose flower buds fully, consistent with the phenotype observed in *Arabidopsis* (Wu *et al.*, 2012) and similar to other reports (Masiero *et al.*, 2004; Liu *et al.*, 2007; Trevaskis *et al.*, 2007; Fornara *et al.*, 2008). The petals contained vegetative features including increased chlorophyll content and stomata, similar to phenotypes observed upon overexpression of potato *StMADS16* and sweet potato *IbMADS3-1* in tobacco (Garcia-Maroto *et al.*, 2000; Shin *et al.*, 2011). Flower development was slower, resulting in significantly delayed anthesis and an increased proportion of sterile flowers, which failed to produce fruit (Fornara *et al.*, 2008; Li *et al.*, 2010). Additional phenotypes included underdeveloped and misshapen fruit, abnormal seed development, reduced seed germination rate, and extended seed germination, suggesting that *SVP3* affected all stages of reproductive development. The severity of the observed phenotypes during reproductive development, combined with normal

vegetative growth, wild-type bud break and flowering time, and relatively high levels of expression in vegetative tissues (Wu *et al.*, 2012), indicate that *SVP3* has a role in the regulatory network of vegetative organogenesis and early stages of floral transition as the key repressor of floral organ development. It is hypothesized that *SVP3* acts to retain vegetative features during plant growth and floral transition, and has to be removed to enable subsequent stages of reproductive development. Ectopic expression phenotypes, combined with constitutive vegetative expression, suggest that there is no involvement of *SVP3* in maintenance of winter dormancy and general growth inhibition. However, *SVP3* may have a role in these processes as an interacting partner of other, rate-limiting proteins, and a loss-of-function phenotype would be required to understand *SVP3* function fully.

SVP3 interferes with anthocyanin biosynthesis in petals via repression of an anthocyanin-related MYB transcription factor

Overexpression of MADS-box genes is often associated with floral abnormalities, including changes in petal pigmentation. Light green sepaloid petals were observed upon overexpression of *Arabidopsis* *AGAMOUS-LIKE 20* (*AGL20* or *SOC-1*) in *Arabidopsis* (Liu *et al.*, 2007) and of the mustard *AGL20* orthologue *MADSA* in short-day tobacco ‘Maryland Mammoth’ (Borner *et al.*, 2008). Similarly, expression of *STMADS* genes such as potato *StMADS16* and sweet potato *IbMADS3-1* in tobacco resulted in development of chlorophyll-enriched petals (Garcia-Maroto *et al.*, 2000; Shin *et al.*, 2011). However, these reports did not investigate anthocyanin biosynthesis and accumulation in the petals of transgenic plants, and very little is known about potential MADS-box protein interference with the regulatory genes of the anthocyanin pathway. This study demonstrated that *SVP3* reduced anthocyanin accumulation in petals of transgenic *A. eriantha* in a quantitative manner. The underlying mechanism was dramatically reduced expression of the key structural




Fig. 3. Biochemical analysis of petal pigmentation. (A) Total chlorophyll content in petals of fully opened flowers. Error bars represent the SE of three independent measurements. (B, C) HPLC assay for cyanidin 3-O-galactoside correlated with visual phenotypes of flower buds (B) and open flowers (C). (D) Quantification of cyanidin 3-O-galactoside using LC-MS. Data represent the means \pm SE ($n=3$).

gene *F3GT1* (Montefiori *et al.*, 2011) and transcriptional repression of regulatory MYB transcription factors, in particular *MYB110a*, the key gene required for kiwifruit petal pigmentation (Fraser *et al.*, 2013). A similar mechanism was observed in transgenic tobacco, where *SVP3* interfered with transcription of the regulatory MYB *NtAn2*.

At this stage, it is unclear if *SVP3* acts by direct or indirect repression of the regulatory MYB transcription. Currently, there is little knowledge about upstream regulators of anthocyanin-related R2R3 MYB expression. Some MADS box genes from the *API/SQUA* class have been implicated in regulation of anthocyanin accumulation. *IbMADS10* expression

correlated with red pigmentation in sweet potato, and ectopic expression resulted in anthocyanin accumulation in transgenic sweet potato callus and transgenic *Arabidopsis* (Lalusin *et al.*, 2006, 2011). *VmTDR4* expression was linked with colour development and anthocyanin-related gene expression in bilberry, while silencing of this gene reduced anthocyanin levels and altered expression of the regulatory R2R3 MYB transcription factor (Jaakola *et al.*, 2010). STMADS proteins have a capacity to interact with other MADS box proteins, and interaction with AP1 affected flowering time and flower development in transgenic *Arabidopsis* (Fornara *et al.*, 2008; Gregis *et al.*, 2008; Wu *et al.*, 2012). Therefore, a similar




Fig. 4. Relative expression of genes responsible for anthocyanin biosynthesis in kiwifruit. (A) Relative expression of kiwifruit flavonoid biosynthetic pathway genes. *CHS*, chalcone synthase; *CHI*, chalcone isomerase; *F3H*, flavonoid 3-hydroxylase; *F3'H*, flavonoid 3'-hydroxylase; *DFR*, dihydroflavonol 4-reductase; *ANS*, anthocyanidin synthase; *F3GT1* and *F3GGT1*; flavonoid 3-O glycosyltransferases. (B) Relative expression of regulatory *MYB* genes. Relative expression of *SVP3* is presented below. The expression was normalized against kiwifruit *actin*. Error bars represent the SE of three replicate reactions.

Table 2. Phenotypic analysis of transgenic 35S:SVP tobacco ‘Maryland Mammoth’ and ‘Samsun’

Flowering time was measured as the number of leaves when visible flower buds were observed. Anthesis time represents days from flower buds to fully open flowers. Total number of flowers includes sterile flowers. Seed germination was recorded as number of days to visible germination on half-strength MS medium.

Cultivar	Genotype	Flowering time (d)	Anthesis time (d)	Total flowers	Sterile flowers	Seed germination
Maryland	35:SVP Line 1	19.5±0.19	7±1.4	27.5±0.5	20.5±2.2	16±0.3
Mammoth	35:SVP Line 2	20.8±1.0	8±0.7	40.7±4.4	28.3±9.2	18±0.2
	Wild-type Line 1	20.9±0.3	5±1.5	27.4±1.6	3.7±0.7	8±0.0
	Wild-type Line 2	22.0±0.6	4±1.2	31.7±3.4	2.5±0.6	8±0.0
Samsun	35:SVP Line 1	33.0±1.1	6±1.0	23.7±1.5	13.2±7.2	9±0.2
	35:SVP Line 2	42.0±0.8	7±1.4	26.7±3.0	12.1±6.3	11±0.4
	Wild-type Line 1	34.0±0.1	4±0.6	21.7±3.7	2.8±0.4	6±0.0
	Wild-type Line 2	36.0±1.5	4±1.6	24.7±3.8	3.0±0.2	6±0.0

Data represent the mean ±SE of six individuals for each genotype.

mechanism might be responsible for repression of *MYB110a* and *NtAn2* transcription and reduced anthocyanin accumulation in transgenic kiwifruit and tobacco flowers. The yeast two-hybrid assays showed that *SVP3* could interact with

kiwifruit AP1/SQUA proteins *FUL* and *FUL*-like, and this interaction may interfere with activation of R2R3 MYB transcription. Even stronger interactions were observed with *AG* and *SEP* MADS box proteins required for normal flower and




Fig. 5. Expression of kiwifruit *SVP3* in transgenic tobacco. (A) Phenotypic analysis of flower buds and open flowers. (B) Floral organ abnormalities: petaloid stamen (black arrow) and ovary position (white arrows). The cross-sections of whole flowers demonstrating colour change are presented in insets. (C) Relative expression of *NtAn1a*, *NtAn1b*, *NtAN2*, and *SVP3* transgene in petals normalized against *Ntα-Tub1*. Error bars represent the SE for three replicate reactions.

fruit development (Varkonyi-Gasic *et al.*, 2011). Therefore, interference with establishment of homeotic MADS box protein complexes, as previously shown in *Arabidopsis* (Fornara *et al.*, 2008; Gregis *et al.*, 2008; Wu *et al.*, 2012), is the likely mechanism by which *SVP3* interferes with various aspects of normal reproductive development.

Further functional studies using transgenic kiwifruit and tobacco generated in the course of this study will aid in establishing direct targets and protein partners of *SVP3*, providing understanding of the pathways and plant processes regulated by *SVP3*. These plants also provide an exciting opportunity for deeper understanding of the regulation of anthocyanin-related MYB expression in flowers.

Supplementary data

Supplementary data are available at *JXB* online.

[Figure S1.](#) *SVP3* in *Actinidia eriantha*.

[Figure S2.](#) LC-MS analysis of flavonoid compounds in petals of 35:*SVP3* and control *A. eriantha*.

[Figure S3.](#) Analysis of transgenic 35:*SVP3* tobacco.

Acknowledgements

The authors wish to thank Andrew Dare for help with HPLC analysis, Ian Hallett and Paul Sutherland for microscopy, Tim Holmes and Darren Snaith for photography, Sakuntala Karunairtnam and Andrew Gleave for cloning support, and Monica Dragulescu and Wade Wadasinghe for maintenance of

plants in the glasshouse. Thanks to Mirco Montefiori and Richard Espley for providing valuable advice and critically reading this manuscript. This work was funded by the New Zealand Foundation for Research, Science and Technology, contract C10X0816 MeriNET.

References

- Aswath CR, Kim SH. 2005. Another story of MADS-box genes—their potential in plant biotechnology. *Plant Growth Regulation* **46**, 177–188.
- Bai Y, Pattanaik S, Patra B, Werkman JR, Xie CH, Yuan L. 2011. Flavonoid-related basic helix–loop–helix regulators, *NtAn1a* and *NtAn1b*, of tobacco have originated from two ancestors and are functionally active. *Planta* **234**, 363–375.
- Bielenberg DG, Wang Y, Li ZG, Zhebentyayeva T, Fan SH, Reighard GL, Scorza R, Abbott AG. 2008. Sequencing and annotation of the evergrowing locus in peach [*Prunus persica* (L.) Batsch] reveals a cluster of six MADS-box transcription factors as candidate genes for regulation of terminal bud formation. *Tree Genetics and Genomes* **4**, 495–507.
- Borner R, Kampmann G, Chandler J, Gleißner R, Wisman E, Apel K, Melzer S. 2008. A MADS domain gene involved in the transition to flowering in *Arabidopsis*. *The Plant Journal* **24**, 591–599.
- Carmona MJ, Ortega N, Garcia-Maroto F. 1998. Isolation and molecular characterization of a new vegetative MADS-box gene from *Solanum tuberosum* L. *Planta* **207**, 181–188.
- Chang SJ, Puryear J, Cairney J. 1993. A simple and efficient method for isolating RNA from pine trees. *Plant Molecular Biology Reporter* **11**, 113–116.
- Cohen O, Borovsky Y, David-Schwartz R, Paran I. 2012. *CaJOINTLESS* is a MADS-box gene involved in suppression of vegetative growth in all shoot meristems in pepper. *Journal of Experimental Botany* **63**, 4947–4957.
- Cooke JEK, Eriksson ME, Junttila O. 2012. The dynamic nature of bud dormancy in trees: environmental control and molecular mechanisms. *Plant, Cell and Environment* **35**, 1707–1728.

- Dare AP, Tomes S, Jones M, McGhie TK, Stevenson DE, Johnson RA, Greenwood DR, Hellens RP.** 2013. Phenotypic changes associated with RNA interference silencing of chalcone synthase in apple (*Malus domestica*). *The Plant Journal* **74**, 398–410.
- Diaz-Riquelme J, Lijavetzky D, Martinez-Zapater JM, Carmona MJ.** 2009. Genome-wide analysis of MIKCC-type MADS box genes in grapevine. *Plant Physiology* **149**, 354–369.
- Fornara F, Gregis V, Pelucchi N, Colombo L, Kater M.** 2008. The rice *StMADS11*-like genes *OsMADS22* and *OsMADS47* cause floral reversions in *Arabidopsis* without complementing the *svp* and *agl24* mutants. *Journal of Experimental Botany* **59**, 2181–2190.
- Fraser LG, Seal AG, Montefiori M, et al.** 2013. An R2R3 MYB transcription factor determines red petal colour in an *Actinidia* (kiwifruit) hybrid population. *BMC Genomics* **14**, 28.
- Garcia-Maroto F, Ortega N, Lozano R, Carmona MJ.** 2000. Characterization of the potato MADS-box gene *STMADS16* and expression analysis in tobacco transgenic plants. *Plant Molecular Biology* **42**, 499–513.
- Gleave AP.** 1992. A versatile binary vector system with a T-DNA organizational-structure conducive to efficient integration of cloned DNA into the plant genome. *Plant Molecular Biology* **20**, 1203–1207.
- Gregis V, Andrés F, Sessa A, et al.** 2013. Identification of pathways directly regulated by SHORT VEGETATIVE PHASE during vegetative and reproductive development in *Arabidopsis*. *Genome Biology* **14**, R56.
- Gregis V, Sessa A, Colombo L, Kater MM.** 2008. *AGAMOUS-LIKE24* and *SHORT VEGETATIVE PHASE* determine floral meristem identity in *Arabidopsis*. *The Plant Journal* **56**, 891–902.
- Gregis V, Sessa A, Dorca-Fornell C, Kater MM.** 2009. The *Arabidopsis* floral meristem identity genes *AP1*, *AGL24* and *SVP* directly repress class B and C floral homeotic genes. *The Plant Journal* **60**, 626–637.
- Hartmann U, Hohmann S, Nettesheim K, Wisman E, Saedler H, Huijser P.** 2000. Molecular cloning of *SVP*: a negative regulator of the floral transition in *Arabidopsis*. *The Plant Journal* **21**, 351–360.
- He C, Tian Y, Saedler R, Efremova N, Riss S, Khan MR, Yephremov A, Saedler H.** 2010. The MADS-domain protein MPF1 of *Physalis floridana* controls plant architecture, seed development and flowering time. *Planta* **231**, 767–777.
- Horsch R, Fry J, Hoffmann N, Eichholtz D, Rogers SG, Fraley R.** 1985. A simple and general method for transferring genes into plants. *Science* **227**, 1229–1231.
- Horvath DP, Sung S, Kim D, Chao W, Anderson J.** 2010. Characterization, expression and function of *DORMANCY ASSOCIATED MADS-BOX* genes from leafy spurge. *Plant Molecular Biology* **73**, 169–179.
- Jaakola L, Poole M, Jones MO, et al.** 2010. A *SQUAMOSA* MADS box gene involved in the regulation of anthocyanin accumulation in bilberry fruits. *Plant Physiology* **153**, 1619–1629.
- Jaudal M, Monash J, Zhang L, Wen J, Mysore KS, Macknight R, Putterill J.** 2013. Overexpression of *Medicago SVP* genes causes floral defects and delayed flowering in *Arabidopsis* but only affects floral development in *Medicago*. *Journal of Experimental Botany* **65**, 429–442.
- Lalusin A, Nishita K, Kim S-H, Ohta M, Fujimura T.** 2006. A new MADS-box gene (*lbMADS10*) from sweet potato (*Ipomoea batatas* (L.) Lam) is involved in the accumulation of anthocyanin. *Molecular Genetics and Genomics* **275**, 44–54.
- Lalusin A, Ocampo E, Fujimura T.** 2011. *Arabidopsis thaliana* plants over-expressing the *lbMADS10* gene from sweet potato accumulate high level of anthocyanin. *Philippine Journal of Crop Science* **36**, 30–36.
- Lee JH, Yoo SJ, Park SH, Hwang I, Lee JS, Ahn JH.** 2007. Role of *SVP* in the control of flowering time by ambient temperature in *Arabidopsis*. *Genes and Development* **21**, 397–402.
- Li ZG, Reighard GL, Abbott AG, Bielenberg DG.** 2009. Dormancy-associated MADS genes from the *EVG* locus of peach [*Prunus persica* (L.) Batsch] have distinct seasonal and photoperiodic expression patterns. *Journal of Experimental Botany* **60**, 3521–3530.
- Li ZM, Zhang JZ, Mei L, Deng XX, Hu CG, Yao JL.** 2010. *PtSVP*, an *SVP* homolog from trifoliolate orange (*Poncirus trifoliata* L. Raf.), shows seasonal periodicity of meristem determination and affects flower development in transgenic *Arabidopsis* and tobacco plants. *Plant Molecular Biology* **74**, 129–142.
- Lichtenthaler HK, Rinderle U.** 1988. The role of chlorophyll fluorescence in the detection of stress conditions in plants. *CRC Critical Reviews in Analytical Chemistry* **19**, 29–85.
- Liu C, Xi WY, Shen LS, Tan CP, Yu H.** 2009. Regulation of floral patterning by flowering time genes. *Developmental Cell* **16**, 711–722.
- Liu C, Zhou J, Bracha-Drori K, Yalovsky S, Ito T, Yu H.** 2007. Specification of *Arabidopsis* floral meristem identity by repression of flowering time genes. *Development* **134**, 1901–1910.
- Mao L, Begum D, Chuang H, Budiman MA, Szymkowiak EJ, Irish EE, Wing RA.** 2000. *JOINTLESS* is a MADS-box gene controlling tomato flower abscission zone development. *Nature* **406**, 910–912.
- Masiero S, Li MA, Will I, Hartmann U, Saedler H, Huijser P, Schwarz-Sommer Z, Sommer H.** 2004. *INCOMPOSITA*: a MADS-box gene controlling prophyll development and floral meristem identity in *Antirrhinum*. *Development* **131**, 5981–5990.
- Mazzitelli L, Hancock RD, Haupt S, et al.** 2007. Co-ordinated gene expression during phases of dormancy release in raspberry (*Rubus idaeus* L.) buds. *Journal of Experimental Botany* **58**, 1035–1045.
- McGhie TK, Hudault SB, Lunken RC, Christeller JT.** 2011. Apple peels, from seven cultivars, have lipase-inhibitory activity and contain numerous urserenoic acids as identified by LC-ESI-QTOF-HRMS. *Journal of Agricultural and Food Chemistry* **60**, 482–491.
- Méndez-Vigo B, Martínez-Zapater JM, Alonso-Blanco C.** 2013. The flowering repressor *SVP* underlies a novel *Arabidopsis thaliana* QTL interacting with the genetic background. *PLoS Genetics* **9**, e1003289.
- Montefiori M, Easley RV, Stevenson D, Cooney J, Datson PM, Saiz A, Atkinson RG, Hellens RP, Allan AC.** 2011. Identification and characterisation of F3GT1 and F3GGT1, two glycosyltransferases responsible for anthocyanin biosynthesis in red-fleshed kiwifruit (*Actinidia chinensis*). *The Plant Journal* **65**, 106–118.
- Murashige T, Skoog F.** 1962. A revised medium for rapid assays with tobacco tissue cultures. *Physiologia Plantarum* **15**, 473–497.
- Pattanaik S, Kong Q, Zaitlin D, Werkman JR, Xie CH, Patra B, Yuan L.** 2010. Isolation and functional characterization of a floral tissue-specific R2R3 MYB regulator from tobacco. *Planta* **231**, 1061–1076.
- Sasaki R, Yamane H, Ooka T, Jotatsu H, Kitamura Y, Akagi T, Tao R.** 2011. Functional and expression analyses of *PmDAM* genes associated with endodormancy in Japanese apricot (*Prunus mume*). *Plant Physiology* **157**, 485–497.
- Shin MR, Seo SG, Kim JS, Joen SB, Kang SW, Lee GP, Kwon SY, Kim SH.** 2011. Alteration of floral organ identity by over-expression of *lbMADS3-1* in tobacco. *Transgenic Research* **20**, 365–376.
- Smaczniak C, Immink RGH, Angenent GC, Kaufmann K.** 2012. Developmental and evolutionary diversity of plant MADS-domain factors: insights from recent studies. *Development* **139**, 3081–3098.
- Trevaskis B, Tadege M, Hemming MN, Peacock WJ, Dennis ES, Sheldon C.** 2007. *Short Vegetative Phase*-like MADS-box genes inhibit floral meristem identity in barley. *Plant Physiology* **143**, 225–235.
- Varkonyi-Gasic E, Moss SM, Voogd C, Wu R, Lough RH, Wang Y-Y, Hellens RP.** 2011. Identification and characterization of flowering genes in kiwifruit: sequence conservation and role in kiwifruit flower development. *BMC Plant Biology* **11**, 72.
- Varkonyi-Gasic E, Moss S, Voogd C, Wang T, Putterill J, Hellens RP.** 2013. Homologs of *FT*, *CEN* and *FD* respond to developmental and environmental signals affecting growth and flowering in the perennial vine kiwifruit. *New Phytologist* **198**, 732–746.
- Wall C, Dozier W, Ebel RC, Wilkins B, Woods F, Foshee W.** 2008. Vegetative and floral chilling requirements of four new kiwi cultivars of *Actinidia chinensis* and *A. deliciosa*. *Hortscience* **43**, 644–647.
- Wang T, Atkinson R, Janssen B.** 2007. Choice of *Agrobacterium* strain for transformation of kiwifruit. *Acta Horticulturae* **753**, 227–232.
- Wang T, Ran Y, Atkinson RG, Gleave AP, Cohen D.** 2006. Transformation of *Actinidia eriantha*: a potential species for functional genomics studies in *Actinidia*. *Plant Cell Reports* **25**, 425–431.

- Wu RM, Walton EF, Richardson AC, Wood M, Hellens RP, Varkonyi-Gasic E.** 2012. Conservation and divergence of four kiwifruit SVP-like MADS-box genes suggest distinct roles in kiwifruit bud dormancy and flowering. *Journal of Experimental Botany* **63**, 797–807.
- Yamane H, Kashiwa Y, Ooka T, Tao R, Yonemori K.** 2008. Suppression subtractive hybridization and differential screening reveals endodormancy-associated expression of an SVP/AGL24-type MADS-box gene in lateral vegetative buds of Japanese apricot. *Journal of the American Society for Horticultural Science* **133**, 708–716.
- Yamane H, Ooka T, Jotatsu H, Hosaka Y, Sasaki R, Tao R.** 2011. Expressional regulation of *PpDAM5* and *PpDAM6*, peach (*Prunus persica*) dormancy-associated MADS-box genes, by low temperature and dormancy-breaking reagent treatment. *Journal of Experimental Botany* **62**, 3481–3488.
- Yu H, Xu YF, Tan EL, Kumar PP.** 2002. *AGAMOUS-LIKE 24*, a dosage-dependent mediator of the flowering signals. *Proceedings of the National Academy of Sciences, USA* **99**, 16336–16341.