



Hearing health geography in England

Document Version

Final published version

[Link to publication record in Manchester Research Explorer](#)

Citation for published version (APA):

Tsimpida, MSc, FHEA, D., Kontopantelis, E., Ashcroft, D., & Panagioti, M. (2020). *Hearing health geography in England: findings from the English longitudinal study of ageing (ELSA) and evidence of a north-south divide*. Poster session presented at Annual Conference of the British Society of Audiology (BSA).

Citing this paper

Please note that where the full-text provided on Manchester Research Explorer is the Author Accepted Manuscript or Proof version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version.

General rights

Copyright and moral rights for the publications made accessible in the Research Explorer are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Takedown policy

If you believe that this document breaches copyright please refer to the University of Manchester's Takedown Procedures [<http://man.ac.uk/04Y6Bo>] or contact uml.scholarlycommunications@manchester.ac.uk providing relevant details, so we can investigate your claim.



D. Tsimpida, PhD (c), Prof E. Kontopantelis, PhD, Prof DM. Ashcroft, PhD, Dr M. Panagioti, PhD
Centre for Primary Care and Health Services Research, Institute for Health Policy and Organisation (IHPO)
Division of Population Health, Health Services Research and Primary Care, School of Health Sciences, The University of Manchester, UK

Aim

To explore the regional patterns and trends of hearing loss (HL) in a representative longitudinal prospective cohort study of the English population aged 50 years old and above.

Methods

Time-series analyses in the full dataset (8 Waves) of the English Longitudinal Study of Ageing (ELSA) [1]. The primary outcome measure was self-reported HL [2].

Local spatial analysis statistical tools used for analysing spatial distributions, patterns, processes and relationships of the geographical data.

The Spatial Join tool used to aggregate the number of cases of self-reported HL to total responses of hearing acuity in each polygon (GOR) to visualise the prevalence rates of HL per GOR.

We computed Adjusted Predictions at the Means (APMs) and the Marginal Effects at the Means (MEMs) [3] of the HL prevalence in each ELSA Wave, with age, gender, education, occupation, income, wealth, IMD and alcohol consumption as the factor variables.

Results

Between 2002-2017 there was an estimated increase of **10.2%** in the total HL prevalence in the English older population. The mean HL prevalence increased from 38.50 (95%CI 37.37-39.14) in Wave 1 to 48.66 (95%CI 47.11-49.54) in Wave 8. (**Figure 1**)

A wide variation in HL prevalence was revealed in representative samples from different regions in England that had significantly equal age, and the increase rate of HL ranged in the 15-year period from 3.2% to 45%. (**Figure 2**)

The highest HL prevalence was observed in GORs with the highest prevalence of participants in the categories of a) most deprived (IMD) quintile, b) those in routine or manual occupations, and c) those with the highest prevalence of alcohol misuse (>14 units/week).

The Hot Spot and Cold Spot analyses showed marked regional variability and evidence of a North-South divide. (**Figure 3**)




Figure 2. Predicted probabilities (%) of hearing loss prevalence at Regions of England in eight Waves (74,699 person-years) of the English Longitudinal Study of Ageing (ELSA)*
*Holding factor variables (age, gender, education, occupation, income, wealth, IMD and alcohol consumption) at their means, for each ELSA Wave.

Summary

- This study is the first that investigated geographical patterns and trends of HL based on a representative cohort of older adults and not through population projections.
- Our study showed that the increase in HL prevalence is not related to the ageing of the population, as widely believed, but potentially due to social and lifestyle changes.
- A socio-spatial approach is crucial for planning sustainable models of hearing care based on the actual needs, and for reducing hearing health inequalities.




Figure 1. Map of England by Government Office Regions, showing prevalence rates of self-reported hearing loss in eight Waves (74,699 person-years) of the English Longitudinal Study of Ageing (ELSA) 2002-2017 (responses collected every 2 years).




Figure 3. Map of England by Government Office Regions showing the spatial clusters of hearing loss prevalence according to Hot Spot and Cold Spot analyses using the Getis-Ord Gi* statistic in eight Waves (74,699 person-years) of the English Longitudinal Study of Ageing (ELSA), 2002-2017.

Conclusions

- HL is highlighted as an increasingly important public health issue in England and globally. A spatial dimension is now being added which shows that the increasing trend in HL prevalence is not age-related, as widely believed [4, 5].
- The Clinical Commissioning Groups (CCGs) who are currently responsible for the NHS audiology services in England should take into consideration socioeconomic factors and modifiable lifestyle behaviours for HL, along with their spatial patterning in England.

References

- [1] Steptoe A. et al. *Int J Epidemiol.* 2013 Dec 1;42(6):1640-8.
- [2] Tsimpida D. et al. *JAMA Network Open* 2020 Aug 3;3(8):e2015009.
- [3] Williams R. *STATA Journal.* 2012; 12:308-31.
- [4] Tsimpida D. et al. *BMJ open.* 2019 Sep 1;9(9):e031030.
- [5] Akeroyd MA et al. *Trends in Hearing.* 2019 Dec;23: 2331216519887614.

Acknowledgments

This research was supported by the NIHR Manchester Biomedical Research Centre (PhD Studentship to DT – personal award reference: NIHR-INF-0551). The views expressed are those of the authors and not necessarily those of the BRC, the NIHR or the Department of Health.