



Queensland University of Technology
Brisbane Australia

This is the author's version of a work that was submitted/accepted for publication in the following source:

Yong, Mostyn R., Savi, Flavia M., Woodruff, Maria A., Saifzadeh, Siamak, Askin, Geoffrey N., Labrom, Robert D., Hutmacher, Dietmar, & Adam, Clayton J.

(2014)

Longitudinal performance of polycaprolactone-based scaffold plus recombinant human morphogenetic protein-2 (rhBMP-2) in large preclinical animal model : 6- versus 12 months. In

Raso, James & Aubin, Carl-Eric (Eds.)

The 10th Meeting of the International Research Society of Spinal Deformities (IRSSD 2014), 29 June - 2 July 2014, Sapporo, Japan.

This file was downloaded from: <http://eprints.qut.edu.au/79636/>

© Copyright 2015 Yong et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

Notice: *Changes introduced as a result of publishing processes such as copy-editing and formatting may not be reflected in this document. For a definitive version of this work, please refer to the published source:*

<http://dx.doi.org/10.1186/1748-7161-10-S1-O58>

LONGITUDINAL PERFORMANCE OF POLYCAPROLACTONE-BASED SCAFFOLD PLUS RECOMBINANT HUMAN MORPHOGENETIC PROTEIN-2 (rhBMP-2) IN LARGE PRECLINICAL ANIMAL MODEL: 6- VERSUS 12 MONTHS

*Yong MR^{1,2}, Savi FM², Woodruff MA², Saifzadeh S², Askin GN^{1,2}, Labrom RD^{1,2}, Hutmacher DW², Adam CJ^{1,2}

1. QUT/Mater Paediatric Spine Research Group, Queensland University of Technology, Mater Misericordiae Health Services, Brisbane, QLD, Australia.

2. Institute of Health and Biomedical Innovation, Queensland University of Technology, Brisbane, QLD, Australia.

Introduction

There is strong current interest in the use of biodegradable scaffolds in combination with bone growth factors as a valuable alternative to the current gold standard autograft in spinal fusion surgery Yong et al. (2013). Here we report on 6- vs 12- month data set evaluating the longitudinal performance of a CaP coated polycaprolactone (PCL) scaffold loaded with recombinant human bone morphogenetic protein-2 (rhBMP-2) as a bone graft substitute within a preclinical ovine thoracic spine.

Methods

Twelve sheep underwent a 3-level (T6/7, T8/9 and T10/11) discectomy with randomly allocated implantation of a different graft substitute at each of the three levels; (i) calcium phosphate (CaP) coated polycaprolactone based scaffold plus 0.54µg rhBMP-2, (ii) CaP coated PCL- based scaffold alone or (iii) autograft (mulched rib head). Fusion assessments were performed via high resolution clinical computed tomography and histological evaluation were undertaken at six (n=6) and twelve (n=6) months post-surgery using the Sucato grading system (Sucato et al. 2004).

Results

The computed tomography fusion grades of the 6- and 12- months in the rhBMP-2 plus PCL-based scaffold group were 1.9 and 2.1 respectively, in the autograft group 1.9 and 1.3 respectively, and in the scaffold alone group 0.9 and 1.17 respectively. There were no statistically significant differences in the fusion scores between 6- and 12- month for the rhBMP plus PCL- based scaffold or PCL – based scaffold alone group however there was a significant reduction in scores in the autograft group. These scores were seen to correlate with histological evaluations of the respective groups.

Conclusions

The results of this study demonstrate the efficacy of scaffold-based delivery of rhBMP-2 in promoting higher fusion grades at 6- and 12- months in comparison to the scaffold alone or autograft group within the same time frame. Fusion grades achieved at six months using PCL+rhBMP-2 are not significantly increased at twelve months post surgery.

References

Sucato, D.J. et al. (2004) Recombinant Human Bone Morphogenetic Protein – 2 Enhances Anterior Spinal Fusion in a Thoracoscopically Instrumented Animal Model. *The Journal of Bone and Joint Surgery American Volume*, 86-A(4): 752-62.

Yong, M.R. et al. (2014) Biological Performance of a Polycaprolactone – based Scaffold plus Recombinant Human Morphogenetic Protein – 2 (rhBMP-2) in an Ovine Thoracic Interbody Fusion Model. *European Spine Journal*, 23: 650-7.