

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Agronomy & Horticulture -- Faculty Publications

Agronomy and Horticulture Department

---

3-15-2020

## Optimum rates of surface-applied coal char decreased soil ammonia volatilization loss

Dinesh Panday

*University of Nebraska-Lincoln*, [dinesh.panday@unl.edu](mailto:dinesh.panday@unl.edu)

Maysoon M. Mikha

*USDA-ARS, Central Great Plains Research Station, Akron, CO*, [maysoon.mikha@usda.gov](mailto:maysoon.mikha@usda.gov)

Harold P. Collins

*USDA-ARS, Grassland Soil and Water Research Laboratory, Temple, TX*, [hal.collins@usda.gov](mailto:hal.collins@usda.gov)

Virginia L. Jin

*USDA-ARS, Agroecosystem Management Research Unit, Lincoln, NE*, [virginia.jin@ars.usda.gov](mailto:virginia.jin@ars.usda.gov)

Michael Kaiser

*University of Nebraska - Lincoln*, [mkaiser6@unl.edu](mailto:mkaiser6@unl.edu)

*See next page for additional authors*

Follow this and additional works at: <https://digitalcommons.unl.edu/agronomyfacpub>



Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), [Agronomy and Crop Sciences Commons](#), [Botany Commons](#), [Horticulture Commons](#), [Other Plant Sciences Commons](#), and the [Plant Biology Commons](#)

---

Panday, Dinesh; Mikha, Maysoon M.; Collins, Harold P.; Jin, Virginia L.; Kaiser, Michael; Cooper, Jennifer; Malakar, Arindam; and Maharjan, Bijesh, "Optimum rates of surface-applied coal char decreased soil ammonia volatilization loss" (2020). *Agronomy & Horticulture -- Faculty Publications*. 1352.  
<https://digitalcommons.unl.edu/agronomyfacpub/1352>

This Article is brought to you for free and open access by the Agronomy and Horticulture Department at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Agronomy & Horticulture -- Faculty Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

---




**Authors**

Dinesh Panday, Maysoon M. Mikha, Harold P. Collins, Virginia L. Jin, Michael Kaiser, Jennifer Cooper, Arindam Malakar, and Bijesh Maharjan

## TECHNICAL REPORTS

## Atmospheric Pollutants and Trace Gases

# Optimum rates of surface-applied coal char decreased soil ammonia volatilization loss

Dinesh Panday<sup>1</sup>  | Maysoon M. Mikha<sup>2</sup> | Harold P. Collins<sup>3</sup> | Virginia L. Jin<sup>4</sup> | Michael Kaiser<sup>1</sup> | Jennifer Cooper<sup>1</sup> | Arindam Malakar<sup>5</sup>  | Bijesh Maharjan<sup>1</sup> 

<sup>1</sup>Dep. of Agronomy and Horticulture, Univ. of Nebraska-Lincoln, Lincoln, NE 68588, USA

<sup>2</sup>USDA-ARS, Central Great Plains Research Station, Akron, CO 80720, USA

<sup>3</sup>USDA-ARS, Grassland Soil and Water Research Laboratory, Temple, TX 76502, USA

<sup>4</sup>USDA-ARS, Agroecosystem Management Research Unit, Lincoln, NE 68583, USA

<sup>5</sup>Nebraska Water Center, Univ. of Nebraska-Lincoln, Lincoln, NE 68588, USA

## Correspondence

Bijesh Maharjan, Dep. of Agronomy and Horticulture, Univ. of Nebraska-Lincoln, Lincoln, NE 68588, USA.

Email: bmaharjan@unl.edu

Assigned to Associate Editor Upendra Sainju.

## Abstract

Fertilizer N losses from agricultural systems have economic and environmental implications. Soil amendment with high C materials, such as coal char, may mitigate N losses. Char, a coal combustion residue, obtained from a sugar factory in Scottsbluff, NE, contained 29% C by weight. A 30-d laboratory study was conducted to evaluate the effects of char addition on N losses via nitrous oxide (N<sub>2</sub>O) emission, ammonia (NH<sub>3</sub>) volatilization, and nitrate (NO<sub>3</sub>-N) leaching from fertilized loam and sandy loam soils. Char was applied at five different rates (0, 6.7, 10.1, 13.4, and 26.8 Mg C ha<sup>-1</sup>; char measured in C equivalent) to soils fertilized with urea ammonium nitrate (UAN) at 200 kg N ha<sup>-1</sup>. In addition, there were two negative-UAN control treatments: no char (no UAN) and char at 26.8 Mg C ha<sup>-1</sup> (no UAN). Treatment applied at 6.7 and 10.1 Mg C ha<sup>-1</sup> in fertilized sandy loam reduced NH<sub>3</sub> volatilization by 26–37% and at 6.7, 10.1, and 13.4 Mg C ha<sup>-1</sup> in fertilized loam soils by 24% compared with no char application. Nitrous oxide emissions and NO<sub>3</sub>-N leaching losses were greater in fertilized compared with unfertilized soil, but there was no effect of char amendment on these losses. Because NO<sub>3</sub>-N leaching loss was greater in sandy loam than in loam, soil residual N was twofold higher in loam than in sandy loam. This study suggests that adding coal char at optimal rates may reduce agricultural reactive N to the atmosphere by decreasing NH<sub>3</sub> volatilization from fertilized soils.

**Abbreviations:** C0N0, no char or urea ammonium nitrate; C0N1, no char and urea ammonium nitrate; C1N1, char rate at 6.7 Mg C ha<sup>-1</sup> and urea ammonium nitrate; C2N1, char rate at 10.1 Mg C ha<sup>-1</sup> and urea ammonium nitrate; C3N1, char rate at 13.4 Mg C ha<sup>-1</sup> and urea ammonium nitrate; C4N1, char rate at 26.8 Mg C ha<sup>-1</sup> and urea ammonium nitrate; C4N0, char rate at 26.8 Mg C ha<sup>-1</sup> and no urea ammonium nitrate; CCR, coal combustion residue; CEC, cation exchange capacity; CV, coefficient of variance; FNR, fertilizer N recovery; OM, organic matter; UAN, urea ammonium nitrate.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2019 The Authors. *Journal of Environmental Quality* published by American Society of Agronomy, Crop Science Society of America, and Soil Science Society of America and Wiley Periodicals, Inc.

## 1 | INTRODUCTION

Fertilizer nitrogen (N) use increased globally at an annual rate of 1.4% from 2014 to 2018 (IFASTAT, 2019). Generally, crop N uptake efficiency is <50% of applied N, which leaves a significant amount of N in soil prone to loss via NH<sub>3</sub> volatilization, NO<sub>3</sub>-N leaching, and/or denitrification as N<sub>2</sub>O emissions (Fageria & Baligar, 2005; Robertson et al., 2013). Nitrogen losses from agricultural systems can be a major limitation for crop production and environmental sustainability.

Numerous management technologies have been proposed to mitigate N losses from agricultural systems, including the proper management of soil C because of its effects on soil properties and processes, including N cycling (Dil, Oelbermann, & Xue, 2014; Ding et al., 2010). Carbon management practices that include amendments with high C content, such as biochar, can boost soil fertility and quality by raising pH and by improving water holding capacity, cation exchange capacity (CEC), and nutrient retention (Bridgwater, 2003; Filiberto & Gaunt, 2013; Singh et al., 2014).

Coal combustion residues (CCRs), such as fly ash, bottom ash, and flue gas desulfurization gypsum, have been used as soil amendments to improve soil health and crop performance (Basu, Pande, Bhadoria, & Mahapatra, 2009; Panday, Ferguson, & Maharjan, 2018; Shaheen, Hooda, & Tsadilas, 2014). However, depending on the composition and nature of CCR, they can enhance mineralization of organic soil N and N losses (Siddaramappa, McCarty, Wright, & Codling, 1994). The CCRs in electric power generating stations obtained from the near-complete combustion of coal during energy production contain very little C. In contrast, coal char (henceforth “char”) resulting from inefficient coal burning can contain up to 29% C by dry weight as well as other essential plant mineral nutrients.

Char stands midway between coal ash and biochar with respect to C content. Biochar and other hydrocarbons are typically produced from pyrolysis of biomass in the presence of little or no oxygen at a range of temperatures and can contain up to 70% of initial biomass C (Atkinson, Fitzgerald, & Hips, 2010; Lehmann, Gaunt, & Rondon, 2006). Biochar can reduce  $\text{NH}_3$  volatilization loss (Steiner, Das, Melear, & Lakly, 2010) and  $\text{NO}_3\text{-N}$  leaching loss (Hagemann, Kammann, Schmidt, Kappler, & Behrens, 2017). However, the beneficial effect of biochar in reducing environmental N losses from fertilized soil is not consistent and depends on sources and production conditions (Ding et al., 2016). Char, which is different from regular CCRs and biochar but has a considerable amount of C, warrants exploration for its potential use in agricultural soil.

The objectives of this study were to evaluate the effects of char on soil N losses in the form of  $\text{NH}_3$  volatilization,  $\text{N}_2\text{O}$  emissions, and  $\text{NO}_3\text{-N}$  leaching from fertilized loam and sandy loam soils. We hypothesized (a) that adding char would reduce N losses from fertilized soil by improving the retention of applied N and (b) that char effectiveness on retaining N would differ by soil type.

## 2 | MATERIALS AND METHODS

The char used in this study was a CCR from a sugar factory in Scottsbluff, NE, and contained 29.3% C and some nutrients (Supplemental Table S1). It also contained heavy metals (As, Cd, Cr, Pb, Hg, and Se), but their concentrations

### Core Ideas

- High C content coal char may reduce environmental N loss from fertilized soil.
- There are implications of using different methods in estimating fertilizer N recovery.
- Must evaluate industrial by-products in agriculture for potential accumulation of trace metals.

were below the USEPA’s ceiling limits for heavy metal soil contamination or phytotoxicity in soil (Cameron, 1992). Char was sieved through a 2-mm sieve. The physical characteristics of char were determined by X-ray diffraction using a PANalytical Empyrean Diffractometer (Malvern Panalytical Ltd.) at the Nebraska Center for Materials and Nanoscience (Supplemental Figure S2). Brunauer-Emmett-Teller surface area of char was analyzed with an ASAP 2460 Surface Area and Porosity Analyzer (Micromeritics Instrument Corporation) at the Nebraska Center for Materials and Nanoscience (Supplemental Table S3).

Two soils were used to evaluate the effects of char on N losses from fertilized soil at the Panhandle Research and Extension Center, University of Nebraska-Lincoln in Scottsbluff, NE. One soil was a Tripp fine sandy loam (coarse-silty, mixed, superactive, mesic Aridic Haplustolls, 0–3% slope) with pH 7.7; 13 g  $\text{kg}^{-1}$  organic matter (OM); and 60, 28, and 12% of sand, silt, and clay contents, respectively. This soil was collected from the Panhandle Research and Extension Center. The other soil was a Duroc loam (fine-silty, mixed, mesic Pachic Haplustolls, 0–1% slope) with pH 7.2; 18 g  $\text{kg}^{-1}$  OM; and 40, 33, and 27% of sand, silt, and clay, respectively. This soil was collected from farmland near the University of Nebraska-Lincoln High Plains Agricultural Laboratory in Cheyenne, NE. Both soils were collected at depths of 0–20 cm in the spring of 2018. Residual inorganic N rates, extracted with 2 M KCl, in loam and sandy loam soils were 5.2 and 3.7 mg  $\text{kg}^{-1}$ , respectively.

Collected soils were air-dried and sieved through a 2-mm mesh. Soils were brought to 10% gravimetric water content (GWC) by applying water and mixing thoroughly, which corresponded to 70 and 50% of field capacity of sandy loam and loam, respectively. Soils were packed in 5-cm-diameter clear acrylic columns (Supplemental Figure S4) to a height of 24 cm with a targeted bulk density of 1400  $\text{kg m}^{-3}$  (Peng et al., 2015). A porous ceramic plate (0.1 MPa strength) was inserted in the bottom of the column and topped with Whatman no. 42 filter paper to prevent soil from clogging the ceramic plate. Soil columns had lid systems at either end. A vacuum port located on the bottom lid allowed suction to be applied during the collection of leachate. The top lid has two parts (lower and

upper). The lower lid part is an elongated connector (height, 5 cm) threaded onto the main column and the upper lid, which was used to install the  $\text{NH}_3$  acid trap. The upper lid part (height, 5 cm) terminates the column with a closed end fitted with a septum port for  $\text{N}_2\text{O}$  sampling from the headspace above the soil.

Char (measured in C equivalent) and UAN were applied to each soil column and mixed in the top 6-cm soil layer. There were seven treatments, each with four replications: (a) C0N0, no char or UAN; (b) C0N1, no char and UAN; (c) C1N1, char rate at  $6.7 \text{ Mg C ha}^{-1}$  and UAN; (d) C2N1, char rate at  $10.1 \text{ Mg C ha}^{-1}$  and UAN; (e) C3N1, char rate at  $13.4 \text{ Mg C ha}^{-1}$  and UAN; (f) C4N1, char rate at  $26.8 \text{ Mg C ha}^{-1}$  and UAN; and (g) C4N0, char rate at  $26.8 \text{ Mg C ha}^{-1}$  and no UAN. All treatments that were fertilized (CxN1) received  $39.5 \times 10^{-3} \text{ g UAN-N}$  that was equivalent to  $200 \text{ kg N ha}^{-1}$ .

After soil columns were prepared, water was periodically added to simulate rainfall (100.8 mm in total) in May 2017 in Scottsbluff, NE (Supplemental Figure S5). Water was added slowly on the surface of soil using a syringe to prevent ponding on the surface. Columns were kept on the laboratory benchtop at constant room temperature ( $25^\circ\text{C}$ ) throughout the 30-d experimental period.

## 2.1 | Sample collection

Ammonia volatilization was measured using an acid trap method (McGinn & Janzen, 1998). The acid trap was made up of a sponge (diameter, 5 cm; thickness, 1.3 cm) with 5 ml of  $\text{H}_3\text{PO}_4$ -glycerol solution (40 ml glycerol, 50 ml  $\text{H}_3\text{PO}_4$  acid, and 910 ml deionized water) placed inside the lower part of the column top lid. The acid traps were installed on Day 0 after all treatments were applied to soil. All  $\text{NH}_3$  traps were exchanged with fresh ones on Days 1, 2, 3, 5, 7, 9, 11, 13, 17, 21, and 25. Each used trap was thoroughly rinsed in 2 M KCl solution and squeezed several times to extract the solution. The collected extracts were analyzed for  $\text{NH}_4\text{-N}$  using a flow injection method (Ahmed, Stalikas, Tzouwara-Karayanni, Karayannis, & Veltsistas, 1997). Cumulative  $\text{NH}_3$  loss was calculated by summing  $\text{NH}_4\text{-N}$  across all collection dates. Cumulative  $\text{NH}_3$  loss was converted to  $\text{kg N ha}^{-1}$  by multiplying the total volatilization loss and the given soil surface area.

Nitrous oxide emissions were measured by collecting gas samples through the septum port on the upper terminal lid. Gas samples were collected on alternate days (Days 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, and 29). During the  $\text{N}_2\text{O}$  sampling period, the  $\text{NH}_3$  trap was removed from the column, which gave a headspace of  $315 \text{ cm}^3$ . Gas samples were collected at 0, 10, and 20 min using a 12-ml syringe. The 0-min samples were collected before closing the lid. At each sampling, gas was transferred to a 10-ml glass sample vial (Wheaton). Samples were analyzed for  $\text{N}_2\text{O}$

concentrations with a gas chromatograph (450-GC, Varian) using an electron capture detector. The  $\text{N}_2\text{O}$  concentration values were converted to mass per volume using the universal gas law equation. Daily gas flux rates ( $\text{mg m}^{-2} \text{ min}^{-1}$ ) were calculated as the linear or quadratic change in headspace  $\text{N}_2\text{O}$  concentration over time (Wagner, Reicosky, & Alessi, 1997) based on regression analysis with the highest  $r^2$  value. Cumulative  $\text{N}_2\text{O}$  emissions ( $\text{kg N ha}^{-1}$ ) were determined by integrating daily  $\text{N}_2\text{O}$  fluxes using the trapezoidal integration method (Dunmola, Tenuta, Moulin, Yapa, & Lobb, 2010).

An attempt was made to collect column leachate on each day after water addition. On each collection date, suction with a 0.25-horsepower air motor (Model 1603007402, Bluffton Motor Works) was applied to the bottom lid of the column to facilitate drainage of water collected at the bottom of soil column through a porous ceramic plate (Peng et al., 2015). Leachate samples were frozen until analysis for  $\text{NO}_3\text{-N}$  using a flow injection method (Ahmed et al., 1997). The total amount of  $\text{NO}_3\text{-N}$  leached in each treatment was calculated by multiplying  $\text{NO}_3\text{-N}$  concentration with leachate volume and summing over collection dates.

All samplings were done in the morning (8:00 a.m.–12:00 p.m.). At the end of the experiment, the porous ceramic plate was removed from the bottom of the soil column, and soil was divided into 6-cm increments. For each increment, 10 g of soil was collected for determination of GWC, and the remaining soil was analyzed for  $\text{NH}_4\text{-N}$  and  $\text{NO}_3\text{-N}$  concentrations. Soil residual inorganic N was calculated as the sum of  $\text{NH}_4\text{-N}$  and  $\text{NO}_3\text{-N}$  concentrations across all soil increments for each column.

## 2.2 | Data analysis

The N losses via  $\text{NH}_3$  volatilization,  $\text{NO}_3\text{-N}$  leaching, and  $\text{N}_2\text{O}$  emissions and soil residual N in unfertilized treatment (C0N0) were subtracted from those in fertilized treatments and divided by the amount of UAN-N applied (i.e., 39.5 mg N) to estimate those losses per applied UAN-N. Fertilizer N recovery (FNR) was estimated by two methods. Equation 1 represents the “N difference” method, where N losses and residual N at the end of the experiment in control treatment (C0N0) were subtracted from those in fertilized treatment to estimate FNR based on “N difference” method ( $\text{FNR}_{\text{CTRL}}$ ) (adapted from Mahal et al. [2019]). Equation 2 estimated FNR based on the initial extractable N ( $\text{FNR}_{\text{ResN}}$ ), which accounted for initial extractable N at the beginning of the experiment (adapted from Li, Hu, Delgado, Zhang, & Ouyang, 2007).

$$\text{FNR}_{\text{CTRL}} = \frac{(\text{N loss}_{\text{Treatment}} - \text{N loss}_{\text{C0N0}}) + (\text{Soil residual N}_{\text{Treatment}} - \text{Soil residual N}_{\text{C0N0}})}{\text{Applied N}} \times 100 \quad (1)$$

$$\text{FNR}_{\text{ResN}} = \frac{\text{N loss}_{\text{Treatment}} + \text{Soil residual N}_{\text{Treatment}}}{(\text{Applied N} + \text{Initial extractable N})} \times 100 \quad (2)$$

The effects of treatment and soil on dependent variables' cumulative values were tested using the PROC MIXED procedure in SAS, with treatment, soil, and their interaction as the fixed effects and rep as random effect (Littell, Milliken, Stroup, Wolfinger, & Schabenberger, 2006; SAS, 2015). When main or interaction effects were significant, means were separated by the LSD test (Littell et al., 2006). Ammonia volatilization and N<sub>2</sub>O emissions data were analyzed using repeated measures in ANOVA to determine the differences among treatments by sampling dates. Statistical significance was evaluated at  $P < .05$  unless otherwise stated.

### 3 | RESULTS

#### 3.1 | Ammonia volatilization

Daily NH<sub>3</sub> volatilization loss with the C4N1 treatment was higher than with other treatments in the first 10 acid trap sample collection dates ( $n = 12$ ) in loamy soil (Figure 1a). The same was true for sandy loam on five different sampling dates (Figure 1b). After Day 17, all treatments showed no or minimal volatilization loss in both soil types. In fertilized treatments, all daily NH<sub>3</sub> losses were >2% of applied N and occurred within the first 2 wk of the experiment, and losses were >1% by the third week in both soil types.

Cumulative NH<sub>3</sub> loss across treatments ranged from 0.2 to 9.1 mg (equivalent to 1.0–46.4 kg N ha<sup>-1</sup>) in loam and from 0.2 to 6.9 mg (equivalent to 1.0–35.2 kg N ha<sup>-1</sup>) in sandy loam soils. There was a significant treatment × soil interaction effect on cumulative NH<sub>3</sub> loss and cumulative NH<sub>3</sub> loss per applied N (Tables 1 and 2). Compared with C0N1, cumulative NH<sub>3</sub> loss (per applied N) was significantly lower for C1N1, C2N1, and C3N1 in loam soil and for C1N1 and C2N1 in sandy loam soil (Table 2). The C3N1 and C4N1 in sandy loam and C4N1 in loam increased NH<sub>3</sub> loss (per applied N) compared with C0N1. The C0N0 and C4N1 had minimal NH<sub>3</sub> losses in both soil types (Figure 1). Among fertilized treatments (C0N1, C1N1, C2N1, C3N1, and C4N1), cumulative NH<sub>3</sub> loss per applied N ranged from 3.2 to 22.3% in loam and from 6.6 to 16.8% in sandy loam soils.

#### 3.2 | Nitrous oxide emissions

Daily N<sub>2</sub>O fluxes varied from 0 to 0.4 mg m<sup>-2</sup> h<sup>-1</sup> in loam and were 0.3 mg m<sup>-2</sup> h<sup>-1</sup> in the sandy loam soil across treatments throughout the experiment (Figure 2). Variability

in daily N<sub>2</sub>O fluxes was high among replications in both loam (coefficient of variance [CV], 32.1–166.1%) and sandy loam (CV, 12.1–176.2%). Of the 15 sampling dates, C0N1 had the highest daily N<sub>2</sub>O flux on the final sampling date in loam and on Days 7 and 9 in sandy loam. Control treatments always had minimal N<sub>2</sub>O fluxes in both soil types.

Cumulative N<sub>2</sub>O emissions differed by treatment but did not differ by soil type or their interaction (Table 1). Emissions were greater in fertilized treatments compared with unfertilized treatments at  $P < .001$ . Cumulative N<sub>2</sub>O emissions among fertilized treatments were not different. Averaged cumulative N<sub>2</sub>O emissions in fertilized treatments were 0.7 kg N ha<sup>-1</sup> in both soil types and 0.03 and 0.05 kg N ha<sup>-1</sup> in unfertilized loam and sandy loam, respectively (Supplemental Figure S6). Among fertilized treatments (C0N1, C1N1, C2N1, C3N1, and C4N1), cumulative N<sub>2</sub>O emissions per applied N ranged from 0.1 to 0.5% in loam and from 0.1 to 0.4% in sandy loam soils.

#### 3.3 | Nitrate leaching

In loam, one leaching event occurred on Day 29 after N fertilization across all treatments. In contrast, three leaching events occurred in sandy loam (Days 20, 21, and 29), with 44.3% of the total NO<sub>3</sub>-N leaching observed on Day 29 (Figure 3).

There was a significant treatment × soil interaction effect on cumulative NO<sub>3</sub>-N leaching (Tables 1 and 2). Cumulative NO<sub>3</sub>-N leaching was consistently greater for all fertilized treatments in sandy loam than in loam (Table 2). Averaged across all treatments, cumulative NO<sub>3</sub>-N leaching was almost fourfold greater for sandy loam ( $17.6 \times 10^{-3}$  g) than for loam ( $4.3 \times 10^{-3}$  g) (Table 1).


Among fertilized treatments, cumulative NO<sub>3</sub>-N leaching per applied N was higher in sandy loam (32.4%) than in loam (2.6%) (Table 1). In sandy loam, C3N1 had lower NO<sub>3</sub>-N leaching ( $16.9 \times 10^{-3}$  g or 21.1% of applied N) than C0N1 ( $24.3 \times 10^{-3}$  g or 39.9% of applied N) (Table 2).

#### 3.4 | Soil residual mineral nitrogen and fertilizer nitrogen recovery

There was a significant treatment × soil interaction effect on soil residual mineral N throughout the column (Table 1). Control treatments (C0N0 and C4N0) had lower soil residual mineral N than fertilized treatments in both soil types (Table 2). Among fertilized treatments, soil residual mineral N was similar in sandy loam but was significantly lower in C4N1 ( $26.4 \times 10^{-3}$  g or 49.8% of applied N) than in the other treatments in loam (Table 2).

When separated by depth, soil residual N was greater in fertilized treatments than in the control treatments at 18–24 cm





**FIGURE 1** Daily and cumulative NH<sub>3</sub> volatilization loss (mean  $\pm$  SE) with different treatments in (a, c) loam and (b, d) sandy loam soils. CON0, no char and no urea ammonium nitrate (UAN); C0N1–C4N1, UAN at 200 kg N ha<sup>-1</sup> and char at 0, 6.7, 10.1, 13.4, and 26.8 Mg C ha<sup>-1</sup>, respectively; C4N0, 26.8 Mg C ha<sup>-1</sup> and no UAN. \*Treatment with significantly higher loss than all other treatments on a given sampling day

in both soil types. Fertilized treatments (C0N1, C1N1, C2N1, C3N1, and C4N1) in loam had greater residual N than the control treatments (CON0 and C4N0) at other depths as well. Soil residual mineral N at 18–24 cm was higher with C1N1 and C3N1 in loam soil than other treatments in both soil types (Figure 4). In loam, C4N1 had lower soil residual N at 18–24 cm than other fertilized treatments. In sandy loam, soil residual N were greater with C3N1 than other treatments

except C1N1. Among fertilized treatments in sandy loam, C4N1 and C0N1 had lower soil residual N than others.

There were no significant differences by treatment or soil in fertilizer N recovery (Table 1). The  $FNR_{CTRL}$  ranged from 67.6 to 77.3% by soil type and from 69.0 to 74.2% by treatment. The UAN-N applied among fertilized treatments (C0N1, C1N1, C2N1, C3N1, and C4N1) that remained unaccounted ranged from 26.3 to 34.4%. However,  $FNR_{ResN}$

**TABLE 1** Analysis of variance results with means for different dependent variables as affected by char, soil, and their interaction

Treatment <sup>a</sup>	NH <sub>3</sub> volatilized		N <sub>2</sub> O emissions		NO <sub>3</sub> -N leached		Soil residual mineral N		FNR <sub>CTRL</sub> <sup>b</sup>	FNR <sub>ResN</sub> <sup>c</sup>
	g (10 <sup>-3</sup> )	% per applied N	g (10 <sup>-3</sup> )	% per applied N	g (10 <sup>-3</sup> )	% per applied N	g (10 <sup>-3</sup> )	% per applied N		
	%									
C0N0	0.4	–	0.01b	–	6.1	–	6.2	–	–	–
C0N1	4.0	9.2	0.12a	0.28	14.3	20.9	22.8	42.1	72.5	97.3
C1N1	2.8	6.2	0.11a	0.28	12.2	15.4	26.4	51.2	73.1	97.8
C2N1	3.0	6.8	0.15a	0.34	14.1	20.4	24.6	46.7	74.2	95.0
C3N1	3.9	8.9	0.10a	0.25	10.8	11.9	25.1	48.0	69.0	94.0
C4N1	7.1	17.1	0.15a	0.36	13.6	19.1	20.9	37.1	73.7	98.3
C4N0	0.3	–	0.01b	–	5.4	–	7.0	–	–	–
Significance	***	***	***	NS	***	NS	***	NS	NS	NS
Soil										
Loam	2.7	8.7	0.10	0.32	4.3	2.6b	25.4	66.1a	77.3	96.2
Sandy loam	3.4	11.0	0.09	0.28	17.6	32.4a	12.5	23.9b	67.6	96.7
Significance	***	***	NS	NS	***	***	***	***	NS	NS
Treatment × soil	***	***	NS	NS	***	NS	***	NS	NS	NS

Note. Means in a column followed by same lowercase letter are not significantly different. When interaction effect was significant, main effect was not reported.

<sup>a</sup>C0N0, no char and no urea ammonium nitrate (UAN); C0N1–C4N1, UAN at 200 kg N ha<sup>-1</sup> and char at 0, 6.7, 10.1, 13.4, and 26.8 Mg C ha<sup>-1</sup>, respectively; C4N0, 26.8 Mg C ha<sup>-1</sup> and no UAN.

<sup>b</sup>Fertilizer N recovery based on “N difference” method.

<sup>c</sup>Fertilizer N recovery based on the initial extractable N.

\*Significant at the .05 probability level.

\*\*Significant at the .01 probability level.

\*\*\*Significant at the .001 probability level. NS, not significant.

**TABLE 2** Interaction effect of treatment and soil on cumulative NH<sub>3</sub> volatilized, NO<sub>3</sub>-N leached, and soil residual N

Treatment <sup>a</sup>	NH <sub>3</sub> volatilized		NO <sub>3</sub> -N leached		Soil residual N	
	Loam	Sandy loam	Loam	Sandy loam	Loam	Sandy loam
	g (10 <sup>-3</sup> )					
C0N0	0.4g	0.3g	3.6cd	8.5c	6.7e	5.6e
C0N1	3.5e	4.5d	4.4cd	24.3a	33.3a	12.4cd
C1N1	2.2f	3.4e	1.5d	22.9ab	36.6a	16.2c
C2N1	2.6f	3.4e	5.8cd	22.5ab	35.1a	14.2c
C3N1	2.3f	5.4c	4.6cd	16.9b	32.8a	17.5c
C4N1	7.6a	6.6b	7.0cd	20.2ab	26.4b	15.2c
C4N0	0.3g	0.3g	3.2cd	7.7cd	7.0de	6.4e

Note. Means for each variable followed by same lowercase letters are not significantly different.

<sup>a</sup>C0N0, no char and no urea ammonium nitrate (UAN); C0N1–C4N1, UAN at 200 kg N ha<sup>-1</sup> and char at 0, 6.7, 10.1, 13.4, 26.8 Mg C ha<sup>-1</sup>, respectively; C4N0, 26.8 Mg C ha<sup>-1</sup> and no UAN.

ranged from 94.0 to 98.3% in treatments and from 96.2 to 96.7% by soil type (Table 1).

## 4 | DISCUSSION


### 4.1 | Ammonia volatilization

Ammonia volatilization loss observed in this study aligned with other studies that reported NH<sub>3</sub> losses from 8 to 13%


(Ma et al., 2010a; Peng et al., 2015; Vaio et al., 2008). Char addition did not enhance or suppress NH<sub>3</sub> volatilization in unfertilized treatments. Fertilization is the major source for NH<sub>3</sub> volatilization loss, as evidenced by a positive correlation between NH<sub>3</sub> volatilization and N fertilization reported in Jantalia et al. (2012) and Jones, Brown, Engel, Horneck, and Olson-Rutz (2013).

The higher clay content and CEC in loam than in sandy loam promoted better retention of NH<sub>4</sub> and subsequently reduced NH<sub>3</sub> loss in loam compared with sandy loam in this





**FIGURE 2** Daily  $\text{N}_2\text{O}$  flux (mean  $\pm$  SE) with different treatments in (a) loam and (b) sandy loam soils. C0N0, no char and no urea ammonium nitrate (UAN); C0N1–C4N1, UAN at  $200 \text{ kg N ha}^{-1}$  and char at 0, 6.7, 10.1, 13.4, and  $26.8 \text{ Mg C ha}^{-1}$ , respectively; C4N0,  $26.8 \text{ Mg C ha}^{-1}$  and no UAN. \*Treatment with significantly higher loss than all other treatments on a given sampling day




**FIGURE 3** Amount of  $\text{NO}_3\text{-N}$  leached (mean;  $n = 4$ ) with different treatments in loam and sandy loam soils at different leaching events. C0N0, no char and no urea ammonium nitrate (UAN); C0N1–C4N1, UAN at  $200 \text{ kg N ha}^{-1}$  and char at 0, 6.7, 10.1, 13.4, and  $26.8 \text{ Mg C ha}^{-1}$ , respectively; C4N0,  $26.8 \text{ Mg C ha}^{-1}$  and no UAN. Leaching events occurred on Days 20, 21, and 29 after fertilization for sandy loam and on Day 29 for loam

study. In addition, a higher sand content would enhance the loss of  $\text{NH}_3$  in sandy loam (McDowell et al., 1958).

Reduction in  $\text{NH}_3$  volatilization observed at lower char rates in both soil types was likely from increased physisorption due to the high surface area ( $82.1 \text{ m}^2 \text{ g}^{-1}$ ) and the high

CEC ( $46.9 \text{ meq } 100 \text{ g}^{-1}$ ) of char. The surface area of char exceeds that of clay-sized particles (Qi & Zhang, 2015) by one or two orders of magnitude and exceeds that of sand particles by three or four orders of magnitude. These results suggest that char functions more like biochar from various sources that have been reported to capture  $\text{NH}_3$  and reduce  $\text{NH}_3$  volatilization loss (Steiner et al., 2010; Taghizadeh-Toosi, Clough, Sherlock, & Condon, 2012). However, the beneficial effect of high-C products, such as char and biochar, in reducing  $\text{NH}_3$  loss depends on their sources, production conditions, containments and quality, and application rates (Ding et al., 2016; Steiner et al., 2008).

Soil pH is another important factor for retention or release of  $\text{NH}_4/\text{NH}_3$  in the soil. At pH below 7.5,  $\text{NH}_4$  is the predominant form, rather than volatile  $\text{NH}_3$  (Fan et al., 1993). As pH increases above 7.5, the  $\text{NH}_3$  form quickly becomes dominant and is susceptible to loss via volatilization (Behera, Sharma, Aneja, & Balasubramanian, 2013). The initial pH of sandy loam in this study was 7.7, which is above the 7.5 pH threshold for  $\text{NH}_3$  volatilization, whereas the loam soil had a pH of 7.2, which is slightly below this threshold. The pH of the char was 7.6, and char contained 19% calcium carbonate. Calcium carbonate aids in increasing soil alkalinity, and hydrolysis of urea to form  $\text{NH}_4$  also raises the pH (Jones et al., 2013). Depending on the nature and composition of CCRs, they could be useful to increase or buffer soil pH (Elsewi, Bingham, & Page, 1978a; Elsewi, Bingham, & Page, 1978b;



**FIGURE 4** Soil residual N (mean;  $n = 4$ ) the 0- to 24-cm depth at different treatments in (a) loam and (b) sandy loam soils. C0N0, no char and no urea ammonium nitrate (UAN); C0N1–C4N1, UAN at  $200 \text{ kg N ha}^{-1}$  and char at 0, 6.7, 10.1, 13.4, and  $26.8 \text{ Mg C ha}^{-1}$ , respectively; C4N0,  $26.8 \text{ Mg C ha}^{-1}$  and no UAN. Means at 18–24 cm with different letters across both soil types are significantly different at  $P < .05$

Phung, Lund, & Page, 1978). There could have been a considerable soil alkalization effect with higher char rates that counteracted and exceeded physiosorption benefits of char.

## 4.2 | Nitrous oxide emissions

The average  $\text{N}_2\text{O}$  emissions rate of  $0.7 \text{ kg N ha}^{-1}$  from fertilized treatments in this study is comparable to the  $0.6 \text{ kg N ha}^{-1}$  emission rate from UAN at  $150 \text{ kg N ha}^{-1}$  in a 28-d field study in eastern Canada (Ma et al., 2010b). In this study, a considerable amount of N moved down the soil profile and/or leached, and char addition would have only facilitated that downward N movement (Basu et al., 2009). A slight increase in  $\text{N}_2\text{O}$  emissions in loam soil compared with sandy loam (Table 1) could be related to anaerobic conditions at some pockets in loam soil, which promotes denitrification (Weier, Doran, Power, & Walters, 1993).

A previous laboratory incubation study documented that  $\text{N}_2\text{O}$  emissions may vary by soil texture (Harrison-Kirk, Beare, Meenken, & Condrón, 2013), but no significant differences in  $\text{N}_2\text{O}$  emissions by soil types were found in our study. Nitrous oxide emissions are primarily driven by N fertilization (Maharjan, Venterea, & Rosen, 2014; Shcherbak, Millar, & Robertson, 2014), as evidenced by greater emissions in fertilized than unfertilized treatments in this study. The high

variability in daily  $\text{N}_2\text{O}$  fluxes among laboratory replicates, which is likely be larger under field conditions, was one reason for the nonsignificant differences and should be kept in mind for evaluation of N losses from agricultural systems because it points toward a highly dynamic pathway. Johnson and Welch (1939) suggested 33% as permissible upper fiducial limit of CV. Although the acceptable range of CV may vary among experiments, the high CV observed in daily fluxes in this study failed to detect differences in treatment means (Patel, Patel, & Shiyani, 2001). Another potential pitfall in this study could be the small headspace used for gas sampling, which reduces minimum detectable flux (De Klein and Harvey, 2012).

## 4.3 | Nitrate leaching

The contrasting effect of C3N1 and C0N1 in sandy loam with respect to  $\text{NH}_3$  loss and  $\text{NO}_3\text{-N}$  leaching underscores the need to account for all possible pathways of N losses in our mitigation efforts. The lower  $\text{NO}_3\text{-N}$  leaching loss in C3N1 than in C0N1 is due to greater soil mineral residual N at the lower bottom of the column (18–24 cm depth) and greater  $\text{NH}_3$  loss in C3N1 than in C0N1. When there are multiple possible pathways for loss, as is the case with mineral N, an effort to reduce N loss via a particular pathway may be undermined or even

outweighed by loss via other pathway(s) (Lam, Suter, Mosier, & Chen, 2016).

The effect of high-C-content amendments on  $\text{NO}_3\text{-N}$  leaching depends on complex physical, chemical, and biological processes. It has been suggested that leaching of soil  $\text{NO}_3\text{-N}$  depends on the ability of biochar to retain  $\text{NO}_3\text{-N}$  and  $\text{NH}_4\text{-N}$  or on the inhibition of nitrification by clay particles (Clough, Condon, Kammann, & Müller, 2013; Liu et al., 2017). Some biochar studies have found decreased  $\text{NO}_3\text{-N}$  leaching depending on fertilizer type, soil type, and leaching conditions, but other studies showed inconsistent effects of biochar on leaching (Fidel, Laird, & Spokas, 2018; Haider, Steffens, Moser, Müller, & Kammann, 2017; Sika & Hardie, 2014).

Ventura, Sorrenti, Panzacchi, George, and Tonon (2013) observed a reduction in  $\text{NO}_3\text{-N}$  leaching only in the second year after biochar application, suggesting an increase in biochar sorption properties over time, possibly due to the oxidation and interaction of biochar and soil particles and an increase in the adsorbing surface due to particle fragmentation with aging (Hagemann et al., 2017; Singh, Hatton, Singh, Cowie, & Kathuria, 2010). In contrast, Gronwald, Don, Tiemeyer, and Helfrich (2015) observed that the adsorption capacity of biochar decreased by 60–80% to less or observed no  $\text{NO}_3/\text{NH}_4\text{-N}$  adsorption after 7 mo of aging in the field compared with the fresh char. A similar trend of decreasing adsorption capacity with biochar from beetroot chips was reported from a laboratory study on loam soil (Bargmann, Martens, Rillig, Kruse, & Kücke, 2014). Possible reasons for decreased adsorption capacity over time can be binding sites of biochar being blocked with organic matter or mineral particles and microbial degradation with subsequent possible changes in surface properties (Cheng, Lehmann, & Engelhard, 2008). In this study, a leaching event was observed on Day 29 after fertilization in loamy soil. The later and lower  $\text{NO}_3\text{-N}$  leaching observed in fertilized loam than in sandy loam in this study may be due to a lower water infiltration rate and greater nutrient retention in loamy soil because of greater clay and OM content (Lehmann & Schroth, 2003). Long-term evaluation is required to understand how char properties might change and affect soil  $\text{NO}_3\text{-N}$  leaching over time.

#### 4.4 | Soil residual mineral nitrogen and fertilizer nitrogen recovery

Lower soil residual mineral N at a depth of 18–24 cm and subsequently lower residual mineral N in the whole soil column with C4N1 compared with other fertilized treatments in loam soil could be the result of higher  $\text{NH}_3$  volatilization loss (cumulative loss of  $7.6 \times 10^{-3}$  g N or 17.1% of applied N) (Figure 1) or a slightly higher  $\text{NO}_3\text{-N}$  leaching loss (Table 1).

In all fertilized treatments, most of N moved down the profile and accumulated at the lower soil layers of the columns 30 d after N addition. This suggests the movement of  $\text{NO}_3\text{-N}$  down the soil profile with water (Bahmani, Nasab, Behzad, & Naseri, 2009; Pierzynski, Vance, & Sims, 2005). Previous research documented that 25.4 mm of irrigation or rainfall can transport soil  $\text{NO}_3\text{-N}$  to 150–200 mm in a loamy sand (Endelman, Keeney, Gilmour, & Saffigna, 1974). During the 30-d experiment, 100.8 mm of water was added. In the case of sandy loam soil, N moved down the profile and leached out of the column; therefore, residual mineral N was overall lower in sandy loam than in loam across fertilized treatments, including C4N1.

The  $\text{FNR}_{\text{CTRL}}$  was much smaller than  $\text{FNR}_{\text{ResN}}$  (Table 1). The  $\text{FNR}_{\text{CTRL}}$  estimate assumes that fertilizer N enhances OM mineralization (Khan, Mulvaney, Ellsworth, & Boast, 2007; Robertson et al., 2013). However, inorganic N inputs can also decrease OM mineralization by decreasing the decomposition of energy-poor OM substrates that are mineralized solely to access N-containing compounds (Craine, Morrow, & Fierer, 2007; Moorhead & Sinsabaugh, 2006). Particularly, in the current study, no crops were grown, and therefore there was no OM to mineralize to make up for potential N deficiency. In a laboratory incubation study with no crops involved, Mahal et al. (2019) demonstrated that fertilizer N suppressed OM mineralization. In contrast, Kaleem Abbasi, Mahmood Tahir, Sabir, and Khurshid (2015) reported that control soil without amendment released a maximum of  $30.9 \text{ mg N kg}^{-1}$  soil on Day 28 compared with  $13.7 \text{ mg kg}^{-1}$  at Day 0 at  $25^\circ\text{C}$  and 58% water filled pore space under laboratory conditions, showing a substantial release of N into the mineral N pool. The wide variation reported in the N mineralization from soils with or without fertilizer N can be affected by applied N rate (Cahill, Osmond, Crozier, Israel, & Weisz, 2007), soil temperature and moisture (Deenik, 2006), amount and type of clay in soil (Breland, 1994; Deenik, 2006). In the current study, mineralization under different treatments were not measured and the long-term effect of char-C in soil N mineralization/immobilization is yet to be explored. Irrespective of the methods of estimating FNR, it did not vary by treatments or soil. However, the differences in  $\text{FNR}_{\text{CTRL}}$  and  $\text{FNR}_{\text{ResN}}$  observed in this study underscores the implications of different methods used in calculating fertilizer N recovery or use efficiency (Mahal et al., 2019) and a critical role that soil OM mineralization might play in soil N availability and N use efficiency.

## 5 | CONCLUSION

In many countries, CCRs have not been properly utilized and still considered as waste products. Benefits of decreasing  $\text{NH}_3$  volatilization loss were observed with optimum rates of char

addition in both coarse and fine-textured soils. There were no adverse effects of adding char on leaching losses or N<sub>2</sub>O emissions. Field research is warranted to evaluate the potential use of char and other similar high C content by-products to improve N management. Further evaluation is warranted to investigate the possible adverse effects of pesticide/herbicide sorption and potential trace metal accumulation in soil, crop tissue, or grains before recommending char for agricultural use.

## ACKNOWLEDGMENTS

This work was supported by the University of Nebraska-Lincoln Agriculture Research Division and Western Sugar Cooperative.

## CONFLICT OF INTEREST

The authors declare no conflict of interest.

## ORCID

Dinesh Panday  <https://orcid.org/0000-0001-8452-3797>

Arindam Malakar  <https://orcid.org/0000-0001-6704-8891>

Bijesh Maharjan  <https://orcid.org/0000-0002-4728-7956>

## REFERENCES

- Ahmed, M. J., Stalikas, P. G., Tzouwara-Karayanni, S. M., Karayannis, M. L., & Veltsistas, P. G. (1997). Simultaneous spectrofluorimetric determination of selenium (IV) and (VI) by flow injection analysis. *Analyst*, *122*, 221–226.
- Atkinson, C. J., Fitzgerald, J. D., & Higgs, N. A. (2010). Potential mechanisms for achieving agricultural benefits from biochar application to temperate soils: A review. *Plant and Soil*, *337*(1–2), 1–18.
- Bahmani, O., Nasab, S. B., Behzad, M., & Naseri, A. A. (2009). Assessment of nitrogen accumulation and movement in soil profile under different irrigation and fertilization regime. *Asian Journal of Agricultural Research*, *3*, 38–46.
- Bargmann, I., Martens, R., Rillig, M. C., Kruse, A., & Kücke, M. (2014). Hydrochar amendment promotes microbial immobilization of mineral nitrogen. *Journal of Plant Nutrition and Soil Science*, *177*(1), 59–67.
- Basu, M., Pande, M., Bhadoria, P. B. S., & Mahapatra, S. C. (2009). Potential fly-ash utilization in agriculture: A global review. *Progress in Natural Science*, *19*(10), 1173–1186.
- Behera, S. N., Sharma, M., Aneja, V. P., & Balasubramanian, R. (2013). Ammonia in the atmosphere: A review on emission sources, atmospheric chemistry and deposition on terrestrial bodies. *Environmental Science and Pollution Research*, *20*(11), 8092–8131.
- Breland, T. A. (1994). Measured and predicted mineralisation of clover green manures at low temperature at different depths in two soils. *Plant and Soil*, *166*, 13–20.
- Bridgwater, A. V. (2003). Renewable fuels and chemicals by thermal processing of biomass. *Chemical Engineering Journal*, *91*, 87–102.
- Cahill, S., Osmond, D., Crozier, C., Israel, D., & Weisz, R. (2007). Winter wheat and maize response to urea ammonium nitrate and a new urea formaldehyde polymer fertilizer. *Agronomy Journal*, *99*(6), 1645–1653.
- Cameron, R. E. (1992). *Guide to site and soil description for hazardous waste site characterization. Vol. 1: Metals* (EPA/600/4-91/029). Washington, DC: USEPA.
- Cheng, C. H., Lehmann, J., & Engelhard, M. H. (2008). Natural oxidation of black carbon in soils: Changes in molecular form and surface charge along a climosequence. *Geochimica et Cosmochimica Acta*, *72*(6), 1598–1610.
- Clough, T., Condon, L., Kammann, C., & Müller, C. (2013). A review of biochar and soil nitrogen dynamics. *Agronomy Journal*, *3*, 275–293.
- Craine, J. M., Morrow, C., & Fierer, N. (2007). Microbial nitrogen limitation increases decomposition. *Ecology*, *88*(8), 2105–2113.
- De Klein, C. A., & Harvey, M. (2012). *Nitrous oxide chamber methodology guidelines*. Wellington, New Zealand: Global Research Alliance on Agricultural Greenhouse Gases, Ministry for Primary Industries.
- Deenik, J. (2006). Nitrogen mineralization potential in important agricultural soils of Hawai'i. *Soil and Crop Management*, *15*.
- Dil, M., Oelbermann, M., & Xue, W. (2014). An evaluation of biochar pre-conditioned with urea ammonium nitrate on maize (*Zea mays* L.) production and soil biochemical characteristics. *Canadian Journal of Soil Science*, *94*, 551–562.
- Ding, Y., Liu, Y., Liu, S., Li, Z., Tan, X., Huang, X., ... Zheng, B. (2016). Biochar to improve soil fertility: A review. *Agronomy for Sustainable Development*, *36*(2).
- Ding, Y., Liu, Y., Wu, W., Shi, D., Yang, M., & Zhong, Z. (2010). Evaluation of biochar effects on nitrogen retention and leaching in multi-layered soil columns. *Water Air and Soil Pollution*, *213*, 47–55.
- Dunmola, A. S., Tenuta, M., Moulin, A. P., Yapa, P., & Lobb, D. A. (2010). Pattern of greenhouse gas emission from a prairie pothole agricultural landscape in Manitoba, Canada. *Canadian Journal of Soil Science*, *90*, 243–256.
- Elsewii, A. A., Bingham, F. T., & Page, A. L. (1978a). Availability of sulfur in fly ash to plants. *Journal of Environmental Quality*, *7*, 69–73. <https://doi.org/10.2134/jeq1978.00472425000700010014x>
- Elsewii, A. A., Bingham, F. T., & Page, A. L. (1978b). Growth and mineral composition of lettuce and Swiss chard grown on flyash amended soils. In D. C. Adriano & I. L. Brisbin (Eds.), *Environmental chemistry and cycling processes* (pp. 568–581). Springfield, VA: U.S. Department of Commerce.
- Endelman, F. J., Keeney, D. R., Gilmour, J. T., & Saffigna, P. G. (1974). Nitrate and chloride movement in the plainfield loamy sand under intensive irrigation. *Journal of Environmental Quality*, *3*, 295–298. <https://doi.org/10.2134/jeq1974.00472425000300030024x>
- Fageria, N. K., & Baligar, V. C. (2005). Enhancing nitrogen use efficiency in crop plants. *Advances in Agronomy*, *88*, 97–185.
- Fan, M. X., & Mackenzie, A. F. (1993). Urea and phosphate interactions in fertilizer microsites: Ammonia volatilization and pH changes. *Soil Science Society of America Journal*, *57*(3), 839–845.
- Fidel, R. B., Laird, D. A., & Spokas, K. A. (2018). Sorption of ammonium and nitrate to biochars is electrostatic and pH-dependent. *Scientific Reports*, *8*, 17627.
- Filiberto, D. M., & Gaunt, J. L. (2013). Practicality of biochar additions to enhance soil and crop productivity. *Agriculture*, *3*, 715–725.
- Gronwald, M., Don, A., Tiemeyer, B., & Helfrich, M. (2015). Effects of fresh and aged chars from pyrolysis and hydrothermal carbonization on nutrient sorption in agricultural soils. *Soil*, *1*(1), 475–489.
- Hagemann, N., Kammann, C. I., Schmidt, H. P., Kappler, A., & Behrens, S. (2017). Nitrate capture and slow release in biochar amended compost and soil. *PLOS ONE*, *12*(2), e0171214.



- Haider, G., Steffens, D., Moser, G., Müller, C., & Kammann, C. I. (2017). Biochar reduced nitrate leaching and improved soil moisture content without yield improvements in a four-year field study. *Agriculture Ecosystems and Environment*, *237*, 80–94.
- Harrison-Kirk, T., Beare, M. H., Meenken, E. D., & Condon, L. M. (2013). Soil organic matter and texture affect responses to dry/wet cycles: Effects on carbon dioxide and nitrous oxide emissions. *Soil Biology and Biochemistry*, *57*, 43–55.
- International Fertilizer Association Statistics (IFASTAT). (2019). *Global consumption of agricultural fertilizer statistics*. Paris: International Fertilizer Association. Retrieved from <https://www.ifastat.org/>
- Jantalia, C. P., Halvorson, A. D., Follett, R. F., Rodrigues Alves, B. J., Polidoro, J. C., & Urquiaga, S. (2012). Nitrogen source effects on ammonia volatilization as measured with semi-static chambers. *Agronomy Journal*, *104*(6), 1595–1603.
- Johnson, N. L., & Welch, B. L. (1939). Applications of the non-central  $t$  distribution. *Biometrika* *31*, 362–389.
- Jones, C., Brown, B. D., Engel, R., Horneck, D., & Olson-Rutz, K. (2013). *Management to minimize nitrogen fertilizer volatilization*. Bozeman: Montana State University Extension.
- Kaleem Abbasi, M., Mahmood Tahir, M., Sabir, N., & Khurshid, M. (2015). Impact of the addition of different plant residues on nitrogen mineralization: Immobilization turnover and carbon content of a soil incubated under laboratory conditions. *Solid Earth*, *6*(1), 197–205.
- Khan, S. A., Mulvaney, R. L., Ellsworth, T. R., & Boast, C. W. (2007). The myth of nitrogen fertilization for soil carbon sequestration. *Journal of Environmental Quality*, *36*(6), 1821–1832. <https://doi.org/10.2134/jeq2007.0099>
- Lam, S. K., Suter, H., Mosier, A. R., & Chen, D. (2016). Using nitrification inhibitors to mitigate agricultural N<sub>2</sub>O emission: A double-edged sword? *Global Change Biology*, *23*(2). <https://doi.org/10.1111/gcb.13338>
- Lehmann, J., & Schroth, G. (2003). Nutrient leaching. In G. Schroth & F. L. Sinclair (Eds.), *Trees, crops and soil fertility* (pp. 151–166). Wallingford, U.K.: CABI Publishing.
- Lehmann, J., Gaunt, J., & Rondon, M. (2006). Bio-char sequestration in terrestrial ecosystems: A review. *Mitigation and Adaptation Strategies for Global Change*, *11*(2), 403–427.
- Li, X., Hu, H. C., Delgado, J. A., Zhang, Y., & Ouyang, Z. (2007). Increased nitrogen use efficiencies as a key mitigation alternative to reduce nitrate leaching in north China plain. *Agricultural Water Management*, *89*, 137–147.
- Littell, R. C., Milliken, G. A., Stroup, W. W., Wolfinger, R. D., & Schabenberger, O. (2006). *SAS for mixed models*. Cary, NC: SAS Institute.
- Liu, Z., He, T., Cao, T., Yang, T., Meng, J., & Chen, W. (2017). Effects of biochar application on nitrogen leaching, ammonia volatilization and nitrogen use efficiency in two distinct soils. *Journal of Plant Nutrition and Soil Science*, *17*, 515–528.
- Ma, B. L., Wu, T. Y., Tremblay, N., Deen, W., McLaughlin, N. B., Morrison, M. J., & Stewart, G. (2010a). On-farm assessment of the amount and timing of nitrogen fertilizer on ammonia volatilization. *Agronomy Journal*, *102*, 134–144.
- Ma, B. L., Wu, T. Y., Tremblay, N., Deen, W., Morrison, M. J., McLaughlin, N. B., ... Stewart, G. (2010b). Nitrous oxide fluxes from corn fields: On-farm assessment of the amount and timing of nitrogen fertilizer. *Global Change Biology*, *16*, 156–170.
- Mahal, N. K., Osterholz, W. R., Miguez, F. E., Poffenbarger, H. J., Sawyer, J. E., Oik, D. C., ... Castellano, M. J. (2019). Nitrogen fertilizer suppresses mineralization of soil organic matter in maize agroecosystems. *Frontiers in Ecology and Evolution*, *7*, Article 59.
- Maharjan, B., Venterea, R. T., & Rosen, C. (2014). Fertilizer and irrigation management effects on nitrous oxide emissions and nitrate leaching. *Agronomy Journal*, *106*, 703–714.
- Major, J., Rondon, M., Molina, D., Riha, S. J., & Lehmann, J. (2012). Nutrient leaching in a Colombian savanna Oxisol amended with biochar. *Journal of Environmental Quality*, *41*, 1076–1086. <https://doi.org/10.2134/jeq2011.0128>
- McDowell, L. L., & Smith, G. E. (1958). The retention and reactions of anhydrous ammonia on different soil types. *Soil Science Society of America Journal*, *22*(1), 38–42.
- McGinn, S. M., & Janzen, H. H. (1998). Ammonia sources in agriculture and their measurement. *Canadian Journal of Soil Science*, *78*(1), 139–148.
- Moorhead, D. L., & Sinsabaugh, R. L. (2006). A theoretical model of litter decay and microbial interaction. *Ecological Monographs*, *76*(2), 151–174.
- Panday, D., Ferguson, R. B., & Maharjan, B. (2018). Flue gas desulfurization (FGD) gypsum as soil amendment. In A. Rakshit, B. Sarkar, & C. Abhilashis (Eds.), *Soil amendments for sustainability: Challenges and perspectives* (pp. 199–208). Boca Raton, FL: CRC Press.
- Patel, J. K., Patel, N. M., & Shiyani, R. L. (2001). Coefficient of variation in field experiments and yardstick thereof: An empirical study. *Current Science*, *81*(9), 1163–1164.
- Peng, X., Maharjan, B., Yu, C., Su, A., Jin, V., & Ferguson, R. B. (2015). A laboratory evaluation of ammonia volatilization and nitrate leaching following nitrogen fertilizer application on a coarse-textured soil. *Agronomy Journal*, *107*, 871–879.
- Phung, H. T., Lund, I. J., & Page, A. L. (1978). Potential use of fly ash as a liming material. In D. C. Driano & I. L. Brisbin (Eds.), *Environmental chemistry and cycling processes* (pp. 504–515). Springfield, VA: U.S. Department of Commerce.
- Pierzynski, G. M., Vance, G. F., & Sims, J. T. (2005). *Soils and environmental quality*. Boca Raton, FL: CRC Press.
- Qi, Y., & Zhang, T. C. (2015). Sorption and desorption of testosterone at environmentally relevant levels: Effects of aquatic conditions and soil particle size fractions. *Journal of Environmental Engineering*, *142*(1), 04015045.
- Robertson, G. P., Bruulsema, T. W., Gehl, R. J., Kanter, D., Mauzerall, D. L., Rotz, C. A., & Williams, C. O. (2013). Nitrogen–climate interactions in U.S. agriculture. *Biogeochemistry*, *114*, 41–70.
- SAS. (2015). *SAS 9.4 in-database products: User's guide* (5th ed.). Cary, NC: SAS Institute.
- Shaheen, S. M., Hooda, P. S., & Tsadilas, C. D. (2014). Opportunities and challenges in the use of coal fly ash for soil improvements: A review. *Journal of Environmental Management*, *145*, 249–267.
- Shcherbak, I., Millar, N., & Robertson, G. P. (2014). Global meta-analysis of the nonlinear response of soil nitrous oxide (N<sub>2</sub>O) emissions to fertilizer nitrogen. *Proceedings of the National Academy of Sciences of the USA*, *111*(25), 9199–9204.
- Siddaramappa, R., McCarty, G. W., Wright, R. J., & Codling, F. E. (1994). Mineralization and volatile loss of nitrogen from soils treated with coal combustion byproducts. *Biology and Fertility of Soils*, *18*(4), 279–284.

- Sika, M. P., & Hardie, A. G. (2014). Effect of pine wood biochar on ammonium nitrate leaching and availability in a South African sandy soil. *European Journal of Soil Science*, *65*, 113–119.
- Singh, B., Macdonald, L. M., Kookana, R. S., van Zwieten, L., Butler, G., Joseph, S., ... Cattle, J. (2014). Opportunities and constraints for biochar technology in Australian agriculture: Looking beyond carbon sequestration. *Soil Research*, *52*, 739–750.
- Singh, B. P., Hatton, B. J., Singh, B., Cowie, A. L., & Kathuria, A. (2010). Influence of biochars on nitrous oxide emission and nitrogen leaching from two contrasting soils. *Journal of Environmental Quality*, *39*(4), 1224–1235. <https://doi.org/10.2134/jeq2009.0138>
- Steiner, C., Glaser, B., Geraldtes Teixeira, W., Lehmann, J., Blum, W. E., & Zech, W. (2008). Nitrogen retention and plant uptake on a highly weathered central Amazonian Ferralsol amended with compost and charcoal. *Journal of Plant Nutrition and Soil Science*, *171*, 893–899.
- Steiner, C., Das, K. C., Melear, N., & Lakly, D. (2010). Reducing nitrogen loss during poultry litter composting using biochar. *Journal of Environmental Quality*, *39*(4), 1236–1242. <https://doi.org/10.2134/jeq2009.0337>
- Taghizadeh-Toosi, A., Clough, T. J., Sherlock, R. R., & Condrón, L. M. (2012). Biochar adsorbed ammonia is bioavailable. *Plant and Soil*, *350*, 57–69.
- Vaio, N., Cabrera, M. L., Kissel, D. E., Rema, J. A., Newsome, J. F., & Calvert, V. H. (2008). Ammonia volatilization from urea-based fertilizers applied to tall fescue pastures in Georgia, USA. *Soil Science Society of America Journal*, *72*, 1665–1671.
- Ventura, M., Sorrenti, G., Panzacchi, P., George, E., & Tonon, G. (2013). Biochar reduces short-term nitrate leaching from a horizon in an apple orchard. *Journal of Environmental Quality*, *42*, 76–82. <https://doi.org/10.2134/jeq2012.0250>
- Wagner, S. W., Reicosky, D. C., & Alessi, R. S. (1997). Regression models for calculating gas fluxes measured with a closed chamber. *Agronomy Journal*, *89*, 279–284. <https://doi.org/10.2134/agronj1997.00021962008900020021x>
- Weier, K. L., Doran, J. W., Power, J. F., & Walters, D. T. (1993). Denitrification and the dinitrogen/nitrous oxide ratio as affected by soil water, available carbon, and nitrate. *Soil Science Society of America Journal*, *57*(1), 66–72.

## SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

**How to cite this article:** Panday D, Mikha MM, Collins HP, et al. Optimum rates of surface applied coal char decreased soil ammonia volatilization loss. *J. Environ. Qual.* 2020;49:256–267. <https://doi.org/10.1002/jeq2.20023>