

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Publications from USDA-ARS / UNL Faculty

U.S. Department of Agriculture: Agricultural
Research Service, Lincoln, Nebraska

2020

Development of nuclear microsatellite markers to facilitate germplasm conservation and population genetics studies of five groups of tropical perennial plants with edible fruits and shoots: rambutan (*Nephelium lappaceum* L.), sapodilla (*Manilkara zapota* (L.) P. Royen), lychee (*Litchi chinensis* Sonn.), mangosteen (*Garcinia mangostana* Linn. and *Garcinia cochinchinensis* (Lour.) Choisy) and bamboo (*Bambusa vulgaris* Schrad. ex J.C. Wendl and *Guadua angustifolia* Kunth)

Renee S. Arias

USDA-ARS National Peanut Research Laboratory & USDA-ARS Plant Germplasm Introduction and Testing Research Unit, renee.arias@usda.gov

Valerie A. Orner

USDA-ARS National Peanut Research Laboratory & USDA-ARS Plant Germplasm Introduction and Testing Research Unit, Valerie.Orner@usda.gov
Follow this and additional works at: <https://digitalcommons.unl.edu/usdaarsfacpub>

Victor S. Sobolev

Arias, Renee S.; Orner, Valerie A.; Sobolev, Victor S.; Martínez-Castillo, Jaime; Ballard, Linda L.; Duke, Mary Y.; Simpson, Sharon A.; Liu, Xiaofen F.; and Scheffler, Brian E., "Development of nuclear microsatellite markers to facilitate germplasm conservation and population genetics studies of five groups of tropical

perennial plants with edible fruits and shoots: rambutan (*Nephelium lappaceum* L.), sapodilla (*Manilkara zapota* (L.) P. Royen), lychee (*Litchi chinensis* Sonn.), mangosteen (*Garcinia mangostana* Linn. and *Garcinia cochinchinensis* (Lour.) Choisy) and bamboo (*Bambusa vulgaris* Schrad. ex J.C. Wendl and *Guadua angustifolia* Kunth)" (2020). *Publications from USDA-ARS / UNL Faculty*. 2355.

<https://digitalcommons.unl.edu/usdaarsfacpub/2355>

USDA-ARS Plant Germplasm Introduction and Testing Research Unit & USDA-ARS Genomics and Bioinformatics Research Unit, Linda.Ballard@usda.gov


This Article is brought to you for free and open access by the U.S. Department of Agriculture: Agricultural Research Service, Lincoln, Nebraska at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Publications from USDA-ARS / UNL Faculty by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.
See next page for additional authors

Authors

Renee S. Arias, Valerie A. Orner, Victor S. Sobolev, Jaime Martínez-Castillo, Linda L. Ballard, Mary V. Duke, Sharon A. Simpson, Xiaofen F. Liu, and Brian E. Scheffler



Development of nuclear microsatellite markers to facilitate germplasm conservation and population genetics studies of five groups of tropical perennial plants with edible fruits and shoots: rambutan (*Nephelium lappaceum* L.), sapodilla (*Manilkara zapota* (L.) P. Royen), lychee (*Litchi chinensis* Sonn.), mangosteen (*Garcinia mangostana* Linn. and *Garcinia cochinchinensis* (Lour.) Choisy) and bamboo (*Bambusa vulgaris* Schrad. ex J.C. Wendl and *Guadua angustifolia* Kunth)

Renée S. Arias · Linda L. Ballard · Mary V. Duke · Sheron A. Simpson · Xiaofen F. Liu · Valerie A. Orner · Victor S. Sobolev · Brian E. Scheffler · Jaime Martinez-Castillo 

Received: 13 March 2020 / Accepted: 15 June 2020
© Springer Nature B.V. 2020

Abstract Simple sequence repeat (SSR) enriched libraries for five groups of tropical perennial plants

with edible fruits and shoots were prepared and sequenced in a GS-FLX Roche 454: sapodilla (*Manilkara zapota* (L.) P. Royen), lychee (*Litchi chinensis* Sonn.), mangosteen (*Garcinia mangostana* Linn. and *G. cochinchinensis* (Lour.) Choisy), rambutan (*Nephelium lappaceum* L.), and bamboo (*Bambusa vulgaris* Schrad. ex J.C. Wendl and *Guadua*

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s10722-020-00965-w>) contains supplementary material, which is available to authorized users.

R. S. Arias · V. A. Orner · V. S. Sobolev
USDA-ARS National Peanut Research Laboratory, 1011
Forrester Dr. S.E, Dawson, GA 39842, USA
e-mail: Renee.Arias@ars.usda.gov

V. A. Orner
e-mail: Valerie.Orner@usda.gov

V. S. Sobolev
e-mail: Victor.Sobolev@usda.gov

R. S. Arias · L. L. Ballard · M. V. Duke ·
S. A. Simpson · X. F. Liu · V. A. Orner ·
V. S. Sobolev · B. E. Scheffler · J. Martinez-Castillo (✉)
USDA-ARS Plant Germplasm Introduction and Testing
Research Unit, 24106 N. Bunn Rd, Prosser,
WA 99350, USA
e-mail: jmartinez@cicy.mx

L. L. Ballard
e-mail: Linda.Ballard@usda.gov

M. V. Duke
e-mail: Mary.Duke@usda.gov

S. A. Simpson
e-mail: Sheron.Simpson@usda.gov

X. F. Liu
e-mail: Fanny.Liu@usda.gov

B. E. Scheffler
e-mail: Brian.Scheffler@ars.usda.gov

L. L. Ballard · M. V. Duke · S. A. Simpson ·
X. F. Liu · B. E. Scheffler
USDA-ARS Genomics and Bioinformatics Research Unit,
141 Experiment Station Rd, Stoneville,
MS 387761, USA

angustifolia Kunth). For SSR development, these species were organized by their common names in five groups. A total of 3870 SSR primer sets were designed, using capillary electrophoresis 1872 nuclear SSRs were tested on 4 to 10 DNA samples within each plant group, that is 384 loci for each of the four groups of fruit trees and 336 loci for the bamboo group. Only 7.9% of the primers tested did not result in amplification. All 1872 SSRs are provided, we highlight 178 SSRs (between 26 and 47 per group) considered top-quality polymorphic SSRs that amplified all the samples, had strong fluorescence signal, presented no stutters and showed minimum non-specific amplification or background fluorescence. A total of 66,057 contig sequences were submitted to GenBank Database. Markers presented here will be useful not only for conservation efforts in banks of germplasm, but also for in-depth analysis of population genetics which usually requires evaluation of large number of loci.

Keywords Germplasm · High throughput · National plant germplasm system (NPGS) · Pyrosequencing · SSR markers · Roche 454 · SSR markers · Tropical trees

Introduction

Approximately 200 plant genomes have been sequenced so far; from those, roughly 50 correspond to ferns, bryophytes and algae, and the rest are mostly temperate climate crops (Mukherjee et al. 2018). Unfortunately, most of the tropical species have yet to be sequenced. This is a worrisome situation since fragmentation and loss of habitat in the tropics is happening at a very fast pace (Aguilar et al. 2018; Cousins 2020; Escobar 2019) and mass extinctions are expected in these regions, even as consequence of habitat disturbance (Alroy 2017). This situation endangers the existence of many species on which man has depended for survival for thousands of years.

One of the biggest challenges for germplasm collections is the molecular characterization of

accessions and their preservation from genetic erosion (Barcaccia 2009). Molecular markers are critical to determine the genetic diversity within collections and in the wild, as well as to select core collections of manageable size that represent the genetic diversity of the collection while maintaining allele specificity and accession rarity (Curry 2017; Reyes-Valdes et al. 2018). Microsatellites (or SSRs—Simple Sequence Repeats—), are one of the most widely used molecular markers in genetic studies, such as population genetics, molecular breeding, and paternity testing (Ellegren 2004). SSRs are abundant, co-dominant, multi-allelic, highly reproducible and easy to use (Richard et al. 2008); and they can be isolated either by data mining of existing sequences (Sharma et al. 2007) or by generating and sequencing SSR-enriched libraries (Kijas et al. 1994; Zane et al. 2002). SSRs are still the markers of choice for many population genetic studies in tropical plants (e. g. Martínez-Castillo et al. 2019a, b; Chaluvadi et al. 2018; Yamanaka et al. 2019).

In the United States of America, the National Plant Germplasm System (NPGS) currently maintains 596,198 accessions from 13,480 species within 239 families (Bretting and Bennet 2007; NPGS 2020). All tropical perennial plants considered in the present study are included in NPGS and they represent an important genetic resource for people living in the tropics: sapodilla [*Manilkara zapota* (L.) P. Royen] Sapotaceae, lychee [*Litchi chinensis* Sonn.] Sapindaceae, rambutan [*Nephelium lappaceum* L.] Sapindaceae, mangosteen [*Garcinia mangostana* Linn.] and false mangosteen [*Garcinia cochinchinensis* (Lour.) Choisy] Clusiaceae, all have edible fruits (Arias et al. 2012; Finocchiaro 2020), whereas the two species of bamboo [*Bambusa vulgaris* Schrad. ex J.C. Wendl and *Guadua angustifolia* Kunth] Poaceae, have edible shoots (Singhal et al. 2013). For these species, the number of SSRs available are not sufficient; since the theoretical quantity of loci for accurate evolutionary inference of populations is greater than 30 (Pollock et al. 1998; Takezaki and Nei 1996). For rambutan, no SSRs have been developed, though transferability of 12 SSRs from lychee has been described (Hock et al. 2005). For lychee, only 4 and 12 SSRs were reported in separate studies (Ekue et al. 2009; Viruel and Hormaza 2004), respectively. For sapodilla, only 8 and 17 SSRs were reported in two studies (Moraes et al. 2013; Silva-Junior et al. 2016), respectively. For

J. Martínez-Castillo
Centro de Investigación Científica de Yucatán (CICY),
Calle 43 No. 130. Col. Chuburná de Hidalgo,
97200 Mérida, Yucatán, Mexico

mangosteen, only 17 SSRs were described (Samsir et al. 2016). For bamboo, only 16 SSRs were developed, but from chloroplast sequencing (Vieira et al. 2016). Our main objective was to develop large sets of nuclear SSR markers for the seven species mentioned earlier with the goal that this information will be a valuable resource for conservation programs in banks of germplasm and for in depth population-genetics studies on these species.

Materials and methods

DNA extraction and preparation of SSR libraries

Leaf samples sapodilla, lychee, mangosteen (two species), rambutan and bamboo (two species) were received from USDA-ARS Tropical Agriculture Research Station (TARS), Mayaguez, Puerto Rico, and organized in five groups for developing simple-sequence repeats (SSRs) to be used in germplasm collection and identification. The list of accessions is shown in Table 1. SSR-enriched libraries were prepared as described in Arias et al. (2015), using the same restriction enzymes adapter 1: SSRLIBF1 and adapter 2: SSRLIBF3 (Techen et al. 2010), and the same biotinylated oligonucleotide repeats and conditions.

Sequencing and SSR primer design

To avoid generating chimeric DNA during PCR reactions in SSR-enriched library preparation, two DNA samples of each of the five groups were processed separately. Then, equal volumes of the two libraries were mixed before proceeding to library preparation for sequencing. DNA quality of pooled pairs of samples was evaluated using Qubit™ fluorometer with the Quant-iT™ PicoGreen® reagent (Invitrogen, Carlsbad, CA) and by Agilent 2100 BioAnalyzer (Agilent Technologies, Santa Clara, CA) equipped with a DNA Ladder 1000 LabChip (Agilent Technologies, New Castle, DE) and its corresponding ladder. The libraries were sequenced using 70 × 75 mm Titanium Pico-Titer Plates (Roche, Branford, CT) on a Roche 454 GS FLX (Roche, Indianapolis, IN) using GS Titanium sequencing kit XLR70 (200 cycles). Read length distribution was analyzed with Roche 454 v.2.0 image/signal

Table 1 List of accessions used for Roche 454 pyrosequencing and SSR development

Common name	Scientific name	Accession name	TARS	A	Group	Total Reads	Bases	Contigs in NCBI	Biosample	Bioproject	Accession	Version
Rambutan	<i>Nephelium lappaceum</i>	Rongren	–	1	NEL	229,908	56,630,415	16,414	SAMNI2545058	PRJNA548147	KDDQ000000000	KDDQ010000000
	<i>Nephelium lappaceum</i>	R 162	–									
Sapodilla	<i>Manilkara zapota</i>	Tikal	17,900	2	MAZ	354,081	94,238,976	19,862	SAMNI2545082	PRJNA559539	KDDP000000000	KDDQ010000000
	<i>Manilkara zapota</i>	Oxkutzcab	17,896									
Lychee	<i>Litchi chinensis</i>	Kai mana	–	2	LIC	268,875	74,609,929	19,746	SAMNI2545059	PRJNA548147	KDDQ000000000	KDDQ010000000
	<i>Litchi chinensis</i>	Brewster	–									
Mangosteen	<i>Garcinia mangostana</i>	–	1202	1	GAM	244,597	67,859,915	8453	SAMNI2545063	PRJNA548147	KDDQ000000000	KDDQ010000000
False mangosteen	<i>Garcinia cochinchinensis</i>	–	1769									
Bamboo	<i>Guadua angustifolia</i>	–	16,914	2	GUA	226,960	51,831,335	1582	SAMNI2545064	PRJNA548147	KDDQ000000000	KDDP010000000
	<i>Bambusa vulgaris</i> var. <i>vittata</i>	–	16,296									

Samples used to develop the SSR markers. TARS: Accession number assigned by the Tropical Agriculture Research Station, Mayaguez, PR. A: adapter used in library, either adapter 1 or 2. Group: letters that identify each group of two samples. Total Reads: number of sequences obtained for each pair of libraries and sequenced in Roche 454. Contigs in NCBI: is the total number of contigs generated in the present work that were submitted to Genbank database in National Center for Biotechnology Information (NCBI). Biosample, Bioproject, Accession and Version: correspond to their assigned identification in NCBI



Fig. 1 Plants used in the present study. Top: sapodilla (left), rambutan (middle), bamboo (right); Bottom: mangosteen (left), lychee (right). Inserts within pictures show fruits cut open.

Photographs, courtesy of USDA-ARS, Peggy Greb (USDA Image Database, open access)

analysis and base caller programs. Contigs were assembled using Roche 454 gsAssembler version 2.0 (Roche, Branford, CT). SSR detection, and primer design followed the same protocol previously described (Arias et al. 2015). When a contig contained more than one repeat, primer sets within the contig were given alphabetical sub-indexes, e.g. “_a”, “_b”, “_c. Designed primer sets were tested on 4 or 10 DNA samples per group in 384-well/clear microtiter plates HSP3811 (Bio-Rad, Hercules, CA) in 5 μ L reactions with 10-ng DNA using Titanium Taq DNA Polymerase (Clontech, Mountain View, CA). Amplicons generated by capillary electrophoresis were analyzed in ABI 3730XL DNA analyzer (Applied Biosystems, Foster City, CA) and data were processed using GeneMapper v. 3.7 (Applied Biosystems, Foster City, CA).

Results

SSR-enriched libraries were prepared for five groups of tropical plant species (Fig. 1), using two accessions in the TARS collection as indicated in Table 1. The libraries were sequenced in a Roche 454 pyrosequencer resulting in 227–354 thousand reads per group. Histograms of the distribution of read number vs. read length showed maximum number of reads for each of the libraries at approximately 300–350 base pairs and reaching up to 600 bp length (Fig. 2). Libraries of bamboo and mangosteen were processed together in the same region of a picotiter plate, then the reads were separated by the sequence of the oligonucleotide adapters used. The number of contigs assembled for each group was between 1582 and 19,862, and their sequences were submitted to GenBank, National Center for Biotechnology Information (NCBI), accession numbers shown in Table 1. The total number of repeats detected by SSR-Finder software in each of the




Fig. 2 Read-length distribution in Roche 454 pyrosequencing of SSR-enriched libraries of *Manilkara zapota* (sapodilla), *Litchi chinensis* (lychee), *Garcinia mangostana* (mangosteen), *Nephelium lappaceum* (rambutan) and *Bambusa vulgaris* or

Guadua angustifolia (bamboo). Mangosteen and bamboo were processed using different adapters and loaded on the same region of the Roche 454 plate. The “y” axis is the number of reads, the “x” axis is the read length in base pairs (bp)

libraries was between 949 (bamboo) and 8084 (rambutan); and the number of unique primers designed were between 353 and 1557, also for bamboo

and rambutan, respectively (Table 2). A total of 384 SSRs were tested for each of the fruit trees, rambutan, sapodilla, lychee, mangosteen, and 336 SSRs were

Table 2 Data of the SSR sets development in seven tropical perennial plants

Common name	Scientific name	Group	SSRs detected	Unique primers	Primers screened	Repeat length (± stdv)	DNA samples tested	Group Primer ID	Repeat Motifs				Top quality polymorphic SSRs
									2-mer	3-mer	4-mer	5-mer	
Rambutan	<i>Nephelium lappaceum</i>	NEL	8084	1557	384	15.3 (± 4.3)	10	(Stv_nel)	422	915	111	49	36 (9.0%)
Sapodilla	<i>Manilkara zapota</i>	MAZ	7484	1389	384	15.7 (± 4.9)	4	(Stv_maz)	530	568	52	22	47 (12.0%)
Lychee	<i>Litchi chinensis</i>	LIC	6262	1161	384	15.6 (± 4.68)	10	(Stv_lic)	389	678	69	25	38 (10.0%)
Mangosteen	<i>Garcinia mangostana</i>	GAM	2235	839	384	15.2 (± 5.50)	4	(Stv_gam)	340	414	65	20	31 (8.0%)
Bamboo	<i>Garcinia cochinchinensis</i>	GUA	949	353	336	16.1 (± 6.77)	10	(Stv_gua)	163	160	22	8	26 (8.0%)
	<i>Bambusa vulgaris</i>												

Summary of simple sequence repeat (SSR) observed for each group of species. Only non-mononucleotide repeats (repeat motif 2–8 bp) were tested and reported, repeats with two-nucleotide motif of higher order are indicated as 2-mer, 3-mer, 4-mer and ≥ 5-mer. Top quality polymorphic SSRs: is the number of the best non-mononucleotide markers that amplified all the samples

t-

tested for bamboo, these 1872 SSRs are provided in Supplementary Table S1. The number of samples used to test the SSRs varied, 10 samples were used for rambutan, lychee and bamboo, whereas only 4 samples of sapodilla and mangosteen were tested. The overall number of repeat motif sizes, whether they were 2, 3, 4 or ≥ 5 nucleotides (nt) is also listed in Table 2.

In the SSR-enriched libraries sequenced of tropical plants, the number of repeat motifs varied from as low as 98 motifs in bamboo to 149 motifs in sapodilla. However, a small group of nine motifs represented from 72 to 83% of all the repeat motifs found in each library, these motifs are shown in Fig. 3. A total of 149 SSR markers did not result in amplification, still leaving 1,723 usable markers generated from this work. In general, less than 10% of the primer sets designed did not produce amplicons, with the lowest values observed in sapodilla, mangosteen and lychee (5.2, 6.3 and 7.8%, respectively), and the highest for rambutan and bamboo (9.4 and 11.6%, respectively) (Fig. 4). The SSRs that resulted in no amplification on the DNAs tested were marked as gray shade cells in Supplementary Table S1. Screening of SSR markers on 4 to 10 individual DNA samples resulted in polymorphism in 30.1 to 52.3% of the markers tested; in the




Fig. 3 Percentage of nuclear simple sequence repeat (SSR) markers that were polymorphic or resulted in no amplification. The total number of SSRs tested was 384 for *Garcinia mangostana* (mangosteen, GAM), 384 for *Litchi chinensis* (lychee, LIC), 384 for *Manilkara zapota* (sapodilla, MAZ), 384 for *Nephelium lappaceum* (rambutan, NEL) and 336 for *Bambusa vulgaris* (bamboo, GUA). One spp. only: indicates that 50% of the markers that were developed using SSR-enriched libraries of *Bambusa vulgaris* and *Guadua angustifolia* amplified only one of these two species. The percentage of polymorphism is probably underestimated given the small number of samples tested




Fig. 4 Percentage of simple sequence repeat (SSR) motifs found in non-monomucleotide nuclear-SSR-enriched libraries over a total of 1093 repeats and 99 motifs of *Garcinia mangostana* (mangosteen), 1948 repeats and 144 motifs of *Litchi chinensis* (lychee), 2135 repeats and 149 motifs of *Manilkara zapota* (sapodilla), 1117 repeats and 114 motifs of *Nephelium lappaceum* (rambutan) and 480 repeats and 98 motifs of *Bambusa vulgaris* (bamboo). A 72–83% of the repeats corresponded to the 9 motifs indicated in the figure legend. Percentage values $\geq 5\%$ are numerically indicated in the colored areas

case of bamboo where SSR-enriched libraries were prepared using two different genera (*Guadua angustifolia* and *Bambusa vulgaris*), 50% of the marker amplified only one of these two species (Fig. 4).

Several criteria were applied to select the top quality SSR markers for each group of tropical plants. These criteria were: amplification of all the samples tested, high fluorescent signal, minimum background amplification (nonspecific amplicons), absence of multiple peaks, and absence of stutter peaks. Application of these criteria to the 1872 SSRs tested resulted in 178 top quality markers reported in Table 3; where 36, 47, 38, 31 and 26 correspond to rambutan, sapodilla, lychee, mangosteen and bamboo, respectively. Examples of SSRs that were chosen as top quality markers are shown in Fig. 5, these are three markers of sapodilla showing amplification of four DNA samples, all of them with high levels of fluorescence (30,000 units scale), and a minimum of background.

Discussion

We reported between 297 and 364 new nuclear SSR markers for five groups of tropical plants studied:

rambutan, sapodilla, mangosteen, lychee and bamboo. Overall, this is a 20-fold higher number of SSR markers than the currently existing in the literature for the five plant groups studied, e.g. 15-fold more for sapodilla and 22-fold more for lychee, respectively. One advantage of using nuclear SSRs, is that the topology of reconstructed phylogenies can be different from the one using plastid data (Lin et al. 2019). This could be an advantage in the case of bamboo (*Guadua angustifolia* and *Bambusa vulgaris*) since to the best of our knowledge the 297 SSRs are the first nuclear SSRs reported for these species; the 16 SSRs previously reported for bamboo are from chloroplast origin (Vieira et al. 2016).

The potential use of the SSRs markers developed in the present study goes beyond their particular use for the seven species considered. One of the characteristics of SSRs is their high level of transferability between closely related species (Ziya et al. 2016). For example, SSRs developed for sapodilla could be useful in the other 64 species that belong to the pantropical genus *Manilkara*, which contains about 30 species in America, about 20 in Africa and about 15 in Asia, Australia and the Pacific; several of them utilized for its timber, fruit and latex (Armstrong 2010). Furthermore, it is possible that the SSRs developed here could be used in species that do not belong to the same genus. For example, SSR markers developed for lychee in two separate studies (Hock et al. 2005; Ekue 2009), have shown transferability to species within different genera of the Sapindaceae family. Sapindaceae is a tropical and subtropical family which contains about 1580 species, several of them with edible fruits (Buerki et al. 2010).

SSRs are being used in multiple conservation efforts to preserve species in the tropics. For example, nine polymorphic SSRs were used to understand the genetic structure and diversity of *Annona cherimola* Mill., to preserve germplasm that could be source of biotic and abiotic stress resistance and to guarantee food security in future generations (Larranaga et al. 2017). In date palm, *Phoenix dactylifera*, 19 SSRs were used to determine the population structure of 195 accessions from Asia and Africa and to understand their vulnerability to diseases and insect pests given sudden changes in climate (Chaluvadi et al. 2018). Also, 46 polymorphic SSRs were used to analyze the genetic structure of mango cultivars from around the world, conservation of germplasm and to facilitate the

Table 3 Selected best quality, polymorphic primer sets for microsatellites of five tropical plant groups

Marker	Forward 5' → 3'	Reverse 5' → 3'	Motif	Range	Amp	Alleles/sample	%H
stv-nel_00221_a	CTTCTCTCTGGAAATTTGGAGGTG	CTGCATCAAAACACGATAAAAACCAC	(GGAGAG) ₅	132–138	2	1.7 ± 0.48	70
stv-nel_00274_a	ATCCAAATCTCAAATCTCAACCACC	TCTAGGGTTTTCTCTGAAGAAATTCG	(TTTA) ₅	160–168	2	1.1 ± 0.31	10
stv-nel_00278_a	GGGTGGAAAATCGGAGAGTAGAAG	ATCCCTCCAATTTCTCTTCCAAAAC	(AGA) ₈	133–136	2	1.1 ± 0.31	10
stv-nel_00333_a	ACTCTGCTGTGTTTTGACCCTTC	CGAGAAAACACACGGTAAAGTGTGAC	(TATG) ₄	158–166	2	1.4 ± 0.51	40
stv-nel_01052_a	CAACCAAGGTATTTTTGCAAGACC	CCTTGTATGAAAAGTATGATGATGC	(ATC) ₅	159–169	2	1.2 ± 0.42	20
stv-nel_01728_a	TTCAATGGATCTTGAATATTTTGTTCG	TTACACATACAAACCTTCGGCAATC	(TTTA) ₅	110–118	2	1.3 ± 0.48	30
stv-nel_02122_a	GTTTTTACCATTGCGATTCGAGAC	AGATTGAGAAAGTCTTAACGGGC	(AAT) ₄	178–188	2	1.1 ± 0.31	10
stv-nel_02811_d	ACCTGACCACAAAACCAACAAG	CACCATATTCGTCTTCCACCAACTTC	(AGA) ₆	121–124	2	1.2 ± 0.42	20
stv-nel_03033_a	TTCCAAGTATTTACTGGCTTTGGC	ATAAATCCCCAAAATGCATCTTC	(TC) ₆	174–176	2	1.6 ± 0.51	60
stv-nel_03406_a	TTGGTGTAGCTAGTGAATAAGGATGAG	CAAAATTAGCATTTACTGGTGGGATG	(ATAC) ₆	164–174	3	1.9 ± 0.31	90
stv-nel_04248_a	TGTTTTTCGGTTTTGTAAGACCACC	CTCCACTGTCCAAAACCTCTCCTC	(GTGTAT) ₄	140–153	3	1.2 ± 0.42	20
stv-nel_04772_a	AGACAGAGAGGTAATGATGGCCC	ATCATCAACAGCAGCAGATCTTTG	(CAT) ₇	163–178	3	2.0 ± 0.00	100
stv-nel_04776_a	TTTGCATGCCAAATCTCTCTTC	AAATATCTATGGTGTCTCAAAAGCAGG	(TTTGT) ₄	210–216	3	1.2 ± 0.42	20
stv-nel_05023_a	GAGAAATTTGATGAAACTCACCGAG	AACAATTCGTCTTGGTTAAAGATGG	(AG) ₇	180–186	4	1.9 ± 0.31	90
stv-nel_05097_a	CGTCACCAAAAGATCTCCAATCTC	AAAAGGGGTGTTTTACAGGCTTAAC	(GAT) ₆	178–184	2	1.6 ± 0.51	60
stv-nel_05277_a	CAGCGCCATTTAGAAGCTGACTAC	AAITGCAACAGCATCAGAAAACCTC	(CTG) ₆	159–162	2	1.6 ± 0.51	60
stv-nel_05295_a	TCCAAATTAATGGTGGGATTTTC	GACCAAAAATAACATAATCGGATGG	(ATT) ₆	171–179	2	1.2 ± 0.42	20
stv-nel_05372_a	TTTTTCGTACGTTTAGTGCCATGTG	GGCTTCCAAGAAAACCACTTTTATC	(CCTC) ₄	177–179	2	1.6 ± 0.51	60
stv-nel_05532_a	TTTTCAAAGGGTTTTGTGAAAATGG	AGTAGAGCTTTCACCGCATCAAAC	(AG) ₈	142–153	4	1.9 ± 0.31	90
stv-nel_05827_a	TCAAATTTGAATGCGGGAAAACCTAGAG	GCAATGCATAACTCTTGTTTTTTGTAAAGG	(ATTA) ₆	198–202	2	1.8 ± 0.42	80
stv-nel_06049_a	TTGCTTTGATCATCACTCTCATCC	TGATGACAAAGGGAGTTTACTGGTG	(CAT) ₆	163–166	2	1.8 ± 0.42	80
stv-nel_06172_a	CACGTGAAAATGACCATAAGGACC	TTCCGATGTGCGATCTCTGTCTTC	(GA) ₆	128–143	2	1.7 ± 0.48	70
stv-nel_06465_a	TTTTCAAAGGCAGATGACAATGATG	TAGATGCTTTCCAAGCACAATCAG	(ATAC) ₇	173–178	3	1.3 ± 0.48	30
stv-nel_07181_a	AGTTCAAAAAGTTCGGATGTCCTG	GATGATCCCCAAAATCGTATTTAGAAG	(AC) ₆	178–182	3	1.7 ± 0.48	70
stv-nel_07204_b	TTCAACAAATTTGCACTGCTCTTTC	GATGGTAATTTTCACCCCGAATTTG	(TTG) ₄	156–159	2	1.2 ± 0.42	20
stv-nel_07374_a	CCAGCCATAATATCAAACACGGTC	CCCCCCCCCAACATTAACAGAAC	(ATCT) ₆	154–162	2	1.2 ± 0.42	20
stv-nel_07884_a	GGAATGGTCTAAGATTAACACCCCC	GGGTTTGTGAAAAGTGTGAGTGTGATG	(TTC) ₇	140–173	4	1.9 ± 0.31	90
stv-nel_08453_a	GCCATTTCTGTACGTGTTCCACAG	AAATAGTAAAACCTCGTTGGGCTCC	(AG) ₇	149–151	2	1.6 ± 0.51	60
stv-nel_08615_a	TGAGGCAACAAAAGGTTTCTTCTC	ACCACCAATTAACCTGCACAAAAAG	(TC) ₇	114–140	3	1.2 ± 0.42	20
stv-nel_08865_a	TTTCACAAAACACCTCTACAGTCCAG	GGACATCTTACAAAACCAAGTGGAG	(ATGT) ₁₀	224–261	5	2.0 ± 0.00	100
stv-nel_09674_a	TTGCGTATTTTCTTTCAAAGTCCG	AAGAAATTCAAATCCCTTCCAATG	(CA) ₇	169–181	2	1.3 ± 0.48	30

Table 3 continued

<i>Nephelium lappaceum</i> (rambutan)						
Marker	Forward 5' → 3'	Reverse 5' → 3'	Motif	Range	Amp	Alleles/sample %H
stv-nel_11131_a	TCAGATCAATGTCAATTTCTTCCCTC	GACAAATGAGAAGAAGATGGAGGC	(TTC) ₄	127–130	2	1.3 ± 0.48
stv-nel_11760_a	CAACAGAGACCTGAGGATTTCCC	AACCCACCTCAATCATAGACATC	(TC) ₆	161–169	2	1.7 ± 0.48
stv-nel_13028_a	GAAAGTTACGCCCTTTTGTCTTTCCG	GAATATGGATGCCACGTTTACAGG	(CAT) ₇	129–138	3	1.3 ± 0.48
stv-nel_13493_a	ATCTGCTCGACTTCAGAATGGC	AACGACGACGAAGAAGAAAGAAAG	(CTT) ₄	165–307	4	3.2 ± 0.42
stv-nel_15792_a	TTCTCTCAGATGTCTTTGGACTTTAGC	TGTATATATGTTGCTTGGATCCTTC	(AG) ₈	125–127	2	1.6 ± 0.51
<i>Manilkara zapota</i> (sapodilla)						
Marker	Forward 5' → 3'	Reverse 5' → 3'	Motif	Range	Amp	Alleles/sample %H
stv-maz_00161_a	ATGGTAGTGGTGATGGCGATAGAG	TTTGTGATCGATATTTTGTGTGGC	(CAT) ₄	166–325	4	2.50 ± 0.57
stv-maz_01419_a	GCGAGACTGAGGATGAAGAAGAAG	CACCTCAAAAACCCAAAGCAAAG	(GAG) ₄	120–124	2	1.25 ± 0.50
stv-maz_01644_a	TGAACAAAGCTTAAGAAAAGTGGCC	AATTAGCACACAGAACTGGGAACC	(GAA) ₅	166–171	2	1.50 ± 0.57
stv-maz_02138_a	GAAAGCAAAAATAGAGCCGGAAAC	TCAATGGTTAGTTCATCGTTTCAAATG	(TCT) ₄	141–143	2	1.75 ± 0.50
stv-maz_02156_b	ACGCTCTTCCCTTGTGTGATCTTC	ACTCGAAGAAGTCTTGAATGGCTG	(CTT) ₅	154–157	2	1.25 ± 0.50
stv-maz_02673_a	ATATTTATGCATTTGATGCGTGGAG	AACCTGCACCTGCTCTTGTTCAC	(CAA) ₄	116–120	2	1.50 ± 0.57
stv-maz_02898_a	ATCTGCAAAATCCACACATACAAG	ACAGCTTGTGTACTTTTGGCCATC	(TTG) ₅	153–157	2	1.25 ± 0.50
stv-maz_03523_a	AGAGCTTCTCCGATAGGATTTCCG	AGGCTGTCACAAAGAAAGAAAAG	(CTT) ₅	172–175	2	1.25 ± 0.50
stv-maz_03685_a	ATGGTATTCAGGTGGATGATGACG	CGGACAAACAGAGTACACAGCCATAC	(TTC) ₆	165–171	2	1.75 ± 0.50
stv-maz_03858_a	TTCCCAAATTCGAGTTTCTCAATTG	TGGTTCCCTTTTCTTTTACCCCTTC	(CT) ₈	154–163	3	1.50 ± 0.57
stv-maz_03945_a	TTGTTCATTTTGGAGTCTTGCTGC	CATGAAAAATGCCAAAATCCTAGC	(AGA) ₅	128–178	3	2.50 ± 0.57
stv-maz_04059_a	TTGTGTGATGAAAAAGGTACAGGC	TTCACAGTCGGCTTAAACAACCTCAG	(TCT) ₅	153–158	2	1.25 ± 0.50
stv-maz_04927_a	CAATATGGAGCTCATGAAAGACCC	CAAACTATGACCATCCCTTTCAGG	(GAC) ₅	106–121	3	2.25 ± 0.50
stv-maz_04989_a	GGCTGAAAAGATGAGTCACTCGAAG	CTGAATGAAAGTTGGTTGATCCTG	(AAG) ₅	163–169	2	1.50 ± 0.57
stv-maz_05004_a	CTGTGATGCAGAAACAAAATTGC	GTTTTCCCTCTCTTCTTCGAC	(TTC) ₄	124–129	2	1.25 ± 0.50
stv-maz_05716_a	TTTCATGTACCATATGCCCTTGCAG	GTTGGTTGGTTGCTTATGAGTGTG	(TTC) ₄	134–164	2	1.25 ± 0.50
stv-maz_05940_a	TGAGATTGATGATTTGCCACAG	GCTCAAGCGATGGGAGTAATAATG	(AGA) ₆	177–180	2	1.50 ± 0.57
stv-maz_05984_a	TTGCCATCGATTTTCTTCTTCTTC	AGCAAAAAGATAGGTCGTGGTGAG	(AGA) ₅	155–212	4	1.75 ± 0.50
stv-maz_06044_a	AGCATATCCTGGTCCCTCTCTTTC	AACAAGTGAGTTTTTGGCCCTCATC	(CTCTTT) ₄	151–179	6	3.00 ± 1.15
stv-maz_06325_a	AGGAAGAGCCATTGGAGTGTATTG	CCTCTGAAAGCCACTAGATTCATACTC	(GAC) ₅	173–181	3	1.25 ± 0.50
stv-maz_06448_a	ACTACGCTTTTACCTCCACGTC	TAAAGGAGCTCAGCCATGGAATAC	(ACAT) ₄	122–131	2	1.50 ± 0.57
stv-maz_06694_a	TTGAGTGGCGACTAGGGTTTAG	CCTGATGATCGCTTAAAGCATTG	(TGT) ₅	157–165	3	1.75 ± 0.50
stv-maz_06769_a	CTTGGCCACCCTCCACTACTAAC	GAATGGTGA CTGAAAGTGGAAAATG	(CAT) ₅	169–172	2	1.50 ± 0.57

Table 3 continued

<i>Manihara zapota</i> (sapidilla)	
Marker	Forward 5' → 3' Reverse 5' → 3' Motif Range Amp Alleles/sample %H
stv-maz_06856_b	AAGGAACTGCTTTTCTTCTC CAGAAATACAAAACCAATGGAATCG (AGAC) ₄ 139–143 3 1.75 ± 0.50 75
stv-maz_06859_a	TCAATTTGGTTCCTTGATTTATGG GGGACCTAATGGCTTACTTCTCTCATC (GAA) ₆ 142–168 4 2.00 ± 0.00 100
stv-maz_06932_a	GAAATGTGTGAATGACAGTACC AGAATCAACATTACCTACAAACCAGG (GAA) ₆ 110–116 3 1.75 ± 0.50 75
stv-maz_07228_a	AATGAGAGGTTGAGGAATGGAGTG AAGAGAACAGCAACACAACAGCAG (TGAT) ₄ 134–137 2 1.50 ± 0.57 50
stv-maz_07725_a	GGAGGACATTTTGAATTTGAAATC GGCGAGGTAACGGTCAGATAAATAC (AAG) ₆ 159–162 2 1.25 ± 0.50 25
stv-maz_08151_b	AAAGCAAGTAATCAGGGTTCCACC TTCATCGTTTGGGTTTCATCTTCTC (AGA) ₅ 138–244 6 2.00 ± 0.00 100
stv-maz_08175_a	TTGATGAAGAGGATGAGGAGAAC CTTAGCCCTCTTTGAGCAAACTG (GA) ₈ 163–178 3 1.75 ± 0.50 75
stv-maz_08303_a	AACCTGTTACAGTAGGACTTGCAC AATCTTTTGAACCCATCTCAGCC (TTG) ₄ 112–153 4 3.50 ± 0.57 100
stv-maz_08356_a	TGAGGATTTTTCATTTCTCCAG TCATGGAAATCAACATGGTAACGG (CTT) ₆ 170–173 2 1.25 ± 0.50 25
stv-maz_08633_a	GATGGCAAAGTGAACAATGGATAG TTTCTTGGCATGTTACAATGATCTG (AG) ₆ 174–177 2 1.25 ± 0.50 25
stv-maz_08689_a	GACTAGTATGGGAGTTCCGATTC ACAAATCATAACCCCTCTTGGCAAC (GA) ₇ 142–148 3 1.50 ± 0.57 50
stv-maz_09208_a	TCAGTACTCAGAAGTTACTAATGTCCGC TCATTTGGTCTTAGTAGTGTCCCTG (GAA) ₅ 159–171 3 1.75 ± 0.50 75
stv-maz_09621_a	TGTCGAACTTAGCAAGACCCCTC AGCACGTGTTCCCATAAAGAAAAG (GA) ₈ 124–128 2 1.50 ± 0.57 50
stv-maz_09673_b	GGTGCCATGTGTTCTATTTCAAGG TGTTGAACAAGCCCAACCCCTG (GAA) ₅ 138–275 4 1.50 ± 0.57 50
stv-maz_10106_a	TCCTCATATCGTTTCCACCACACTC AAAGATTCTGATATTTCCATTGTTG (CTT) ₆ 105–118 3 1.75 ± 0.50 75
stv-maz_10490_a	GGGATCTGCATTTTCTCGGTAAG GTAGAATAACCCACACAACCTCCGC (TGG) ₄ 182–551 3 2.00 ± 0.81 75
stv-maz_11051_a	AGGATTAATGCAATGAGGGAAAGTTG CCAGGGATGTGATACAAAGTGAATC (ATT) ₄ 165–178 2 1.75 ± 0.50 75
stv-maz_12181_b	CTGCTACATTTGCTGATAGCCCTTG TTCTCAATCACATTTGCTGCTTTTC (TCT) ₅ 151–157 2 1.25 ± 0.50 25
stv-maz_12205_c	AAGCACCTCATGATTAGAACTGC GTGCTGCACTATTGCTCATCTCAG (TC) ₇ 171–183 4 1.75 ± 0.50 75
stv-maz_12995_a	AGAGGTGCAAAAAGAAATGGATGTC TTATACCAGCCATATGCCCTTC (TC) ₈ 135–139 2 1.25 ± 0.50 25
stv-maz_13002_a	TTTTCTCCTTTTACATAGCCCTAGTTG GGAAACACCAAGGGTACACAAAAC (TTAT) ₄ 109–112 2 1.75 ± 0.50 75
stv-maz_13536_a	TGGTACCTTAGTTAATTTGGCATCG TGGATGTTGGACATGTTG (TGA) ₄ 124–127 2 1.25 ± 0.50 25
stv-maz_13697_a	TACAAAAGTAGAAGGAGCTCAGCC GTCACGTCCTTAACCCACAGAG (GAT) ₄ 158–170 3 1.50 ± 0.57 50
stv-maz_14601_a	TGAGTGGAGCAGATCTCAGAACAC TTATAGCTTAGCTTACACAGCAGC (TC) ₆ 165–169 2 1.25 ± 0.50 25
<i>Litchi chinensis</i> (lychee)	
Marker	Forward 5' → 3' Reverse 5' → 3' Motif Range Amp Alleles/sample %H
stv-lic_00007_a	TCGCTTTAGGGTTTTCTTCTGCTG CGAAACCACCGTATTATTCCATTTTC (GA) ₆ 178–188 4 1.9 ± 0.31 90
stv-lic_00456_a	GTGTAAAACACAACGACGACGGAAG AAAACAGTAAACGAAAGCCAAAACCTGTG (CTCA) ₄ 135–169 4 2.0 ± 0.47 90
stv-lic_00551_a	GTTTGGCACTATCTCGTAACCACC ATGATGTGAATCGGGTTCAAAGAAAG (CAT) ₄ 149–158 2 1.6 ± 0.51 60
stv-lic_00661_a	GTTCCGAGTCTGTCAATTTCCCTC TCAAAGAGAGGTTGTTGTTGTTG (CAAA) ₄ 147–172 2 1.7 ± 0.48 70

Table 3 continued

Litchi chinensis (lychee)

Marker	Forward 5' → 3'	Reverse 5' → 3'	Motif	Range	Amp	Alleles/sample	%H
stv-lic_00878_a	TATGGACCGAATTCTCCTTCATTG	CCAATCTTCACAACCCAAAATAGC	(AGA) ₈	174–326	4	2.7 ± 0.94	90
stv-lic_01270_a	ATCACTCTATGCATCACTTGCAGC	TTCTAACACCAATTCTCTGTCTCAGG	(ACA) ₅	164–167	3	1.4 ± 0.51	40
stv-lic_01347_a	GAAGCCACAAGAGAAAGATTGACG	AAACACAACAACCCCAATTACCCAC	(TGG) ₅	173–180	3	1.5 ± 0.52	50
stv-lic_02612_a	CGCAGATTGACAGAACAGAGATTG	ACCCAAGTACGCCCTTTTCCTTTTAG	(AG) ₆	124–140	3	1.9 ± 0.31	90
stv-lic_03732_a	GAAGTCCAAAACCCAGTTTCATTTC	TTATCGATGTAAACCGCTTTGTTTG	(ATCA) ₄	174–178	2	1.2 ± 0.42	20
stv-lic_04081_b	TCGAAATATGCCAGCCTTATAACC	TGGTCATAITTCATGATGTGTCTGC	(ACA) ₅	126–128	2	1.2 ± 0.42	20
stv-lic_04717_a	GTCAGGTTGGTTCGATGTGTTG	CGCTGTAGGCTTTTCTTAGCTG	(TTTTG) ₄	98–122	5	2.1 ± 0.87	70
stv-lic_05050_a	CACCAGAAAGGATGATTTACAGAG	TATTTGGCAATTGGACTTTTCTC	(AAAT) ₄	131–135	2	1.2 ± 0.42	20
stv-lic_05155_a	CGACAAATGCATTCACATACAG	TCTGGTCAACTTTCTTCACAATCG	(ATAC) ₄	115–137	3	1.9 ± 0.31	90
stv-lic_05167_a	AACGTTCCAAACATGAAACCAAGAC	TAGGGGGCTTTTATAATCAGGACG	(TACA) ₅	104–132	4	1.9 ± 0.31	90
stv-lic_05730_a	TCGTGTTGGGTTACATAAAAGTTG	AGCTTGTAGGAAAATAAGGGTGGG	(GA) ₈	125–150	4	1.9 ± 0.31	90
stv-lic_05952_a	CACTTTGAAGAGACAGAAGCCACC	CACGAGCATTTGTTTGAGTTGAAG	(ATC) ₄	158–161	2	1.5 ± 0.52	50
stv-lic_06125_a	ATGGAGAATGAATCAGTCGGAGAC	AAACAGGCAAATAATGAGAAAAGCG	(CAT) ₇	132–139	3	1.3 ± 0.48	30
stv-lic_06240_a	CCACCATTAATTAACACTTTCGCC	AACTGGATGATGAGGATCAGGAAG	(CT) ₆	138–140	2	1.5 ± 0.52	50
stv-lic_06505_a	ACACAGGAGATGAGGCAAGTTAATG	CCAGATAAGTTTCTCTGTGCTCG	(TGTT) ₄	181–185	2	1.6 ± 0.51	60
stv-lic_06578_a	GACCAATCCTTCAGAGAAAAGAAC	TCAGTTGATATGCACCAATTAAGC	(AAC) ₆	163–183	4	1.8 ± 0.63	70
stv-lic_06771_a	GCGAGACTCAAAAATTACATATCACTCC	TTCAGTTTCAGACCAAAGGTTTATCTC	(ATAC) ₄	163–179	2	1.5 ± 0.52	50
stv-lic_06871_a	ATCCAAAAGAGGAATCAGAAAGACC	GTTCCGATGACATGCTTCTCTCTC	(GA) ₆	140–143	2	1.2 ± 0.42	20
stv-lic_06873_a	TGGTTCCATGGAGAATAATAATACGAG	GTAGCGCAATGAAACCAAAAGAATC	(TTG) ₄	131–138	3	1.8 ± 0.42	80
stv-lic_07043_a	ATAACGACATCCAAAGTGGAGAAG	ACCTGTCAACAAGAACCCGAATAG	(GGT) ₇	139–150	3	2.2 ± 0.91	70
stv-lic_07417_a	ACCATTTCAGTAAACTATGGGTGGTC	CCACACATCAATTTCTAAGAACAATATCG	(TCA) ₈	167–181	3	1.7 ± 0.48	70
stv-lic_07423_a	TTCTGATTTTAAATTTGTCAGGTG	ACAGAAGACCAAGATTGCAGAGAG	(CT) ₇	131–152	2	1.6 ± 0.51	60
stv-lic_07476_a	CCTCGTTTGGCAATTTGTAATGAAAC	CCATCATCACTTAATCTTTTGCCC	(GGT) ₄	151–154	2	1.4 ± 0.51	40
stv-lic_07889_a	AGTAGAACCCACCACTTTGGTCTG	CCAAGGCTGTCTTTCGGATTTAG	(TGT) ₄	150–153	2	1.2 ± 0.42	20
stv-lic_08181_a	TATAAATTTCCACCCTGCTTTGTGTG	CTCGTTTTAAAGCAAAAAGCCTAGC	(TA) ₆	156–162	3	1.7 ± 0.48	70
stv-lic_08315_a	GAATGAAGATAAAAACCCAGATAACAGACG	TCTTGATGCACCAAGAAAAGTTAG	(GAA) ₅	138–141	2	1.7 ± 0.48	70
stv-lic_09862_a	TTGAGTGTGGTGTGTTTGACAATG	TCCATCTCTGTTTTTGTAACTGGC	(TGACA) ₄	172–179	2	1.2 ± 0.42	20
stv-lic_10478_a	TTTTAATGTGGAGATGGTTTTTGGG	GCGGTTGGTGTTCGCAATTTTATAC	(GGTT) ₄	175–179	2	1.7 ± 0.48	70
stv-lic_10896_a	AACCAGAGATGGTGGAGTGGAG	AGTAAACACGAAACGAGAAITGGG	(AG) ₇	145–173	4	1.7 ± 0.48	70
stv-lic_13448_a	ACCGTACTCTCCATTACAACGCTC	TTGGGAACATAATTTCTCCACAC	(GAT) ₅	164–167	2	1.5 ± 0.52	50
stv-lic_14104_b	AGTAAATGCGTCACTCATGGATCG	TGACAGATGACTGAAGAGGCTGAG	(GGAC) ₅	140–144	2	1.6 ± 0.51	60

Table 3 continued

Forward 5' → 3'		Reverse 5' → 3'		Marker	Range	Amp	Alleles/sample	%H
<i>Litchi chinensis</i> (lychee)								
Marker	Forward 5' → 3'	Reverse 5' → 3'	Marker	Range	Amp	Alleles/sample	%H	
stv-lic_16470_b	CTTCGTCAGTACAAGGAGGAGGAG	AACCACCTTCAATGCCATAGAGCC	(AGT) ₄	153–168	3	1.9 ± 0.31	90	
stv-lic_18234_a	TGAGCTTAAAGGCATGATACCTTTTCG	CCTTTTAGAGATGCTCAAAGTCTGC	(TACA) ₆	108–116	3	1.9 ± 0.31	90	
stv-lic_19633_a	CCCCATCTTCATTTTATTATTGTTG	ATGTGGGTATCTTTCTTTTCAGCC	(TTGT) ₇	152–169	3	1.9 ± 0.31	90	
<i>Garcinia mangostana</i> and <i>Garcinia cochinchinensis</i> (mangosteen)								
Marker	Forward 5' → 3'	Reverse 5' → 3'	Marker	Range	Amp	Alleles/sample	%H	
stv-gam_00231_a	GTTGCACCTCCTCCGAGGTCAG	TTCTTTTTGATTTCTTGCAGGTGG	(TTG) ₄	107–188	6	1.50 ± 0.58	50	
stv-gam_00278_a	TTTGGAGTAGCACTTACCAAAAGGG	GATTTGAATCTTCACCACAACCCCTC	(GAT) ₅	168–256	4	1.25 ± 0.50	25	
stv-gam_00546_b	ATACACCTCATACAACCTCCGGCTC	CACAGGGATAGGGATAGGGATAGG	(CAT) ₄	143–218	9	3.25 ± 0.50	100	
stv-gam_00560_a	GATAAAAGAGGCAATGTGTGAGGG	TGCAACAAAGAAACAACACCACCTC	(ACT) ₄	135–141	3	1.50 ± 0.57	50	
stv-gam_00645_a	AGAAAGCTCAAGTCTGCTTGGTG	ACTCAGAAGAAAGGAATTTCCACGC	(TGT) ₅	174–191	8	2.00 ± 1.41	50	
stv-gam_00660_a	CTAGCCACTCATGGTGGTAAAGTG	TAAAAGCCAGAAAGGAGACTCGAC	(GAA) ₄	108–228	4	1.25 ± 0.50	25	
stv-gam_01553_a	TGAACCTGCTGTGCTGCTCTG	TCGAACTGGTGTAGAGGTAGAGG	(CCTCTA) ₄	98–271	7	2.25 ± 1.25	75	
stv-gam_01788_b	TCCCCATTTCCATCTCTTAAACATC	TTGGATTAATAAAAATGGGTGGTTC	(TTC) ₄	172–214	4	1.25 ± 0.50	25	
stv-gam_01820_a	AGAGAAGACCTGTGCGACATAGG	AGCGACTTGTTAGGGAAAAGGC	(TCT) ₆	166–301	5	1.25 ± 0.50	25	
stv-gam_01864_a	GTTACAACATCTTTAGGTTCCGCCG	TTAGAGGTGACAAGGGAGGATGAG	(TATT) ₄	178–377	4	1.00 ± 0.00	0	
stv-gam_01886_a	AAGAATAGACCGATTGCCGATATG	CCTACCTAAAAATGGACCCAGCTTC	(GGA) ₄	138–274	2	1.25 ± 0.50	25	
stv-gam_01984_a	TGAGTAAAGAAAAGAGGTGCTCGC	ATGGATTTTCGAAAGGTTTCATGC	(ACA) ₄	158–233	2	1.00 ± 0.00	0	
stv-gam_02195_a	AGAAACAGACCAGAAATGTGAGGG	TTTGTGTGATTTGCTAGTGTGGATTG	(TTC) ₄	160–344	7	1.75 ± 0.95	50	
stv-gam_02824_a	CAGTGGTAGCTCGCTCCTAGAAATG	ATCTCATCTCTGATCCCTCTGGGGTG	(AAT) ₄	164–180	3	1.50 ± 0.57	50	
stv-gam_02895_a	ACAGCCACAATAGTCACTCCCTCCTC	TTTGGTTGTTTTGTAGTAGGTTTCTG	(AAC) ₁₀	156–185	6	1.75 ± 1.50	25	
stv-gam_03207_a	AAATGATCACAAAATTCACCCCCAC	GGATGGAATTAACAACGTTACAACATTAC	(GAT) ₄	159–231	5	1.75 ± 0.50	75	
stv-gam_03342_a	ACCTACCTCCAGCTGCTGATTTG	AGATTGCAACCTCAAGAAGACTGCTC	(CCA) ₄	111–130	2	1.25 ± 0.50	25	
stv-gam_03495_a	TTCGAGGAAGGATAAGTTGTTTG	CATAAACCAAAACCATCAAGAACC	(GGA) ₆	163–203	7	1.75 ± 0.95	50	
stv-gam_03790_a	CTTCTCAATGATCCCCATGTTTG	AAGGTTCTTTTGCCTTTTGTTC	(CCA) ₄	97–304	5	2.00 ± 1.41	50	
stv-gam_03796_a	GGATGTGAGTGAAGTTAGTGACCG	TATAATCCATCATCACCCATGACG	(TATG) ₄	97–388	11	2.75 ± 1.70	75	
stv-gam_04008_a	TTCTTTGGTTTCTTGACGCTTAGG	TCATCAACCCCACTAAAACCTCCAC	(TTC) ₄	148–176	6	1.75 ± 0.95	50	
stv-gam_04053_a	TAGACAAGGACAAGTGCAAAGTCCC	CTAAGCACTACTTCTGCCAGCCAC	(AGA) ₇	140–168	6	2.00 ± 1.41	50	
stv-gam_04292_a	ATCATGATCTGCAGCAATATGCC	AGTTACATGAATATGACGACGGGG	(CTC) ₄	147–163	5	1.75 ± 0.95	50	
stv-gam_04801_a	CATCATCTTCTCTCCCTCTTGTCC	AGACATGCTTGCAGTTCTAGTCCC	(CTT) ₄	108–111	2	1.00 ± 0.00	0	
stv-gam_05115_a	TTGATGGTAAATGTGGGATTTGATG	GAGTCTGTCTCACATCTGCAACC	(ATG) ₆	112–245	9	3.25 ± 1.50	100	

Table 3 continued

<i>Garcinia mangostana</i> and <i>Garcinia cochinchinensis</i> (mangosteen)									
Marker	Forward 5' → 3'	Reverse 5' → 3'	Motif	Range	Amp	Alleles/sample	%H		
stv-gam_05136_a	GTGGTCCATGTATAGGTGGGATG	ATTACCCATGGCAGTTGGCTC	(TA) ₆	126–445	4	1.00 ± 0.00	0		
stv-gam_05237_a	CAACAGCCATGCCTCGTTACTAC	CAAGAGACGGCGTTAGGAAATTAC	(TC) ₈	151–170	7	2.00 ± 0.81	75		
stv-gam_05474_a	CAAAGCCACCAACTTACCACAAAAC	TGGTTTTAGAGGATGACGTGTGAG	(TCT) ₅	113–283	4	1.50 ± 0.57	50		
stv-gam_05662_a	TGGATTTGTTAGGGTTAGGGTTTG	ACCCCTCCATTACTCCCTCTAC	(TG) ₈	123–333	6	1.75 ± 1.50	25		
stv-gam_05897_a	CTCTCACTTCCCTCTCTTTGGATGG	ATGATGATGACGATGATGACAAATG	(TCG) ₄	147–291	4	1.75 ± 0.95	50		
stv-gam_06047_a	GAAAGTAGATATGTGGAGCAAGCC	AAATTGAGAGTTTCCCTTTTGAGC	(ACA) ₄	177–481	5	1.50 ± 0.57	50		
<i>Guadua angustifolia</i> and <i>Bambusa vulgaris</i> (bamboo)									
Marker	Forward 5' → 3'	Reverse 5' → 3'	Motif	Range	Amp	Alleles/sample	%H		
stv-gua_00022_a	AAAAGAAGGGAAAGAGGAAAGGAGG	CTCTCTCCCTTCTGGACTGACAAG	(GA) ₄	122–326	3	2.0 ± 1.18	50		
stv-gua_00085_a	TCTCCCTACCCTATCTCTCTCTCG	CTTGGAGTTTCATGCACCACCTGTA	(TC) ₄	151–160	2	1.0 ± 0.00	0		
stv-gua_00089_a	TAATCGAGCTGGTTACGAGGAAAA	CCGTGTTCCCTCCCGTACTCT	(ATAA) ₄	177–180	2	1.1 ± 0.70	30		
stv-gua_00145_a	TCTCCAAAAGTTGGCTTTCTGATTC	TAGGCCTCAAGCAAAATCAATCTTC	(CT) ₅	144–150	3	1.1 ± 0.53	20		
stv-gua_00273_a	CCTTGGAGTAGAGGAGGGCAATAG	CCCTCTCCTTCTTCTCTCTCTCGCT	(AG) ₄	96–249	3	1.7 ± 0.45	70		
stv-gua_00307_a	CTCTACCTCGACGTTTCATGTCA	GAAGGAGATATTCAAACGGTGTATGG	(TC) ₄	111–440	8	2.7 ± 1.41	90		
stv-gua_00323_a	AGTCCAGAGAACTCAGACCAGAGC	GCAGTTTGGCACAACATTTGTTTA	(CT) ₄	173–181	2	1.0 ± 0.77	30		
stv-gua_00354_a	ATGGAGGAGATGAATACGGAAAGAA	AAGGTTGTAGTTTTTGGATGGGA	(AC) ₄	163–165	2	0.9 ± 0.70	20		
stv-gua_00445_a	ATTTAGGTCAGAAATGGATCCAGGG	TGTGTGTTCTGACTCTATTGGATCAAG	(AGT) ₅	174–180	2	1.0 ± 0.00	0		
stv-gua_00461_a	GCCAAATCCAAACAATGTAAACAGA	GTTTGTGGCAAGGATGTGACAATA	(CA) ₈	163–173	3	1.6 ± 0.80	60		
stv-gua_00511_a	AAATCAATCGACGGGATGAGAAAT	ATAAGCCGTCGGATCCCTATCTAC	(AG) ₁₀	168–180	2	1.4 ± 0.80	60		
stv-gua_00558_a	TCTCGCAGACTAATAATGGCAGGT	CCCTAGTAATTCAGAGAAACGGGA	(TCT) ₄	136–159	5	2.0 ± 0.77	70		
stv-gua_00578_b	CCACTCTTTGTTCTCCAAATCTCCA	GAAGAGGAGAATGAGACGGAGCTA	(CG) ₄	178–370	3	1.9 ± 1.22	50		
stv-gua_00638_a	CACCTCCAGCAATCTTCTTCAACCT	ATCGAGGAGTTGGGTATAACCGT	(CTT) ₄	159–181	2	1.1 ± 0.30	10		
stv-gua_00670_a	GACTAGACATGCTCCGATTTGACA	AGCATTTGCTTCTTCCCTCACAC	(AG) ₇	101–182	3	2.3 ± 1.18	60		
stv-gua_00709_a	ACGCAAAAACGAGGACGGTATAGTA	CAAAAGCAACTAAAAGCAAAAGGGGA	(TC) ₄	121–131	3	2.0 ± 0.77	70		
stv-gua_00841_a	TATTTGGAAGCTAGTGCCACAACAA	CGACATGGACAACATTTGACTGATTT	(TGCAC) ₄	143–145	2	1.4 ± 0.66	30		
stv-gua_00878_a	TCAGGATATAGGCGACAGAGCGGA	GACTCGATCCGACCGACGAT	(GTG) ₄	111–128	4	2.2 ± 0.87	80		
stv-gua_00917_a	CAACGATGCTAGCCCTTCTATTCCG	CTACTCCGGTACTACTCTCCACGCC	(AC) ₄	112–467	2	1.7 ± 0.45	70		
stv-gua_00918_a	GCTGATGCTGCTGCTGTACTCTTT	CCACCACGCAAAAACCTCTATAAAA	(AGCT) ₄	171–174	2	0.7 ± 0.45	0		
stv-gua_01020_a	ATCACCCGATCTGTACTCTGAAGC	GCAGATCCAGTTGTTTCGTTTCTT	(ACC) ₆	115–124	3	1.1 ± 0.53	20		
stv-gua_01140_a	ATCTTGTTCCTCCAACTCTCTCCAC	TTTTCTTCTCTCTCTCTCTCTCTCT	(GA) ₈	138–160	2	2.1 ± 0.30	100		

Table 3 continued

Marker	Forward 5' → 3'	Reverse 5' → 3'	Motif	Range	Amp	Alleles/sample	%H
stv-gua_01211_a	GGGTCTGGTCAGCCATCTTACTTT	TCCAGTTTGAGTTTCCATCCATTT	(TC) ₈	102–155	6	2.4 ± 1.56	60
stv-gua_01473_a	TTACAAAGGCACAATCTACACTCC	GGACCAAGAGCCGAAAGAGC	(TC) ₈	113–138	2	2.5 ± 0.67	100
stv-gua_01515_a	ACCTTCCTACTGCCTCCTCCTTT	GAGGATGGGGTAGGTGAAAGCTC	(CT) ₄	158–187	3	1.3 ± 0.45	30
stv-gua_01550_a	GAGTCAACAATCCAAACATCTCCC	CTTTGAGACGTCGTGTGATTCAGT	(AG) ₆	145–156	4	1.9 ± 0.70	90

%H: Percentage of heterozygosity of the samples at each locus

use of genetic resources for breeding purposes (Yamanaka et al. 2019). Even in current times when genotyping by sequencing (GBS) has become inexpensive, SSRs are still the preferred effective, robust, reproducible and simple to use tool to determine genetic diversity of landraces of maize and preserve rare allele sources; as SSRs do not require large bioinformatics infrastructure and expertise for data analysis (Hayano-Kanashiro et al. 2017).

Germplasm conservation and genetic population studies can be performed with a small number of markers, many have used between 8 and 17 SSRs (Amici et al. 2019; Ekue et al. 2009; Moraes et al. 2013; Samsir et al. 2016; Silva-Junior et al. 2016; Viruel and Hormaza 2004). Indeed, the most common number of loci that had been used in population studies of wild species was six and usually no more than twelve (Koskinen et al. 2004). However, for the estimation of the population-genetic parameter θ ($4N\mu$) a linear gain in accuracy occurs when increasing the count of loci from 1 to 100 (Carling and Brumfield 2007), and the theoretical quantity of loci for accurate evolutionary inference was estimated between 30 and hundreds of loci (Pollock et al. 1998; Takezaki and Nei 1996). Thus, the number of nuclear SSR markers provided in the present work for rambutan, sapodilla, lychee, mangosteen (two species) and bamboo (two species) would allow to meet those theoretical ideal figures for each of these groups. In addition, we report 26–47 top quality polymorphic markers for each for the species, which are sufficient for screening large number of samples and facilitate their correct identification and conservation in banks of germplasm; whereas for more in depth characterization of population-genetic parameters we provide hundreds of SSRs. Regarding the level of polymorphism of the markers reported here, 30–50%, is probably underestimated given the small number of accessions (4–10 per group) used for testing these markers.

Conclusions

The seven perennial plant species considered in the present study; rambutan, sapodilla, mangosteen, lychee and bamboo, represent an important genetic resource for the people living in the tropics. The markers reported here will help to generate




Fig. 5 Examples of what was considered good quality simple sequence repeat (SSR) markers in the present work. Three markers of *Manilkara zapota* (Stv_maz_13697; Stv_maz_06044; Stv_maz_06859) tested on four DNA samples

and showing discrimination of all the samples tested. “x” axis corresponds to amplicon sizes in base pairs (bp), “y” axis for all the markers was set at a maximum of 30,000 fluorescent units

information in relation to conservation genetics and breeding programs for these species.

Acknowledgements This work was supported by USDA-ARS project number 6044-21000-005-00D and the U.S. National Plant Germplasm System (NPGS). Any library/sequence information requirements can be addressed to Dr. Renee S. Arias at Renee.Arias@usda.gov. Mention of trade names or commercial products in this article is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the US Department of Agriculture.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest directly or indirectly and informed consent to publish this study.

References

- Aguilar R, Calvino A, Ashworth L, Aguirre-Acosta N, Carbone LM, Albrieu-Llinas G, Nolasco M, Ghilardi A, Cagnolo L (2018) Unprecedented plant species loss after a decade in fragmented subtropical Chaco Serrano forests. *PLoS ONE* 13(11):e0206738. <https://doi.org/10.1371/journal.pone.0206738>
- Alroy J (2017) Effects of habitat disturbance on tropical forest biodiversity. *P Natl Acad Sci USA* 114(23):6056–6061. <https://doi.org/10.1073/pnas.1611855114>
- Amici AA, Nadkarni NM, DiBlasi E, Seger J (2019) Contrasting effects of host tree isolation on population connectedness in two tropical epiphytic bromeliads. *Am J Bot* 106(12):1602–1611. <https://doi.org/10.1002/ajb2.1391>
- Armstrong K (2010) Systematics and biogeography of the pantropical genus *Manilkara* Adans. (Sapotaceae) Dissertation. University of Edinburgh
- Arias RS, Borrone JW, Tondo CL, Kuhn DN, Schnell RJ (2012) Genomics of tropical fruit tree crops. In: Schnell RJ, Privadharshan PM (eds) *Genomics of tree crops*. Springer, New York, p 369

- Arias RS, Martínez-Castillo J, Sobolev VS, Blancarte-Jasso NH, Simpson SA, Ballard LL, Duke MV, Liu XF, Irish BM, Scheffler BE (2015) Development of a large set of microsatellite markers in Zapote mamey [*Pouteria sapota* (Jacq.) H.E. Moore & Stearn] and their potential use in the study of the species. *Molecules* 20(6):11400–11417. <https://doi.org/10.3390/molecules200611400>
- Barcaccia G (2009) Molecular markers for characterizing and conserving crop plant germplasm. In: Mohan Jain S, Brar DS (eds) *Molecular Techniques in Crop Improvement*, vol Part 2. Springer, Netherlands, pp 231–254
- Bretting P, Bennet R (2007) The national plant germplasm system: an overview. *Phytopathology* 97:S150
- Buerki S, Lowry PP II, Alvarez N, Razafimandimbison SG, K'pfer P, Callmander MW (2010) Phylogeny and circumscription of Sapindaceae revisited: molecular sequence data, morphology and biogeography support recognition of a new family Xanthoceraceae. *Plant Ecol Evol* 143:148–159. <https://doi.org/10.5091/plecevo.2010.437>
- Carling MD, Brumfield RT (2007) Gene sampling strategies for multi-locus population estimates of genetic diversity (theta). *PLoS ONE* 2(1):e160. <https://doi.org/10.1371/journal.pone.0000160>
- Chaluvadi SR, Young P, Thompson K, Bahri BA, Gajera B, Narayanan S, Krueger R, Bennetzen JL (2018) *Phoenix* phylogeny, and analysis of genetic variation in a diverse collection of date palm (*Phoenix dactylifera*) and related species. *Plant Divers* 41(5):330–339. <https://doi.org/10.1016/j.pld.2018.11.005>
- Cousins S (2020) Bushfires expose weaknesses in Australia's health system. *Lancet* 395(10219):175–176. [https://doi.org/10.1016/S0140-6736\(20\)30096-9](https://doi.org/10.1016/S0140-6736(20)30096-9)
- Curry HA (2017) From working collections to the World Germplasm Project: agricultural modernization and genetic conservation at the Rockefeller Foundation. *Hist Philos Life Sci* 39(2):5. <https://doi.org/10.1007/s40656-017-0131-8>
- Ekue MR, Gailing O, Finkeldey R (2009) Transferability of Simple Sequence Repeat (SSR) Markers Developed in *Litchi chinensis* to *Blighia sapida* (Sapindaceae). *Plant Mol Biol Report* 27:570–574. <https://doi.org/10.1007/s11105-009-0115-2>
- Ellegren H (2004) Microsatellites: simple sequences with complex evolution. *Nat Rev Genet* 5(6):435–445. <https://doi.org/10.1038/nrg1348>
- Escobar H (2019) Amazon fires clearly linked to deforestation, scientists say. *Science* 365(6456):853. <https://doi.org/10.1126/science.365.6456.853>
- Finocchiaro A (2020) Australian Tropical Fruits. <https://www.australian-tropical-foods.com/index.php/exotic-fruits/>. Accessed 10 Feb 2020
- Hayano-Kanashiro C, Martínez de la Vega O, Reyes-Valdés MH, Pons-Hernández J-L, Hernández-Godínez F, Alfaro-Laguna E, Herrera-Ayala JL, Vega-Sánchez MC, Carrera-Valtierra JA, Simpson J (2017) An SSR-based approach incorporating a novel algorithm for identification of rare maize genotypes facilitates criteria for landrace conservation in Mexico. *Ecol Evol* 7(6):1680–1690. <https://doi.org/10.1002/ece3.2754>
- Hock S, Mahani M, Choong C, Salma I (2005) Transferability of SSR markers from lychee (*Litchi chinensis* Sonn.) to pulasan (*Nephelium ramboutan-ake* L.). *Fruits* 60:379–385. <https://doi.org/10.1051/fruits:2005043>
- Kijas JM, Fowler JC, Garbett CA, Thomas MR (1994) Enrichment of microsatellites from the citrus genome using biotinylated oligonucleotide sequences bound to streptavidin-coated magnetic particles. *Biotechniques* 16(4):656–660
- Koskinen MT, Hirvonen H, Landry PA, Primmer CR (2004) The benefits of increasing the number of microsatellites utilized in genetic population studies: an empirical perspective. *Hereditas* 141(1):61–67. <https://doi.org/10.1111/j.1601-5223.2004.01804.x>
- Larranaga N, Albertazzi FJ, Fontecha G, Palmieri M, Rainer H, van Zonneveld M, Hormaza JI (2017) A Mesoamerican origin of cherimoya (*Annona cherimola* Mill.): implications for the conservation of plant genetic resources. *Mol Ecol* 26(16):4116–4130. <https://doi.org/10.1111/mec.14157>
- Lin H-Y, Hao Y-J, Li J-H, Fu C-X, Soltis PS, Soltis DE, Zhao Y-P (2019) Phylogenomic conflict resulting from ancient introgression following species diversification in *Stewartia* s.l. (Theaceae). *Mol Phylogenet Evol* 135:1–11. <https://doi.org/10.1016/j.ympev.2019.02.018>
- Martínez-Castillo J, Arias RS, Andueza-Noh RH, Ortiz-García MM, Irish BM, Scheffler BE (2019a) Microsatellite markers in Spanish lime (*Melicoccus bijugatus* Jacq., Sapindaceae), a neglected Neotropical fruit crop. *Genetic Res Crop Evol* 66(7):1371–1377. <https://doi.org/10.1007/s10722-019-00815-4>
- Martínez-Castillo J, Blancarte-Jasso NH, Chepe-Cruz G, Nah-Chan NG, Ortiz-García MM, Arias RS (2019b) Structure and genetic diversity in wild and cultivated populations of Zapote mamey (*Pouteria sapota*, Sapotaceae) from southeastern Mexico: its putative domestication center. *Tree Genet Genomes* 15(4):ARTN61. <https://doi.org/10.1007/s11295-019-1368-z>
- Moraes RC, Vivas CV, Oliveira FA, Menezes IP, van den Berg C, Gaiotto FA (2013) Microsatellite markers for an endemic Atlantic Forest tree, *Manilkara multifida* (Sapotaceae). *AoB PLANTS* 5:plt006. <https://doi.org/10.1093/aobpla/plt006>
- Mukherjee S, Stamatis D, Bertsch J, Ovchinnikova G, Katta HY, Mojica A, Chen I-MA, Kyripides NC, Reddy T (2018) Genomes OnLine database (GOLD) vol 7: updates and new features. *Nucleic Acids Res* 47(D1):D649–D659. <https://doi.org/10.1093/nar/gky977>
- NPGRS (2020) U.S. National Plant Germplasm System. <https://npgsweb.ars-grin.gov/gringlobal/query/summary.aspx>. Accessed 10 Feb 2020
- Pollock DD, Bergman A, Feldman MW, Goldstein DB (1998) Microsatellite behavior with range constraints: parameter estimation and improved distances for use in phylogenetic reconstruction. *Theor Popul Biol* 53(3):256–271. <https://doi.org/10.1006/tpbi.1998.1363>
- Reyes-Valdés MH, Burgueno J, Singh S, Martínez O, Sansaloni CP (2018) An informational view of accession rarity and allele specificity in germplasm banks for management and conservation. *PLoS ONE* 13(2):e0193346. <https://doi.org/10.1371/journal.pone.0193346>

- Richard GF, Kerrest A, Dujon B (2008) Comparative genomics and molecular dynamics of DNA repeats in eukaryotes. *Microbiol Mol Biol Rev* 72(4):686–727. <https://doi.org/10.1128/MMBR.00011-08>
- Samsir SA, Bunawan H, Yen CC, Noor NM (2016) Dataset of SSR markers for ISSR-Suppression-PCR to detect genetic variation in *Garcinia mangostana* L. in Peninsular Malaysia. *Data Brief* 8:1438–1442. <https://doi.org/10.1016/j.dib.2016.08.016>
- Sharma PC, Grover A, Kahl G (2007) Mining microsatellites in eukaryotic genomes. *Trends Biotechnol* 25(11):490–498. <https://doi.org/10.1016/j.tibtech.2007.07.013>
- Silva-Junior JA, de Souza FD, Moraes RC, Gaiotto FA (2016) Development of microsatellite markers for *Manilkara maxima* T.D. Penn. (Sapotaceae) and their use in conservation genetics. *Mol Biol Rep* 43(6):451–455. <https://doi.org/10.1007/s11033-016-3981-3>
- Singhal P, Bal LM, Satya S, Sudhakar P, Naik SN (2013) Bamboo shoots: a novel source of nutrition and medicine. *Crit Rev Food Sci Nutr* 53(5):517–534. <https://doi.org/10.1080/10408398.2010.531488>
- Takezaki N, Nei M (1996) Genetic distances and reconstruction of phylogenetic trees from microsatellite DNA. *Genetics* 144(1):389–399
- Techen N, Arias RS, Glynn NC, Pan Z, Khan I, Scheffler BE (2010) Optimized construction of microsatellite-enriched libraries. *Mol Ecol Res* 10:508–515. <https://doi.org/10.1111/j.1755-0998.2009.02802.x>
- Vieira LD, dos Anjos KG, Faoro H, Fraga HPD, Greco TM, Pedrosa FD, de Souza EM, Rogalski M, de Souza RF, Guerra MP (2016) Phylogenetic inference and SSR characterization of tropical woody bamboos tribe Bambuseae (Poaceae: Bambusoideae) based on complete plastid genome sequences. *Curr Genet* 62(2):443–453. <https://doi.org/10.1007/s00294-015-0549-z>
- Viruel MA, Hormaza JI (2004) Development, characterization and variability analysis of microsatellites in lychee (*Litchi chinensis* Sonn., Sapindaceae). *Theor Appl Genet* 108(5):896–902. <https://doi.org/10.1007/s00122-003-1497-4>
- Yamanaka S, Hosaka F, Matsumura M, Onoue-Makishi Y, Nashima K, Urasaki N, Ogata T, Shoda M, Yamamoto T (2019) Genetic diversity and relatedness of mango cultivars assessed by SSR markers. *Breed Sci* 69(2):332–344. <https://doi.org/10.1270/jsbbs.18204>
- Zane L, Bargelloni L, Patarnello T (2002) Strategies for microsatellite isolation: a review. *Mol Ecol* 11(1):1–16. <https://doi.org/10.1046/j.0962-1083.2001.01418.x>
- Ziya ME, Kafkas S, Khodaeiaminjan MNÇ, Gözel H (2016) Genome survey of pistachio (*Pistacia vera* L.) by next generation sequencing: development of novel SSR markers and genetic diversity in *Pistacia* species. *Bmc Genom* 17:998. <https://doi.org/10.1186/s12864-016-3359-x>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.