ISSN: 2322 - 0902 (P) ISSN: 2322 - 0910 (O)



International Journal of Ayurveda and Pharma Research

Research Article

EVALUATION OF EFFICACY OF *JALOUKAVACHARANA* AND *SIRAVYADHANA* IN THE MANAGEMENT OF *VATARAKTA*: A COMPARATIVE STUDY

Rajeshwari^{1*}, Deenaprakash Bharadwaj², Harshavardhana K³

*1PG Scholar, ²Professor Emeritus, ³Professor & HOD, Department of Shalyatantra, K.V.G.A.M.C Sullia Dakshina Kannada, Karnataka, India.

ABSTRACT

Vatarakta is the illness that exhibits different signs and symptoms based upon the *Dosha, Utthana* and *Gambheera Avastha*. Dietary habits and life style modalities play a major role in the causation of *Vatarakta*. Though various remedies are there to treat it, *Raktamokshana* is said to be an effective, and standard technique. Hence it is claimed to be an important affective tool in the management of *Vatarakta*. This study is undertaken to explore and compare the efficacy of *Jaloukavacharana* and *Siravyadhana* in *Vatarakta*.

Randomized comparative clinical trial was adopted in this study. 40 subjects fulfilling diagnostic and inclusion criteria were selected and divided into two groups. The day on which the procedure was conducted was considered as first day of the trail. Follow up was on 8th & 15th day. Total duration of Study was 15 days. In both the group there is a significant improvement of subjective and objective symptoms except on *Vaivarnya*. The collected data is statistically analysed *Jaloukavacharana* is found to be more effective in the features *Daha and Shopha*. *Siravydhana* is more effective in *Shoola* and *Vivarnya*. Both the procedure shows equal effect on *Sparshaasahatva*. The overall results of Group A are 82.11% and Group B is 82.76%. Both the treatments are equally effective in *Vatarakta*. Hence alternative hypothesis H3 is proved. This simple and cost-effective treatments are painless do not require any anaesthesia. Hence it can be easily performed in OPD level on day-care basis.

KEYWORDS: Jaloukavacharana, Raktamokshana Siravydahana, Vatraakta.

INTRODUCTION

Vatarakta is very distressing disease due to its continued relapsing and remittent nature^[1]. The factors said in the classics which are responsible for the *Vatarakta* are *Viruddha Ahara* and *Mithya Vihara*. These leads to the vitiation of *Vayu* and *Rakta*. Aggravated *Vata* paves for vitiation of *Rakta* in turn leading to further aggravation of *Vata*, thus evolves condition *Vatarakta*.

In this context the modalities under Siravvadhana^[2] scrutiny are and Ialouka*vacharana*^[3] which annihilate the *Doshic* vitiation in the aforesaid condition. *Raktamokshana* is more effectual in quick action than administering Snehana and Alepa^[4]. Expulsion or removal of vitiated Rakta (as well as other vitiated Doshas) from the body is known as *Raktamokshan*^[5]. This can be done either through *Siravyadhana* (vene puncture) or with Jaloukavacharana (Leech therapy). Hereby, the study tries to expound the effectiveness of these two procedures on two groups of subjects.

Sushruta Acharya while explaining the *Shastiupakrama* mentioned that *Rakta Visravana* is

very useful in *Vedana Upashamana* and to relive *Achirottita Shopa*^[6]. So, this study is also aimed to understand which among the two procedures are more capable of removing the *Avarana*.

Objectives of the Study

- 1. To study in detail about Vatarakta
- 2. To study the clinical efficacy of *Jaloukavacharana* on *Vatarakta*
- 3. To study the clinical efficacy of *Siravyadhana* on *Vatarakta*. To compare the results of *Jaloukavacharana* as well as *Siravyadhana* and analyse the results statically.

Source of Data

Patients who attended the OPD and IPD of K.V.G. Ayurvedic Medical College and Hospital, Sullia, with signs and symptoms of *Vatarakta* were selected for this study.

Method of Collection of Data

Data regarding history, duration of illness, findings on clinical examination and other relevant information including Lab investigations were recorded in a case proforma specially designed for this study, based on the clinical features of the disease *Vatarakta*.

Inclusion criteria

- ✓ Patients with clinical features of *Vatarakta*
- ✓ Patients with age ranging from 20-60 years

Exclusion criteria

- ✓ Patients of age below 20 years and above 60 years.
- ✓ Patients with other systemic diseases
- ✓ Patients contraindicated for *Siravyadhana* and *Jaloukavacaharana*
- ✓ Patient with the history of bleeding diathesis
- ✓ Pregnant women

Diagnostic criteria

The clinical features of *"Vatarakta"* as mentioned in Classics were taken as the diagnostic criteria.

Selected subjective and objective parameters are graded accordingly.

Pathological investigation

Routine haematological investigations like Hb%, total count, differential count, ESR, bleeding time, clotting time, RBS were done to rule out any other disease pathology as well as to evaluate safety and efficacy of the trial therapy.

Grouping and sampling

Total 40 patients of *Vatarakta* were included in this study. They were randomly divided into two groups.

Study duration

- ✓ Jaloukavacharana- 1 sitting, on the first day only
- ✓ Siravyadhana- 1 sitting, on the first day only
- ✓ Follow up- On 8th day & 15th day
- ✓ Total duration-15 days

Observation and recording of data

- ✓ First Recording-Pre-treatment examination (1st day BT)
- ✓ Second Recording-Soon after the treatment (1st day AT)
- ✓ Follow up observation- On 8th day and 15th day

Assessment Criteria

Post procedural observation of the assessment criteria was done and recorded.

Subjective parameter

- ✓ Shoola
- ✓ Daaha

Objective parameter

- ✓ Sparshaasahatva
- ✓ Shopha
- ✓ Vaivarnya

Table 1: Shoola

| Shoola 🦉 | Grade |
|--|-------|
| Absent - no pain | 0 |
| <i>Avara</i> - pain of bearable in nature, but no difficulty in movement of joints, appears frequently | 1 |
| <i>Madhyama</i> - slight difficulty in joint movements due to severe pain requires medication and pain remain throughout the day | 2 |
| Pravara- Difficult to move the joints due to pain. Affects the sleep also | 3 |

Table 2: Daha

| Daha | Grade |
|--|-------|
| Absent - no burning sensation | 0 |
| Avara- occasionally burning sensation | 1 |
| Madhyama- frequently burning sensation | 2 |
| Pravara - continuous burning sensation | 3 |

Table 3: Sparshaasahatva

| Sparshaasahatva | Grade |
|-------------------------------|-------|
| Absent- no tenderness | 0 |
| Avara- slight tenderness | 1 |
| Madhyama- moderate tenderness | 2 |
| Pravara- severe tenderness | 3 |

Table 4: Shopha

| Shopha | Grade |
|-----------------------------|-------|
| Absent- no swelling | 0 |
| Avara- slight swelling | 1 |
| Madhyama- moderate swelling | 2 |
| Pravara- severe swelling | 3 |

Table 5: Vaivarnya

| Vaivarnya | Grade |
|--------------------------|-------|
| Absent- no discoloration | 0 |
| Avara | 1 |
| Madhyama | 2 |
| Pravara | 3 |

Observation and Results

Table 6: Effect of Jaloukavacharana on Daha in Group A

| Symptom | Mean | score | | | % | S D (+) | S E (+) | tualua | p value |
|---------|------|-------|------|-------|-----|---------|---------|---------|---------|
| | BT | | | BT-AT | 70 | S.D (±) | S.E (±) | t value | p value |
| Daha | 1.05 | AT | 0.45 | 0.60 | 57 | 0.503 | 0.112 | 2.76 | < 0.05 |
| | | FU1 | 0.30 | 0.75 | 71 | 0.444 | 0.008 | 3.76 | < 0.05 |
| | | FU2 | 0.00 | 1.05 | 100 | 0.759 | 0.011 | 6.19 | < 0.05 |

Statistical analysis showed that the mean score which was 1.05 before the treatment was reduced to 0.45 after the treatment and after follow up it became 0.00 with 100% improvement and this is a statistically significant (P<0.05).

| Symptom | Mean | score | | | 0/ | 6 D (L) | | tualua | p value | |
|---------|------|-------|------|-------|---------|---------|---------|---------|---------|--|
| | BT | | | BT-AT | % | S.D (±) | S.E (±) | t value | | |
| Daha | 0.80 | AT | 0.55 | 0.25 | 31 | 0.444 | 0.099 | 0.89 | >0.05 | |
| | | FU1 | 0.45 | 0.35 | 44urved | 0.489 | 0.016 | 1.33 | >0.05 | |
| | | FU2 | 0.10 | 0.70 | 87.5 | 0.923 | 0.021 | 2.98 | < 0.05 | |

Table 7: Effect of Siravyadhana on Daha in Group B

Statistical analysis showed that the mean score which was 0.80 before the treatment was reduced to 0.55 after the treatment and after follow up it became 0.10 with 87.5% improvement and this is statistically significant (P<0.05).

Table 8: Effect of Jaloukavacharana in Group-A on Shoola

| Symptom | Mean | score | | 92, F3, | % | S.D (±) | S.E (±) | t value | p value | |
|---------|------|-------|------|---------|------|---------|---------|---------|---------|--|
| Symptom | BT | | | BT-AT | JAPR | 3.D (±) | 3.E (±) | t value | | |
| | | AT | 1.50 | 0.65 | 30 | 0.489 | 0.109 | 3.44 | < 0.05 | |
| Shoola | 2.15 | FU1 | 1.00 | 1.15 | 53 | 0.489 | 0.016 | 5.88 | <0.05 | |
| | | FU2 | 0.65 | 1.50 | 70 | 0.513 | 0.010 | 8.08 | < 0.05 | |

The mean score from 2.15 to 1.50 after the treatment and after follow up it became 0.65 with 70% improvement. It is found to be statistically significant (P<0.05).

Table 9: Effect of Siravyadhana in Group-B on Shoola

| Symptom | Mean | score | | | % | S D (+) | S E (+) | t value | n value |
|---------|------|-------|------|-------|----|---------|---------|---------|---------|
| | BT | | | BT-AT | | S.D (±) | S.E (±) | tvalue | p value |
| | | AT | 1.45 | 0.80 | 36 | 0.410 | 0.092 | 4.07 | < 0.05 |
| Shoola | 2.25 | FU1 | 1.25 | 1.00 | 44 | 0.324 | 0.009 | 5.75 | < 0.05 |
| | | FU2 | 0.65 | 1.60 | 71 | 0.503 | 0.011 | 8.89 | < 0.05 |

The mean score from 2.25 to 1.45 after the treatment and after follow up it became 0.65 with 71% improvement. It is found to be statistically significant (P<0.05).

Table 10: Effect on Jaloukavacharana in Group-A on Sparshaasahatva

| Symptom | Mean | score | | | 0/ | S D (+) | S E (+) | t | n valuo |
|---------------------|------|-------|------|-------|----|---------|---------|-------|---------|
| | BT | | | BT-AT | % | S.D (±) | S.E (±) | value | p value |
| Sparshaa sahatva | 1.50 | AT | 0.95 | 0.55 | 37 | 0.605 | 0.135 | 2.03 | < 0.05 |
| | | FU1 | 0.50 | 1.00 | 67 | 0.795 | 0.022 | 3.98 | < 0.05 |
| | | FU2 | 0.10 | 1.40 | 93 | 0.940 | 0.014 | 5.98 | < 0.05 |

Int. J. Ayur. Pharma Research, 2020;8(10):47-55

The mean score which was 1.50 before treatment reduced to 0.95 after the treatment and after follow up it became 0.10 with 93% improvement it is statistically significant (P<0.05).

| Symptom | Means | score | | | % | S.D (±) | SE(I) | tualua | p value |
|-----------------|-------|-------|------|-------|----|---------|---------|---------|---------|
| | BT | | | BT-AT | | 3.D (±) | S.E (±) | t value | |
| Sparshaasahatva | 1.50 | AT | 0.70 | 0.80 | 53 | 0.410 | 0.092 | 4.66 | < 0.05 |
| | | FU1 | 0.50 | 1.00 | 67 | 0.324 | 0.006 | 5.63 | < 0.05 |
| | | FU2 | 0.10 | 1.40 | 93 | 0.598 | 0.009 | 9.20 | < 0.05 |

Table 11: Effect of Siravyadhana in Group-B on Sparshaasahatva

The mean score which was 1.50 before treatment reduced to 0.70 after the treatment and after follow up it become 0.10 with 93% improvement study is statistically significant (P<0.05).

| Symptom | | Mea | n score | | 0/ | C D (+) | S.E (±) | t value | р | | |
|---------|------|-----|---------|-------|----|---------|-----------------|---------|--------|--|--|
| | BT | | | BT-AT | % | S.D (±) | 5. с (т) | | value | | |
| Shopha | 1.75 | AT | 1.00 | 0.75 | 43 | 0.444 | 0.099 | 3.00 | < 0.05 | | |
| | | FU1 | 0.75 | 1.00 | 57 | 0.795 | 0.019 | 4.02 | < 0.05 | | |
| | | FU2 | 0.05 | 1.70 | 97 | 0.923 | 0.016 | 8.11 | < 0.05 | | |

Table 12: Effect of Jaloukavacharana in Group-A on Shopha

The mean score which was 1.75 before treatment reduced to 1.00 after the treatment and after follow up it became 0.05 with 97% improvement study is statistically significant (P<0.05).

| Table 15: Effect of Shavyaanana in Group-B on Shopha | | | | | | | | | | | |
|--|------|-------|------|-------|----|---------|---------|---------|---------|--|--|
| Commentant | Mean | score | | | % | S.D (±) | S.E (±) | t value | p value | | |
| Symptom | BT | | | BT-AT | | | | | | | |
| | | AT | 0.95 | 0.60 | 39 | 0.503 | 0.112 | 2.14 | < 0.05 | | |
| Shopha | 1.55 | FU1 | 0.60 | 0.95 | 61 | 0.510 | 0.013 | 3.97 | < 0.05 | | |

FU2

0.15

1.40

Table 13: Effect of Siravyadhana in Group-B on Shopha

The mean score which was 1.55 before treatment reduced to 0.95 after the treatment and after follow up it become 0.15 with 90% improvement, study is statistically significant (P<0.05).

Table 14: Effect of *Jaloukavacharana* over *Vaivarnya* in Group-A

90

0.754

0.012

6.93

< 0.05

| Sumatom | Mean | score | | 241 II | % | S.D (±) | S.E (±) | t value | p value |
|-----------|------|-------|------|--------|----|---------|---------|---------|---------|
| Symptom | BT | | | BT-AT | | | | | |
| Vaivarnya | 0.95 | AT | 0.85 | 0.10 | 11 | 0.308 | 0.069 | 0.33 | >0.05 |
| | | FU1 | 0.70 | 0.25 | 26 | 0.444 | 0.042 | 0.85 | >0.05 |
| | | FU2 | 0.60 | 0.35 | 37 | 0.489 | 0.019 | 1.21 | >0.05 |

The mean score which was 0.95 before treatment reduced to 0.85 after the treatment and after follow up it become 0.60 (37%) with non-significant improvement (P>0.05).

| Symptom | Mean | score | | | % | S.D (±) | S.E (±) | t value | p value | |
|-----------|------|-------|------|-------|----|---------|---------|---------|---------|--|
| | BT | | | BT-AT | | | | | | |
| Vaivarnya | 0.50 | AT | 0.45 | 0.05 | 10 | 0.224 | 0.050 | 0.20 | >0.05 | |
| | | FU1 | 0.40 | 0.10 | 20 | 0.308 | 0.031 | 0.42 | >0.05 | |
| | | FU2 | 0.30 | 0.20 | 40 | 0.410 | 0.021 | 0.89 | >0.05 | |

Table 15: Effect of Siravyadhana over Vaivarnya in Group-B

Study showed statistically Non significant improvement (P>0.05). The mean score which was 0.50 before treatment reduced to 0.45 after the treatment and after follow up it became 0.30 with 40% improvement.




Fig No:1 Procedure of Jaloukavacharana 1- Jalouka in fresh water 2- Jalouka in Haridra Jala 3- Covered the body of Jalouka with Pichu 4horse-shoe shape neck of Jalouka showing proper sucking 5,6- Chardana of Jalouka





Fig.No:2 Procedure of Siravyadhana 1- Sthanika Snehana 2- Sthanika Patta Sweda 3- Puncturing the vein 4- Flowing of blood 3

Assessment of Total Effect of Therapy Result on Group A



Result on Group B




Table 16: Comparative results of Group-A and Group-B

| Characteristics | Group-A | | | Group-B | | | |
|-----------------|---------|-------|------------|---------|------------|-----------|--|
| Signs and | Mean | score | Percentage | Mean | Percentage | | |
| Symptoms | ВТ | FU | of relief | ВТ | FU | of relief | |
| Daha | 1.05 | 0.00 | 100 | 0.80 | 0.10 | 87.5 | |
| Shoola | 2.15 | 0.65 | 70 | 2.25 | 0.65 | 71 | |
| Sparshaasahatva | 1.50 | 0.10 | 93 | 1.50 | 0.10 | 93 | |
| Shopha | 1.75 | 0.05 | 97 | 1.55 | 0.15 | 90 | |
| Vaivarnya | 0.95 | 0.60 | 37 | 0.50 | 0.30 | 40 | |

DISCUSSION

Effect of Jaloukavachara on Daha [7]

Vitiated *Rakta* and *Pitta* causes severe burning sensation, *Jalouka* is claim to be *Seetha* and *Madhura* there by *Jalouka* decreases the *Ushna*, *Teekshna Guna* and cures the *Daha*. The present study with the statistical analysis shows 100% eradication of *Daha* by *Jaloukavacharana*. Process is highly significant.

Effect of Siravyadhana on Daha

In the present study 87.5% improvement was seen by *Siravyadhana*. This procedure removes the *Dooshita Rakta* and decreases the *Rakta Dusti* hence this statistically significant is the result.

Effect of Jaloukavacharana and Siravyadhana on Shoola

Shoola is purely due to vitiated *Vata*. due to the *Vishamagni Agnimandhya* occurs and it develops the *Aama*. The *Aama* eases the blockage of *Srotas*

which in turn aggravates the *Vata* causes the *Shoola*. So, the vitiated *Vata* and *Aama* blocks the *Rakta*, by *Anyonya avarana* and the disease originates.^[8]

By doing *Raktamokshana Avarana* is removed there by it decreases the pain. So, in this study both *Jaloukavacharna* and *Siravyadhana* have shown almost equal result.

The results in group A on *Shoola* says that mean score is reduced from 2.15 to 0.65 i.e. 70% improvement. The P value is less than 0.05 and treatment is statistically significant.

In *Siravydhana* group result is 71% there the mean score is reduced from 2.25 to 0.65. P value is less than 0.05 and the treatment is statistically significant.

Effect of Jaloukavacharana and Siravyadhana on Sparshasahatva

Sparsaasahatva is mainly the symptom of *Shonitavrta* condition^[9], that too when *Vata* is predominant. When *Avarana* is removed tenderness reduces. Statistically both treatments don't show any difference. Mean score of both the treatment was 1.50 and reduced to 0.10 and the treatment shows significant result i.e., 93% improvement.

Effect on Shopha

Shopha is a Tridoshaja condition when aggravated Vata affects the Kapha, Rakta and Pitta obstruction in the Srotas takes place.

Jaloukavacharana

As it is a localised condition in a *Vata Rakta*, when *Jalouka* is applied it does *Shamana* of vitiated *Dosha* and removes *Dooshita Rakta*. In the present study mean score before the treatment was 1.75 and it reduced to 0.05 (P value is less than 0.05) i.e., 97% improvement was seen and treatment is statistically significant.

Siravyadhana

Visravana is helpful in removing *Achirothitta Shopha*. By *Visravana Avarodha* of *Srotas* is removed there by decrease in the *Shotha*.^[10] the present study revealed that mean score before the treatment was 1.55 and after the treatment it was 0.15, and P value less than 0.05 with 90% improvement and it is statistically significant.

Effect of Jaloukavacharana and Siravyadhana on Vaivarnya

Prakupitha Vata and vitiated Rakta get Sangha/ Avarodha in the Srotas. Here lakshanas of Vruddhi of Vata as well as Rakta favoring Vaivarnya of skin. Along with this the Pitta, Rakta in Ashrayaashrayi Bhava and Bhrajaka Pitta in the Twak also responsible for altered Varna.^[11] Here in Vatarakta Rakta Dhatu does Avarana to Pitta and Vata which causes Vaivarnya. By Raktamokshana these Dooshita *Rakta* and *Avarana* is removed there by *Vaivarnya* is decreased.

This study statistically shows that there is no significant improvement in *Vaivarnya* in *Vatarakta* by both procedures. *Jaloukavacharanaa* shows 37% improvement and *Siravyadhana* shows 40% improvement.

Overall effect of Jaloukavacharanaa on Vatarakta

Out of 20 subjects in group A 14 subjects show good improvement 1 shows moderate improvement and 5 subject show complete relief from the symptoms.

Overall effect of Siravyadhana on Vatarakta

Out of 20 subjects 11 show good improvement 1 moderate improvement and 8 got complete relief from the symptoms.

Comparative result of group A and group B Daha

In the present study statistical analysis of subjective and objective parameter shows that *Jaloukavacharana* on the *Daha* shows good results (100%) and *Siravyadhana* shows 87.5 % results. *Jaloukavacharana* shows good results than *Siravyadhana* may be because *Jalouka* is *Madhura* and *Sita* which has good effect on *Daha* of *Pitta* and *Rakta*.

Shoola

Statistical analysis of result of Jaloukavacharana 70% on Shoola shows improvement and Siravyadhana shows 71% improvement i.e., almost equal results. It may be because, by doing both procedure Avarana of Vata and *Rakta* gets removed.

Sparshasahatva

Statistical analysis shows 93% improvement in both the groups. Both procedures are equally effective in decreasing *Avarana* of *Vata* and *Rakta* there by it reduces *Sparshashatva*.

Shopha

Statistical analysis of Jaloukavacharana shows 97% result and Siravyadhana shows 90% improvement in Shopha. Jaloukavacharana shows more improvement than Siravaydhana because Visravana is good in subsiding the Achirothita Shotha, Shows good effect. It is because in the local action Jalouka are more powerful than Siravyadhana.

Vaivarnya

Statistical analysis of treatment by *Jaloukavacharana* shows 37% improvement and by *Siravyadhana* 40% improvement. But both treatments are statistically not significant. It may be because one sitting of procedures is not sufficient to remove the *Vaivarnya*.

Comparative results of overall effect in Group A and Group B

Comparative analysis of the overall effect of the treatment in the both the group was done statistically with unpaired t test. It shows the treatments are statistically insignificant. Both the treatments are equally effective in *Vatarakta*.

Raktavisravana is the standard mode of treatment in *Vatarakta*. *Rakta* is the main *Dooshya* in this disease so by removing *Dooshitha Rakta* it gives good results.

Probable mode of action of Jaloukavacharanaa

Jalouka is one among the Anushastra and it is Sheetala and Madhura hence they may be used to decrease the Pitta.^[12]

Vatarakta is a disease of Prakupitha Vata and Dooshitha Rakta. According to the texts the main treatment modality of Vatarakta is Raktamokshana. Rakta is having similar Gunas of Pitta. By doing Jaloukavacharanaa it will remove the Dooshitha Rakta and Pitta Shamana takes place. Prakupitha Vata also gets reduced by this treatment. The Pitta Shamaka effect of Jaloukavacharana results in the reduction of Daha, Shopha and Sparsaasahatva.

Probable mode of Action of Siravyadhana

Siravyadhana is indicated in Sarvadehanusari Dooshita Rakta. And Raktamokshana can treat almost all disorders resulted due to the derangement in Mamsa, Meda, Asthi and Majja. It purifies the Dhamani and corrects the vitiated blood. It also purifies Rasa and Swedavaha srotas.

As it is a systemic disease and thereby *Raktamokshana* reduces the symptoms of *Vatarakta*.. **CONCLUSION**

Based on the literature, observation and results of this clinical Study following conclusions are drawn.

Present lifestyle and dietary habits are the main contributing factors in the establishment of *Vatarakta.*

Most of the subjects were in Uttana Avasta.

Jaloukavacharana and *Siravyadhana* are equally effective in the case of *Vatarakta*. Thus, alternative hypothesis is proved in this study.

While considering the symptom *Daha* and *Shopha*, *Jaloukavacharana* is found to be more effective. Likewise, in *Shoola* and *Vivarnya Siravydhana* is more effective. In *Sparshaasahatva* both the procedure is equally effective.

For both procedures' materials required are easily available. They are simple and cost effective. Because they are painless do not require any anaesthesia. Hence can be easily performed in OPD level on daycare basis.

Repeated bloodletting periodically by *Jaloukavacharana* and *Siravyadhana* definitely gives good results.

All the symptoms except *Vaivarnya* responded to the treatment very well. *Vaivarnya* remained further May be because *Raktamoshana* is done only once which is not sufficient to reverse the pathology completely in such a short span.

REFERENCES

- 1. Agnivesha. Charaka Samhitha, with Ayurveda Dipika commentary, edited by Vaidya Yadavji Trikamji Acharya, Chikitsasthana -29 verse no.18, Varanasi, Choukhamba Surabharathi Prakashan, reprint-2011, p.628.
- 2. Acharya Sushruta. Sushruta Samhita with Nibandha Sangraha Commentary by Dalhana, edited by Vaidya Jadavji Trikamji, Sharirasthan-8, verse no.1-25, Varanasi, Chaukhamba Krishnadas Academy, reprint 2014, p. 379-383.
- Acharya Sushruta. Sushruta Samhita with Nibandha Sangraha Commentary by Dalhana, edited by Vaidya Jadavji Trikamji, Sutrasthan-13, verse no.19, Varanasi, Chaukhamba Krishnadas Academy, reprint 2014, p. 57.
- 4. Acharya Sushruta.Sushruta Samhita with Nibandha Sangraha Commentary by Dalhana, edited by Vaidya Jadavji Trikamji, Sharirasthan-8, verse no.22, Varanasi, Chaukhamba Krishnadas Academy, reprint 2014, p. 383.
- 5. Acharya Sushruta. Sushruta Samhita with Nibandha Sangraha Commentary by Dalhana, edited by Vaidya Jadavji Trikamji, Sutrasthan-14, verse no.34, Varanasi, Chaukhamba Krishnadas Academy, reprint 2014, p. 65.
- 6. Acharya Sushruta. Sushruta Samhita with Nibandha Sangraha Commentary by Dalhana, edited by Vaidya Jadavji Trikamji, Chiktsasthan-1, verse no.27-28, Varanasi, Chaukhamba Krishnadas Academy, reprint 2014, p. 399
- 7. Agnivesha. Charaka Samhitha, with Ayurveda Dipika commentary, edited by Vaidya Yadavji Trikamji Acharya, Sutrasthana -12 verse no.11, Varanasi, Choukhamba Surabharathi Prakashan, reprint-2011,p.80.
- 8. Acharya Sushruta. Sushruta Samhita with Nibandha Sangraha Commentary by Dalhana, edited by Vaidya Jadavji Trikamji, Chiktsasthan-1, verse no.27, Varanasi, Chaukhamba Krishnadas Academy, reprint 2014, p. 399.
- 9. Acharya Sushruta. Sushruta Samhita with Nibandha Sangraha Commentary by Dalhana, edited by Vaidya Jadavji Trikamji, Sutrasthan-13,

verse no.6, Varanasi, Chaukhamba Krishnadas Academy, reprint 2014, p. 55.

- 10. K.R.Sreekanthamurthy. Madhava Nidana, Chapter-26, verse no. 2-5, Varanasi, Chaukhambha Subharathi Prakashan, Reprint 1995, p.3.
- 11. Acharya Sushruta. Sushruta Samhita with Nibandha Sangraha Commentary by Dalhana,

Cite this article as:

Rajeshwari, Deenaprakash Bharadwaj, Harshavradhana K Evaluation of Efficacy of Jaloukavacharana And Siravyadhana In The Management of Vatarakta; A Comparative Study. International Journal of Ayurveda and Pharma Research. 2020;8(10):47-55.

Source of support: Nil, Conflict of interest: None Declared

edited by Vaidya Jadavji Trikamji, Nidanasthan-1, verse no.33, Varanasi, Chaukhamba Krishnadas Academy, reprint 2014, p. 263.

12. Acharya Sushruta. Sushruta Samhita with Nibandha Sangraha Commentry by Dalhana, edited by Vaidya Jadavji Trikamji, Sutrasthan-13, verse no.6, Varanasi, Chaukhamba Krishnadas Academy, reprint 2014, p. 55.

> *Address for correspondence Dr Rajeshwari PG Scholar, Department of Shalyatantra, K.V.G.A.M.C Sullia Dakshina Kannada, Karnataka, India. Email: aryapavithran92@gmail.com

Disclaimer: IJAPR is solely owned by Mahadev Publications - dedicated to publish quality research, while every effort has been taken to verify the accuracy of the content published in our Journal. IJAPR cannot accept any responsibility or liability for the articles content which are published. The views expressed in articles by our contributing authors are not necessarily those of IJAPR editor or editorial board members.

