

Ateneo de Manila University

Archivum Ateneo

Mathematics Faculty Publications

Mathematics Department

2015

Cauchy Extension of M_α -Integral and Absolute M_α -Integrability

Ian June L. Garces

Abraham P. Racca

Follow this and additional works at: <https://archivum.ateneo.edu/mathematics-faculty-pubs>

 Part of the [Mathematics Commons](#)

International Journal of Mathematical Analysis
Vol. 9, 2015, no. 27, 1323 - 1329
HIKARI Ltd, www.m-hikari.com
<http://dx.doi.org/10.12988/ijma.2015.5368>

Cauchy Extension of M_α -Integral and Absolute M_α -Integrability

I. J. L. Garces

Department of Mathematics
Ateneo de Manila University
1108 Loyola Heights, Quezon City
The Philippines

Abraham P. Racca

Mathematics and Computer Science Department
Adventist University of the Philippines
4118 Silang, Cavite, The Philippines

Copyright © 2015 I. J. L. Garces and Abraham P. Racca. This article is distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Park, Ryu, and Lee recently introduced a Henstock-type integral, which lies between the Mcshane and the Henstock integrals. This paper proves the closure property of this new integral under Cauchy extension, and presents a characterization on absolute M_α -integrability.

Mathematics Subject Classification: 26A39

Keywords: M_α -integral, Cauchy extension, absolute M_α -integrable

1 Introduction

Park, Ryu, and Lee [4] recently defined a Henstock-type integral, which they call M_α -integral. Several properties and convergence theorems of the integral were established in [1] and [4]. Most of them paralleled the usual properties

of Henstock integral, including the Saks-Henstock Lemma [4, Lemma 2.5]. Moreover, by providing examples, it was also shown in [4] that M_α -integral lies strictly between the McShane integral (which is known to be equivalent to the Lebesgue integral (see [2])) and the Henstock integral.

Let $\alpha > 0$ be a constant, and $I = [a, b]$ a non-degenerate closed and bounded interval in \mathbb{R} .

- (1) A *partial division* D of I is a finite collection of interval-point pairs $([u, v], \xi)$ such that the closed intervals $[u, v]$ are non-overlapping, $\bigcup [u, v] \subseteq I$, and $\xi \in I$. If $\bigcup [u, v] = I$, we call the division D simply a *division*.
- (2) A positive function defined on I is called a *gauge* on I .
- (3) Let δ be a gauge on I , and $D = \{([u, v], \xi)\}$ a partial division of I . If $[u, v] \subseteq (\xi - \delta(\xi), \xi + \delta(\xi))$ for all $([u, v], \xi) \in D$, then we say that D is a δ -*fine McShane partial division*. Moreover, if D is a McShane partial division such that $\xi \in [u, v]$ for all $([u, v], \xi) \in D$, then D is called a δ -*fine Henstock partial division*.
- (4) A McShane division $D = \{([u, v], \xi)\}$ of I is said to be an M_α -*division* if

$$\sum_{([u,v],\xi) \in D} \text{dist}(\xi, [u, v]) < \alpha,$$

where $\text{dist}(x, J) = \inf\{|y - x| : y \in J\}$.

- (5) Let $D = \{([u, v], \xi)\}$ be a partial division on I , and f a real-valued function defined on I . We write

$$S(f, D) = \sum_{([u,v],\xi) \in D} f(\xi)(v - u).$$

With these terms and notations, the definition of M_α -integrability can now be presented.

Definition 1.1 ([4, Definition 2.1]). A function $f : [a, b] \rightarrow \mathbb{R}$ is M_α -*integrable* if there exists a real number A such that, for each $\epsilon > 0$, there is a gauge δ on $[a, b]$ such that

$$|S(f, D) - A| < \epsilon$$

for each δ -fine M_α -division D of $[a, b]$. Here, A is called the M_α -*integral* of f on $[a, b]$, and we write $A = \int_a^b f$.

Moreover, f is M_α -*integrable on* $E \subseteq [a, b]$ if $f\chi_E$ is M_α -integrable on $[a, b]$, where χ_E is the characteristic function over E , and we write $\int_a^b f\chi_E = \int_E f$.

This paper proves the closure property of this new integral under Cauchy extension (Theorem 2.1) and a characterization on absolute M_α -integrability (Theorem 3.3).

2 Cauchy Extension

The following theorem shows that M_α -integral is closed under Cauchy extension, a property that is not valid for McShane integral.

Theorem 2.1 (Cauchy Extension). *Let $f : [a, b] \rightarrow \mathbb{R}$ be M_α -integrable on $[c, b]$ for each $c \in (a, b)$. Then f is M_α -integrable on $[a, b]$ if and only if $\lim_{c \rightarrow a^+} \int_c^b f$ exists. In this case, we have*

$$\int_a^b f = \lim_{c \rightarrow a^+} \int_c^b f.$$

Proof. Suppose that f is M_α -integrable on $[a, b]$, and let $\epsilon > 0$ be given. Then there exists a gauge δ on $[a, b]$ such that if D is a δ -fine M_α -division of $[a, b]$, then

$$\left| S(f, D) - \int_a^b f \right| < \epsilon.$$

Moreover, for each $c \in (a, b)$, there exists a gauge δ_c on $[c, b]$ such that if D_c is a δ_c -fine M_α -division of $[c, b]$, then

$$\left| S(f, D_c) - \int_c^b f \right| < \epsilon.$$

For each $c \in (a, b)$, we may assume that $\delta_c(x) \leq \delta(x)$ for all $x \in [c, b]$.

Choose $s \in (a, a + \delta(a))$ such that $|f(a)|(s - a) < \epsilon$. Let D_s be a δ_s -fine M_α -division of $[s, b]$. Then $D = D_s \cup \{([a, s], a)\}$ is a δ -fine M_α -division of $[a, b]$. Hence, we have

$$\left| \int_a^b f - \int_s^b f \right| \leq \left| \int_a^b f - S(f, D) \right| + \left| S(f, D_s) - \int_s^b f \right| + |f(a)|(s - a) < 3\epsilon.$$

Since s is chosen arbitrarily close to a , we obtain $\lim_{c \rightarrow a^+} \int_c^b f$ exists and is equal to $\int_a^b f$.

Conversely, suppose that $\lim_{c \rightarrow a^+} \int_c^b f = A$ exists, and let $\{x_0, x_1, x_2, \dots\}$ be a strictly decreasing sequence of real numbers such that $x_0 = b$ and $x_n \rightarrow a$. Then f is M_α -integrable on $[x_n, x_{n-1}]$ for every $n \in \mathbb{Z}^+$.

Let $\epsilon > 0$ be given. For each integer $n \geq 1$, choose a gauge δ_n on $[x_n, x_{n-1}]$ such that if D_n is a δ_n -fine M_α -division of $[x_n, x_{n-1}]$, then

$$\left| S(f, D_n) - \int_{x_n}^{x_{n-1}} f \right| < \frac{\epsilon}{2^n}.$$

Furthermore, choose an integer $N > 0$ such that

$$\left| \int_t^b f - A \right| < \epsilon \quad \text{and} \quad |f(a)|(t - a) < \epsilon$$

for every $t \in (a, x_N]$.

Define a gauge δ on $[a, b]$ as follows:

$$\delta(x) = \begin{cases} x_N - x & \text{if } x = a \\ \min\{\delta_1(x), x - x_1\} & \text{if } x \in (x_1, x_0] \\ \min\{\delta_n(x), x - x_n, x_{n-1} - x\} & \text{if } x \in (x_n, x_{n-1}), n \geq 2 \\ \min\{\delta_n(x), \delta_{n+1}(x), x_{n-1} - x, x - x_{n+1}\} & \text{if } x = x_n, n \geq 1. \end{cases}$$

Let D be a δ -fine M_α -division of $[a, b]$. Then

$$D = \{([a, t], a)\} \cup D_t \cup \bigcup_{i=1}^m D_i,$$

where $t \in (x_{m+1}, x_m]$ for some integer $m \geq N$, D_t a δ_{m+1} -fine M_α -division of $[t, x_m] \subset (x_{m+1}, x_m]$, and D_i a δ_i -fine M_α -division of $[x_i, x_{i-1}]$. Thus, by Saks-Henstock Lemma, we have

$$\begin{aligned} |S(f, D) - A| &\leq |f(a)|(t - a) + \left| S(f, D_t) - \int_t^{x_m} f \right| \\ &\quad + \sum_{i=1}^m \left| S(f, D_i) - \int_{x_i}^{x_{i-1}} f \right| + \left| \int_t^b f - A \right| \\ &< \epsilon + \frac{\epsilon}{2^{m+1}} + \sum_{i=1}^{\infty} \frac{\epsilon}{2^i} + \epsilon \\ &< 4\epsilon. \end{aligned}$$

This ends the proof of the theorem. \square

Corollary 2.2. *Let $f : [a, b] \rightarrow \mathbb{R}$ be M_α -integrable on every $[c, d] \subset (a, b)$. Then f is M_α -integrable on $[a, b]$ if and only if $\lim_{\substack{c \rightarrow a^+ \\ d \rightarrow b^-}} \int_c^d f$ exists. In this case, we have*

$$\int_a^b f = \lim_{\substack{c \rightarrow a^+ \\ d \rightarrow b^-}} \int_c^d f.$$

3 Absolute M_α -Integrability

Recall that an integral theory is *absolute* if a function f is integrable (in the sense of that theory) if and only if $|f|$ is also integrable. Both Riemann and McShane integrals are absolute (see [2] or [5]), but Henstock integral is known to be not (see [3]).

Example 3.1. Let $\sum_{n=1}^{\infty} a_n$ be a convergent series. Define a function $f : [0, 1] \rightarrow \mathbb{R}$ as follows:

$$f(x) = \begin{cases} 2^n a_n & \text{if } x \in \left(\frac{1}{2^n}, \frac{1}{2^{n-1}}\right], n \in \mathbb{Z}^+ \\ 0 & \text{if } x = 0. \end{cases}$$

Clearly, f is M_α -integrable on $[c, 1]$ for each $c \in (0, 1)$. Thus, by Theorem 2.1, we have

$$\int_0^1 f = \lim_{c \rightarrow 0^+} \int_c^1 f = \sum_{n=1}^{\infty} a_n.$$

However, it is not difficult to see that $|f|$ would be M_α -integrable if and only if the given series is absolutely convergent.

In general, due to the closure under Cauchy extension, it is possible for a function f to be M_α -integrable on $[a, b]$, but $|f|$ is not. We have the following proposition.

Proposition 3.2. *The M_α -integral is not absolute.*

Let a function $f : [a, b] \rightarrow \mathbb{R}$ be M_α -integrable on $[a, b]$. Then the function $F : [a, b] \rightarrow \mathbb{R}$, defined as $F(a) = 0$ and $F(x) = \int_a^x f$ for $x \in (a, b]$, is called the *primitive* of f on $[a, b]$.

Recall that a function $F : [a, b] \rightarrow \mathbb{R}$ is said to be of *bounded variation* on $[a, b]$ if

$$V(F, [a, b]) = \sup \left\{ \sum_P |F(u, v)| \right\} < \infty,$$

where the supremum runs over all partitions P of $[a, b]$ (that is, a finite collection of non-overlapping intervals $[u, v]$ such that $\bigcup [u, v] = [a, b]$), and $F(u, v) = F(v) - F(u)$.

The following theorem answers the question on exactly when $|f|$ is M_α -integrable on $[a, b]$, given that f is M_α -integrable on $[a, b]$.

Theorem 3.3. *Let a function $f : [a, b] \rightarrow \mathbb{R}$ be M_α -integrable on $[a, b]$ with primitive F . Then $|f|$ is M_α -integrable on $[a, b]$ if and only if F is of bounded variation over $[a, b]$. In this case, we have*

$$\int_a^b |f| = V(F, [a, b]).$$

Proof. Suppose that $|f|$ is M_α -integrable on $[a, b]$. Let $P = \{[u, v]\}$ be a partition of $[a, b]$. Then

$$\sum_P |F(u, v)| = \sum_P \left| \int_u^v f \right| \leq \sum_P \int_u^v |f| = \int_a^b |f|,$$

which implies $V(F, [a, b]) < \infty$. Thus, F is of bounded variation over $[a, b]$.

Conversely, suppose that F is of bounded variation over $[a, b]$. Let $\epsilon > 0$ be given. Then there exists a partition $P = \{[x_{i-1}, x_i] : i = 1, 2, \dots, n\}$ of $[a, b]$ such that

$$V(F, [a, b]) - \frac{\epsilon}{2} < \sum_{i=1}^n |F(x_{i-1}, x_i)| \leq V(F, [a, b]).$$

Moreover, by the M_α -integrability of f , there exists a gauge δ_0 on $[a, b]$ such that

$$|S(f, D) - F(a, b)| < \frac{\epsilon}{4}$$

for any δ_0 -fine M_α -division D of $[a, b]$.

Define

$$\delta(x) = \begin{cases} \min\{\delta_0(x), x - x_{i-1}, x_i - x\} & \text{if } x \in (x_{i-1}, x_i) \text{ for some } i \\ \min\{\delta_0(x), x - x_{i-1}, x_{i+1} - x\} & \text{if } x = x_i, 1 \leq i \leq n - 1 \\ \min\{\delta_0(x), x_1 - x\} & \text{if } x = x_0 = a \\ \min\{\delta_0(x), x - x_{n-1}\} & \text{if } x = x_n = b, \end{cases}$$

and let $D = \{([u_k, v_k], \xi_k) : k = 1, 2, \dots, m\}$ be a δ -fine M_α -division of $[a, b]$. By the definition of δ , D has $x_0 = a, x_1, x_2, \dots, x_n = b$ as among its tags. Further, if $([u, v], \xi) \in D$, where $\xi \in \{x_0, x_1, \dots, x_n\}$, then $\xi \in [u, v]$, so that these tags can be made as endpoints and

$$V(F, [a, b]) - \frac{\epsilon}{2} < \sum_{i=1}^n |F(x_{i-1}, x_i)| \leq \sum_{k=1}^m |F(u_k, v_k)| \leq V(F, [a, b]).$$

Thus, applying Saks-Henstock Lemma and triangle inequality, we have

$$\begin{aligned} |S(|f|, D) - V(F, [a, b])| &\leq \left| \sum_{k=1}^m (|f(\xi_k)|(v_k - u_k) - |F(u_k, v_k)|) \right| \\ &\quad + \left| \sum_{k=1}^m |F(u_k, v_k)| - V(F, [a, b]) \right| \\ &< \sum_{k=1}^m |f(\xi)(v_k - u_k) - F(u_k, v_k)| + \frac{\epsilon}{2} \\ &< \epsilon. \end{aligned}$$

This ends the proof of the theorem. □

Corollary 3.4. *Let the functions $f, g : [a, b] \rightarrow \mathbb{R}$ be M_α -integrable on $[a, b]$ with $|f(x)| \leq g(x)$ for all $x \in [a, b]$. Then $|f|$ is M_α -integrable on $[a, b]$, and*

$$\left| \int_a^b f \right| \leq \int_a^b |f| \leq \int_a^b g.$$

References

- [1] I. J. L. Garces and A. P. Racca, Characterizing Convergence Conditions for the M_α -Integral, *J. Chungcheong Math. Soc.* 24 (3) (2011) 469-480.
- [2] R. A. Gordon, *The Integrals of Lebesgue, Denjoy, Perron, and Henstock*, American Mathematical Society, 1994. <http://dx.doi.org/10.1090/gsm/004>
- [3] P. Y. Lee, *Lanzhou Lectures on Henstock Integration*, World Scientific, 1989. <http://dx.doi.org/10.1142/0845>
- [4] J. M. Park, H. W. Ryu and H. K. Lee, The M_α -Integral, *J. Chungcheong Math. Soc.* 23 (1) (2010) 99-108.
- [5] H. L. Royden and P. M. Fitzpatrick, *Real Analysis*, Pearson, 2010.

Received: March 19, 2015; Published: April 30, 2015