

Ateneo de Manila University

**Archium Ateneo**

---

Mathematics Faculty Publications

Mathematics Department

---

2020

## The sigma chromatic number of the Sierpinski gasket graphs and the Hanoi graphs

Agnes Garciano

Reginaldo M. Marcelo

Mari-Jo P. Ruiz

Mark Anthony C. Tolentino

Follow this and additional works at: <https://archium.ateneo.edu/mathematics-faculty-pubs>

 Part of the [Mathematics Commons](#)


---

PAPER • OPEN ACCESS

## The sigma chromatic number of the Sierpiński gasket graphs and the Hanoi graphs

To cite this article: A D Garciano *et al* 2020 *J. Phys.: Conf. Ser.* **1538** 012002

View the [article online](#) for updates and enhancements.



**IOP | ebooks™**

Bringing together innovative digital publishing with leading authors from the global scientific community.

Start exploring the collection—download the first chapter of every title for free.

# The sigma chromatic number of the Sierpiński gasket graphs and the Hanoi graphs

A D Garciano, R M Marcelo, M J P Ruiz and M A C Tolentino

Department of Mathematics, School of Science and Engineering, Loyola Schools,  
Ateneo de Manila University, Philippines

E-mail: [agarciano@ateneo.edu](mailto:agarciano@ateneo.edu), [rmarcelo@ateneo.edu](mailto:rmarcelo@ateneo.edu), [mruiz@ateneo.edu](mailto:mruiz@ateneo.edu),  
[mtolentino@ateneo.edu](mailto:mtolentino@ateneo.edu)

**Abstract.** A vertex coloring  $c : V(G) \rightarrow \mathbb{N}$  of a non-trivial connected graph  $G$  is called a *sigma coloring* if  $\sigma(u) \neq \sigma(v)$  for any pair of adjacent vertices  $u$  and  $v$ . Here,  $\sigma(x)$  denotes the sum of the colors assigned to vertices adjacent to  $x$ . The *sigma chromatic number* of  $G$ , denoted by  $\sigma(G)$ , is defined as the fewest number of colors needed to construct a sigma coloring of  $G$ . In this paper, we determine the sigma chromatic numbers of the Sierpiński gasket graphs and the Hanoi graphs. Moreover, we prove the uniqueness of the sigma coloring for Sierpiński gasket graphs.

## 1. Introduction

In [4], Chartrand, Okamoto, and Zhang introduced a new kind of vertex coloring called a sigma coloring. It is defined as follows.

**Definition 1.1** (Chartrand, Okamoto, Zhang [4]). *For a non-trivial connected graph  $G$ , let  $c : V(G) \rightarrow \mathbb{N}$  be a vertex coloring of  $G$ . For each  $v \in V(G)$ , the **color sum** of  $v$ , denoted by  $\sigma(v)$ , is defined to be the sum of the colors of the vertices adjacent to  $v$ . If  $\sigma(u) \neq \sigma(v)$  for every two adjacent  $u, v \in V(G)$ , then  $c$  is called a **sigma coloring** of  $G$ . The minimum number of colors required in a sigma coloring of  $G$  is called its **sigma chromatic number** and is denoted by  $\sigma(G)$ .*

Sigma coloring is an example of a neighbor-distinguishing coloring, the most studied example of which is the proper vertex coloring. Over the years, various neighbor-distinguishing colorings have also been introduced and discussed in literature such as in [3] and [5]. The notion of sigma coloring is closely related to the vertex colorings/labellings, discussed in [1], [9], [12] that also use the sum of the colors/labels of a vertex's neighbors.

Sigma colorings of different families of graphs have already been studied. For instance, Chartrand et al. determined the sigma chromatic numbers of paths, cycles, bipartite, and complete multipartite graphs in [4]. More recently, in [10], Luzon, Ruiz, and Tolentino have determined the sigma chromatic numbers of some families of circulant graphs. The complexity of the sigma coloring problem has also been studied in [6].

In this paper, we determine the sigma chromatic numbers of the Sierpiński gasket graphs and the Hanoi graphs.

The **Sierpiński gasket graph**  $S_n$ ,  $n \geq 1$ , is the graph whose vertices are the intersection points of the finite Sierpiński gasket and whose edges are the line segments of the gasket [13].



Content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](https://creativecommons.org/licenses/by/3.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

An alternative description of  $S_n$  can also be found in [8]. The construction of  $S_n$  is shown in Fig. 1 while a sigma coloring of  $S_2$  is shown in Fig. 2.




Figure 1. Sierpiński gasket graphs.




Figure 2. A sigma coloring of  $S_2$ .

On the other hand, the **Hanoi graphs** ([2], [7], [8]) are derived from the states of the Tower of Hanoi problem. Like the Sierpiński gasket graphs, the Hanoi graphs are also constructed in an iterative manner as shown in Fig. 3. A sigma 2-coloring of  $H_2$  is shown in Fig. 4.




Figure 3. Hanoi graphs.

The (ordinary) chromatic numbers of  $S_n$  and  $H_n$  have already been determined by other researchers. For the Hanoi graphs, Parisse has constructed a natural coloring that leads to the following result.

**Theorem 1.2** (Parisse, [11]). *For all  $n \geq 1$ , the chromatic number of  $H_n$  is 3.*

For the Sierpiński gasket graphs, Tegui and Godbole proved the following:




Figure 4. A sigma 2-coloring of  $H_2$ .

**Theorem 1.3** (Teguaia, Godbole [13]). *For all  $n \geq 1$ , the chromatic number of  $S_n$  is 3.*

Moreover, Klavžar proved the following stronger result.

**Theorem 1.4** (Klavžar [8]). *The proper 3-coloring of  $S_n$  is unique for any  $n \geq 1$ .*

More precisely, Theorem 1.4 states that if  $c_1$  and  $c_2$  are two distinct proper 3-colorings of  $S_n$ , then it is possible to transform  $c_1$  to  $c_2$  by performing rotations, reflections, and/or a change of colors used.

In [4], Chartrand et al. also showed the following:

**Theorem 1.5** (Chartrand, Okamoto, Zhang [4]). *For every graph  $G$ ,*

$$\sigma(G) \leq \chi(G).$$

Hence, the sigma chromatic numbers of the Sierpiński gasket graphs and the Hanoi graphs are at most three. In this paper, we determine the exact sigma chromatic number of these graphs. Moreover, we also prove the uniqueness, in the same sense as in Theorem 1.4, of the sigma coloring of  $S_n$  ( $n \geq 2$ ) that uses the minimum number of colors. The following are our main results.

**Theorem 1.6.**

- (1) *For all  $n \geq 3$ , the sigma chromatic number of  $H_n$  is 3.*
- (2) *For all  $n \geq 2$ , the sigma chromatic number of  $S_n$  is 2.*
- (3) *For all  $n \geq 2$ , the sigma 2-coloring of  $S_n$  is unique.*

In Section 2, we prove statement (1) by showing that  $H_n$ ,  $n \geq 3$ , does not have a sigma 2-coloring. In Section 3, we prove statement (2) by constructing iteratively a sigma 2-coloring of  $S_n$ ,  $n \geq 2$ . Finally, in Section 4, we prove the uniqueness of this sigma 2-coloring.

## 2. The Sigma Chromatic Number of the Hanoi Graphs

In this section, we prove that the sigma chromatic number of the Hanoi graph  $H_n$  is 3 for all  $n \geq 3$ . We begin with the following proposition.

**Proposition 2.1.** *Let  $n \geq 3$ . If  $H_n$  is not sigma 2-colorable, then neither is  $H_{n+1}$ .*

*Proof.* Suppose that  $H_{n+1}$  has a sigma 2-coloring  $c$  that uses the colors  $a$  and  $b$ . Moreover, assume that  $a$  and  $b$  have been chosen so that the sets  $\{3a, 2a + b, 2b + a, 3b\}$  and  $\{2a, a + b, 2b\}$  are disjoint.


Let  $S$  be one of the three  $H_n$  subgraphs of  $H_{n+1}$  and define the coloring  $c'$  to be the restriction of  $c$  to  $S$ . We want to show that  $c'$  is a sigma 2-coloring of  $S$ .

To this end, let  $x$  and  $y$  be the two corner vertices of  $S$  that originally have degree three as vertices of  $H_{n+1}$ . Note that  $c$  and  $c'$  induce the same color sums for all vertices in  $S \setminus \{x, y\}$ . Hence, to show that  $c'$  is a sigma coloring of  $S$ , we need to focus only on the color sums of  $x$  and  $y$ . Now, since  $x$  and  $y$  are vertices with degree two in  $S$ , their color sums with respect to  $c'$  are in the set  $\{2a, a + b, 2b\}$ . Hence, by the assumption, these color sums cannot be equal to the color sum of any of the neighbors of  $x$  or  $y$  since these neighbors have degree three. This completes the proof.  $\square$

Thus, to prove that  $\sigma(H_n) = 3, n \geq 3$ , it is sufficient to prove that  $H_3$  is not sigma 2-colorable. We show this in the following lemma.

**Lemma 2.2.** *The sigma chromatic number of  $H_3$  is 3.*

*Proof.* Suppose there is a sigma 2-coloring  $c : V(H_3) \rightarrow \{a, b\}$  of  $H_3$ . We show that  $c$  produces a contradiction. Let us label the vertices of  $H_3$  as shown in Fig. 5.



**Figure 5.** Vertex labels of the Hanoi graph  $H_3$ .

We consider cases based on the possible values of  $c(v_4), c(v_6)$ , and  $c(v_7)$ . There are four cases, in each of which we show how the contradiction arises.

**Case 1**  $(c(v_4), c(v_6), c(v_7)) = (a, a, a)$

Refer to Fig. 6. In this case, the partial color sums of  $v_4, v_6$ , and  $v_7$  are all equal to  $2a$ . Since  $c$  is a sigma coloring, the actual color sums of the three vertices must be different from each other. This implies that  $c(v_2), c(v_8)$ , and  $c(u_1)$  must all have different values. Clearly, this is not possible since there are only two colors.

**Case 2**  $(c(v_4), c(v_6), c(v_7)) = (b, a, a)$

In this case, the partial color sums of  $v_6$  and  $v_7$  are both equal to  $a + b$ . Since  $c$  is a sigma coloring, the actual color sums of  $v_6$  and  $v_7$  must be distinct. Hence,  $c(u_1) \neq c(v_8)$ .

(2.a) Suppose  $(c(u_1), c(v_8)) = (a, b)$ . Refer to Fig. 7. In this case,  $\sigma(v_6) = 2a + b$  and  $\sigma(v_7) = 2b + a$ . Since  $\sigma(v_6)$  cannot be equal to  $\sigma(u_1)$ , we must have  $\{c(u_2), c(u_3)\} \neq \{a, b\}$ . Moreover, Case 1 implies that  $(c(u_2), c(u_3)) \neq (a, a)$ . Hence,  $(c(u_2), c(u_3))$  must be  $(b, b)$ . This implies that  $\sigma(u_1) = 2b + a$  and that  $u_2$  and  $u_3$  both have partial color sums equal to  $a + b$ . Since the actual color sums of  $u_2$  and  $u_3$  cannot be equal to  $\sigma(u_1)$ , there are no possible values for  $c(u_4)$  and  $c(u_5)$  that do not violate the assumption that  $c$  is a sigma coloring.




Figure 6. Case 1.




Figure 7. Case 2.a.

(2.b) Suppose  $(c(u_1), c(v_8)) = (b, a)$ . This case can be treated in a similar way as Case 2.1.

**Case 3**  $(c(v_4), c(v_6), c(v_7)) = (a, a, b)$

In this case, the partial color sums of  $v_4$  and  $v_6$  are both equal to  $a + b$ . Since  $c$  is a sigma coloring, the actual color sums of  $v_4$  and  $v_6$  must be distinct. Hence,  $c(v_2) \neq c(u_1)$ .

(3.a) Suppose  $(c(v_2), c(u_1)) = (b, a)$ . Then  $\sigma(v_4) = 2b + a$  and  $\sigma(v_6) = 2a + b$ . This proceeds in the same manner as Case (2.a).

(3.b) Suppose  $(c(v_2), c(u_1)) = (a, b)$ . Then  $\sigma(v_6) = 2b + a$  and  $\sigma(v_4) = 2a + b$ . It follows that  $c(v_8) = a$  and  $\sigma(v_7) = 3a$ .

- (i) Suppose  $(c(v_5), c(v_9)) = (a, a)$ . By Case 1, this leads to a contradiction.
- (ii) Suppose  $(c(v_5), c(v_9)) = (b, a)$ . By Case 2, this leads to a contradiction as well.
- (iii) Suppose  $(c(v_5), c(v_9)) = (a, b)$ . Refer to Fig. 8. In this case, the partial color sums of  $v_2$  and  $v_3$  are both equal to  $2a$ . Since their actual color sums must be distinct, there is no possible value for  $c(v_1)$ .




Figure 8. Case 3.b.iii.




Figure 9. Case 3.b.iv.




Figure 10. Case 4.

(iv) Suppose  $(c(v_5), c(v_9)) = (b, b)$ . If  $c(v_3) = b$ , then the partial color sums of  $v_2$  and  $v_3$  are equal and there is no possible value for  $c(v_1)$ . Hence,  $c(v_3) = a$  and  $\sigma(v_5) = 2a + b$ . It follows that  $c(w_1) = b$  and  $\sigma(v_9) = 2b + a$ . This case then proceeds similarly as Case (2.a). Refer to Fig. 9

**Case 4**  $(c(v_4), c(v_6), c(v_7)) = (a, b, a)$

Refer to Fig. 10. In this case, the partial color sums of  $v_4$  and  $v_7$  are both equal to  $a + b$ . Since  $c$  is a sigma coloring, the actual color sums of  $v_4$  and  $v_7$  must be distinct. Hence,  $c(v_2) \neq c(v_8)$ . This implies that  $\{\sigma(v_4), \sigma(v_7)\} = \{2a + b, 2b + a\}$ ; hence,  $c(u_1) = a$  so that  $\sigma(v_6) = 3a$ . By Cases 1, 2, and 3, there are no possible values for  $c(u_2)$  and  $c(u_3)$ . Therefore, there is no sigma 2-coloring of  $H_3$ . Since  $\chi(H_3) = 3$  and  $\sigma(H_3) \leq \chi(H_3)$ , we must have  $\sigma(H_3) = 3$ .  $\square$

**3. The Sigma Chromatic Number of the Sierpiński Gasket Graphs**


In this section, we prove that the sigma chromatic number of the Sierpiński gasket graph  $S_n$  is 2 for all  $n \geq 2$ . Throughout this section, we use  $a, b$  to denote distinct positive integers for which the sets  $\{a + b, 2a, 2b\}$  and  $\{2a + 2b, 3a + b, 3b + a, 4a, 4b\}$  are disjoint. In Fig. 11, we present a sigma 2-coloring of  $S_2$  using  $a$  and  $b$  as colors. Our proof consists of the following steps:

- (i) Construct a sigma 2-coloring of  $S_3$  using the sigma 2-coloring of  $S_2$  in Fig. 11
- (ii) Construct a sigma 2-coloring of  $S_4$  using the obtained sigma 2-coloring of  $S_3$
- (iii) Generalize to any  $S_n$  by induction


*3.1. Construction of a sigma 2-coloring of  $S_3$*

Denote the coloring in Fig. 11 by LL. Perform two operations on LL to obtain two new colorings LR and U:


- (i) To obtain LR, we rotate LL 120° clockwise. LR is shown in Fig. 12.
- (ii) To obtain U, we reflect LL along the angle bisector of its lower left corner. We then interchange the colors  $a$  and  $b$ . U is shown in Fig. 13.



**Figure 11.** LL.



**Figure 12.** LR.



**Figure 13.** U.

Notice that the above operations ensure that the colorings in LR and U are sigma colorings as well. Using LL, LR, and U, construct a sigma coloring of  $S_3$  as shown in Fig. 14. In this figure, a double arrow between a pair of corner vertices signifies that these two vertices are to be identified with each other. The resulting sigma coloring of  $S_3$  is shown in Fig. 15.


*3.2. Construction of a sigma 2-coloring of  $S_4$*

Denote by  $LL_2$  the coloring in Fig. 15. As in the previous construction, perform operations on  $LL_2$  to obtain two new colorings  $LR_2$  and  $U_2$ :


- (i) To obtain  $LR_2$ , reflect  $LL_2$  along the angle bisector of its lower left corner, then interchange the colors  $a$  and  $b$ .
- (ii) To obtain  $U_2$ , rotate  $LL_2$  120° counterclockwise.

Using  $LL_2, U_2$ , and  $LR_2$ , construct the sigma 2-coloring of  $S_4$  as done for  $S_3$ . The constructed sigma coloring is shown in Fig. 16.






**Figure 14.** Constructing a sigma coloring of  $S_3$  using LL, LR, and U.



**Figure 15.** The constructed sigma coloring of  $S_3$ .



**Figure 16.** The constructed sigma 2-coloring of  $S_4$ .

3.3. Induction

In Fig. 17, a simplified version of Fig. 16 is shown. This version shows only the vertices of the three corner triangles, their colors, and their color sums.

Fig. 16 shows the important similarities between the sigma 2-colorings of  $S_2$  and  $S_4$ :

- Corresponding corner vertices have the same colors and color sums.
- The corresponding neighbors of corresponding corner vertices have the same colors and color sums.

Due to these similarities, it is possible to repeat the operations performed previously to obtain sigma 2-colorings of  $S_5$  and  $S_6$ . This process can then be repeated to complete the proof. Therefore, the sigma chromatic of  $S_n, n \geq 2$ , is 2. □




Figure 17. A simplified version of Fig. 16.

The preceding proof leads to the following result, which is more precise than statement (2) of Theorem 1.6.

**Theorem 3.1.** *Let  $n \geq 2$  be an integer.*

- (i) *If  $n$  is even, then  $S_n$  has a sigma 2-coloring of the form shown in Fig. 18.*
- (ii) *If  $n$  is odd, then  $S_n$  has a sigma 2-coloring of the form shown in Fig. 19.*




Figure 18. Sigma 2-coloring for  $S_n$ ,  $n$  even.




Figure 19. Sigma 2-coloring for  $S_n$ ,  $n$  odd.

#### 4. Uniqueness of the Sigma 2-coloring of Sierpiński Gasket Graphs

As in Section 3, use  $a, b$  to denote distinct positive integers for which the sets  $\{a + b, 2a, 2b\}$  and  $\{2a + 2b, 3a + b, 3b + a, 4a, 4b\}$  are disjoint. We now make the following definitions.

**Definition 4.1.** *Let  $n \geq 2$  be an integer and let  $c$  be a sigma 2-coloring, using the colors  $a$  and  $b$ , of  $S_n$ . Define the operations  $r, s$ , and  $t$  as follows:*

- (i) *The coloring  $rc$  is obtained by rotating  $c$   $120^\circ$  clockwise.*
- (ii) *The coloring  $sc$  is obtained by reflecting  $c$  along the angle bisector of the lower left corner vertex.*
- (iii) *The coloring  $tc$  is obtained by interchanging the colors  $a$  and  $b$ .*

Also, define  $G$  to be the group  $\langle r, s, t \rangle$  and  $Gc$  to be the set  $\{gc : g \in G\}$ .

Clearly,  $G$  is isomorphic to  $D_3 \times \mathbb{Z}_2$ ; hence, it has 12 elements. Moreover, all the elements of  $Gc$  are also sigma 2-colorings of  $S_n$ .

**Definition 4.2.** *Let  $n \geq 2$  be an integer and let  $c$  be a sigma 2-coloring of  $S_n$ . Define the sigma 2-colorings  $LL(c), U(c), LR(c)$  of  $S_{n-1}$  as follows:*

- (i)  *$LL(c)$  is the restriction of  $c$  to the lower left  $S_{n-1}$  subgraph of  $S_n$ .*

(ii)  $U(c)$  is the restriction of  $c$  to the upper  $S_{n-1}$  subgraph of  $S_n$ .

(iii)  $LR(c)$  is the restriction of  $c$  to the lower right  $S_{n-1}$  subgraph of  $S_n$ .

We also write  $c = (\alpha, \beta, \gamma)$  to signify that  $\alpha, \beta$ , and  $\gamma$  are sigma 2-colorings of  $S_{n-1}$  for which  $(LL(c), U(c), LR(c)) = (\alpha, \beta, \gamma)$ . For instance, the coloring  $c_3$  shown in Fig. 15 can be written as  $c_3 = (c_2, tsc_2, rc_2)$ , where  $c_2$  is the coloring shown in Fig. 11.

For  $S_n, n \geq 2$ , we denote by  $c_n$  the sigma 2-coloring constructed in the proof in Section 3. Recall that  $c_n$  follows the forms shown in Figures 18 and 19. In order to prove statement (3) of Theorem 1.6, we prove the following equivalent statement.

**Lemma 4.3.** *Let  $a, b$  be fixed distinct positive integers for which the sets  $\{a + b, 2a, 2b\}$  and  $\{2a + 2b, 3a + b, 3b + a, 4a, 4b\}$  are disjoint. Then any sigma 2-coloring  $c$  of  $S_n, n \geq 2$ , is in  $Gc_n$ .*

*Proof.* Our proof is by induction. The case  $n = 2$  can be verified easily; that is, the totality of sigma 2-colorings of  $S_2$  is given by  $Gc_2$ , where  $c_2$  is the sigma 2-coloring shown in Fig. 11.

Now, suppose the statement holds for  $n$  and let  $c$  be a sigma 2-coloring of  $S_{n+1}$ . By the assumption on the colors  $a$  and  $b$ ,  $LL(c), U(c)$ , and  $LR(c)$  must be sigma 2-colorings of  $S_n$ ; hence, by the inductive hypothesis, they belong to the set  $Gc_n$ .

**Case 1.** Suppose  $n$  is even.

By applying  $t$  to  $c$  at most once, we can guarantee that the resulting coloring  $c'$  has the lower left corner vertex colored  $a$ ; that is,  $LL(c')$  is in  $\{c_n, sc_n, trc_n, strc_n, tr^2c_n, str^2c_n\}$ . Moreover, by applying  $s$  to  $c'$  at most once, we can guarantee that the resulting coloring  $c''$  has  $LL(c'')$  that is in  $\{c_n, trc_n, tr^2c_n\}$ . Therefore, we can simply assume without loss of generality that  $LL(c)$  is in  $\{c_n, trc_n, tr^2c_n\}$ .

*Case 1.1.* Suppose  $LL(c) = c_n$ . Since there are only 12 possible choices each for  $U(c)$  and  $LR(c)$ , it is easy to verify that  $c = (c_n, tsc_n, rc_n)$ . In this case,  $c = c_{n+1}$ , which is the sigma 2-coloring constructed in Section 3. Naturally,  $c = c_{n+1} \in Gc_{n+1}$ .

*Case 1.2.* Suppose  $LL(c) = trc_n$ . In this case, we must have  $c = (trc_n, sr^2c_n, sc_n)$ . Furthermore, we have

$$c_{n+1} = (c_n, tsc_n, rc_n) \implies rsc_{n+1} = (trc_n, rsc_n, rsrc_n).$$

By applying the property that  $(sr)^2$  is the identity,  $rsc_{n+1}$  simplifies to  $c$ ; that is,  $c \in Gc_{n+1}$ .

*Case 1.3.* Suppose  $LL(c) = tr^2c_n$ . In this case, we must have  $c = (tr^2c_n, trc_n, sr^2c_n)$ . Furthermore, we have

$$c_{n+1} = (c_n, tsc_n, rc_n) \implies trc_{n+1} = (tr^2c_n, trc_n, rsc_n).$$

As in Case 1.2.,  $trc_{n+1}$  simplifies to  $c$ ; that is,  $c \in Gc_{n+1}$ .

**Case 2.** Suppose  $n$  is odd.

Similar to Case 1, we can simply assume without loss of generality that  $LL(c)$  is in  $\{c_n, r^2c_n, trc_n\}$ .

*Case 2.1.* Suppose  $LL(c) = c_n$ . In this case,  $c = (c_n, r^2c_n, tsc_n)$ . In this case,  $c = c_{n+1}$ , which is the sigma 2-coloring constructed in Section 3. Naturally,  $c = c_{n+1} \in Gc_{n+1}$ .

*Case 2.2.* Suppose  $LL(c) = r^2c_n$ . In this case,  $c = (r^2c_n, tsc_n, tsrc_n)$ . Furthermore,

$$c_{n+1} = (c_n, r^2c_n, tsc_n) \implies srtc_{n+1} = (rsrc_n, tsc_n, tsrc_n).$$

Hence,  $srtc_{n+1}$  is equal to  $c$ ; that is,  $c \in Gc_{n+1}$ .

*Case 2.3.* Suppose  $LL(c) = trc_n$ . In this case,  $c = (trc_n, src_n, tr^2c_n)$ . Furthermore,

$$c_{n+1} = (c_n, r^2c_n, tsc_n) \implies r^2tc_{n+1} = (trc_n, r^2sc_n, tr^2c_n).$$

Hence,  $r^2tc_{n+1}$  is equal to  $c$ ; that is,  $c \in Gc_{n+1}$ .

This completes the proof of the lemma and, consequently, of statement (3) of Theorem 1.6.  $\square$

## 5. Conclusion

In this paper, we have determined the sigma chromatic numbers of two families of graphs: the Sierpiński gasket graphs and the Hanoi graphs. For the Hanoi graphs, our approach involved showing that, for  $n \geq 3$ , the sigma 2-colorability of  $H_{n+1}$  implies the sigma 2-colorability of  $H_n$ . On the other hand, for the Sierpiński gasket graphs, we employed a recursive construction of a sigma 2-coloring of  $S_n$ . Moreover, we have proven that this sigma 2-coloring is unique up to rotations, reflections, and choice of colors. This uniqueness result is the sigma coloring analog of Klavžar's uniqueness result for proper 3-colorings of  $S_n$ .

## Acknowledgment

The authors would like to thank the Office of the Vice President for the Loyola Schools and the Department of Mathematics BCA fund of Ateneo de Manila University for supporting our attendance to this conference. Moreover, we would like to thank the organizers of ICCGANT 2019 for warmly welcoming us and hosting a wonderful conference. Finally, we would like to thank the referees of this paper for their valuable comments and suggestions.

## References

- [1] Akbari S, Ghanbari M, Manaviyat R and Zare S 2013 On the lucky choice number of graphs *Graphs and Combinatorics* **29** 157–163
- [2] Arett D and Dorée S 2010 Coloring and counting on the tower of Hanoi graphs *Mathematics Magazine* **83** 200–209
- [3] Chartrand G, Okamoto F, Rasmussen C W and Zhang P 2009 The set chromatic number of a graph *Discussiones Mathematicae Graph Theory* **29** 545–561
- [4] Chartrand G, Okamoto F and Zhang P 2010 The sigma chromatic number of a graph *Graphs and Combinatorics* **26** 755–773
- [5] Czerwiński S, Grytczuk J and Żelazny W 2009 Lucky labelings of graphs *Information Processing Letters* **109** 1078–1081
- [6] Dehghan A, Sadeghi M R and Ahadi A 2014 The complexity of the sigma chromatic number of cubic graphs [arXiv:1403.6288v1](https://arxiv.org/abs/1403.6288v1) [math.CO]
- [7] Hinz A 1999 The tower of Hanoi in algebra and combinatorics *ICAC '97, Hongkong, Lecture Notes in Computer Science* 277–289
- [8] Klavžar S 2008 Coloring Sierpiński graphs and Sierpiński gasket graphs *Taiwanese Journal of Mathematics* **12** 513–522
- [9] Kristiana A I, Dafik, Utoyo M I, Slamain, Alfarisi R, Agustin I H and Venkatachalam M 2019 Local irregularity vertex coloring of graphs *International Journal of Civil Engineering and Technology*, **10** 1606–1616
- [10] Luzon P A D, Ruiz M J P and Tolentino M A C 2016 The sigma chromatic number of the circulant graphs  $C_n(1, 2)$ ,  $C_n(1, 3)$ , and  $C_{2n}(1, n)$  *Lecture Notes in Computer Science* **9943** 216–227
- [11] Parisse D 2009 On some metric properties of the Sierpiński graphs  $s(n, k)$  2009 *Ars Combinatoria* **90** 145–160
- [12] Slamain 2017 On distance irregular labelling of graphs *Far East Journal of Mathematical Sciences* **102** 919–932
- [13] Tegui A and Godbole A 2006 Sierpiński gasket graphs and some of their properties *Australasian Journal of Combinatorics* **35** 181–192