

# Gene Section

## Short Communication

# ID2 (inhibitor of DNA binding 2, dominant negative helix-loop-helix protein)

Menno C van Zelm

Dept Immunology, Erasmus MC, University Medical Center, Rotterdam, the Netherlands (MCv)

Published in Atlas Database: June 2014

Online updated version : <http://AtlasGeneticsOncology.org/Genes/ID2ID436ch2p25.html>  
DOI: 10.4267/2042/56406

This work is licensed under a Creative Commons Attribution-Noncommercial-No Derivative Works 2.0 France Licence.  
© 2015 Atlas of Genetics and Cytogenetics in Oncology and Haematology

## Abstract

Review on ID2, with data on DNA/RNA, on the protein encoded and where the gene is implicated.

## Identity

**Other names:** GIG8, ID2A, ID2H, bHLHb26

**HGNC (Hugo):** ID2

**Location:** 2p25.1


## DNA/RNA

### Description

The gene spans 5608 bp containing 5 exons of which 2 (exons 3 and 4) are protein-encoding.

### Transcription

The ID2 gene has 4 transcripts, of which 3 can generate functional protein: ID2-001 (2041bp) contains all 5 exons, ID2-201 (1362 bp) contains 3 exons and ID2-002 contains 2 exons. ID2-003 (474 bp) contains a retained intron.



## Pseudogene

A pseudogene of ID2 is located on chromosome 3.

## Protein

### Description

ID2 belongs to the helix-loop-helix (HLH) protein family. It is composed by 134 aa and belongs to a subgroup HLH family members (ID1, ID2, ID3, ID4) that lack a basic DNA-binding domain. ID proteins form heterodimers with class I basic HLH-group members such as MyoD (Langlands et al., 1997), NEDD9 (Law et al., 1999), and E2A gene products E12 and E47. ID2 has a main domain located on 38-79 aa responsible for the helix-loop-helix conformation. In addition, Id2 contains a 10 aa motif that is responsible for the nuclear export signaling.

### Expression

Expression of ID2 is found in the brain, ovary, liver, lung, thyroid gland and prostate and several subsets of leukocytes. ID2 expression is especially high in Natural Killer (NK-) cells, but is also found in CD4+ T cells, CD8+ T cells, monocytes and precursor B cells.

### Localisation

Nucleus.

### Function

Although it does not bind directly to DNA, by binding basic helix-loop-helix transcription factors through its HLH motif, ID2 may control tissue-specific genes related to cell growth, proliferation and differentiation (Hara et al., 1994; Iavarone et al., 1994). ID2 functions in cell fate decisions in early leukocyte development. Specifically, ID2 is required for NK-cell, innate lymphoid and lymphoid tissue inducer cells (Boos et al., 2007; Moro et al., 2010; Yokota et al., 1999). Furthermore, ID2 functions in the development of several dendritic cell subsets: Langerhans cells, cutaneous dendritic cells and splenic CD8a+ dendritic cells (Hacker et al., 2003). Although ID2 seems redundant for T-cell development in thymus, ID2 promotes NKT-cell development (Verykokakis et al., 2013), and it is involved in effector differentiation (Masson et al., 2013), as well as  $\gamma\delta$ T cell homeostasis (Zhang et al., 2014). Finally, ID2 inhibits progression of precursor-B-cell development (Hara et al., 1997; Jensen et al., 2013), as well as activation-induced deaminase expression during B-cell responses (Gonda et al., 2003), likely through inhibition of E47 (Sayegh et al., 2003).

Besides leukocytes, ID2 has been found to function in erythrocyte development (Ji et al., 2008), enterocyte precursor and lung epithelial cell differentiation in mice (Rawlins et al., 2009).

Furthermore, female mice lacking ID2 show lactation defects (Mori et al., 2000), and male mice have impaired spermatogenesis (Sablitzky et al., 1998).

### Homology

ID2 is highly conserved in vertebrates, including mammals, reptiles, and fish.

## Mutations

### Germinal

No germinal mutations have been reported.

### Somatic

No somatic mutations have been reported.

## Implicated in

### Neuroblastoma

#### Note

ID2 functions as a key regulator in the phenotypic transition of neuroblastoma tumor cells (Chakrabarti et al., 2013). Anchorage-dependent (AD) neuroblastoma cells express much higher levels of ID2 than anchorage-independent (AI) cells. Moreover, knockdown of ID2 in AD cells induces an AI phenotype, whereas the opposite is seen upon forced expression of ID2 in AI cells. The function of ID2 in this process is at least in part via negative regulation of the TGF $\beta$ /Smad pathway.

### Colon carcinoma

#### Note

ID2 expression is upregulated by enhanced beta-catenin signaling and subsequent beta-catenin /TCF mediated transcription.

The induction of ID2 expression increases anchorage-independent survival of these cells (Rockman et al., 2001).

### Melanoma

#### Note

The transition of melanoma to a more aggressive malignancy is associated with the resistance to growth inhibition by TGF- $\beta$  (Javelaud et al., 2008). In susceptible cells, TGF- $\beta$  suppresses ID2 expression and allows p15<sup>Ink4b</sup> to induce a cell cycle arrest (Schlegel et al., 2009). Upon obtaining resistance to TGF- $\beta$ , the tumor cells overexpress ID2 and remain in cycle.

### Retinoblastoma

#### Note

ID2 is overexpressed due to transcriptional activation by oncoproteins of the Myc family in retinoblastoma, where it is thought to inhibit Retinoblastoma protein family members (Lasorella et al., 2000; Lasorella et al., 2005).

## Hodgkin's lymphoma

### Note

The majority of Hodgkin's lymphomas are derived from germinal center B cells. Still, Hodgkin-Reed/Sternberg (HRS) cells of classical Hodgkin's lymphoma have a very atypical phenotype.

This is the result of overexpression of ABF-1 and Id2, which inhibit the function of the B cell-determining transcription factor E2A (Mathas et al., 2006; Renne et al., 2006).

The mechanism resulting in ID2 overexpression, nor the impact of ID2 on cell cycle progression in HRS cells have been demonstrated experimentally yet (Cotta and Medeiros, 2008).

## References

Hara E, Yamaguchi T, Nojima H, Ide T, Campisi J, Okayama H, Oda K. Id-related genes encoding helix-loop-helix proteins are required for G1 progression and are repressed in senescent human fibroblasts. *J Biol Chem.* 1994 Jan 21;269(3):2139-45

Iavarone A, Garg P, Lasorella A, Hsu J, Israel MA. The helix-loop-helix protein Id-2 enhances cell proliferation and binds to the retinoblastoma protein. *Genes Dev.* 1994 Jun 1;8(11):1270-84

Hara E, Hall M, Peters G. Cdk2-dependent phosphorylation of Id2 modulates activity of E2A-related transcription factors. *EMBO J.* 1997 Jan 15;16(2):332-42

Langlands K, Yin X, Anand G, Prochownik EV. Differential interactions of Id proteins with basic-helix-loop-helix transcription factors. *J Biol Chem.* 1997 Aug 8;272(32):19785-93

Sablitzky F, Moore A, Bromley M, Deed RW, Newton JS, Norton JD. Stage- and subcellular-specific expression of Id proteins in male germ and Sertoli cells implicates distinctive regulatory roles for Id proteins during meiosis, spermatogenesis, and Sertoli cell function. *Cell Growth Differ.* 1998 Dec;9(12):1015-24

Law SF, Zhang YZ, Fashena SJ, Toby G, Estojak J, Golemis EA. Dimerization of the docking/adaptor protein HEF1 via a carboxy-terminal helix-loop-helix domain. *Exp Cell Res.* 1999 Oct 10;252(1):224-35

Yokota Y, Mansouri A, Mori S, Sugawara S, Adachi S, Nishikawa S, Gruss P. Development of peripheral lymphoid organs and natural killer cells depends on the helix-loop-helix inhibitor Id2. *Nature.* 1999 Feb 25;397(6721):702-6

Lasorella A, Noseda M, Beyna M, Yokota Y, Iavarone A. Id2 is a retinoblastoma protein target and mediates signalling by Myc oncoproteins. *Nature.* 2000 Oct 5;407(6804):592-8

Mori S, Nishikawa SI, Yokota Y. Lactation defect in mice lacking the helix-loop-helix inhibitor Id2. *EMBO J.* 2000 Nov 1;19(21):5772-81

Rockman SP, Currie SA, Ciavarella M, Vincan E, Dow C, Thomas RJ, Phillips WA. Id2 is a target of the beta-catenin/T cell factor pathway in colon carcinoma. *J Biol Chem.* 2001 Nov 30;276(48):45113-9

Gonda H, Sugai M, Nambu Y, Katakai T, Agata Y, Mori KJ, Yokota Y, Shimizu A. The balance between Pax5 and Id2

activities is the key to AID gene expression. *J Exp Med.* 2003 Nov 3;198(9):1427-37

Hacker C, Kirsch RD, Ju XS, Hieronymus T, Gust TC, Kuhl C, Jorgas T, Kurz SM, Rose-John S, Yokota Y, Zenke M. Transcriptional profiling identifies Id2 function in dendritic cell development. *Nat Immunol.* 2003 Apr;4(4):380-6

Sayegh CE, Quong MW, Agata Y, Murre C. E-proteins directly regulate expression of activation-induced deaminase in mature B cells. *Nat Immunol.* 2003 Jun;4(6):586-93

Lasorella A, Rothschild G, Yokota Y, Russell RG, Iavarone A. Id2 mediates tumor initiation, proliferation, and angiogenesis in Rb mutant mice. *Mol Cell Biol.* 2005 May;25(9):3563-74

Mathas S, Janz M, Hummel F, Hummel M, Wollert-Wulf B, Lusatis S, Anagnostopoulos I, Lietz A, Sigvardsson M, Jundt F, Jöhrens K, Bommert K, Stein H, Dörken B. Intrinsic inhibition of transcription factor E2A by HLH proteins ABF-1 and Id2 mediates reprogramming of neoplastic B cells in Hodgkin lymphoma. *Nat Immunol.* 2006 Feb;7(2):207-15

Renné C, Martin-Subero JI, Eickernjäger M, Hansmann ML, Küppers R, Siebert R, Bräuninger A. Aberrant expression of ID2, a suppressor of B-cell-specific gene expression, in Hodgkin's lymphoma. *Am J Pathol.* 2006 Aug;169(2):655-64

Boos MD, Yokota Y, Eberl G, Kee BL. Mature natural killer cell and lymphoid tissue-inducing cell development requires Id2-mediated suppression of E protein activity. *J Exp Med.* 2007 May 14;204(5):1119-30

Cotta CV, Medeiros LJ. Expression of helix-loop-helix proteins in classical Hodgkin lymphoma: a possible explanation for a characteristic immunophenotype. *Adv Anat Pathol.* 2008 Mar;15(2):97-104

Javelaud D, Alexaki VI, Mauviel A. Transforming growth factor-beta in cutaneous melanoma. *Pigment Cell Melanoma Res.* 2008 Apr;21(2):123-32

Ji M, Li H, Suh HC, Klarmann KD, Yokota Y, Keller JR. Id2 intrinsically regulates lymphoid and erythroid development via interaction with different target proteins. *Blood.* 2008 Aug 15;112(4):1068-77

Rawlins EL, Clark CP, Xue Y, Hogan BL. The Id2+ distal tip lung epithelium contains individual multipotent embryonic progenitor cells. *Development.* 2009 Nov;136(22):3741-5

Schlegel NC, Eichhoff OM, Hemmi S, Werner S, Dummer R, Hoek KS. Id2 suppression of p15 counters TGF-beta-mediated growth inhibition of melanoma cells. *Pigment Cell Melanoma Res.* 2009 Aug;22(4):445-53

Moro K, Yamada T, Tanabe M, Takeuchi T, Ikawa T, Kawamoto H, Furusawa J, Ohtani M, Fujii H, Koyasu S. Innate production of T(H)2 cytokines by adipose tissue-associated c-Kit(+)/Sca-1(+) lymphoid cells. *Nature.* 2010 Jan 28;463(7280):540-4

Chakrabarti L, Wang BD, Lee NH, Sandler AD. A mechanism linking Id2-TGFβ crosstalk to reversible adaptive plasticity in neuroblastoma. *PLoS One.* 2013;8(12):e83521

Jensen K, Rother MB, Brusletto BS, Olstad OK, Dalsbotten Aass HC, van Zelm MC, Kierulf P, Gautvik KM. Increased ID2 levels in adult precursor B cells as compared with children is associated with impaired Ig locus contraction and decreased bone marrow output. *J Immunol.* 2013 Aug

1;191(3):1210-9

Masson F, Minnich M, Olshansky M, Bilic I, Mount AM, Kallies A, Speed TP, Busslinger M, Nutt SL, Belz GT. Id2-mediated inhibition of E2A represses memory CD8+ T cell differentiation. *J Immunol*. 2013 May 1;190(9):4585-94

Verykokakis M, Krishnamoorthy V, Iavarone A, Lasorella A, Sigvardsson M, Kee BL. Essential functions for ID proteins at multiple checkpoints in invariant NKT cell development. *J Immunol*. 2013 Dec 15;191(12):5973-83

Zhang B, Lin YY, Dai M, Zhuang Y. Id3 and Id2 act as a dual safety mechanism in regulating the development and population size of innate-like  $\gamma\delta$  T cells. *J Immunol*. 2014 Feb 1;192(3):1055-63

---

*This article should be referenced as such:*

van Zelm MC. ID2 (inhibitor of DNA binding 2, dominant negative helix-loop-helix protein). *Atlas Genet Cytogenet Oncol Haematol*. 2015; 19(3):168-171.

---