

University of Texas Rio Grande Valley

ScholarWorks @ UTRGV

Mathematical and Statistical Sciences Faculty
Publications and Presentations

College of Sciences

2020

Screening Potential Citrus Rootstocks for Phytophthora nicotianae Tolerance

Madhurababu Kunta

Sandy Chavez

Zenaida Vilorio

Hilda S. del Rio

Madhavi Devanaboina

The University of Texas Rio Grande Valley

See next page for additional authors

Follow this and additional works at: https://scholarworks.utrgv.edu/mss_fac



Part of the [Mathematics Commons](#), and the [Plant Sciences Commons](#)

Recommended Citation

Kunta, Madhurababu; Chavez, Sandy; Vilorio, Zenaida; del Rio, Hilda S.; Devanaboina, Madhavi; Yanev, George; Park, Jong-Won; and Louzada, Eliezer S., "Screening Potential Citrus Rootstocks for Phytophthora nicotianae Tolerance" (2020). *Mathematical and Statistical Sciences Faculty Publications and Presentations*. 61.

https://scholarworks.utrgv.edu/mss_fac/61

This Article is brought to you for free and open access by the College of Sciences at ScholarWorks @ UTRGV. It has been accepted for inclusion in Mathematical and Statistical Sciences Faculty Publications and Presentations by an authorized administrator of ScholarWorks @ UTRGV. For more information, please contact justin.white@utrgv.edu, william.flores01@utrgv.edu.

Authors

Madhurababu Kunta, Sandy Chavez, Zenaida Vilorio, Hilda S. del Rio, Madhavi Devanaboina, George Yanev, Jong-Won Park, and Eliezer S. Louzada

Screening Potential Citrus Rootstocks for *Phytophthora nicotianae* Tolerance

Madhurababu Kunta and Sandy Chavez

Texas A&M University-Kingsville Citrus Center, 312 N. International Boulevard, Weslaco, TX 78599

Zenaida Viloría

UK Research and Education Center at Princeton, Entomology, 1205 Hopkinsville Street, Princeton, KY 42445

Hilda S. del Río

Texas A&M University-Kingsville Citrus Center, 312 N. International Boulevard, Weslaco, TX 78599

Madhavi Devanaboina

The University of Texas Rio Grande Valley, School of Mathematics and Statistical Sciences, 1201 W. University Drive, Edinburg, TX 78539

George Yanev

The University of Texas Rio Grande Valley, School of Mathematics and Statistical Sciences, 1201 W. University Drive, Edinburg, TX 78539; and Institute of Mathematics and Informatics, Bulgarian Academy of Sciences, 8 Acad. G. Bonchev Str., 1113 Sofia, Bulgaria

Jong-Won Park and Eliezer S. Louzada

Texas A&M University-Kingsville Citrus Center, 312 N. International Boulevard, Weslaco, TX 78599

Additional index words. citrus, *Phytophthora*, root rot, rootstock, tolerance

Abstract. Seeds from four citrus rootstocks including sour orange, Bitters-C22 citrandarin, Sarawak pummelo × Rio Red grapefruit, and Sarawak pummelo × Bower mandarin were exposed to high inoculum levels of *Phytophthora nicotianae* to screen for tolerance. Inoculation of pregerminated seeds (PGIS) and non-PGIS was carried out. The average *P. nicotianae* propagule counts from the soil samples where these seedlings were raised ranged from 424 to 1361 colony forming units/cm³. The proportion of live to dead plants was recorded at 11 months postinoculation, which showed that Sarawak × Bower performed significantly better than other rootstocks. Evaluation of the rootstocks 18 months postinoculation resulted in only one surviving sour orange plant, which suggests potential rootstock resistance.

Citrus trees are susceptible to a variety of diseases and can be combating more than one at the same time. In the Lower Rio Grande Valley (LRGV) of south Texas, a common oomycete causing major crop losses in citrus is *Phytophthora nicotianae* (Kunta et al., 2007). Two species of *Phytophthora*, *P. nicotianae* Breda de Haan (synonymous with *P. parasitica* Dast.) and *P. citrophthora* (Sm. & Sm.) Leonian are the most prevalent and highly destructive pathogens affecting citriculture worldwide (Graham and Menge,

2000). Another species, *P. palmivora*, causes serious damage to fibrous roots and causes citrus brown rot in Florida and Japan (Tashiro et al., 2012; Zitko and Timmer, 1994). Only *P. nicotianae* has been confirmed in Texas (Chaudhary, 2018; Kunta et al., 2007). *Phytophthora* can reproduce asexually through sporangia and zoospores as well as sexually through oospores (Meng et al., 2014). The thick-walled long-term survival spores called chlamydozoospores germinate during wet weather to produce sporangiophores bearing sporangia. The sporangia will develop short-lived motile zoospores with flagella that can swim in water and infect a new host plant. The chlamydozoospore will remain in the soil for several years and can serve as an inoculum source until the environmental conditions are optimal.

Phytophthora causes both above and below ground diseases, such as foot rot and gummosis of the trunk, brown rot of fruit, and fibrous root rot. Fibrous root rot may go unnoticed by growers until the canopy shows

stunting, premature defoliation and branch dieback (Naqvi, 2004). Foot rot infection begins above the soil surface where a lesion on the bark can extend from the scion to the base of the rootstock (Savita and Nagpal, 2012). Gum will exude from diseased bark, serving as a clear characteristic of *P. nicotianae* infection and is referred to as gummosis (Graham and Menge, 1999). Gumming is more noticeable during dry weather vs. wet seasons when rain may potentially rinse away obvious gumming evidence (Savita and Nagpal, 2012). Plants infected with *P. nicotianae* may have a decayed root system leaving roots sloughed of all outer layers. The production of new fibrous roots is apparently outpaced by root death in heavily infected mature trees (Graham, 1995). Root death will ultimately decline the overall health of the host plant as seen in tobacco, ornamental plants, tomato, and citrus cultivars (Lamour et al., 2003).

The economic losses for the citrus industry worldwide exclusively caused by *P. nicotianae* are difficult to estimate as most trees are often affected by more than one disease at the same time (Ludowici et al., 2013). In California, the largest citrus producer in the United States, brown rot and root rot caused by *Phytophthora* accounted for \$12.9 million annual loss (Savita and Nagpal, 2012). In Florida, the second top U.S. citrus producer, *Phytophthora* damage has been estimated to be ≈\$30 to \$60 million annual yield loss, without control treatments (Graham and Menge, 1999). In Texas, there are no reports on the impact of *Phytophthora*; however, because of the use of flood irrigation, *Phytophthora* is endemic.

Phytophthora nicotianae disease management strategies include use of resistant rootstocks, biological control, fungicides, clean material for propagation, and irrigation with adequate drainage (Cacciola and di San Lio, 2008). Metalaxyl and phosphite fungicides have been reported to be effective in combating *Phytophthora* diseases when applied at proper dosages and at right time frame (Graham and Feichtenberger, 2015). *Phytophthora* can be controlled with fungicides, but overuse may render them ineffective through resistance development (Timmer et al., 1998). Biological control efforts using *Penicillium funiculosum* and *Chaetomium globosum* on citrus cultivars (Fang and Tsao, 1995; Hung et al., 2015) have been proven effective; however, it is not preferred as a large-scale control strategy due to specific temperature, soil pH, and host preference for different biological control agents.

Use of resistant rootstocks is an ideal and long-term solution to fight *Phytophthora* diseases. Sour orange is one of the few rootstock that can produce high yields and yet tolerate the high pH calcareous soils common in the LRGV, and therefore, it is the predominant rootstock planted in this area. *Phytophthora* infected trees with typical symptoms of foot rot, root rot, and gummosis are commonly present. Field trials with *Phytophthora*-infected rootstocks in Florida showed Swingle citrumelo and Carrizo citrange having high

Received for publication 6 Feb. 2020. Accepted for publication 22 Apr. 2020.

Published online 28 May 2020.

This research was possible because of financial support from Texas Citrus Producers Board. E.S.L. is the corresponding author. E-mail: eliezer.louzada@tamuk.edu.

This is an open access article distributed under the CC BY-NC-ND license (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

seedling mortality whereas sour orange did not have significant seedling mortality (Bowman et al., 2002, 2003). Graham (1990) tested seven rootstocks including trifoliolate orange (*Poncirus trifoliata*), Ridge Pineapple sweet orange, Carrizo citrange, Swingle citrumelo, sour orange, Cleopatra mandarin, and Volkameriana lemon; the results revealed that trifoliolate orange and Swingle citrumelo were among the most tolerant to root rot. Similarly, trifoliolate orange and Swingle citrumelo performed better than the other rootstocks in their ability to regenerate roots under *Phytophthora* infection (Graham, 1995). In Florida, Carrizo citrange and Swingle citrumelo were considered tolerant and moderately resistant rootstocks, respectively (Graham and Feichtenberger, 2015); however, the same rootstocks failed to grow under field trials conducted in the LRGV (Louzada et al., 2008) due to soil characteristics.

There is an urgent need for a superior rootstock that can provide resistance to *Phytophthora* diseases, tolerate alkaline soil and the weather conditions in the LRGV area, and yet produce high quality fruit. Sour orange rootstock plants show genetic diversity (Lamine and Mliki, 2015) among them which may reflect in differences in relative tolerance to *Phytophthora* diseases. To find an alternative rootstock possessing all the qualities needed for *P. nicotianae* tolerance and successfully grow in the LRGV; four citrus rootstocks including sour orange rootstock were subjected to high concentrations of *P. nicotianae* inoculation.

Materials and Methods

Plant material. Four citrus rootstocks were screened for *P. nicotianae* tolerance including sour orange (*Citrus aurantium*), Bitters-C22 citrandarin (*C. sunki* × Swingle *Poncirus trifoliata*), and hybrids Sarawak pummelo (*C. maxima*) × Rio Red (*C. paradisi* Macfadyen) grapefruit and Sarawak pummelo (*C. maxima*) × Bower mandarin (*C. reticulata* Blanco). This last hybrid was included because it is a similar cross that resulted sour orange even though it was bred as a scion. The Sarawak × Rio Red cross was originally performed for scion improvement. Grapefruit is a pummelo × sweet orange hybrid, whereas sweet orange is a pummelo × mandarin with 75% of the mandarin parent, so we decided to test Sarawak × Rio Red hybrid. Seeds were thoroughly washed with Golden-Glo soap (Spartan, Maumee, OH) and surface-sterilized with 10% commercial bleach (5.25% NaOCl) for 20 min and rinsed three times with autoclaved reverse osmosis water.

Preparation of *P. nicotianae* stock and seed inoculation. Pure *P. nicotianae* culture, previously isolated from LRGV soil and citrus fibrous root and maintained in our laboratory, was used in this study (Chaudhary, 2018). Identity for this culture was previously confirmed by morphological and culture characteristics, and polymerase

chain reaction (PCR). V-8 agar plugs with 7-d-old cultures were placed in 1 L of clarified V8 broth (50 mL V8 juice, 0.5 g CaCO₃, 949.5 mL of water), and incubated in the dark for 7 d at room temperature (24 °C), after which the cultures were poured out into beakers topped with strainers to separate the agar plugs from V8 broth. Zoospores and chlamydospores were produced using standard methods that were previously described (Dhingra and Sinclair, 1995). The zoospore concentration was adjusted to 10⁶ zoospores/mL to inoculate the seeds. The mycelium, zoospores, and chlamydospores were mixed with sterile Metro Mix soil (Sun Gro Horticulture, Agawam, MA) which was moistened to field capacity to produce soil with high concentrations of *P. nicotianae* inoculum (Colburn and Graham, 2007).

Inoculation of PGIS. A total of 408 seeds of each rootstock were inoculated. The seeds were placed in autoclaved paper towels moistened with autoclaved double distilled (DD) water and stored in a tightly closed tray. Pregermination was completed under laboratory conditions at room temperature until radicles emerged from most seeds. After 2 weeks, PGIS were placed into petri dishes (150 × 15 mm) with 50 mL of zoospore suspension per dish and placed under light for 24 h.

Inoculation of NPGIS. A total of 408 seeds of each rootstock were inoculated. NPGIS were immersed in a chlamydospore solution for 5 min and sown into Metro Mix soil infested with *P. nicotianae* as previously described. Two trays were designated for the planting of each rootstock treatment—NPGIS and PGIS—for a total of 16 trays, with 204 seeds per tray.

Afterward, NPGIS and PGIS trays were re-inoculated with *P. nicotianae* by adding the suspension into the soil and moistened to field capacity. Plastic trays with no holes for drainage were chosen and metal-net liners were placed inside. The metal-net liners allowed drainage of excess water and root to grow through, thus facilitate visual identification of root rot and continuous inoculation of the germinated seeds. Roots were clipped to accelerate *P. nicotianae* continuous infection at two and four months after initial planting.

Plant evaluation. Eleven months postinoculation, the plants from each tray were removed and classified as dead or alive. Live plants were evaluated and rated on plant decay from a level of 1 to 3 (Fig. 1). Level 1 represented low decay, level 2 represented moderate decay, and level 3 represented high decay. Plants were characterized level 1 if the root system showed no signs of root decay, the stem remained green, and leaves did not exhibit yellowing. Plants characterized at level 2 had some thinning roots, partially yellow leaves, and slight browning on the stem. Plants characterized at level 3 had yellowing leaves, browning stem, and roots with extreme thinning. Plants were classified as dead or alive by uprooting each plant from each individual tray and analyzing its overall

condition. Plants in each tray classified as alive were individually photographed and measured for plant height and root length. Plants classified as dead from each tray were recorded, but no further measurements were taken. To test for the presence and viability of *P. nicotianae*, root samples were taken from both the live and dead plants. Additionally, soil samples from each tray were collected in Ziploc bags to measure *P. nicotianae* propagule density. Once plants were classified, the remaining surviving plants were replanted in the same soil and left to grow for an additional 7 months after which dead plants were discarded and remaining plants were evaluated for root rot tolerance. The total postinoculation time was 18 months.

Detection of *P. nicotianae* in infected plants and soil. The propagule densities were measured using previously described methods (Bright et al., 2004; Timmer et al., 1988). Root samples were surface sterilized with 10% commercial bleach (5.25% NaOCl) for 15 min, rinsed three times with DD water, and cut into 1-cm pieces. Both root and soil samples were placed on CMA-PARPH medium plates (Jeffers, 2006) and analyzed for *P. nicotianae* propagule densities, expressed as colony forming units (cfu) per cm³ soil.

Conventional PCR assay. DNA was isolated from a small piece of mycelium using the Qiagen DNeasy Plant Mini Kit (Qiagen, Valencia, CA). To detect *P. nicotianae*, conventional PCR (cPCR) assay (Grote et al., 2002) was performed on 2 µL of DNA in a 25-µL reaction using the r-Gradient Thermoblock (Biometra GmbH, Göttingen, Germany). The PCR products were analyzed by electrophoresis on 1% agarose gels stained with ethidium bromide, in TAE buffer and visualized on a BioSpectrum Darkroom (Ultraviolet Products, Upland, CA).

Statistical analysis. The test results given in Table 1 imply no statistically significant difference between the lengths of the live seedlings subject to NPGIS and PGIS treatments. As a result, in the further analysis, we combined the data for NPGIS and PGIS. To determine whether the rootstocks differ with respect to live or dead status and to determine if there is a difference among the rootstocks at the conditional levels 2 and 3, chi-square tests were performed. Level 1 was not included in the testing due to insufficient data. Post hoc tests for pairwise comparisons of proportions were carried out to determine which of the four rootstocks stood out among the others, having most different proportion of Live/Total counts after 11 months. All statistical analyses were performed using JMP 13.0.0 (2016), SAS Institute Inc. (Cary, NC).

Results

Plant decay, height, and root length for each citrus rootstock. The plant height and root length measurements for the seedlings derived from NPGIS and PGIS seeds at 11-month postinoculation with *P. nicotianae* are shown in Fig. 2 and Table 1. Plants were



Fig. 1. Citrus plant decay showed three levels of disease symptoms, levels 1, 2, and 3, at 11 months postinoculation with *Phytophthora nicotianae*.

Table 1. Plant height and root length (cm) from seedlings derived from non-pregerminated inoculated seeds (NPGIS) were compared with the seedlings derived from pregerminated inoculated seeds (PGIS) by rootstock to evaluate their performance at 11-months postinoculation with *Phytophthora nicotianae*.

Rootstock	Shoot						Root					
	L1		L2		L3		L1		L2		L3	
	NPGIS	PGIS	NPGIS	PGIS	NPGIS	PGIS	NPGIS	PGIS	NPGIS	PGIS	NPGIS	PGIS
Sarawak × Bower	12.5	11.2	11.5	11.6	7.9	8.9	14.1	11.1	11.4	11.1	7.81	12.8
<i>P</i> value		0.39		0.83		0.51		0.11		0.77		0.01*
Sour orange	—	9.30	7.00	9.50	7.50	7.75	—	6.50	6.00	5.60	4.78	4.05
<i>P</i> value				0.13		0.69				0.73		0.38
Bitters-C22	—	—	12.00	10.30	9.59	8.90	—	—	5.00	7.50	5.77	5.90
<i>P</i> value						0.43						0.94
Sarawak × Rio Red	—	12.30	—	7.60	—	6.00	—	11.10	—	7.00	—	4.50

L1, L2, and L3: levels 1, 2, and 3 of plant decay symptoms.

*Significantly different.




Fig. 2. The boxplots display the range in plant height and root length measurements for the seedlings derived from non-pregerminated (NPGIS) and pregerminated (PGIS) seeds at 11 months postinoculation with *Phytophthora nicotianae*. The dots represent outliers.

classified into levels based on visual cues of the root rot symptoms shown by each plant. An analysis was run for each rootstock between treatment groups and level classification to determine if there was a statistical difference ($P < 0.05$) between each level (Table 1).

Sour orange. There were no surviving sour orange NPGIS plants at 11 months postinoculation that could be classified as level 1; therefore, there was no analysis done between the NPGIS group and the four PGIS plants classified as level 1. Level 2 plant height for the PGIS treatment group averaged to 2.5 cm taller than the NPGIS group but still was not significantly different. The root measurements between NPGIS and PGIS for level 2 differed by less than 0.5 cm. A level

3 category was designated for the most damaged plants and sour orange NPGIS group had nine plants collected while PGIS group had 10 plants. Level 3 NPGIS and PGIS displayed fairly equal overall plant height and root length with means differing less than 1 cm for both measurement averages. Analysis of variance (ANOVA) performed for the shoot length of the NPGIS did not provide significant results between levels ($P = 0.782$) nor did the shoot length analysis of the PGIS group ($P = 0.157$). The root length analysis for the NPGIS treatment group did not provide a significant difference ($P = 0.705$); however, the root length analysis for the PGIS treatment group did show statistical differences between classification levels ($P = 0.048$). Only one PGIS sour orange plant

survived 18 months postinoculation with *P. nicotianae*.

Sarawak pummelo × Bower mandarin. The Sarawak pummelo × Bower mandarin level 1 NPGIS treatment group ($n = 5$) resulted in a plant height with a mean of 12.50 cm, whereas the PGIS group had a slightly lower mean of 11.20 cm ($n = 20$) but did not provide a significant difference ($P = 0.39$). The corresponding root measurements of level 1 Sarawak pummelo × Bower mandarin NPGIS and PGIS treatment groups also did not provide a significant difference in lengths ($P = 0.11$). Level 2 plant height and root length measurements of NPGIS and PGIS groups were almost identical, with means having a difference of less than 0.5 cm, therefore did not have a significant

Table 2. Contingency table: status by rootstock.

Count total %	C22	Sarawak × Bower	Sarawak × RioRed	Sour orange	Total
Dead	112 15.36	89 12.21	133 18.24	174 23.87	508 69.68
Live	28 3.84	96 13.17	24 3.29	73 10.01	221 30.32
Total	140 19.20	185 25.38	157 21.54	247 33.88	729
Tests					
N	729	df	-LogLike	R ² (U)	
		3	32.014891	0.0322	
Likelihood ratio			χ ²	P > χ ²	
Pearson			64.030	<0.0001*	
			64.675	<0.0001*	




Fig. 3. Contingency analysis of rootstock by status (dead/live).




Fig. 4. Contingency analysis of rootstock by level of plant decay.

difference. Level 3 shoot heights for NPGIS and PGIS groups were not significantly different ($P = 0.51$), but the root lengths were significantly different with PGIS seeds growing 5 cm longer. An ANOVA performed

between levels of the NPGIS group determined there was a significant difference for both shoot and root measurements, with P values of 0.002 and <0.001 , respectively. The pregerminated treatment group did not show

significant differences between the PGIS group for either shoot length ($P = 0.079$) or root length ($P = 0.584$).

Bitters-C22 citrandarin. There were no seedlings that could be classified as level 1 for NPGIS and PGIS; therefore, no analysis was performed for this level. An analysis was not conducted between Level 2 NPGIS and PGIS treatment groups due to the large difference in sample sizes. One plant from the NPGIS treatment group displayed moderate symptoms enough to be classified at Level 2 with a height of 12 cm and root length of 5 cm. PGIS had 11 plants that were classified as Level 2 with a mean shoot height of 10.3 cm and root length of 7.5 cm. Plants classified at a Level 3 for this rootstock did not show a significant difference in average plant height or root length between NPGIS and PGIS treatment groups where in both cases the differences were less than one centimeter and P values were 0.43 and 0.94, respectively. The NPGIS treatment group had 11 plants collected for evaluation. There were no plants classified as level 1, one plant classified as level 2 and 11 plants classified as level 3. There were no significant differences calculated for either treatment group between any levels. A greater sample size would have provided a more accurate analysis to determine the statistical difference between classification levels.

Sarawak pummelo × RioRed grapefruit. There were no Sarawak pummelo × Rio Red grapefruit NPGIS group plants that survived at 11 months postinoculation collection. Because no measurements could be taken, comparison analyses were not performed between NPGIS and PGIS groups by level. The NPGIS group did not provide any surviving plants 11 months postinoculation. The PGIS treatment group did have 24 surviving plants in total, and there were significant differences between means for both the plant height ($P = 0.008$) and root measurements ($P < 0.001$).

Confirmation of P. nicotianae in plants and soil. The presence of *P. nicotianae* was confirmed in each of the PARPH media plates by observing colony morphology, sporangium, hyphal, and oospore characteristics (Jeffers, 2006). The propagules in the infested soil where the plants were raised ranged from 424 to 1361 cfu/cm³ at 11-month post inoculation. The soil sample taken from the tray where there was only one surviving sour orange rootstock plant at 18-month post inoculation showed propagule count of 890 cfu/cm³. The root pieces consistently produced characteristic *P. nicotianae* colonies on CMA-PARPH medium plates. An expected PCR amplification product of 737 bp was obtained with all the DNA samples derived from all the morphologically characterized colonies.

Phytophthora tolerance in different citrus rootstocks. Plants collected from each individual tray displayed symptoms corresponding to *P. nicotianae* infection; leaves exhibited yellowing, roots were thinning, and stems were fragile and brown. A contingency table (Table 2) and a mosaic graph

(Fig. 3) were developed for the four rootstocks by the status dead (D) or live (L). The R^2 test had a P value of 0.0322; hence, the rootstocks differed in status D or L. Mosaic graph and contingency table for the four rootstocks by level 2 and level 3 for the category live are shown in Fig. 4 and Table 3, respectively. The R^2 test had a P value 0.0216. Therefore, given that the plants are alive, there is difference between the rootstocks at the conditional levels 2 and 3. Pairwise comparisons of proportions for the four rootstocks indicated that the proportion of Sarawak \times Bower is significantly different from the other three rootstocks. Indeed, the proportion of live for Sarawak \times Bower ($96/185 = 0.52$) is much higher than the proportions of the others. The results given in Table 4 identified the percent of live Sarawak \times Bower seedlings as significantly higher than that for all other rootstocks.

Only surviving sour orange plant. After 18 months postinoculation, only one PGIS sour orange plant out of all treatments survived, and it did not display any symptoms corresponding to *P. nicotianae* infection. Leaves on the sour orange plant were not discolored, the stem was sturdy and visibly healthy (Fig. 5A), and the root system showed no signs of thinning or root decay (Fig. 5B).

Discussion

The economic losses due to *Phytophthora* diseases in the LRGV citrus orchards and unavailability of a resistant rootstock led to the screening of different citrus rootstocks for tolerance. Rootstocks such as Swingle citrumelo, Carrizo citrange, and Troyer citrange are tolerant to *P. nicotianae*-induced diseases; however, they are not adapted to the LRGV soils.

Rootstocks in the LRGV should be tolerant to *P. nicotianae*, high pH calcareous soils, and preferentially cold hardy. The most urgent need is tolerance/resistance to *Phytophthora* and adaptability to our soil conditions; we decided to start looking at potential variation in seedlings of rootstocks already tested in the LRGV and also at population of hybrids that we recently produced. One important characteristic of any citrus rootstock is that they should present a high degree of nucellar embryony to maintain uniformity of propagated scion varieties. Bitters-C22 citrandarin was screened due to its past performance exhibiting all favorable traits, including high degree of polyembryony. Bitters-C22 has shown tolerance to freezing temperatures as well as to calcareous soils, while still producing quality fruit (Federici et al., 2009). After a 6-year field trial in the LRGV, the Bitters-C22 rootstock produced the highest yield over six other rootstocks, which included sour orange (Louzada et al., 2008). The Bitters-C22 rootstock was susceptible to *P. nicotianae* infection, whereas trifoliate orange was described as resistant after field and greenhouse trials (Graham, 1995). Federici et al. (2009) described

Table 3. Contingency table: level by rootstock

Count total %	C22	Sarawak \times Bower	Sarawak \times RioRed	Sour Orange	Total
Level 2	12 6.09	61 30.96	14 7.11	50 25.38	137 69.54
Level 3	16 8.12	19 9.64	6 3.05	19 9.64	60 30.46
Total	28 14.21	80 40.61	20 10.15	69 35.03	197
Tests					
N		df	-LogLike		R^2 (U)
197		3	5.2920186		0.0216
			χ^2		$P > \chi^2$
Likelihood ratio			10.584		0.0142*
Pearson			11.393		0.0098*

Table 4. Pairwise comparisons of proportions using Holm-Bonferroni adjustment method.

P value	C22	Sarawak \times Bower	Sarawak \times RioRed
Sarawak \times Bower	<0.0001		
Sarawak \times RioRed	0.3607	<0.0001	
Sour orange	0.1057	<0.0001	0.0048



Fig. 5. The sour orange rootstock plant derived from pregerminated seeds was actively growing at 18 months postinoculation with *Phytophthora nicotianae*. It had a strong root system with no signs of thinning or root rot.

Bitters-C22 as moderately tolerant to *P. nicotianae*, and therefore in this study, Bitters-C22 was expected to perform well. However, after the evaluation of rootstocks 11 months postinoculation in this study, Bitters-C22 did not perform well, and only 28 of 816 plants survived. Even though the survival rate was low, the surviving plants did have among the highest plant height and equivalent root length compared with the other three rootstocks in this study.

Sour orange is believed to be a hybrid of *C. reticulata* (mandarin) and *C. grandis* (pummelo) according to nursery stock evaluations and molecular marker analyses (Barrett and Rhodes, 1976; Nicolosi et al., 2000). Because sour orange is believed to be

a hybrid of these two species, similar qualities of sour orange hybrids could be achieved by crossing mandarins and pummelos (Grosser et al., 2004). Crosses between these two species have provided rootstocks with tolerance to *P. nicotianae* in greenhouse conditions as well as field conditions (Grosser et al., 2007). In this study, Sarawak pummelo \times Bower mandarin rootstock hybrid showed the highest seed survival rate with 96 plants at 11 months postinoculation. Plants exhibited leaf drop, stem yellowing, and fibrous root rot at lower degrees than the other rootstocks in this study. The root length and plant height were the highest of all rootstocks measured and had only moderate root damage. However, 18 months

postinoculation the tolerance of these plants was eventually overcome by *P. nicotianae* induced diseases. Pummelo does not present nucellar embryony, producing only monoembryonic zygotic seed; therefore, it should be expected that the many of its offspring would also be monoembryonic. However, there is still chance of obtaining hybrids that have some degree of polyembryony, such as sour orange, which is an offspring of the same kind of cross. If any of the tested hybrid shows high resistance to *Phytophthora* and is monoembryonic, propagation by tissue culture would be an option.

Grapefruit was described as susceptible (Hutchison and Grimm, 1972; Klotz, 1978) while pummelo was noted as resistant to *P. nicotianae* induced diseases (Ollitrault et al., 2006). Rio Red grapefruit provides high-quality fruit with visually appealing red flesh color, and field observation in Texas show Rio Red to have some tolerance to *Phytophthora* (unpublished data). Hybrid seeds of these two rootstocks were selected for *P. nicotianae* tolerance screening with the objective of combining some fruit favorable traits and develop more tolerance to *P. nicotianae*. In this study, Sarawak pummelo × Rio Red grapefruit seedlings showed the lowest survival rate of 24 plants total, which included both PGIS and NPGIS inoculated seed treatment groups. The NPGIS group did not have any surviving plants 11 months postinoculation. The surviving plants were on average the shortest plants recovered from the screening.

Sour orange was a candidate for screening against *P. nicotianae* diseases because it is the most commonly used rootstock in the LRGV. Seedlings variation in sour orange was first reported by Webber (1932) from experiments started in 1914 and conducted for more than 15 years. Xiang and Roose (1988) reported percentage of zygotic seeds greater than 30% in Taiwanica sour orange. Regardless of the percentage of zygotic seeds, sour orange has been the best ever rootstock for high pH calcareous soils and when high-quality scion production is desired. It seems that this does not cause enough variation to hinder this rootstock. Castle et al. (1993) described sour orange as tolerant to foot rot, cold temperatures, and capable of producing high-quality yields. Hutchison and Grimm (1972) had some success with sour orange clones and determined they were moderately resistant to *P. nicotianae* in Florida. However, when Graham (1995) tested the ability of root regeneration of sour orange, it was found to be intolerant to *P. nicotianae* infection. The difference in tolerance could be due to the amount of *P. nicotianae* inoculum present in the soil or the developmental stage of the seedlings. In this study, PGIS and NPGIS seeds were used for each rootstock, and both were exposed to high amounts of *P. nicotianae* inoculum. Eleven months postinoculation, sour orange had the second highest survival rating with more than 70 plants. Most plants did exhibit the typical *P. nicotianae* disease symptoms

and had the shortest root length and plant height compared with the other rootstocks. However, after an additional 7 months in the same *P. nicotianae* infested soil, the sole surviving rootstock was sour orange and it was visibly healthy. Aboveground the leaves were uniformly green with no yellowing, and below ground there were no signs of fibrous root decay. The sour orange plant shown to be highly tolerant/resistant to infection by *P. nicotianae*. This plant has continued growing for more than 2 years in heavily infested soil while not showing any symptoms as of this writing. The sour orange plant was propagated by cuttings and will be evaluated under field conditions.

To investigate the origin of *Phytophthora*-resistant sour orange (PRSO) plant, simple sequence repeat (SSR) marker analysis (Barkley et al., 2006; Kijas et al., 1997) was conducted using this plant and six sour orange lines (T1, T2, and T3 are standard sour orange ID: VI 95 from UCR Citrus Clonal Protection Program, T25 = Brazilian sour orange, T21 = standard sour orange, and T20 = Regular sour orange) that are currently used for the propagation of sour orange rootstock in the Texas A&M University–Kingsville Citrus Center. PRSO had two TAA1 alleles between 700 and 800 bp DNA size markers that are polymorphic among four parental lines except T20 and T2, both of which had the same two alleles (Supplemental Fig. 1). This may suggest that PRSO could be derived from a nucellar embryo of either of T20 and T2. However, the possibility of nucellar origin of PRSO from either of the lines were excluded because T20 and T2 had no or multiple CAGG9 alleles, whereas PRSO had single allele (Supplemental Fig. 1). Although SSR marker analysis failed to pinpoint two parental lines from which PRSO was originated, the data indicated that PRSO could be derived from a hybridization between T20 and T21 considering their physical proximity in the field.

Conclusion


In summary, among the four citrus rootstocks evaluated for *Phytophthora* tolerance, Sarawak × Bower hybrid has significantly performed well at 11 months postinoculation. A resistant sour orange plant was obtained that was the only plant to survive at 18 months postinoculation. This rootstock plant may offer a viable solution to *Phytophthora*-associated problems in citrus production under the Lower Rio Grande Valley area of South Texas.

Literature Cited

- Barrett, H.C. and A.M. Rhodes. 1976. A numerical taxonomic study of affinity relationships in cultivated citrus and its close relatives. *Syst. Bot.* 1(2):105–136.
- Barkley, N.A., M.L. Roose, R.R. Krueger, and C.T. Federici. 2006. Assessing genetic diversity and population structure in a citrus germplasm collection utilizing simple sequence repeat markers (SSRs). *Theor. Appl. Genet.* 112: 1519–1531.

- Bowman, K.D., J.P. Albano, and J.D. Graham. 2002. Greenhouse testing of rootstocks for resistance to *Phytophthora* species in flatwoods soil. *Proc. Fl. State Hort. Soc.* 115:10–13.
- Bowman, K.D., J.H. Graham, and R.C. Adair, Jr. 2003. Young tree growth in a flatwoods rootstock trial with *Diaprepes* weevil and *Phytophthora* diseases. *Proc. Fl. State Hort. Soc.* 116:249–251.
- Bright, D.B., J.H. Graham, M.S. Irey, and L.E. Baucum. 2004. Soil, rootstock, and climatic factors affect populations of *Phytophthora nicotianae* in south Florida citrus plantings. *Proc. Fl. State Hort. Soc.* 117:148–151.
- Cacciola, S.O. and G.M. di San Lio. 2008. Management of citrus diseases caused by *Phytophthora* spp., p. 61–84. In: A. Ciancio and K.G. Mukerji (eds.). *Integrated management of diseases caused by fungi, phytoplasma and bacteria. Integrated Management of Plant Pests and Diseases*, vol. 3. Springer, Dordrecht, the Netherlands.
- Castle, W.S., D.P.H. Tucker, A.H. Krezdorn, and C.O. Youtsey. 1993. *Rootstocks for Florida citrus* (Univ. Florida Publ. SP-42), 2nd ed. 15 May 2020. <https://crec.ifas.ufl.edu/extension/citrus_rootstock/Rootstock-Literature/Rootstocks%20for%20Florida%20Citrus.pdf>.
- Chaudhary, S. 2018. Incidence, distribution, and epidemiology of phytophthora diseases of citrus in south Texas and their interaction with Huanglongbing. Texas A&M University, College Station, PhD diss. 15 May 2020. <<http://hdl.handle.net/1969.1/174478>>.
- Colburn, G.C. and J.H. Graham. 2007. Protection of citrus rootstocks against *Phytophthora* spp. with a hypovirulent isolate of *Phytophthora nicotianae*. *Phytopathology* 97:958–963.
- Dhingra, O. and J.B. Sinclair. 1995. *Basic plant pathology methods*. 2nd ed. CRC Press, Boca Raton, FL.
- Fang, J.G. and P.H. Tsao. 1995. Efficacy of *Penicillium funiculosum* as a biological agent against *Phytophthora* rots of azalea and citrus. *Phytopathology* 85:871–878.
- Federici, C.T., R.S. Kupper, and M.L. Roose. 2009. ‘Bitters’, ‘Carpenter’ and ‘Furr’ trifoliolate hybrids: Three new citrus rootstocks. 15 May 2020. <<http://plantbiology.ucr.edu/faculty/new%20citrus%20rootstocks%202009.pdf>>.
- Graham, J.H. 1995. Root regeneration and tolerance of citrus rootstocks to root rot caused by *Phytophthora nicotianae*. *Phytopathology* 85: 111–117.
- Graham, J.H. and J.A. Menge. 1999. Root health: Fungal diseases, p. 126–135. In: L.W. Timmer and L.W. Duncan (eds.). *Citrus health management*. APS Press, St. Paul, MN.
- Graham, J.H. 1990. Evaluation of tolerance of citrus rootstocks to *Phytophthora* root rot in chlamydospore-infested soil. *Plant Dis.* 74: 743–746.
- Graham, J.H. and J.A. Menge. 2000. *Phytophthora*-induced diseases, p. 12–15. In: *Compendium of citrus diseases*. 2nd ed. APS Press, St. Paul, MN.
- Grosser, J.W., V.M. Urrutia, G. Ananthkrishnan, and P. Serrano. 2004. Building a replacement sour orange rootstock: Somatic hybridization of selected mandarin + pummelo combinations. *J. Amer. Soc. Hort. Sci.* 129:530–534.
- Grosser, J.W., J.H. Graham, A. Hoyte, H.M. Rubio, D.B. Bright, J. Gmitter, C.X. Chen, and F.G. Gmitter, Jr. 2007. Continued development of rootstocks tolerant of the *Diaprepes/Phytophthora* complex via greenhouse screening. *Proc. Annu. Meet. Fla. State Hort. Soc.* 120:103–109.

- Graham, J. and E. Feichtenberger. 2015. Citrus *Phytophthora* diseases: Management challenges and successes. *J. Citrus Pathol.* 2:1–11.
- Grote, D., A. Olmos, A. Kofoet, J.J. Tuset, E. Bertolini, and M. Cambra. 2002. Specific and sensitive detection of *Phytophthora nicotianae* by simple and nested-PCR. *Eur. J. Plant Pathol.* 108:197–207.
- Hung, P.M., P. Wattanachai, S. Kasem, and S. Poeaim. 2015. Efficacy of *Chaetomium* species as biological control agents against *Phytophthora nicotianae* root rot in Citrus. *Mycobiology* 43:288–296.
- Hutchison, D.J. and G.R. Grimm. 1972. Variation in *Phytophthora* resistance of Florida rough lemon and sour orange clones. *Proc. Annu. Meet. Fla. State Hort. Soc.* 85:38–39.
- Jeffers, S.N. 2006. Identifying species of *Phytophthora*. Department of Entomology, Clemson University, Clemson, SC. 15 May 2020. <https://www.fs.fed.us/foresthealth/htm/sp/sod/misc/culturing_species_phytophthora.pdf>.
- Kijas, J.M.H., M.R. Thomas, J.C.S. Fowler, and M.L. Roose. 1997. Integration of trinucleotide microsatellites into a linkage map of citrus. *Theor. Appl. Genet.* 94:701–706.
- Klotz, L.J. 1978. Fungal, bacterial and non-parasitic diseases and injuries originating in the seedbed, nursery and orchard, p. 1–66. In: *The citrus industry: Vol. 4. Crop protection.* W. Reuther, E.C. Calavan, and G.F. Carman (eds.). University of California Agricultural Sciences Publications, Richmond, CA.
- Kunta, M., J.V. da Graça, and M. Skaria. 2007. Molecular detection and prevalence of citrus viroids in Texas. *HortScience* 42:600–604.
- Lamour, K.H., M.L. Daughtrey, D.M. Benson, J. Hwangand, and M.K. Hausbeck. 2003. Etiology of *Phytophthora drechsleri* and *P. nicotianae* (= *P. parasitica*) diseases affecting floriculture crops. *Plant Dis.* 87:854–858.
- Lamine, M. and A. Mliki. 2015. Elucidating genetic diversity among sour orange rootstocks: A comparative study of the efficiency of RAPD and SSR markers. *Appl. Biochem. Biotechnol.* 175:2996–3013.
- Louzada, E.S., H.S. del Rio, M. Sétamou, J.W. Watson, and D.M. Swietlik. 2008. Evaluation of citrus rootstocks for the high pH, calcareous soils of South Texas. *Euphytica* 164:13–18.
- Ludowici, V.A., W. Zhang, L.M. Blackman, and A.R. Hardham. 2013. *Phytophthora nicotianae*, p. 113–123. In: K. Lamour (ed.). *Phytophthora— a global perspective.* CAB International, Wallingford, Oxfordshire, UK.
- Meng, Y., Q. Zhang, W. Ding, and W. Shan. 2014. *Phytophthora parasitica*: A model oomycete plant pathogen. *Mycology* 5:43–51.
- Naqvi, S.A.M.H. 2004. Diagnosis and management of certain important fungal diseases of citrus, p. 247–290. In: S.A.M.H. Naqvi (ed.). *Diseases of fruits and vegetables: Vol. I. Diagnosis and management.* Kluwer Academic, Dordrecht, the Netherlands.
- Nicolosi, E., Z.N. Deng, A. Gentile, S. La Malfa, G. Continella, and E. Tribulato. 2000. Citrus phylogeny and genetic origin of important species as investigated by molecular markers. *Theor. Appl. Genet.* 100:1155–1166.
- Ollitrault, P., Y. Froelicher, D. Dambier, F. Luro, and M. Yamamoto. 2006. Seedlessness and ploidy manipulations. In: I.A. Khan (ed.). *Citrus genetics, breeding and biotechnology.* CAB International, Wallingford, UK; Cambridge, MA.
- Savita, G.S. and A. Nagpal. 2012. Citrus diseases caused by *Phytophthora* species. *GERF Bull. Biosci.* 3:18–27.
- Tashiro, N., S. Uematsu, and Y. Ide. 2012. First report of *Phytophthora palmivora* as a causal pathogen of citrus brown rot in Japan. *J. Gen. Plant Pathol.* 78:233–236.
- Timmer, L.W., H.A. Sandler, J.H. Graham, and S.E. Zitko. 1988. Sampling citrus orchards in Florida to estimate populations of *Phytophthora parasitica*. *Phytopathology* 78:940–944.
- Timmer, L.W., J.H. Graham, and S.E. Zitko. 1998. Metalaxyl-resistant isolates of *Phytophthora nicotianae*: Occurrence, sensitivity, and competitive parasitic ability on citrus. *Plant Dis.* 82:254–261.
- Webber, H.J. 1932. Variation in citrus seedlings and their relation to rootstock selection. *Hilgardia* 7:1–77.
- Xiang, C. and M.L. Roose. 1988. Frequency and characteristics of nucellar and zygotic seedlings of rootstocks. *Scientia Hort.* 37:47–59.
- Zitko, S.E. and L.W. Timmer. 1994. Competitive parasitic abilities of *Phytophthora parasitica* and *P. palmivora* on fibrous roots of citrus. *Phytopathology* 84:1000–1004.



Supplemental Fig. 1. Comparison of simple sequence repeat (SSR) marker profiles between *Phytophthora*-resistant sour orange (PSRO) and sour orange parental lines. Marker loci are indicated on the bottom. > indicated polymorphic markers. T1, T2, and T3 are standard sour orange ID: VI 95 from UCR Citrus Clonal Protection Program, T25 = Brazilian sour orange, T21 = standard sour orange, and T20 = Regular sour orange.