

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

5,000

Open access books available

125,000

International authors and editors

140M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.

For more information visit www.intechopen.com



Application and Mechanisms of Plant Growth Promoting Fungi (PGPF) for Phytostimulation

Md. Motaher Hossain and Farjana Sultana

Abstract

Plant growth-promoting fungi (PGPF) constitute diverse genera of nonpathogenic fungi that provide a variety of benefits to their host plants. PGPF show an effective role in sustainable agriculture. Meeting increasing demand for crop production without damage to the environment is the biggest challenge nowadays. The use of PGPF has been recognized as an environmentally friendly way of increasing crop production. These fungi have proven to increase crop yields by improving germination, seedling vigor, plant growth, root morphogenesis, photosynthesis, and flowering through either a direct or indirect mechanism. The mechanisms of PGPF involve solubilizing and mineralizing nutrients for easy uptake by plants, regulating hormonal balance, producing volatile organic compounds and microbial enzyme, suppressing plant pathogens and ameliorating abiotic stresses. Successful colonization is an intrinsic factor for most PGPF to exert their beneficial effects on plants. A certain level of specificity exists in the interactions between plant species and PGPF for root colonization and growth promoting effects. There is a gap between the number of reported efficacious PGPF and the number of PGPF as biofertilizer. Efforts should be strengthened to improve the efficacy and commercialization of PGPF. Hence, this chapter summarizes valuable information regarding the application and mechanisms of PGPF in sustainable agriculture.

Keywords: seed germination, seedling vigor, root morphogenesis, yield, root colonization, formulation

1. Introduction

The world's population exceeded ~7 billion just after 2010, and still continues to grow fast. Roughly, 83 million people are added to the world's population every year and with this pace of growth, the global population is projected to reach around 9.7 billion by 2050, ~24% higher than today [1]. In order to feed this large population, crop production must increase by approximately 25–70% above current production levels [2]. Intensification of agriculture is considered a potential solution. By relying on intensive use of fertilizers, pesticides and other inputs, agricultural intensification increases the productivity of existing farmland and delivers more food to the added population. However, the chemical-based crop

intensification produces more food in a way that the future production potential of farmland is being undermined and the environment is being affected. An increasingly degraded soil, overwhelming health hazards from soil and water pollution, disturbed natural microbial populations are a few of the direct implications in chemical-intensive agriculture. To avoid these potentially harmful effects of agrochemicals in agriculture, alternative approaches must be persuaded. An ecocentric approach that provides both environmental and economic benefits is increasingly needed. Organic farming is one of many such approaches that promote agroecosystem health, ensuring sustainable intensification in agriculture.

The uniqueness of microorganisms and the dynamic part played by them in sustaining agricultural ecosystems have made them likely candidates for playing a central role in organic-based modern agriculture. Fortunately, plant roots harbor an abundant association of beneficial microorganisms. Root exudates are the largest source of carbon that attracts the microbial populations and allow them to forge an intimate association with host plants [3]. In response, the rhizosphere microbial populations play versatile roles in transforming, mobilizing and solubilizing soil nutrients, which are crucial for plant growth and development. Among the diverse rhizosphere microbial population, fungi known as plant growth promoting fungi (PGPF) are receiving a growing attention in recent days. Over the decades, varieties of PGPF have been studied including those belong to genera *Trichoderma*, *Penicillium*, *Phoma* and *Fusarium* [4]. Studies have shown that PGPF modulate plant growth and enhance resilience to plant pathogens without environmental contamination [5]. The positive effects of PGPF on plant and environment make them well fitted to organic agriculture.

The course of plant growth promotion by PGPF is a complex process and often cannot be attributed to a single mechanism. A variety of direct and indirect mechanisms, including solubilization of minerals, synthesis of phytohormones, production of volatile organic compounds, exploitation of microbial enzymes, increases in nutrient uptake, amelioration of abiotic stresses and suppression of deleterious phytopathogens are involved. These wide arrays of interconnected mechanisms help PGPF maintaining rhizosphere competence and stability in host performance. Compared to the large number of PGPF identified in the laboratory, only a small fraction of them is in agricultural practice worldwide. Inconsistent performance of the inoculated PGPF under field conditions limits the commercial application of them. Development of appropriate formulation could improve the performance in the field and pave the way for commercialization of the PGPF. An ideal formulation of PGPF should fit with existing application technologies, protect biological actives from stress, ensure viability, remains unaffected after storage under ambient conditions, ensure microbial actives in the field and be cost effective [6].

Considering the aspects discussed above, the need for superior PGPF to supplement inorganic chemical fertilizers as one of the crucial steps of moving toward organic farming practices has been highlighted. Inclusion of new techniques in these processes has been vital to the development of novel PGPF applications. This review will therefore attempt to shed light on the recent findings related to the impact of PGPF on plant growth and yield, duration of their effects, host specificity of the cooperation, root colonization mechanisms, their modes of action and commercial formulation for enhancement of plant growth and yield. The knowledge produced from this review could be very useful to those who are apprehensive about environmental protection and agricultural sustainability.

2. Plant growth promoting fungi (PGPF)

Plants have intricate relationships with an array of microorganisms, particularly rhizosphere fungi and bacteria, which can lead to an increase in plant vigor, growth and development as well as changes in plant metabolism [7]. The group of rhizosphere fungi that colonize plant roots and enhance plant growth is referred to as PGPF [4]. PGPF are heterogeneous group of nonpathogenic saprotroph fungi. They can be separated into endophytic, whereby they live inside roots and exchange metabolites with plants directly, and epiphytic, whereby they live freely on the root




Figure 1. Beneficial interaction between plant and plant growth promoting fungi (PGPF). PGPF can modulate plant growth and development through the production of phytohormones and volatile compounds. PGPF also influence plant nutrition via solubilization of phosphorus and mineralization of organic substrates. PGPF modify plant functioning against biotic and abiotic stresses by negating their harmful effects.

surface and free-living PGPF, which live outside plant cells, i.e., in the rhizosphere [5]. PGPF establish a non-obligate mutualism with a broader range of host plants. That is why symbiotic mycorrhizal fungi are not considered as PGPF, although they are known to improve growth of the plants [8]. Moreover, PGPF encompass a diverse taxonomic group in comparison to mycorrhiza. They are often involved in a range of complex interactions with plants and develop distinct strategies to mediate improvements in seed germination, seedling vigor, plant growth, flowering and productivity of host plants (**Figure 1**). PGPF are not only associated with the root to mediate positive effects on plant growth and development but also have beneficial effects on suppressing phytopathogenic microorganisms [9]. Not every organism identified as PGPF will improve plant growth under all conditions or in association with all plant hosts [10]. Some PGPF biocontrol inoculants usually contain necrotrophic mycoparasites such as *Trichoderma* spp. [11], while a limited number such as *Sphaerodes mycoparasitica* is biotrophic mycoparasitic agent [12]. Therefore, PGPF are considered one of the potential active ingredients in both biofertilizer and mycofungicide formulation.

3. The nature and composition of PGPF

PGPF are common root-associated and soil-borne fungi from diverse genera. Fungi reported as PGPF include Ascomycetes, Basidiomycetes and Oomycetes [5]. Some strains of hypovirulent binucleate *Rhizoctonia* (HBNR) are known to be PGPF [13]. PGPF also include isolates of mycelial fungi that do not produce any spores, generally known as sterile black fungus (SBF), sterile dark fungus (SDF) and sterile red fungus (SRF) [14]. The non-sporulating PGPF are often difficult to identify and mostly lack formal taxonomic status. Among the PGPF *Aspergillus*, *Fusarium*, *Penicillium*, *Phoma* and *Trichoderma* have a wide distribution and are, by far, the most extensively reported (**Table 1**). Each of the genera has a variety of species. *Aspergillus*, *Fusarium*, *Penicillium* and *Phoma* were frequently found in the rhizosphere or in the roots of plants. Instead, *Trichoderma* were mostly isolated from soil. Among the rhizosphere population, PGPF have a high relative abundance. A total of 619 (44%) out of 1399 fungal isolates collected from rhizosphere of six different plants were PGPF, while frequency of occurrence of PGPF in zoysiagrass, wheat, corn and eggplant rhizosphere were 46, 47, 38 and 10%, respectively [4]. This indicates that abundance of PGPF varies largely according to the host rhizosphere. Similarly, the dominating fungal genus is not necessarily the dominating PGPF in the rhizosphere population. The order of the frequency of the main genera among 1399 fungal isolates was *Fusarium* > *Trichoderma* > sterile fungi > *Penicillium* > *Pythium* > *Rhizoctonia* > *Mucor*, while that of PGPF from each plant genus was: *Trichoderma* (~82%) > *Pythium* (~75%) > *Penicillium* (~69%) > *Alternaria* (~63%) > *Fusarium* (~44%) > sterile fungi (40%) > *Mucor* (~38%) [4]. The important characteristics of these fungi are their high rhizosphere competence and ability to promote plant growth.

Initial search for identification of PGPF was concentrated to rhizosphere fungi. Recent studies have demonstrated the potential of phyllosphere fungi as PGPF. The phyllosphere, which consists of the above ground surfaces of plants, is one of the most prevalent microbial habitats on earth. Phyllosphere fungi can act as mutualists promoting plant growth and tolerance of environmental stressors [53]. A few of other fungi isolated from tree bark, decorticated wood and water damaged building functioned as PGPF [43, 49]. More interestingly, the fungal entomopathogens also show potential to be PGPF and promote plant growth [54]. PGPF seem to have a cosmopolitan occurrence.

PGPF	Original source of isolation	References
<i>Alternaria</i> sp.	<i>Zoysia tenuifolia</i> , <i>Rosa rugosa</i> , <i>Camellia japonica</i> , <i>Delonix regia</i> , <i>Dianthus caryophyllus</i> , <i>Rosa hybrid</i>	[4, 15]
<i>Aspergillus</i> sp., <i>As. fumigatus</i> , <i>As. niger</i> , <i>As. terreus</i> , <i>As. ustus</i> , <i>As. clavatus</i>	<i>Capsicum annuum</i> , <i>Glycine max</i> , <i>Cicer arietinum</i> , <i>Elymus mollis</i> , <i>Solanum tuberosum</i> , <i>Nymphoides peltata</i>	[16–21]
<i>Aureobasidium pullulans</i>	Dark chestnut soil	[22]
<i>Chaetomium globosum</i>	<i>Capsicum annuum</i>	[23]
<i>Cladosporium</i> sp., <i>Cladosporium sphaerospermum</i>	<i>Cucumis sativus</i> , <i>Glycine max</i>	[24, 25]
<i>Colletotrichum</i> sp.	<i>Rosa rugosa</i> , <i>Camellia japonica</i> , <i>Delonix regia</i> , <i>Dianthus caryophyllus</i> , <i>Rosa hybrid</i>	[15]
<i>Exophiala</i> sp.	<i>Cucumis sativus</i>	[26]
<i>Fusarium</i> sp., <i>F. equiseti</i> , <i>F. oxysporum</i> , <i>F. verticillioides</i>	<i>Cynodon dactylon</i> , <i>Lygeum spartum</i> , <i>Zoysia tenuifolia</i> , <i>Musa</i> sp. and other environment	[27–32]
Non-sporulating sterile fungi	<i>Zoysia tenuifolia</i>	[14]
<i>Penicillium</i> sp., <i>Pe. chrysogenum</i> , <i>Pe. citrinum</i> , <i>Pe. kloeckeri</i> , <i>Pe. menonorum</i> , <i>Pe. resedanum</i> , <i>Pe. simplicissimum</i> , <i>Pe. janthinellum</i> , <i>Pe. viridicatum</i>	Halophyte, <i>Ixeris repenes</i> , <i>Cicer arietinum</i> , <i>Elymus mollis</i> , <i>Capsicum annuum</i> , <i>Zoysia tenuifolia</i>	[9, 16, 22, 33–40]
<i>Phoma</i> sp., <i>Phoma herbarum</i> , <i>Phoma multirostrata</i>	<i>G. max</i> , <i>Rosa rugosa</i> , <i>Camellia japonica</i> , <i>Delonix regia</i> , <i>Dianthus caryophyllus</i> , <i>Rosa hybrid</i> , <i>Zoysia tenuifolia</i>	[4, 14, 15, 34, 41, 42]
<i>Phomopsis</i> sp., <i>Phomopsis liquidambari</i>	<i>Rosa rugosa</i> , <i>Camellia japonica</i> , <i>Delonix regia</i> , <i>Dianthus caryophyllus</i> , <i>Rosa hybrid</i> , <i>Bischofia polycarpa</i> bark	[15, 43]
<i>Purpureocillium lilacinum</i>	Soil	[44]
<i>Rhizoctonia</i> spp.	Orchid, <i>Lycopersicon lycopersicum</i> , and soil	[13, 45, 46]
<i>Rhodotorula mucilaginosa</i>	Soil	[22]
<i>Talaromyces wortmannii</i>	Soil	[40]
<i>Trichoderma asperellum</i> , <i>T. atroviride</i> , <i>T. hamatum</i> , <i>T. harzianum</i> , <i>T. longibrachiatum</i> , <i>T. pseudokoningii</i> , <i>T. viride</i> , <i>T. virens</i>	Soil, wood and damaged building	[34, 47–52]

Table 1. Different fungi reported as plant growth promoting fungi (PGPF) with their original source of isolation.

4. Impact of PGPF on plant growth promotion

PGPF exhibit traits beneficial to plant and as such, their capacity to enhance plant growth and development is well founded. PGPF mediate both short- and long-term effects on germination and subsequent plant performance. Improvement in germination, seedling vigor, shoot growth, root growth, photosynthetic efficiency, flowering, and yield are the most common effects decreed by PGPF. A particular PGPF may condition plant growth by exerting all or one or more of these effects.

4.1 Impact of PGPF on seed germination and seedling vigor

Seed germination and germinant growth are critical developmental periods of the young plantlet until it begins producing its own food by photosynthesis. Treatment with PGPF, particularly of the genus *Aspergillus*, *Alternaria*, *Trichoderma*, *Penicillium*, *Fusarium*, *Sphaerodes* and *Phoma* has been reported to improve seed germination and seedling vigor in different agronomic and horticultural crops (Table 2). Scarified seeds inoculated with spores from *Aspergillus* and *Alternaria* had significant increases in germination of Utah milkvetch (*Astragalus utahensis*) *in vitro*, and in greenhouse and fall-seeded plots near Fountain Green and Nephi [55]. The *Aspergillus*-treated seeds performed out seeds inoculated with *Alternaria*. An increase of 30% in seedling emergence was observed in cucumber plant raised upon the treatment of *T. harzianum* [47]. Application of *T. harzianum* also significantly increased seed germination, emergence index, seedling vigor and successful transplantation percentage in muskmelon compared to the untreated controls [59]. Early seedling emergence and enhanced vigor were observed in bacterial wilt susceptible tomato cultivar treated with *T. harzianum*, *Phoma multirostrata*, and *Penicillium chrysogenum* compared to untreated controls [34]. The culture filtrate of *Penicillium* was as effective as the living inocula in improving seed germination of tomato [70]. Significantly, higher germination and vigor index were observed in Indian spinach, when seeds were sown in sterilized field soil amended with wheat grain inoculum of *Fusarium* spp. PPF1 [27]. *Sphaerodes mycoparasitica*, a biotrophic mycoparasite of *Fusarium* species, improved wheat seed germination and seedling growth *in vitro* compared to *T. harzianum*, while under phytotron conditions, both *S. mycoparasitica* and *T. harzianum* had positive impact on wheat seedlings growth in the presence of *F. graminearum* [12]. These results show the positive impact of PGPF on seed germination and seedlings growth of a wide arrays of hosts.

4.2 Impact of PGPF on shoot growth

The most common form of growth promotion by PGPF is the augmented shoot in colonized plants. Shoot growth promotion has been shown by a great diversity of PGPF across a large number of plant species. Isolates of *Aspergillus*, *Trichoderma*, *Penicillium*, and *Fusarium* were capable of enhancing the shoot growth in model plant *Arabidopsis* [9, 20, 28, 33, 48]. Different species of *Aspergillus* are known to support shoot growth in chickpea [16], Chinese cabbage [56], cucumber [17], soybean [18, 65] and wheat [76]. Species of nonpathogenic *Fusarium* were reported to stimulate shoot growth in Indian spinach [27] and banana [29]. Application of barley grain inoculum of *Penicillium viridicatum* GP15-1 to the potting medium resulted in 26–42% increase in stem length, 37–46% increase in shoot fresh weight and 100–176% increase in shoot dry weight of cucumber plants [35]. Similarly, inoculation of cucumber plants with *Pe. menonorum* KNU3 increased cucumber shoot dry biomass by as much as 52% [36]. Stimulated shoot growth by *Penicillium* spp. was also reported in tomato [69], Waito-c rice [37, 38], chili [23, 39] and sesame [74]. Application of *T. longipile* and *T. tomentosum* increased shoot dry weight of cabbage seedlings by 91–102% in glasshouse trials [57]. Likewise, cottonseeds pretreated with *T. viride* showed four-fold increases in shoot length elongation and an almost 40-fold increase in plant dry weight compared to the control [66]. Augmented shoot growth by *Trichoderma* has also been reported in chickpea [16], wheat [79], maize [78], cucumber [60] and other plant species (Table 2). Isolates of *Phoma* were found to be an efficient stimulator of plant shoot [15, 41, 62]. A few hypovirulent *Rhizoctonia* isolates were able to induce significantly higher fresh leaves and stems weights in tomato plants grown in greenhouse [13]. Enhancement of shoot growth was also observed

Test crop	PGPF strain	Improvement	References
<i>Arabidopsis thaliana</i>	<i>Trichoderma virens</i> Gv. 29-8	Biomass, lateral root development	[48]
	<i>Penicillium janthinellum</i> GP16-2	Shoot biomass, leaf number	[33]
	<i>Pe. simplicissimum</i> GP17-2	Shoot biomass, leaf number	[9]
	<i>Fusarium oxysporum</i> NRRL 38499, NRRL 26379 and NRRL 38335,	Shoot-root growth	[28]
	<i>Aspergillus ustus</i>	Shoot growth, lateral root, root hair numbers	[20]
<i>Astragalus utahensis</i>	<i>Aspergillus</i> spp., <i>Alternaria</i> spp.	Seed germination	[55]
<i>Basella alba</i>	<i>Fusarium</i> spp. PPF1	Germination, seedling vigor, shoot-root growth, leaf area, leaf chlorophyll content	[27]
<i>Brassica campestris</i>	<i>Talaromyces wortmannii</i> FS2	Shoot fresh weight	[40]
<i>B. chinensis</i>	<i>A. niger</i> 1B and 6A	Plant dry weight, N and P content	[56]
<i>B. oleracea</i> var. <i>capitata</i>	<i>T. longipile</i> , <i>T. tomentosum</i>	Shoot dry weight, leaf area	[57]
<i>Capsicum annuum</i>	<i>Pe. resedanum</i> LK6	Shoot length, biomass, chlorophyll content, photosynthesis	[39]
	<i>Chaetomium globosum</i> CAC-1G	Plant biomass, root-shoot growth	[23]
<i>Cicer arietinum</i>	<i>A. niger</i> BHUAS01, <i>Pe. citrinum</i> BHUPC01, <i>T. harzianum</i>	Plant growth	[16]
	<i>T. harzianum</i> T-75	Yield	[58]
<i>Cucumis melo</i>	<i>T. harzianum</i> Bi	Germination, seedling health, vigor	[59]
<i>Cucumis sativus</i>	<i>Pe. simplicissimum</i> GP17-2	Root-shoot growth	[4]
	<i>Pe. viridicatum</i> GP15-1	Root-shoot length, biomass	[35]
	<i>T. harzianum</i> GT3-2	Root-shoot growth	[60]
	<i>F. equiseti</i> GF19-1	Root-shoot growth	[61]
	<i>Aspergillus</i> spp. PPA1	Root-shoot length, biomass, leaf area, chlorophyll content	[17]
	<i>Exophiala</i> sp. LHL08	Plant growth under drought and salinity	[26]
	<i>Phoma</i> sp.	Root-shoot growth, yield in the field	[62]
GiSeLa6® (<i>Prunus cerasus</i> × <i>P. canescens</i>)	<i>Phoma</i> sp. GS8-2, GS8-3	Root-shoot growth	[63]
	<i>T. harzianum</i> T-22	Root growth, development	[64]
<i>Glycine max</i>	<i>A. fumigatus</i> HK-5-2	Shoot growth, biomass, leaf area, chlorophyll contents, photosynthetic rate	[65]
	<i>A. fumigatus</i> LH02	Shoot growth, biomass, leaf area, chlorophyll contents, photosynthetic rate	[18]
	<i>Phoma herbarum</i> TK-2-4	Plant length, biomass	[41]
<i>Gossypium arboreum</i> L	<i>T. viride</i>	Root-shoot length, plant dry weight	[66]

Test crop	PGPF strain	Improvement	References
<i>Helianthus annuus</i>	<i>Trichoderma</i> sp., <i>Aspergillus</i> sp., <i>Penicillium</i> sp., <i>Phoma</i> sp., <i>Fusarium</i> sp.	Seed germination, seedling vigor	[67]
<i>Lactuca sativus</i>	<i>F. oxysporum</i> MSA 35	Root-shoot growth, chlorophyll content	[68]
<i>Lycopersicon lycopersicum</i>	<i>T. harzianum</i> TriH_JSB27, <i>Phoma multirostrata</i> PhoM_JSB17, <i>T. harzianum</i> TriH_JSB36, <i>Pe. chrysogenum</i> PenC_JSB41	Seedling emergence, vigor	[34]
	<i>T. harzianum</i> T-22	Seed germination under stress	[69]
	<i>Penicillium</i> spp.	Seed germination, root-shoot growth	[70]
	<i>F. equiseti</i> GF19-1	Plant biomass, root-shoot growth	[71]
<i>Musa</i> sp.	<i>F. oxysporum</i> V5W2, Eny 7.11o, Emb 2.4o	Yield	[29]
<i>Nicotiana tabacum</i>	<i>Alternaria</i> sp., <i>Phomopsis</i> sp., <i>Cladosporium</i> sp., <i>Colletotrichum</i> sp., <i>Phoma</i> sp.	Root-shoot growth, chlorophylls, soluble sugars, plant biomass	[15]
<i>Pinus sylvestris</i> var. <i>mongolica</i>	<i>T. harzianum</i> E15, <i>T. virens</i> ZT05	Seedling biomass, root structure, soil nutrients, soil enzyme activity	[72]
<i>Saccharum officinarum</i>	<i>T. viride</i>	Yield	[73]
<i>Sesamum indicum</i>	<i>Penicillium</i> spp. NICS01, DFC01	Root-shoot growth, chlorophylls, proteins, amino acids, lignans	[74]
<i>Solanum tuberosum</i>	<i>A. ustus</i>	Root-shoot growth, lateral root, root hair numbers	[20]
<i>Spinacia oleracea</i>	<i>F. equiseti</i>	Plant biomass, root-shoot growth	[75]
<i>Suaeda japonica</i>	<i>Penicillium</i> sp. Sj-2-2	Plant length	[38]
	<i>Cladosporium</i> sp. MH-6	Shoot length	[24]
	<i>Pe. citrinum</i> IR-3-3	Root-shoot length	[37]
	<i>Phoma herbarum</i> TK-2-4	Plant length	[41]
<i>Triticum aestivum</i>	<i>T. harzianum</i> , <i>T. koningii</i>	Plant biomass, root-shoot growth.	[4]
	<i>Sphaerodes mycoparasitica</i>	Seed germination, seedling growth	[12]
	<i>A. niger</i> NCIM	Shoot and total plant length ratio	[76]
<i>Vinca minor</i>	<i>T. harzianum</i>	Flowering, plant height, weight	[77]
<i>Zea mays</i>	<i>T. harzianum</i> T22	Shoot growth, area and size of main and secondary roots	[78]

Table 2. Effect of different plant growth promoting fungi (PGPF) on seed germination, plant growth and yield in various plants.

by *Talaromyces wortmannii* in cabbage [40], *Chaetomium globosum* in chili [23], *Colletotrichum* sp. in tobacco and *Exophiala* sp. in cucumber [26]. The results from these studies are consistent with numerous field and growth chamber experiments that have shown that PGPF inoculants can mediate shoot growth improvement.

4.3 Impact of PGPF on photosynthesis

The plant growth promotion in some plant-PGPF interaction is occasionally associated with improvement in state and function of the photosynthetic apparatus of plants. Treatment with *T. longipile* and *T. tomentosum* increased leaf area of cabbage by 58–71% in glasshouse trials [57]. Tomato plants grown with HBNR isolates had significantly higher leaf fresh weight than control plants in greenhouse [13]. Arabidopsis grown in soil amended with *Pe. simplicissimum* GP17-2 and *Pe. janthinellum* GP16-2 were more greener and had approximately 1 more leaflet per plant than control plants 4 weeks after treatment [9]. *Penicillium* spp. also enhanced leaf chlorophyll content in cucumber and chili [36, 39]. Soil amendment with *Aspergillus* spp. PPA1 and *Fusarium* spp. PPF1 significantly increased leaf area and leaf chlorophyll content in cucumber and Indian spinach, respectively [27]. Improvement in leaf number, leaf area and leaf chlorophyll levels would contribute to increases in photosynthesis rate and net accumulation of carbohydrate in plants.

4.4 Impact of PGPF on root growth and architecture

Roots are vital plant organs that remain below the surface of the soil. The root system is important for plant fitness because it facilitates the absorption of water and nutrients, provides anchorage of the plant body to the ground and contributes to overall growth of plants. Root functions as the major interface between the plant and the microbes in the soil environment. The bulk of previous studies have evidenced the immense ability of PGPF in enhancement of root growth in different plants (Table 2). Plants forming association with PGPF show faster and larger root growth resulting in a rapid increase in the root biomass [27, 35, 50, 57]. Moreover, root length, root surface area, root diameter and branch number are under direct influence of intimate interaction with PGPF. Application of *T. vires* ZT05 increased root length, root surface area, average root diameter, root tip number and root branch number of pines by 25.11, 98.19, 5.66, 45.89 and 74.42%, respectively [72]. *A. ustus* is known to cause alterations in the root system architecture by promoting the formation of secondary roots in Arabidopsis and potato [20]. In maize (*Zea mays*), *Trichoderma* inoculation enhanced root biomass production and increased root hair development [78]. The abundance in root hair formation significantly increases root surface area, suggesting that PGPF inoculants could enhance the potential for plant roots to acquire nutrients under nutrient-limited conditions.

4.5 Impact of PGPF on flowering

The application of PGPF may influence the number, size and timing of flower in flowering plants. *Tagetes* (marigolds) grown with companion of *Pe. simplicissimum* flowered earlier and had greater flower size and weight [80]. Steamed or raw soil infested with *T. harzianum* hastened flowering of periwinkle and increased the number of blooms per plant on chrysanthemums [77]. Under greenhouse conditions, *T. harzianum* TriH_JSB27 and *Pe. Chrysogenum* PenC_JSB41 accelerated the flowering time in tomato [34]. Similarly, root colonization by the nematophagous fungus *Pochonia chlamydosporia* hurried flowering in *Arabidopsis thaliana* [81]. Root colonization by *Piriformospora indica* also results in early flowering in *Coleus forskohlii*, bottle gourd and *Nicotiana tabacum* [82]. Flowering time has commercial significance for crops and ornamental plants by shortening crop duration and improving productivity. A short duration crop would have several advantages over a long duration crop, even with equal total yields such as require less water, expose less to stresses and

increase the availability of the land for subsequent cropping. This indicates that PGPF improve the plasticity of complex plant traits.

4.6 Impact of PGPF on yield

PGPF show promising ability to promote growth through extensive improvements and betterment of fundamental processes operating in the plants, all of which directly and indirectly contributes to the crop yield increase. Inoculation of banana (cv. Giant Cavendish and Grand Nain) with *F. oxysporum* resulted in 20–36% yield increase in the field [29]. Soil treatment with *T. harzianum* alone or in combination with organic amendment and fungicide significantly improved seed yield in pea [83] and chickpea [58]. Similarly, soil treatment with *T. viride* produced significantly the highest number of fruits per plant, number of seeds per fruit, fruit weight and dry weight of 100 seeds as compared to untreated control [84]. The beneficial association of plants with nonpathogenic binucleate *Rhizoctonia* spp. resulted in increase in yield of carrot, lettuce, cucumber, cotton, radish, wheat, tomato, Chinese mustard and potato [13, 45, 46]. These results demonstrate that PGPF hold great promise in the improvement of agriculture yields.

5. Duration of sustained plant growth promotion effect by PGPF

The duration of biofunctional activities of PGPF in plants is a key factor for their effective application in the field. Naturally, a legitimate question may arise whether PGPF isolates that have shown promising effects on early growth stage of plants, could also affect the middle or late ontogenetic stages and ultimately contribute to yield increases at harvest. As for potato, an increase in leaf, shoot, and tuber weight was observed by a nonpathogenic isolate (No. 521, AG-4) of *Rh. solani* 63–70 days after planting, while it was not expressed in yield at harvest [85]. Conversely, increased growth responses of wheat plants treated with PGPF were observed during seedling (2 weeks after sowing), vegetative (4 weeks), pre-flowering (6 weeks), flowering (10 weeks) and seed maturation stages (14 weeks) [4]. The isolates of *Phoma* sp. (GS6-1, GS7-4) and non-sporulating fungus (GU23-3), increased plant height, ear-head length and weight, seed number and plant biomass at harvest [79]. Again, isolates of *Phoma* sp. and non-sporulating fungus significantly increased plant length, dry biomass, leaf number and fruit number of cucumber cv. Jibai until 10 weeks post planting in greenhouse trials [62]. These isolates were equally effective in promoting growth and increasing yield of cucumber at 6 and 10 weeks post planting in the field [62]. There are other PGPF, which as well have shown the ability to confer long-term growth benefits to different plants. Rice and pea plants inoculated with *Westerdykella aurantiaca* FNBR-3, *T. longibrachiatum* FNBR-6, *Lasiodiplodia* sp. FNBR-13 and *Rhizopus delemar* FNBR-19 showed a stimulatory increase of growth for 8 weeks in the greenhouse [86]. Similarly, a single inoculation with inoculum of *Penicillium* and *Pochonia* affected the whole life cycle of tomato and Arabidopsis, respectively, accelerating the growth rate, shortening their vegetative period and enhancing seed maturation [34, 81]. As such, majority of PGPF strains are able to induce sustained beneficial effects on plant growth. The basis of sustained effects of PGPF on plants is not fully understood. One possibility is that the fungus continues to colonize the root system and establishes a life-long colonization with crop roots. The ability of PGPF to confer sustained benefit to plant is of great agriculture importance in terms of improving crop yield.

6. Host specificity of the plant growth-promoting cooperation

Although plants harbor a diverse community of fungi, a preferential interaction exists between certain PGPF and a particular host. Once a particular host mutualizes this fungus, it undergoes host-specific adaptations. The outcome of such adaptations is a highly specialized and finely tuned mutualism, leading to improved responsiveness to each other needs. Evidences show that PGPF that induce growth in one plant species do not necessarily have the same effect in other species [5]. Some PGPF exert general growth promotion effects in several plant species, other fungi only do so in specific host plant. A field study showed that most of eight non-sporulating PGPF isolates enhanced the growth of one wheat variety, whereas a few isolates enhanced the growth of the other variety [87]. Moreover, at least four isolates increased yields of both varieties. Thus, the efficacy of the PGPF isolates depended upon the wheat variety in addition to their inherent growth promoting abilities. Similarly, many of the zoysiagrass PGPF isolates promoted growth of bentgrass [4], in contrast to a few isolates enhanced growth in soybean [88]. Similarly, nine isolates belonging to *Phoma* sp. and one non-sporulating fungus caused consistent plant length enhancement in cucumber cv. Shogoin fusharii compared to nine isolates except the non-sporulating fungus in cv. Aodai Kyuri. Again, plant length enhancement in cv. Jibai was shown by eight *Phoma* sp. and one non-sporulating fungus compared to five *Phoma* sp. isolates in cv. Ociai fushinari [62]. Identically, *Pe. simplicissimum* GP17-2 and *F. equiseti* 19–1 demonstrated sufficient growth-promoting effects on different host plants [4, 9, 60], but did not have effect on *Lotus japonicas* [89]. The outcome of the plant-PGPF interaction, therefore, depends on the plant and PGPF species. It is likely that the specific interaction develops during long-term co-evolution, as it has been observed for compatible and incompatible interactions of pathogens with plants [90]. Moreover, certain components of root exudates may attract and interact microbe specifically and allow it colonize the roots.

7. Mechanisms of plant growth promotion

The course of plant growth promotion by PGPF is complex and often cannot be attributed to a single mechanism. Various mechanisms that are known to modulate plant growth and development can be either direct or indirect. Direct growth promotion occurs when substances produced by the fungi or nutrient available by them facilitate plant growth. On the other hand, the ability of fungi to suppress plant pathogens and to ameliorate stress are considered major indirect mechanisms of plant growth promotion by PGPF. A particular PGPF may affect growth and development of plants using one or more of these mechanisms (**Table 3**).

7.1 Phosphate solubilization

Phosphorus is the second most important and frequently limiting macronutrient for plant growth and productivity. It is an important component of the key macromolecules in living cells and thereby, required for wide array of functions necessary for the survival and growth of living organisms. Despite the abundance of phosphorus in agricultural soils, the majority occurs in an insoluble form. Phosphorus forms complex compounds by reacting with iron, aluminum or calcium depending on the soil types and becomes insoluble and unavailable to plants [102]. To circumvent this problem, phosphate-solubilizing PGPF can play an important role dissolving insoluble P into the soluble form and making it available for plants. PGPF produce

Mechanisms	Specific activities	PGPF strain	References
Phosphate solubilization	Solubilized P by acid phosphatase and alkaline phosphatase	<i>F. verticillioides</i> RK01, <i>Humicola</i> sp. KNU01	[30]
	Solubilized P from rock phosphate and Ca-P by organic acid	<i>A. niger</i> 1B and 6A	[56]
	Solubilize P from tricalcium phosphate (TCP)	<i>A. niger</i> BHUAS01, <i>Pe. citrinum</i> BHUPC01, <i>T. harzianum</i>	[16]
	Solubilized P by organic acid activities	<i>Pe. oxalicum</i> NJDL-03, <i>Aspergillus niger</i> NJDL-12	[91]
	Phytase-mediated improvement in phytate phosphorus	<i>A. niger</i> NCIM	[76]
	Increased HCO ₃ -extractable P (23% increase)	<i>Pe. bilaiae</i> RS7B-SD1	[92]
Mineralization of organic substrate	Increased production of NH ₄ -N and NO ₂ -N in soil	<i>T. harzianum</i> GT2-1, <i>T. harzianum</i> GT3-1	[4]
	Increased availability of ammonium nitrogen from barley grain	<i>Phoma</i> sp.GS6-1, GS6-2, GS7-3, GS7-4, GS8-6, GS10-1, GS10-2, sterile fungus GU23-3	[87]
	Solubilize minerals such as MnO ₂ and metallic zinc	<i>T. harzianum</i> Rifai 1295-22	[93]
	Increased availability of ammonium nitrogen from barley grain	<i>Phoma</i> sp. GS8-1, GS8-2, GS8-3, Sterile fungus GU21-1	[62]
	Increased concentration of Cu, P, Fe, Zn, Mn and Na in roots Increased concentration of Zn, P and Mn in shoot	<i>T. harzianum</i> strain T-203	[47]
	Increased soil organic carbon, N, P and K content	<i>T. viride</i>	[73]
	Increased availability of macro and micronutrients and organic carbon	<i>T. harzianum</i> strain Th 37	[94]
Phytohormone and enzyme production	Auxin-related compounds (indole-3-acetic acid, IAA)	<i>T. virens</i> Gv. 29-8	[48]
	Gibberellins (GA1 and GA4) production	<i>A. fumigatus</i> HK-5-2	[65]
	GAs production	<i>Pe. resedanum</i> LK6	[39]
	GAs production	<i>Penicillium</i> sp. Sj-2-2	[38]
	GAs production	<i>Cladosporium</i> sp.MH-6	[24]
	GAs production	<i>Pe. citrinum</i> IR-3-3	[37]
	GAs and IAA production	<i>Chaetomium globosum</i> CAC-1G	[23]
	GAs production	<i>Exophiala</i> sp. LHL08	[26]
	GAs production	<i>Phoma herbarum</i> TK-2-4	[41]
	GAs production	<i>A. fumigatus</i> HK-5-2	[65]
	GAs production	<i>A. fumigatus</i> LH02	[18]
	IAA production	<i>T. harzianum</i> T-22	[64]
Zeatin (Ze), IAA, 1-aminocyclopropane-1-carboxylic acid (ACC)	<i>T. harzianum</i>	[95]	

Mechanisms	Specific activities	PGPF strain	References
Suppression of deleterious pathogens	Suppressed damping off caused by <i>Pythium irregular</i> , <i>Pythium</i> sp., <i>Pythium parocandrum</i> , <i>Pythium aphanidermatum</i> and <i>Rhizoctonia solani</i> AG4	Sterile fungus GSP102, <i>T. harzianum</i> GT3-2, <i>F. equiseti</i> GF19-1, <i>Pe. simplicissimum</i> GP17-2	[4]
	Induced systemic resistance against <i>Colletotrichum graminicola</i>	<i>T. harzianum</i> T22	[78]
	Suppressed bacterial wilt disease caused by <i>Ralstonia solanacearum</i>	<i>T. harzianum</i> TriH_JSB27, <i>Phoma multirostrata</i> PhoM_JSB17, <i>T. harzianum</i> TriH_JSB36, <i>Pe. chrysogenum</i> PenC_JSB41	[34]
	Suppressed Fusarium wilt caused by <i>Fusarium oxysporum</i> f. sp. <i>ciceris</i>	<i>T. harzianum</i> T-75	[58]
	Suppressed <i>Fusarium graminearum</i>	<i>Sphaerodes mycoparasitica</i>	[12]
	Suppressed damping off caused by <i>Rhizoctonia solani</i> AG4	<i>Pe. viridicatum</i> GP15-1	[35]
	Suppressed nematodes <i>Pratylenchus goodeyi</i> and <i>Helicotylenchus multicinctus</i>	<i>F. oxysporum</i> V5W2, Eny 7.11o and Emb 2.4o	[29]
	Suppressed seedling mortality by <i>Rhizoctonia solani</i>	<i>T. harzianum</i> isolate T-3	[83]
Amelioration of abiotic stress	Increased tolerance to salt stress	<i>T. harzianum</i> T-22	[69]
	Mitigation of oxidative stress due to NaOCl and cold stress	<i>T. harzianum</i> Rifai strain 1295-22	[96]
	Enhanced maize seedling copper stress tolerance	<i>Chaetomium globosum</i>	[97]
	Minimized Cu-induced electrolytic leakage and lipid peroxidation	<i>Pe. funiculosum</i> LHL06	[98]
	Increased tolerance to drought stress	<i>T. atroviride</i> ID20G	[99]
Volatile organic compounds (VOCs)	Produced abundant classes of VOCs (sesquiterpenes and diterpenes)	<i>F. oxysporum</i> NRRL 26379, NRRL 38335	[28]
	Produced mainly terpenoid-like volatiles including β -caryophyllene	<i>Talaromyces wortmannii</i> FS2	[40]
	Produced 2-methyl-propanol and 3-methyl-butanol	<i>Phoma</i> sp. GS8-3	[100]
	Produced abundant amount of isobutyl alcohol, isopentyl alcohol, and 3-methylbutanal	<i>T. viride</i>	[101]

Table 3. Different mechanisms of plant growth promotion used by various plant growth promoting fungi (PGPF).

phosphate-solubilizing enzymes such as phytases and phosphatases and organic acids, which liberate P from insoluble phosphates. The most efficient phytase and phosphatase producing PGPF belong to the genera *Aspergillus*, *Trichoderma*, and *Penicillium* [103]. The order in terms of phytate hydrolysis efficacy was *Aspergillus* > *Penicillium* > *Trichoderma* [104]. *Fusarium verticillioides* RK01 and *Humicola* sp. KNU01 solubilized phosphate by increasing activities of acid phosphatase and alkaline phosphatase, and promoted soybean growth significantly [30]. The phosphate solubilizing fungi possess greater phosphorus solubilization ability than bacteria,

especially under acidic soil conditions [105]. The main reason is most fungi are eosinophilic, and have relatively higher growth in acidic environments than bacteria [106]. The acidity has significant influence on organic acid-mediated phosphate solubilizing activities of *Pe. oxalicum* NJDL-03 and *A. niger* NJDL-12 [91]. However, acidification is not always the major mechanism of P solubilization by *T. harzianum* Rifai 1295-22 (T-22), where pH of cultures never fell below 5.0 and no organic acids were detected [93]. Some of the reported PGPF such as *Aspergillus niger* has twin abilities of P mineralization and solubilization [104]. The fungus releases P both from organic and inorganic sources. These suggests that specific PGPF may have specific activity in solubilizing phosphate and making it available for crop growth.

7.2 Substrate degradation (mineralization)

Microorganisms primarily mediate soil nutrient pathways. Microbial mineralization of nutrients from organic matter is crucial for plant growth. Some PGPF promote plant growth, but do not produce plant hormones or solubilize fixed phosphate. Among *Pe. radicum*, *Pe. bilaiae* (strain RS7B-SD1) and *Penicillium* sp. strain KC6-W2, the strongest growth promotion in wheat, medic, and lentil was shown by *Penicillium* sp. KC6-W2, while the only significant P increase (~23% increase) was found in *Pe. bilaiae* RS7B-SD1-treated plants [92]. Similarly, seven *Trichoderma* isolates significantly improved the growth of bean seedlings; despite some of them do not possess any of the assessed growth-promoting traits such as soluble P, indole acetic acid (IAA) and siderophores [107]. These PGPF are believed to encourage plant growth by accelerating mineralization in the soil. Fungi have better substrate assimilation efficiency than any other microbes and are able to break down complex polyaromatic compounds such lignin and humic or phenolic acids [108]. A close relationship was found between the cellulose and starch degradation activity of PGPF for decomposing barley grain and their subsequent growth promotion effect in plants [109]. Application of *T. harzianum* strain Th 37 increased the availability of macro and micronutrients and organic carbonate in the ratoon initiation stage in sugarcane [94]. Colonization of *T. harzianum* in cucumber roots enhanced the availability and uptake of nutrients by the plants [47]. Cucumber plants grew better and produced more marketable fruits due to an increase in soil nutrients caused by PGPF, and accumulated more inorganic minerals like Ca, Mg, and K in aerial shoots [62]. PGPF are also directly involved the degradation of the nitrogenous organic materials through ammonization and nitrification. Formation of $\text{NH}_4\text{-N}$ and $\text{NO}_2\text{-N}$ in soil was accelerated during soil amendment with PGPF-infested barley grains [109]. More interestingly, the fungal entomopathogen *Metarhizium robertsii*, when established as a root endophyte, was shown to translocate nitrogen from a dead insect to a common bean plant host, suggesting this PGPF's potential to acquire mineral nutrients from organic matter and promote plant growth [54]. Nutrient release by mineralization could explain why PGPF other than mycorrhizae improve plant growth when added to soil.

7.3 Phytohormone production

Phytohormones are involved in many forms of plant-microbe interactions and also in the beneficial interactions of plants with PGPF. The commonly recognized classes of phytohormones produced by PGPF are the auxins (IAA) and gibberellins (GAs) (Table 3). IAA, the most studied auxin, regulates many aspects of plant growth, in particular, root morphology by inhibiting root elongation, increasing lateral root production, and inducing adventitious roots [48]. The *T. harzianum* T-22-mediated root biomass production and root hair development in maize is

believed to operate through a classical IAA response pathway [78]. Similarly, a direct correlation exists between increased levels of fungal IAA and lateral root development in *Arabidopsis* seedlings inoculated with *T. virens* [48].

GAs are well known for their role in various developmental processes in plants, including stem elongation. Shoot elongation of waito-c rice seedlings by culture filtrates of *Pe. citrinum* IR-3-3 and *A. clavatus* Y2H0002 was attributed to the activity of physiologically active GAs existing in the culture filtrates [19, 37]. Biochemical analyses of *Penicillium* sp. LWL3 and *Pe. glomerata* LWL2 culture filtrates that enhanced the growth of Dongjiin beyo rice cultivar and in GA-deficient mutant *Waito-C* revealed the presence of IAA and various GAs [110]. Similarly, production of bioactive GAs correlated with enhanced growth of *Waito-C* under salinity by *Penicillium* sp. Sj-2-2 [38]. GA also played key roles during root colonization by *P. indica* in pea roots [111].

Another phytohormone through which PGPF mediate plant growth is cytokinin, especially the Zeatin. Zeatin production has been documented in *Piriformospora indica*, *T. harzianum* and *Phoma* sp., and the fungi that also produce other phytohormones [95, 112, 113]. *P. indica* produces low amounts of auxins, but high levels of cytokinins. *Trans*-Zeatin cytokinin biosynthesis was found crucial for *P. indica*-mediated growth stimulation in *Arabidopsis* [112]. This evidence suggests that PGPF often mediate the various growth and developmental processes in plants by influencing the balance of various plant hormones.

7.4 Microbial ACC deaminase

PGPF produces a crucial enzyme ACC (1-aminocyclopropane-1-carboxylic acid) deaminase. ACC deaminase cleaves the ethylene precursor, 1-aminocyclopropane-1-carboxylic acid (ACC), into NH_3 (ammonia) and α -ketobutyrate [114]. The ACC deaminase regulates the plant growth by cleaving ACC produced by plants and thereby minimizing the ethylene level in the plant, which when present in high concentrations can lead to a reduced plant growth [115]. ACC deaminase is an inducible enzyme encoded by *acdS* genes of fungi and bacteria [116]. ACC deaminase appears to be central to the functional interactions of some plant-PGPF. *T. asperellum* T203 produced high levels of ACC deaminase and showed an average 3.5-fold induction of the *acds* gene [117]. When ACC deaminase expression is impaired in the fungus *T. asperellum* T203, the plant growth promotion abilities of this organism are also decreased [51]. The root colonizing bacteria *T. harzianum* T22 no longer promote canola root elongation after its *acdS* gene is knocked out [64]. Production of ACC deaminase was reported in some other fungi, which include *Issatchenkia occidentalis* [118], and *Penicillium citrinum* and a stramnopile, *Phytophthora sojae* [119, 120]. The ACC deaminase-producing microbes have competitive advantages in the rhizosphere over nonproducing microorganisms because the enzyme acts as a nitrogen source for them [116]. Moreover, bacteria and fungi that express ACC deaminase can lower the impact of a range of different stresses that affect plant growth and development [114]. These show that ACC deaminase is not only related to plant growth promotion abilities of the microbes, but also play additional roles in the rhizosphere.

7.5 Suppression of deleterious microorganisms by PGPF

The key indirect mechanism of PGPF-mediated plant growth promotion is through their activities as biocontrol agents. PGPF protect and empower plants to resist harmful pathogens and ensure their better growth. The mechanisms by which PGPF suppress growth or activity of invading pathogens in crop plants

include antibiosis, competition for nutrient and space, mycoparasitism and induced systemic resistance (ISR) [121]. PGPF of diverse genera promoted growth of field-soil grown cucumber by counteracting damping off pathogen *Pythium* sp. through microbial antagonism [4]. Banana plants inoculated with PGPF *F. oxysporum* significantly suppressed nematode pathogens *Pratylenchus goodeyi* and *Helicotylenchus multincinctus* resulting in up to ~20 to 36% increase in banana yields [29]. The mycoparasite *Sphaerodes retispora* has been reported to improve the plant dry weight and to decrease plant mortality in the presence of *F. oxysporum* [122]. Similarly, under phytotron conditions, seed germination, root biomass, total biomass, root length, and total length of *F. graminearum*-infected wheat were noticeably increased with the treatments of *S. mycoparasitica* and *T. harzianum*, as compared to inoculation with *F. graminearum* alone. Both mycoparasites prevented colonization and reduction in root growth by the pathogen [12]. PGPF compete with the pathogen for colonization niche on roots [79]. Other mechanisms of disease suppression by PGPF are, therefore, likely to include competition with pathogens for infection sites on the root surface. Moreover, there is a long and growing list of PGPF such as *Trichoderma*, *Penicillium*, *Fusarium*, *Phoma*, and non-sporulating fungi, which can protect crop plants against pathogens by eliciting ISR [14, 31, 123, 124]. Although many fungal strains to act as PGPF and elicit ISR, it is not clear how far both mechanisms are connected. These microbes may use some of the same mechanisms to promote plant growth and control plant pathogens.

7.6 Rhizoremediation and stress control

The microbial association of plants has a major influence on plant adaptation to abiotic stresses such as salinity, drought, heavy metal toxicity, extreme temperatures and oxidative stress. Recent studies indicate that fitness benefits conferred by certain PGPF contribute plant adaption to stresses [125]. There are reports of enhanced plant growth because of the association of PGPF with plants, even when plants are under suboptimal conditions [126]. Root colonization by *T. atroviride* ID20G increased fresh and dry weight of maize roots under drought stress [99]. Supplementation of *T. harzianum* to NaCl treated mustard seedlings showed elevation by 13.8, 11.8, and 16.7% in shoot, root length and plant dry weight, respectively as compared to plants treated with NaCl (200 mM) alone [127]. The fungus *Pe. funiculosum* significantly increased the plant biomass, root physiology and nutrients uptake to soybean under copper stress [98]. These fungi have been known to produce plant growth regulators (like GAs and auxins) and extend plant tolerance to abiotic and biotic stresses [23, 125]. Recurrently, *T. harzianum* T22 has little effect upon seedling performance in tomato, however, under stress; treated seeds germinate consistently faster and more uniformly than untreated seeds [69]. A few other fungi like *Microsphaeropsis*, *Mucor*, *Phoma*, *Alternaria*, *Peyronellaea*, *Steganosporium*, and *Aspergillus* are known to grow well in polluted medium and protect plants from adverse effects of metal stress [128]. There are numerous similar examples of PGPF ameliorating abiotic stresses and promoting plant growth. Despite significant differences between different stresses, cellular responses to them share common features. Enhanced resistance of PGPF-treated plants to abiotic stresses is explained partly due to higher capacity to scavenge ROS and recycle oxidized ascorbate and glutathione [99, 127]. The increase in proline content is found to be very useful in providing tolerance to these plants under stress [129]. Both enzymatic (peroxidase, catalase, superoxide dismutase, ascorbate peroxidase, monodehydroascorbate reductase, dehydroascorbate reductase, glutathione reductase, glutathione S-transferase and glucoyl peroxidase), and non-enzymatic (ascorbic acid, reduced glutathione, oxidized glutathione) antioxidants are induced by PGPF further

enhance the synthesis of these phytoconstituents and defend the plants from further damage [127].

7.7 Production of volatile organic compounds (VOCs)

Microorganisms produce various mixtures of gas-phase, carbon-based compounds called volatile organic compounds (VOCs) as part of their normal metabolism. The comparative analysis of experimental data has shown that volatile metabolites make a much greater contribution to the microbial interactions than non-volatile ones [130]. Recent studies reveal that VOC emission is indeed a common property of a wide variety of soil fungi, including PGPF. Some of these VOCs produced by PGPF exert stimulatory effects on plants. A PGPF, *Talaromyces wortmannii* emits a terpenoid-like volatile, β -caryophyllene, which significantly promoted plant growth and induced resistance in turnip [40]. The identified VOCs emitted by *Phoma* sp. GS8-3 belonged to C4-C8 hydrocarbons, where 2-methyl-propanol and 3-methyl-butanol formed the main components and promoted the growth of tobacco seedlings [100]. These two components were also extracted from PGPR [131]. On the other hand, 3-methyl-butanol has been reported from *T. viride* [101]. The other most abundant VOCs from *T. viride* were isobutyl alcohol, isopentyl alcohol, farnesene and geranylacetone. *Arabidopsis* cultured in petri plates in a shared atmosphere with *T. viride*, without direct physical contact was taller with more lateral roots, bigger with augmented total biomass (~45%) and earlier flowered with higher chlorophyll concentration (~58%) [101]. Moreover, volatile blends showed better growth promotion than individual compounds [132]. Volatile compounds produced by PGPF are also heavily involved in induce systemic resistance toward pathogens [100].

8. Pattern and process of root colonization by PGPF

Root colonization is considered as an important strategy of PGPF for plant growth promotion. Root colonization is the ability of a fungus to survive and proliferate along growing roots in the presence of the indigenous microflora over a considerable period [35]. The fungus that colonizes plant root effectively is more rhizosphere competent than others [107]. Rhizosphere competence is a necessary condition for a fungus to be an efficient PGPF. Re-isolation frequency of the fungus from the colonized roots is an indirect measure of its root colonizing ability and thereby, its rhizosphere competence. In such studies, *Pe. simplicissimum* GP17-2 and *Pe. viridicatum* GP15-1 were re-isolated from *Arabidopsis* Col-0 roots 3 weeks after planting at high frequencies which were found to be >90% (**Figure 2**). Similarly, the re-isolation frequency of *Pe. janthinellum* GP16-2 from the roots of Col-0 plants was recorded to be, on average, 85% [33]. *Aspergillus* spp. PPA1 was re-isolated from the roots of cucumber plants at a frequency of 95–100% 3 weeks after planting [17], indicating a rapid and efficient root colonization by the PGPF. However, a slow root colonization by PGPF was also reported, as it was the case with *Phoma* sp. GS8-2, which achieved maximum colonization on cucumber roots at 10 weeks [62]. The relative growth rate of the fungi and roots seems to determine the length of time required for maximum root colonization.

Some PGPF selectively colonize host roots and promote growth. Isolates of *Phoma* and sterile fungi showed poor ability to colonize the soybean roots and were unable to enhance the growth of soybean [79]. Similarly, *T. koningi* colonized roots and enhanced growth of *Lotus japonicas*, but *Pe. simplicissimum* and *F. equiseti* did not [89]. It was observed that *T. koningi* induced a transient and decreased level of defense gene expression in *L. japonicas* during its entry into the roots, while a



Figure 2. Re-isolation of *Penicillium simplicissimum* GP17-2 and *Penicillium viridicatum* GP15-1 at higher frequencies from colonized roots of *Arabidopsis thaliana* ecotype Col-0 3 weeks after sowing.

stimulated expression of these genes was induced by *Pe. simplicissimum* and *F. equiseti* [89]. *T. koningi* resembles symbiotic fungi, while *Pe. simplicissimum* and *F. equiseti* act similar to fungal pathogens in activating host defense. This shows that legumes selectively avoid some PGPF and thus allow only specific PGPF to interfere.

There are also PGPF, in particular, the non-sporulating sterile fungi that lack root colonization ability, but they are able to promote growth and yield of plants [62, 133]. This indicates that root colonization is not an indispensable condition for growth promotion by all PGPF. Some chemical factor(s) produced by them might be responsible for growth promotion.

The colonization of the root system of by PGPF is not always homogenous; the density of PGPF varies in different parts of the root system. The colonization of roots by the majority of PGPF appears to be higher in the upper than in the middle and lower root parts of roots, [35, 133]. The lower part was always less colonized by PGPF, especially during first 2 weeks of colonization. This is probably due to the faster growth of the roots than of the hyphae. Moreover, the main zone of root exudation is located behind the apex [134]. However, some PGPF can keep up with root growth and colonize the entire root system [35]. Only fungi with large nutrient reserves can move to the root and along the root over larger distances [135].

Anatomical data show that PGPF may colonize root tissues internally and establish a mutualistic relationship with host. *F. equiseti* GF19-1 produced abundant hyphal growth on the root surface, formed appressoria-like structures and grew in the intercellular space, not inside the cell [31]. *T. harzianum* CECT 2413 exhibited profuse adhesion of hyphae to the tomato roots and colonized the epidermis and cortex. Intercellular hyphal growth and the formation of plant-induced papilla-like hyphal tips were also observed [136]. Hyphae of *T. koningi* penetrated the epidermis and entered the intercellular inner cortex tissues [89]. Sterile red fungus has been also demonstrated to invade the inner root regions that helped plants derive nutrients from the soil and protected roots from pathogens [137].

9. Formulation of PGPF

PGPF, especially *Trichoderma*, have many success stories as plant growth promoting agents and appear to have much potential as a commercial formulation. Different organic and inorganic carrier materials have been studied for effective delivery of bioinoculants. A talc-based formulation was developed for *T. harzianum* to supply concentrated conidial biomass of the fungus with high colony forming units (CFU) and long shelf life [138]. The concentrated formulation provided an extra advantage of smaller packaging for storage and transportation, and low

product cost as compared to other carriers such as charcoal, vermiculite, sawdust and cow dung. Seed application of the formulation recorded significant increase in growth promotion in chickpea [138]. Corn and sugarcane bagasse were used as potential carriers for *Trichoderma* sp. SL2 inoculants. The corn formulation of SL2 significantly enhanced rice seedlings root length, wet weight and biomass compared to inoculum mixed with sugarcane bagasse and control [139]. A spray-dried flowable powder formulation was developed for biostimulant *Trichoderma* strains using a CO₂ generating dispersant system, based on polyacrylic acid, citric acid and sodium bicarbonate, polyvinyl alcohol as adhesives and lecithin as wetting agent [140]. Hydrolytic amino acids derived from pig corpses were used in the preparation of *T. harzianum* T-E5-containing bioorganic fertilizer. The resulting bioorganic fertilizer supported higher densities of *T. harzianum* T-E5 and substantially enhanced plant growth when applied as a soil amendment [141]. A composted cattle manure-based *Trichoderma* biofertilizer was developed and tested in the field. Plots fertilized with biofertilizer had the greatest aboveground biomass of any treatment and were significantly more productive than non-amended plots and plots fertilized with any rate of organic fertilizer [142]. Effective formulation of *P. indica* was prepared in talcum powder or vermiculite with 20% moisture. The talcum-based formulations performed significantly better as bioinoculant over vermiculite-based formulations in glasshouse experiments [143]. These show the feasibility of commercial level production and applicability of different PGPF formulations for plant growth promotion in the field.

10. Conclusions

Because of current concerns over the adverse effects of agrochemicals, there is a growing interest in improving our understanding of the role and application of beneficial microbes in agriculture. The plant-associated growth promoting fungi show excellent potential for wider use in sustainable agriculture as they improve plant growth and yield in an ecofriendly and cost-effective manner. However, the PGPF continue to be greatly underutilized, primarily due to some practical problems such as the inconsistency in field performance, which appears to be the greatest challenge in the development of microbial inoculants for plant growth until now and well into the future. If our understanding of complex rhizosphere environment, of the mechanisms of action of PGPF and of the practical aspects of mass production, inoculant formulation and delivery increase, more PGPF products will become available. Knowledge of multiple microbial interaction with different or complementary mode of actions is also of extreme value for development of bio-formulation.

Recent advances in biotechnological tools and reliable transformation system could be useful in engineering of the PGPF to confer improved benefits to the crop. Genetic transformation and overexpression of one or more of the plant growth promoting traits that act synergistically may lead to enhanced performance by the inoculant. Research may be required periodically in order to evaluate the genetic stability and ecological persistence of the genetically modified strain. Efforts should be strengthened to foster linkage between investigators and entrepreneurs in facilitating technology transfer, promotion and acceptance by end users.

IntechOpen

Author details

Md. Motaher Hossain^{1*} and Farjana Sultana²

1 Department of Plant Pathology, Bangabandhu Sheikh Mujibur Rahman Agricultural University, Gazipur, Bangladesh

2 College of Agricultural Sciences, International University of Business Agriculture and Technology, Dhaka, Bangladesh

*Address all correspondence to: hossainmm@bsmrau.edu.bd

IntechOpen

© 2020 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] WPP. World Population Prospect: 2019. New York: Population Division, Department of Economic and Social Affairs, United Nations; 2019. p. 2
- [2] Hunter MC, Smith RG, Schipanski ME, Atwood LW, Mortensen DA. Agriculture in 2050: Recalibrating targets for sustainable intensification. *BioScience*. 2017;**67**(4):386-391
- [3] Vacheron J, Desbrosses G, Bouffaud ML, Touraine B, Moëne-Loccoz Y, Muller D, et al. Plant growth-promoting rhizobacteria and root system functioning. *Frontiers in Plant Science*. 2013;**4**:356
- [4] Hyakumachi M. Plant-growth-promoting fungi from turf grass rhizosphere with potential for disease suppression. *Soil Microorganisms*. 1994;**44**:53-68
- [5] Hossain MM, Sultana F, Islam S. Plant growth-promoting fungi (PGPF): Phytostimulation and induced systemic resistance. In: Singh D, Singh H, Prabha R, editors. *Plant-Microbe Interactions in Agro-Ecological Perspectives*, Volume 2: Microbial Interactions and Agro-Ecological Impacts. Singapore: Springer; 2017. pp. 135-191. DOI: 10.1007/978-981-10-6593-4
- [6] Lyn ME, Burnett D, Garcia AR, Gray R. Interaction of water with three granular biopesticide formulations. *Journal of Agricultural and Food Chemistry*. 2010;**58**:1804-1814
- [7] Aly AH, Debbab A, Proksch P. Fungal endophytes: Unique plant inhabitants with great promises. *Applied Microbiology and Biotechnology*. 2011;**90**:1829-1845
- [8] Bent E. Induced systemic resistance mediated by plant growth-promoting rhizobacteria (PGPR) and fungi (PGPF). In: Tuzun S, Bent E, editors. *Multigenic and Induced Systemic Resistance in Plants*. New York: Springer; 2006. pp. 225-258
- [9] Hossain MM, Sultana F, Kubota M, Koyama H, Hyakumachi M. The plant growth-promoting fungus *Penicillium simplicissimum* GP17-2 induces resistance in *Arabidopsis thaliana* by activation of multiple defense signals. *Plant & Cell Physiology*. 2007;**48**(12):1724-1736
- [10] Hyakumachi M, Kubota M. Fungi as plant growth promoter and disease suppressor. In: Arora DK, editor. *Mycology Series*. Vol. 21. *Fungal Biotechnology in Agricultural, Food, and Environmental Applications*. New York: Marcel Dekker; 2004. pp. 101-110
- [11] Kaewchai S, Soyong K, Hyde KD. Mycofungicides and fungal bio-fertilizers. *Fungal Diversity*. 2009;**38**: 25-50
- [12] Vujanovic V, Goh YK. qPCR quantification of *Sphaerodes myco-parasitica* biotrophic mycoparasite interaction with *Fusarium graminearum*: *in vitro* and *in planta* assays. *Archives of Microbiology*. 2012;**194**(8):707-717
- [13] Muslim A, Horinouchi H, Hyakumachi M. Biological control of *Fusarium* wilt of tomato with hypovirulent binucleate *Rhizoctonia* in greenhouse conditions. *Mycoscience*. 2003;**44**:77-84
- [14] Sultana F, Hossain MM, Kubota M, Hyakumachi M. Elicitation of systemic resistance against the bacterial speck pathogen in *Arabidopsis thaliana* by culture filtrates of plant growth-promoting fungi. *Canadian Journal of Plant Pathology*. 2008;**30**(2):196-205
- [15] Zhou Z, Zhang C, Zhou W, Li W, Chu L, Yan J, et al. Diversity and plant

- growth-promoting ability of endophytic fungi from the five flower plant species collected from Yunnan, Southwest China. *Journal of Plant Interactions*. 2014;**9**(1):585-591
- [16] Yadav J, Verma JP, Tiwari KN. Plant growth promoting activities of fungi and their effect on chickpea plant growth. *Asian Journal of Biological Sciences*. 2011;**4**:291-299
- [17] Islam S, Akanda AM, Sultana F, Hossain MM. Chilli rhizosphere fungus *Aspergillus* spp. PPA1 promotes vegetative growth of cucumber (*Cucumis sativus*) plants upon root colonisation. *Archives of Phytopathology and Plant Protection*. 2014;**47**:1231-1238
- [18] Khan AL, Hamayun M, Kim YH, Kang SM, Lee JH, Lee IJ. Gibberellins producing endophytic *Aspergillus fumigatus* sp. LH02 influenced endogenous phytohormonal levels, plant growth and isoflavone biosynthesis in soybean under salt stress. *Process Biochemistry*. 2011;**46**:440-447
- [19] You YH, Kwak TW, Kang SM, Lee MC, Kim JG. *Aspergillus clavatus* Y2H0002 as a new endophytic fungal strain producing gibberellins isolated from *Nymphoides peltata* in fresh water. *Mycobiology*. 2015;**43**:87-91
- [20] Salas-Marina MA, Silva-Flores MA, Cervantes-Badillo MG, Rosales-Saavedra MT, Islas-Osuna MA, Casas-Flores S. The plant growth-promoting fungus *Aspergillus ustus* promotes growth and induces resistance against different lifestyle pathogens in *Arabidopsis thaliana*. *Journal of Microbiology and Biotechnology*. 2011;**21**(7):686-696
- [21] Waqas M, Khan AL, Hamayun M, Shahzad R, Kim YH, Choi KS, et al. Endophytic infection alleviates biotic stress in sunflower through regulation of defence hormones, antioxidants and functional amino acids. *European Journal of Plant Pathology*. 2015;**141**:803-824
- [22] Ignatova LV, Brazhnikova YV, Berzhanova RZ, Mukasheva TD. Plant growth-promoting and antifungal activity of yeasts from dark chestnut soil. *Microbiological Research*. 2015;**175**:78-83
- [23] Khan AL, Shinwari ZK, Kim YH, Waqas M, Hamayun M, Kamran M, et al. Role of endophyte *Chaetomium globosum* LK4 in growth of *Capsicum annum* by production of gibberellins and indole acetic acid. *Pakistan Journal of Botany*. 2012;**44**:1601-1607
- [24] Hamayun M, Khan SA, Khan AL, Rehman G, Kim YH, Iqbal I, et al. Gibberellin production and plant growth promotion from pure cultures of *Cladosporium* sp. MH-6 isolated from cucumber (*Cucumis sativus* L.). *Mycologia*. 2010;**102**:989-995
- [25] Hamayun M, Khan SA, Ahmad N, Tang D-S, Kang S-M, et al. *Cladosporium sphaerospermum* as a new plant growth-promoting endophyte from the roots of *Glycine max* (L.) Merr. *World Journal of Microbiology and Biotechnology*. 2009;**25**(4):627-632
- [26] Khan AL, Hamayun M, Ahmad N, Waqas M, Kang SM, Kim YH, et al. *Exophiala* sp. LHL08 reprograms *Cucumis sativus* to higher growth under abiotic stresses. *Physiologia Plantarum*. 2011;**143**(4):329-343
- [27] Islam S, Akanda AM, Prova A, Sultana F, Hossain MM. Growth promotion effect of *Fusarium* spp. PPF1 from Bermuda grass (*Cynodon dactylon*) rhizosphere on Indian spinach (*Basella alba*) seedlings are linked to root colonization. *Archives of Phytopathology and Plant Protection*. 2014;**47**:2319-2331
- [28] Bitas V, McCartney N, Li N, Demers J, Kim JE, Kim HS, et al.

Fusarium oxysporum volatiles enhance plant growth via affecting auxin transport and signaling. *Frontiers in Microbiology*. 2015;**6**:1248

[29] Waweru B, Turoop L, Kahangi E, Coyne D, Dubois T. Non-pathogenic *Fusarium oxysporum* endophytes provide field control of nematodes, improving yield of banana (*Musa* sp.). *Biological Control*. 2014;**74**:82-88

[30] Radhakrishnan R, Khan AL, Kang SM, Lee IJ. A comparative study of phosphate solubilization and the host plant growth promotion ability of *Fusarium verticillioides* RK01 and *Humicola* sp. KNU01 under salt stress. *Annales de Microbiologie*. 2015;**65**(1):585-593

[31] Kojima H, Hossain MM, Kubota M, Hyakumachi M. Involvement of the salicylic acid signaling pathway in the systemic resistance induced in *Arabidopsis* by plant growth-promoting fungus *Fusarium equiseti* GF19-1. *Journal of Oleo Science*. 2013;**62**(6):415-426

[32] Maciá-Vicente JG, Jansson HB, Talbot NJ, Lopez-Llorca LV. Real-time PCR quantification and live-cell imaging of endophytic colonization of barley (*Hordeum vulgare*) roots by *Fusarium equiseti* and *Pochonia chlamydosporia*. *New Phytologist*. 2009;**182**:213-228

[33] Hossain MM, Sultana F, Kubota M, Hyakumachi M. Differential inducible defense mechanisms against bacterial speck pathogen in *Arabidopsis thaliana* by plant-growth-promoting-fungus *Penicillium* sp. GP16-2 and its cell free filtrate. *Plant and Soil*. 2008;**304**:227-239

[34] Jogaiah S, Abdelrahman M, Tran LSP, Shin-ichi I. Characterization of rhizosphere fungi that mediate resistance in tomato against bacterial wilt disease. *Journal of Experimental Botany*. 2013;**64**:3829-3842

[35] Hossain MM, Sultana F, Miyazawa M, Hyakumachi M. The plant growth promoting fungi *Penicillium* spp. GP15-1 enhances growth and confers protection against damping-off and anthracnose in the cucumber. *Journal of Oleo Science*. 2014;**63**(4):391-400

[36] Babu AG, Kim SW, Yadav DJ, Hyum U, Adhikari M, Lee YS. *Penicillium menonorum*: A novel fungus to promote growth and nutrient management in cucumber plants. *Mycobiology*. 2015;**43**:49-56

[37] Khan SA, Hamayun M, Yoon H, Kim HY, Suh SJ, Hwang SK, et al. Plant growth promotion and *Penicillium citrinum*. *BMC Microbiology*. 2008;**8**:231. DOI: 10.1186/1471-2180-8-231

[38] You YH, Yoon H, Kang SM, Shin JH, Choo YS, Lee IJ, et al. Fungal diversity and plant growth promotion of endophytic fungi from six halophytes in Suncheon Bay. *Journal of Microbiology and Biotechnology*. 2012;**22**:1549-1556

[39] Khan AL, Waqas M, Lee IJ. Resilience of *Penicillium resedanum* LK6 and exogenous gibberellin in improving *Capsicum annuum* growth under abiotic stresses. *Journal of Plant Research*. 2015;**128**(2):259-268

[40] Yamagiwa Y, Toyoda K, Inagaki Y, Ichinose Y, Hyakumachi M, Shiraishi T. *Talaromyces wortmannii* FS2 emits β -caryophyllene, which promotes plant growth and induces resistance. *Journal of General Plant Pathology*. 2011;**77**:336-341

[41] Hamayun M, Khan SA, Khan AL, Rehman G, Sohn EY, Shah AA, et al. *Phoma herbarum* as a new gibberellin-producing and plant growth-promoting fungus. *Journal of Microbiology and Biotechnology*. 2009;**19**:1244-1249. DOI: 10.4014/jmb.0901.030

[42] Hamayun M, Khan SA, Khan AL, Tang DS, Hussain J, Ahmad N, et al.

Growth promotion of cucumber by pure cultures of gibberellin-producing *Phoma* sp. GAH7. *World Journal of Microbiology and Biotechnology*. 2010;**26**:889-894

[43] Siddikee MA, Zereen MI, Li CF, Dai CC. Endophytic fungus *Phomopsis liquidambari* and different doses of N-fertilizer alter microbial community structure and function in rhizosphere of rice. *Scientific Reports*. 2016;**6**:32270

[44] Cavello IA, Crespo JM, García SS, Zapiola JM, Luna MF, Cavalitto SF. Plant growth promotion activity of keratinolytic fungi growing on a recalcitrant waste known as “Hair Waste”. *Biotechnology Research International*. 2015;**2015**:952921. DOI: 10.1155/2015/952921

[45] Pascual CB, Raymundo AD, Hayakumachi M. Efficacy of hypovirulent binucleate *Rhizoctonia* sp. to control banded leaf and sheath blight in corn. *Journal of General Plant Pathology*. 2000;**66**:95-102

[46] Jiang J, Tam S, Toda T, Chen L. Controlling *Rhizoctonia* damping-off of Chinese mustard by using endomycorrhizal *Rhizoctonia* spp. isolated from orchid mycorrhizae. *Plant Disease*. 2015;**100**:85-91

[47] Yedidia I, Srivastva AK, Kapulnik Y, Chet I. Effect of *Trichoderma harzianum* on microelement concentrations and increased growth of cucumber plants. *Plant and Soil*. 2001;**235**:235-242

[48] Contreras-Cornejo HA, Macías-Rodríguez LI, Cortés-Penagos C, López-Bucio J. *Trichoderma virens*, a plant beneficial fungus, enhances biomass production and promotes lateral root growth through an auxin-dependent mechanism in *Arabidopsis*. *Plant Physiology*. 2009;**149**:1579-1592

[49] Lee S, Yap M, Behringer G, Hung R, Bennett JW. Volatile organic compounds emitted by *Trichoderma* species mediate

plant growth. *Fungal Biology and Biotechnology*. 2016;**3**:7

[50] Zhang S, Gan Y, Xu B. Application of plant-growth-promoting fungi *Trichoderma longibrachiatum* t6 enhances tolerance of wheat to salt stress through improvement of antioxidative defense system and gene expression. *Frontiers in Plant Science*. 2016;**7**:1405

[51] Brotman Y, Landau U, Cuadros-Inostroza Á, Takayuki T, Fernie AR, et al. *Trichoderma*-plant root colonization: Escaping early plant defense responses and activation of the antioxidant machinery for saline stress tolerance. *PLoS Pathogens*. 2013;**9**(3):e1003221

[52] Studholme DJ, Harris B, Le Cocq K, Winsbury R, Perera V, Ryder L, et al. Investigating the beneficial traits of *Trichoderma hamatum* GD12 for sustainable agriculture—Insights from genomics. *Frontiers in Plant Science*. 2013;**4**:258

[53] Stone BWG, Weingarten EA, Jackson CR. The role of the phyllosphere microbiome in plant health and function. *Annual Plant Reviews*. 2018;**1**:1-24. DOI: 10.1002/9781119312994.apr0614

[54] Behie SW, Zelisko PM, Bidochka MJ. Endophytic insect-parasitic fungi translocate nitrogen directly from insects to plants. *Science*. 2012;**336**:1576-1577

[55] Eldredge SD, Geary B, Jensen SL. Seed isolates of *Alternaria* and *Aspergillus* fungi increase germination of *Astragalus utahensis*. *Native Plants*. 2016;**17**(2):89-94

[56] Chuang CC, Kuo YL, Chao CC, Chao WL. Solubilization of inorganic phosphates and plant growth promotion by *Aspergillus niger*. *Biology and Fertility of Soils*. 2007;**43**:575-584

- [57] Rabeendran N, Moot DJ, Jones EE, et al. Inconsistent growth promotion of cabbage and lettuce from *Trichoderma* isolates. New Zealand Plant Protection. 2000;**53**:143-146
- [58] Hossain MM, Hossain N, Sultana F, Islam SMN, Islam S, Bhuiyan MKA. Integrated management of *Fusarium* wilt of chickpea (*Cicer arietinum* L.) caused by *Fusarium oxysporum* f.sp. *ciceris* with microbial antagonist, botanical extract and fungicide. African Journal of Biotechnology. 2013;**12**(29):4699-4706
- [59] Kaveh H, Vatandoost S, Aroiee H, Mazhabi M. Would *Trichoderma* affect seed germination and seedling quality of two muskmelon cultivars, Khatooni and Qasri and increase their transplanting success? Journal of Biological and Environmental Sciences. 2011;**5**:169-175
- [60] Chandanie WA, Kubota M, Hyakumachi M. Interactions between the arbuscular mycorrhizal fungus *Glomus mosseae* and plant growth-promoting fungi and their significance for enhancing plant growth and suppressing damping-off of cucumber (*Cucumis sativus* L.). Applied Soil Ecology. 2009;**41**:336-341
- [61] Saldajeno MGB, Hyakumachi M. The plant growth-promoting fungus *Fusarium equiseti* and the arbuscular mycorrhizal fungus *Glomus mosseae* stimulate plant growth and reduce severity of anthracnose and damping-off diseases in cucumber (*Cucumis sativus*) seedlings. The Annals of Applied Biology. 2011;**159**:28-40
- [62] Shivanna MB, Meera MS, Kubota M, Hyakumachi M. Promotion of growth and yield in cucumber by zoysiagrass rhizosphere fungi. Microbes and Environments. 2005;**20**(1):34-40
- [63] Chandanie WA, Kubota M, Hyakumachi M. Interaction between arbuscular mycorrhizal fungus *Glomus mosseae* and plant growth promoting fungus *Phoma* sp. on their root colonization and growth promotion of cucumber (*Cucumis sativus* L.). Mycoscience. 2005;**46**:201-204
- [64] Sofu A, Tataranni G, Xiloyannis C, Dichio B, Scopa A. Direct effects of *Trichoderma harzianum* strain T-22 on micropropagated shoots of GiSeLa6® (*Prunus cerasus* × *Prunus canescens*) rootstock. Environmental and Experimental Botany. 2012;**76**:33-38
- [65] Hamayun M, Khan SA, Khan MA, Khan AL, Kang SM, Kim SK, et al. Gibberellin production by pure cultures of a new strain of *Aspergillus fumigates*. World Journal of Microbiology and Biotechnology. 2009;**25**:1785-1792
- [66] Shanmugaiah V, Mathivanan N, Varghese B. Purification, crystal structure and antimicrobial activity of phenazine-1-carboxamide produced by a growth-promoting biocontrol bacterium, *Pseudomonas aeruginosa* MML2212. Journal of Applied Microbiology. 2010;**108**:703-711. DOI: 10.1111/j.1365-2672.2009.04466.x
- [67] Nagaraju A, Murali M, Sudisha J, Amruthesh KN, Murthy SM. Beneficial microbes promote plant growth and induce systemic resistance in sunflower against downy mildew disease caused by *Plasmopara halstedii*. Current Botany. 2012;**3**(5):12-18
- [68] Minerdi D, Bossi S, Maffei ME, Gullino ML, Garibaldi A. *Fusarium oxysporum* and its bacterial consortium promote lettuce growth and expansin A5 gene expression through microbial volatile organic compound (MVOC) emission. FEMS Microbiology Ecology. 2011;**76**:342-351
- [69] Mastouri F, Björkman T, Harman GE. Seed treatment with *Trichoderma harzianum* alleviates biotic, abiotic, and physiological stresses in germinating seeds and seedlings. Phytopathology. 2010;**100**:1213-1221

- [70] Mushtaq S, Nasim G, Khokhar I, Mukhtar I. Effects of *Penicillium* extracts on germination vigour in subsequent seedling growth of tomato (*Solanum lycopersicum* L.). Archives of Phytopathology and Plant Protection. 2012;**45**(8):932-937
- [71] Horinouchi H, Katsuyama N, Taguchi Y, Hyakumachi M. Control of *Fusarium* crown and root rot of tomato in a soil system by combination of a plant growth-promoting fungus, *Fusarium equiseti*, and biodegradable pots. Crop Protection. 2008;**27**:859-864
- [72] Halifu S, Deng X, Song X, Song R. Effects of two *Trichoderma* strains on plant growth, rhizosphere soil nutrients, and fungal community of *Pinus sylvestris* var. *mongolica* annual seedlings. Forests. 2019;**10**(9):758
- [73] Yadav RL, Shukla SK, Suman A, Singh PN. *Trichoderma* inoculation and trash management effects on soil microbial biomass, soil respiration, nutrient uptake and yield of ratoon sugarcane under subtropical conditions. Biology and Fertility of Soils. 2009;**45**:461-468
- [74] Radhakrishnan R, Kang S, Baek I, Lee I. Characterization of plant growth-promoting traits of *Penicillium* species against the effects of high soil salinity and root disease. Journal of Plant Interactions. 2014;**9**(1):754-762
- [75] Horinouchi H, Muslim A, Hyakumachi M. Biocontrol of *Fusarium* wilt of spinach by the plant growth promoting fungus *Fusarium equiseti* GF183. Journal of Plant Pathology. 2010;**92**(1):249-254
- [76] Gujar PD, Bhavsar KP, Khire JM. Effect of phytase from *Aspergillus niger* on plant growth and mineral assimilation in wheat (*Triticum aestivum* Linn.) and its potential for use as a soil amendment. Journal of the Science of Food and Agriculture. 2013;**93**:2242-2247
- [77] Chang YC, Baker R, Kleifeld O, Chet I. Increased growth of plants in the presence of the biological control agent *Trichoderma harzianum*. Plant Disease. 1986;**70**:145-148
- [78] Harman GE, Petzoldt R, Comis A, Chen J. Interactions between *Trichoderma harzianum* strain T22 and maize inbred line Mo17 and effects of this interaction on diseases caused by *Pythium ultimum* and *Colletotrichum graminicola*. Phytopathology. 2004;**94**:147-153
- [79] Shivanna MB, Meera MS, Hyakumachi M. Role of root colonization ability of plant growth promoting fungi in the suppression of take-all and common root rot of wheat. Crop Protection. 1996;**15**(6):497-504
- [80] Hameed K, Couch HB. Effects of *Penicillium simplicissimum* on growth, chemical composition, and root exudation of axenically grown marigolds. Phytopathology. 1971;**62**:669
- [81] Zavala-Gonzalez EA, Rodríguez-Cazorla E, Escudero N, Aranda-Martínez A, Martínez-Laborda A, Ramírez-Lepe M, et al. *Arabidopsis thaliana* root colonization by the nematophagous fungus *Pochonia chlamydosporia* is modulated by jasmonate signaling and leads to accelerated flowering and improved yield. The New Phytologist. 2016;**213**(1):351-364. DOI: 10.1111/nph.14106
- [82] Varma A, Bakshi M, Lou B, et al. *Piriformospora indica*: A novel plant growth-promoting Mycorrhizal fungus. Agricultural Research. 2012;**1**:117-131
- [83] Akhter W, Bhuiyan MKA, Sultana F, Hossain MM. Integrated effect of microbial antagonist, organic amendment and fungicide in controlling seedling mortality (*Rhizoctonia solani*) and improving yield in pea (*Pisum sativum* L.). Comptes Rendus Biologies. 2015;**338**:21-28

- [84] Hammad R, Elbagory M. Using plant growth-promoting fungi (PGPF), as a biofertilizer and biocontrol agents against *Tetranychus cucurbitacearum* on Nubian watermelon (*Citrullus lanatus* L.). *Journal of Advances in Microbiology*. 2019;**16**(2):1-15
- [85] Sneh B, Ichielevich-Auster M, Barash I, Koltin Y. Increased growth response induced by a nonpathogenic *Rhizoctonia solani*. *Canadian Journal of Botany*. 1986;**64**:2372-2378
- [86] Srivastava PK, Shenoy BD, Gupta M, Vaish A, Mannan S, Singh N, et al. Stimulatory effects of arsenic-tolerant soil fungi on plant growth promotion and soil properties. *Microbes and Environments*. 2012;**27**(4):477-482
- [87] Shivanna MB, Meera MS, Hyakumachi M. Sterile fungi from zoysiagrass rhizosphere as plant growth promoters in spring wheat. *Canadian Journal of Microbiology*. 1994;**40**:637-644
- [88] Shivanna MB, Meera MS, Kageyama K, Hyakumachi M. Influence of zoysiagrass rhizosphere fungal isolates on growth and yield of soybean plants. *Mycoscience*. 1995;**36**:25-30
- [89] Masunaka A, Hyakumachi M, Takenaka S. Plant growth-promoting fungus, *Trichoderma koningi* suppresses isoflavonoid phytoalexin vestitol production for colonization on/in the roots of *Lotus japonicus*. *Microbes and Environments*. 2011;**26**(2):128-134
- [90] Desender S, Andrivon D, Val F. Activation of defence reactions in Solanaceae: Where is the specificity? *Cellular Microbiology*. 2007;**9**:21-30
- [91] Li Z, Bai T, Dai L, Wang F, Tao J, Meng S, et al. A study of organic acid production in contrasts between two phosphate solubilizing fungi: *Penicillium oxalicum* and *Aspergillus niger*. *Scientific Reports*. 2016;**6**:25313
- [92] Wakelin SA, Gupta VVSR, Harvey PR, Ryder MH. The effect of *Penicillium* fungi on plant growth and phosphorus mobilization in neutral to alkaline soils from southern Australia. *Canadian Journal of Microbiology*. 2007;**53**:106-115
- [93] Altomare C, Norvell WA, Björkman T, Harman GE. Solubilization of phosphates and micronutrients by the plant-growth-promoting and biocontrol fungus *Trichoderma harzianum* Rifai 1295-22. *Applied and Environmental Microbiology*. 1999;**65**:2926-2933
- [94] Singh V, Singh PN, Yadav RL, Awasthi SK, Joshi BB, Singh RK, et al. Increasing the efficacy of *Trichoderma harzianum* for nutrient uptake and control of red rot in sugarcane. *Journal of Horticulture and Forestry*. 2010;**2**(4):66-71
- [95] Martínez-Medina A, Roldán A, Albacete A, Pascual JA. The interaction with arbuscular mycorrhizal fungi or *Trichoderma harzianum* alters the shoot hormonal profile in melon plants. *Phytochemistry*. 2011;**72**:223-229
- [96] Björkman T, Blanchard LM, Harman GE. Growth enhancement of shrunken-2 sweet corn with *Trichoderma harzianum* 1295-22: Effect of environmental stress. *Journal of the American Society for Horticultural Science*. 1998;**123**:35-40
- [97] Alhamed MFA, Shebany YM. Endophytic *Chaetomium globosum* enhances maize seedling copper stress tolerance. *Plant Biology*. 2012;**14**:859-863
- [98] Khan AL, Lee IJ. Endophytic *Penicillium funiculosum* LHL06 secretes gibberellin that reprograms *Glycine max* L. growth during copper stress. *BMC Plant Biology*. 2013;**13**:86
- [99] Guler NS, Pehlivan N, Karaoglu SA, Guzel S, Bozdeveci A. *Trichoderma*

atroviride ID20G inoculation ameliorates drought stress-induced damages by improving antioxidant defence in maize seedlings. *Acta Physiologiae Plantarum*. 2016;**38**:132. DOI: 10.1007/s11738-016-2153-3

[100] Naznin HA, Kiyohara D, Kimura M, Miyazawa M, Shimizu M, Hyakumachi M. Systemic resistance induced by volatile organic compounds emitted by plant growth-promoting fungi in *Arabidopsis thaliana*. *PLoS One*. 2014;**9**(1):e86882

[101] Hung R, Lee S, Bennett JW. *Arabidopsis thaliana* as a model system for testing the effect of *Trichoderma* volatile organic compounds. *Fungal Ecology*. 2013;**6**:19-26

[102] Islam MT, Hossain MM. Plant probiotics in phosphorus nutrition in crops with special reference to rice. In: Maheshwari DK, editor. *Bacteria in Agrobiolgy: Plant Probiotics*. Berlin, Heidelberg: Springer; 2012. pp. 325-363

[103] Rajankar PN, Tambekar DH, Wate SR. Study of phosphate solubilization efficiencies of fungi and bacteria isolated from saline belt of Purna river basin. *Research Journal of Agriculture and Biological Sciences*. 2007;**3**(6):701-703

[104] Gaind S, Nain L. Soil-phosphorus mobilization potential of phytate mineralizing fungi. *Journal of Plant Nutrition*. 2015;**38**(14):2159-2175. DOI: 10.1080/01904167.2015.1014561

[105] Nahas E. Factors determining rock phosphate solubilization by microorganisms isolated from soil. *World Journal of Microbiology and Biotechnology*. 1996;**12**:567-572

[106] Rousk J, Brookes PC, Bååth E. Contrasting soil pH effects on fungal and bacterial growth suggests functional redundancy in carbon mineralisation. *Applied*

and Environmental Microbiology. 2009;**75**:1589-1596

[107] Hoyos-Carvajal L, Orduz S, Bissett J. Growth stimulation in bean (*Phaseolus vulgaris* L.) by *Trichoderma*. *Biological Control*. 2009;**51**:409-416

[108] Ruess L, Ferris H. Decomposition pathways and successional changes. In: *Proceeding of the Fourth International Congress of Nematology*. Vol. 2. 2004. pp. 547-556

[109] Hyakumachi M. *Microorganism Resources: Its Characteristics and Utilization*. Tokyo: Soft Science Inc.; 2000. pp. 81-92

[110] Waqas M, Khan AL, Kamran M, Hamayun M, Kang SM, Kim YH, et al. Endophytic fungi produce gibberellins and indole acetic acid and promotes host-plant growth during stress. *Molecules*. 2012;**17**:10754-10773

[111] Foo E, Ross JJ, Jones WT, Reid JB. Plant hormones in arbuscular mycorrhizal symbioses: An emerging role for gibberellins. *Annals of Botany*. 2013;**111**:769-779

[112] Vadassery J, Ritter C, Venus Y, Camehl I, Varma A, Shahollari B, et al. The role of auxins and cytokinins in the mutualistic interaction between *Arabidopsis* and *Piriformospora indica*. *Molecular Plant-Microbe Interactions*. 2008;**21**:1371-1383

[113] Saxena S. *Applied Microbiology*. India: Springer Pvt. Ltd; 2015. p. 190

[114] Nascimento FX, Rossi MJ, Soares CRFS, McConkey BJ, Glick BR. New insights into 1-aminocyclopropane-1-carboxylate (ACC) deaminase phylogeny, evolution and ecological significance. *PLoS One*. 2014;**9**(6):e99168

[115] Glick BR, Cheng Z, Czarny J, Duan J. Promotion of plant growth by ACC deaminase-producing soil bacteria.

European Journal of Plant Pathology.
2007;**119**:329-339

[116] Glick BR. Bacteria with ACC deaminase can promote plant growth and help to feed the world. *Microbiological Research*. 2014;**169**:30-39

[117] Viterbo A, Landau U, Kim S, Chernin L, Chet I. Characterization of ACC deaminase from the biocontrol and plant growth-promoting agent *Trichoderma asperellum* T203. *FEMS Microbiology Letters*. 2010;**305**:42-48

[118] Palmer C, Golden K, Danniels L, Ahmad H. ACC deaminase from *Issatchenkia occidentalis*. *Journal of Biological Sciences*. 2007;**7**:188-193

[119] Jia YJ, Kakuta Y, Sugawara M, Igarashi T, Oki N, Kisaki M, et al. Synthesis and degradation of 1-aminocyclopropane-1-carboxylic acid by *Penicillium citrinum*. *Bioscience, Biotechnology, and Biochemistry*. 1999;**63**:542-549

[120] Singh N, Kashyap S. *In-silico* identification and characterization of 1-aminocyclopropane-1-carboxylate deaminase from *Phytophthora sojae*. *Journal of Molecular Modeling*. 2012;**18**:4101-4111

[121] Whipps JM. Microbial interactions and biocontrol in rhizosphere. *Journal of Experimental Botany*. 2001;**52**:487-511

[122] Harveson RM, Kimbrough JW, Hopkins DL. Novel use of a Pyrenomycetous mycoparasite for management of *Fusarium* wilt of watermelon. *Plant Disease*. 2002;**86**(9):1025-1030

[123] Koike N, Hyakumachi M, Kageyama K, Tsuyumu S, Doke N. Induction of systemic resistance in cucumber against several diseases by plant growth-promoting fungi: Lignification and superoxide

generation. *European Journal of Plant Pathology*. 2001;**107**:523-533

[124] Hossain MM, Sultana F. Genetic variation for induced and basal resistance against leaf pathogen *Pseudomonas syringae* pv. *tomato* DC3000 among *Arabidopsis thaliana* accessions. *Springerplus*. 2015;**4**:296. DOI: 10.1186/s40064-015-1070-z

[125] Khan AL, Hamayun M, Kim YH, Kang SM, Lee IJ. Ameliorative symbiosis of endophyte (*Penicillium funiculosum* LHL06) under salt stress elevated plant growth of glycine max L. *Plant Physiology and Biochemistry*. 2011;**49**:852-862

[126] Bae H, Sicher RC, Kim MS, Kim SH, Strem MD, Melnick RL, et al. The beneficial endophyte *Trichoderma hamatum* isolate DIS 219b promotes growth and delays the onset of the drought response in *Theobroma cacao*. *Journal of Experimental Botany*. 2009;**60**:3279-3295

[127] Ahmad P, Hashem A, Abd-Allah EF, Alqarawi AA, John R, Egamberdieva D, et al. Role of *Trichoderma harzianum* in mitigating NaCl stress in Indian mustard (*Brassica juncea* L.) through antioxidative defense system. *Frontiers in Plant Science*. 2015;**6**:868

[128] Li B, Li Q, Xiong L, Kronzucker HJ, Krämer U, Shi W. *Arabidopsis* plastid AMOS1/EGY1 integrates abscisic acid signaling to regulate global gene expression response to ammonium stress. *Plant Physiology*. 2012;**160**:2040-2051

[129] Rasool S, Ahmad A, Siddiqi TO, Ahmad P. Changes in growth, lipid peroxidation and some key antioxidant enzymes in chickpea genotypes under salt stress. *Acta Physiologiae Plantarum*. 2013;**35**:1039-1050

[130] Kanchiswamy CN, Malnoy M, Maffei ME. Chemical diversity of

- microbial volatiles and their potential for plant growth and productivity. *Frontiers in Plant Science*. 2015;**6**:151. DOI: 10.3389/fpls.2015.00151
- [131] Farag MA, Ryu CM, Sumner LW, Pare PW. GC-MS SPME profiling of rhizobacterial volatiles reveals prospective inducers of growth promotion and induced systemic resistance in plants. *Phytochemistry*. 2006;**67**:2262-2268
- [132] Naznin HA, Kimura M, Miyazawa M, Hyakumachi M. Analysis of volatile organic compounds emitted by plant growth-promoting fungus *Phoma* sp. GS8-3 for growth promotion effects on tobacco. *Microbes and Environments*. 2013;**28**(1):42-49
- [133] Meera MS, Shivanna MB, Kageyama K, Hyakumachi M. Plant growth promoting fungi from zoysiagrass rhizosphere as potential inducers of systemic resistance in cucumbers. *Phytopathology*. 1994;**84**:1399-1406
- [134] Rovira AD, Newman EI, Bowen HJ, Campbell R. Quantitative assessment of the rhizoplane microflora by direct microscopy. *Soil Biology and Biochemistry*. 1974;**6**:211-216
- [135] Bowen GD. Microbial dynamics in the rhizosphere: Possible strategies in managing rhizosphere populations. In: Keister DL, Cregan PB, editors. *The Rhizosphere and Plant Growth*. Dordrecht: Kluwer; 1991. pp. 25-32
- [136] Chacón MR, Rodríguez-Galán O, Benítez T, Sousa S, et al. Microscopic and transcriptome analyses of early colonization of tomato roots by *Trichoderma harzianum*. *International Microbiology*. 2007;**10**(1):19-27
- [137] Dewan MM, Sivasithamparam K. Effect of colonization by a sterile fungus on viability of seed and growth and anatomy of roots. *Mycological Research*. 1990;**94**:553-557
- [138] Singh PC, Nautiyal CS. A novel method to prepare concentrated conidial biomass formulation of *Trichoderma harzianum* for seed application. *Journal of Applied Microbiology*. 2012;**113**:1142-1450. DOI: 10.1111/j.1365-2672.2012.05426.x
- [139] Doni F, Isahak A, Zain CRCM, Ariffin SM, Mohamad WNW, Yusoff WMW. Formulation of *Trichoderma* sp. SL2 inoculants using different carriers for soil treatment in rice seedling growth. *Springerplus*. 2014;**3**:532. DOI: 10.1186/2193-1801-3-532
- [140] Oancea F, Raut J, Şesan TE, Cornea PC. Dry flowable formulation of biostimulants *Trichoderma* strains. *Agriculture and Agricultural Science Procedia*. 2016;**10**:494-502
- [141] Zhang FG, Meng XH, Feng CL, Ran W, Yu GH, Zhang YJ, et al. Hydrolytic amino acids employed as a novel organic nitrogen source for the preparation of PGPF-containing bio-organic fertilizer for plant growth promotion and characterization of substance transformation during BOF production. *PLOS One*. 2016;**11**:e0149447
- [142] Zhang FG, Huo YQ, Cobb AB, Luo GW, Zhou JQ, Yang GW, et al. *Trichoderma* biofertilizer links to altered soil chemistry, altered microbial communities, and improved grassland biomass. *Frontiers in Microbiology*. 2018;**9**:848
- [143] Sarma MVRK, Kumar V, Saharan K, Srivastava R, Sharma AK, Prakash A, et al. Application of inorganic carrier-based formulations of fluorescent pseudomonads and *Piriformospora indica* on tomato plants and evaluation of their efficacy. *Journal of Applied Microbiology*. 2011;**111**:456-466