

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

5,000

Open access books available

125,000

International authors and editors

140M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



# L-Arginine/Nitric Oxide Pathway and KCa Channels in Endothelial Cells: A Mini-Review

Marcelo González and José Carlos Rivas

## Abstract

The endothelium is an organ with a key role in the maintenance of cardiovascular health through the regulation of vascular tone, vascular resistance, blood flow, and arterial pressure. These functions are related with the synthesis and release of vasoactive molecules, mainly vasodilators like nitric oxide (NO) and endothelium-derived hyperpolarizing factor (EDHF). Both factors are released and diffused from endothelial cells to the smooth muscle cells, where there is a subsequent activation of signaling pathways that finally decrease the intracellular calcium to induce the vascular relaxation. The study of the molecular mechanisms that underlie the endothelial function still is in development, but from the evidence obtained from the endothelial cells *in vitro* studies are possible to partially describe the pathways to regulate the physiological endothelial function and the disturbances in pathological conditions. In this mini-review, we describe the main mechanisms for NO synthesis and the role of potassium channels related with EDHF. We include schemes and graphical summaries for better understanding of the molecular regulation of vascular tone in the human cardiovascular system.

**Keywords:** L-arginine, nitric oxide, potassium channels, endothelium


## 1. Characteristics of the endothelium

The endothelial cells (ECs) have mesenchymal origin, length of 25–50  $\mu\text{m}$  and form a flat epithelium called endothelium. The endothelium in a human adult is composed of approximately  $1-6 \times 10^{13}$  cells, constituting an organ that weighs approximately 1 kg and covers a surface area of approximately  $1-7 \text{ m}^2$  [1]. For decades, the endothelium was considered as a simple barrier between blood and the rest of the body's tissues. However, since the early 1980s, this vision changed radically [2] and, today, the endothelium is considered a true organ that fulfills multiple functions in the physiology and pathophysiology of vascular system, including autocrine, paracrine, and endocrine actions and the regulation of coagulation and fibrinolysis processes [3].

One of the most important functions of endothelial cells is their participation in the regulation of vascular tone. In the classic article of Furchgott and Zawadzki in 1980, it was demonstrated that the presence of the endothelium is essential for the vasodilator effect induced by acetylcholine in isolated blood vessels pre-constricted with norepinephrine. In those years, it was proposed that the vasodilation was

produced through a factor that was released by the endothelium in response to agonists [4]. This factor was called the endothelial-derived relaxing factor (EDRF) [5]. Between 1986 and 1990, it was concluded that this factor corresponded to nitric oxide (NO) [6, 7]. The endothelium responds to mechanical stimuli such as pressure and flow stress (“shear stress”), hormonal stimuli, and vasoactive substances that regulate the vascular tone. The endothelial cells release molecules that regulate vasomotor function, inflammation, and hemostasis. Vasodilators agents include NO, prostacyclin, and endothelium-derived hyperpolarizing factor (EDHF). Vasoconstrictors agents include endothelin 1, angiotensin II, thromboxane A<sub>2</sub>, and reactive oxygen-derived species (ROS). Inflammatory mediators include NO, intercellular adhesion molecule 1 (ICAM-1), vascular adhesion molecule 1 (VCAM-1), E-selectin, and NFκB (Figure 1 [8]).

Since the discovery of NO, the mechanisms of endothelial cell activation and endothelial dysfunction have been studied. In this way, the quiescent endothelial cells express a vasodilator, anticoagulant, and anti-adhesive phenotype, while the activated endothelial cell expresses procoagulant, pro-adhesive, and vasoconstrictive properties [9]. It has been considered that the decrease in the capacity of the vascular endothelium to stimulate vasodilation generates endothelial dysfunction, a phenomenon that is observed in several pathological conditions such as hypertension, hypercholesterolemia, diabetes mellitus, hyperhomocysteinemia, chronic kidney failure, chronic heart failure, etc. Although the molecular basis for endothelial dysfunction is not fully understood, numerous studies point to decreased biosynthesis and/or NO activity as a central mechanism [10–13].




**Figure 1.**

*Vascular tone regulation. The vascular tone is partially regulated by the local factors secreted by endothelial cells (ECs) in response to physical factors like shear stress and humoral and chemical factors like hormones and oxygen levels. The changes in blood flow are detected by membrane proteins, mainly receptors (Rs), transporters (Ts), and ion channels (ICs). There is a network connecting the activities of these proteins through signaling pathways that induce the release of different mediators like thromboxane A<sub>2</sub> (TxA<sub>2</sub>), endothelin 1, reactive oxygen species (ROS), nitric oxide (NO), endothelium-derived hyperpolarizing factor (EDHF), or prostacyclin (PGI<sub>2</sub>), among others. The equilibrium between the vasoconstrictors and vasodilators factors maintains the endothelial function and vascular health.*

## 2. Synthesis of nitric oxide in the endothelium

NO is synthesized from the semi-essential cationic amino acid L-arginine, which must be transported from the extracellular space into the endothelial cell by a family of cationic amino acid transporters (CATs) [14]. This amino acid is the substrate in a reaction where the metabolic product corresponds to L-citrulline in an equimolar proportion with the coproduct NO [15, 16]. This reaction is catalyzed by the enzyme NO synthase (NOS), which can be classified into their constitutive forms (cNOS) and their inducible form (iNOS) [17]. The cNOS includes the endothelial isoform (eNOS) and the neuronal isoform (nNOS), both producing NO in short bursts at low concentrations (nM) and in a calcium-dependent manner to fulfill the physiological functions of NO. The physiological activity of eNOS is dependent on several cofactors and is regulated by signaling pathways that induce phosphorylation in different sites for activation (serine 1177) or inhibition (threonine 495) [17]. NO diffuses from endothelial cells to smooth muscle cells (SMCs) and activates the soluble guanylate cyclase (sGC) pathway, to reduce the intracellular calcium and induce vasodilation (**Figure 2**). iNOS is mainly expressed in cells that participate in the inflammatory response after induction by cytokines and other inflammatory mediators, producing NO in high concentrations ( $\mu\text{M}$ ) and independently of calcium [18–20].

The availability of NO *in vivo* is regulated by a combination of NO synthesis and inactivation. The decrease in the availability of NO may be due to a lower expression



**Figure 2.**

*L-arginine transport and nitric oxide synthesis in endothelial cells. hCAT-1 is a protein expressed in plasma membrane of endothelial cells, mainly in plasma membrane invagination called caveolae. The L-arginine enters to the cell from blood and is used by eNOS to synthesize L-citrulline and nitric oxide (NO). The eNOS needs different cofactors to maintain its function, which include tetrahydrobiopterin (BH<sub>4</sub>), nicotinamide adenine dinucleotide phosphate (NADPH), and heat shock protein 90 (Hsp90). Nitric oxide diffuses through the cell membranes and enters the smooth muscle cells to activate the soluble guanylate cyclase (sGC). The sGC synthesizes cyclic GMP (cGMP), which activates protein kinase G and, after subsequent steps, the intracellular calcium decreases to induce the vasodilation.*

or activity of eNOS, as a result of the action of endogenous and exogenous inhibitors or due to the lower availability of the substrate L-arginine [8, 14]. The availability of NO can also be diminished by the rapid reaction between NO and reactive oxygen-derived species (ROS) [13].

### 3. Reactive oxygen-derived species (ROS) in endothelium

Endothelial cells generate ROS, including the superoxide radical ( $O_2^{\cdot-}$ ), hydrogen peroxide ( $H_2O_2$ ), peroxynitrite ( $ONOO^-$ ), hydroxyl radical ( $\cdot OH$ ), among others [15, 16]. In endothelial cells, the main sources of ROS are the enzymatic complex xanthine oxidoreductase (XOR) [17], the complex of membrane nicotinamide adenine dinucleotide phosphate oxidase (NADPH oxidase) [18], eNOS itself when it is “uncoupled” due to lack of tetrahydrobiopterin ( $BH_4$ ) or L-arginine [19], mitochondrial cytochromes [20], and hemoglobin [21].

Among all endothelial ROS sources, NADPH oxidases are enzymes whose primary function is the generation of ROS and they play an important role in redox signaling [22]. On the other hand, the activity of NADPH oxidase can cause the uncoupling of eNOS by the oxidative degradation of  $BH_4$ , leading to the eNOS-dependent synthesis of  $O_2^{\cdot-}$  and detriment of the synthesis of NO [18, 23]. Once  $O_2^{\cdot-}$  is synthesized, it can act as a precursor to other ROS due to its use by superoxide dismutase (SOD) to generate  $H_2O_2$  that has greater stability and capacity to cross biological membranes, and it therefore can act as a modulator of signal transduction pathways [24]. Furthermore,  $O_2^{\cdot-}$  reacts quickly with NO to generate  $ONOO^-$ , a powerful oxidizing agent that causes DNA fragmentation and lipid oxidation [25].

It is currently postulated that the mechanism by which  $O_2^{\cdot-}$  “kidnaps” NO would play a central role in the development of endothelial dysfunction that is seen in pathologies such as diabetes mellitus [26–28], preeclampsia [29, 30], and hypertension [31].

### 4. L-arginine transport in human fetal endothelium

The amino acid L-arginine is taken up by endothelial cells through the transporter systems  $y^+$ ,  $y^+L$ ,  $b^{0,+}$ , and  $yB^{0,+}$  [32–35]. Of these systems, there are two that have been described in HUVEC, that is,  $y^+$  system [36–38] and  $y^+L$  system [39]. The  $y^+$  system family is currently known to include at least five cationic amino acid transporters (CATs) called CAT-1, CAT-2A, CAT-2B, CAT-3, and CAT-4. CAT-1 is expressed ubiquitously, CAT-2A and CAT-3 are constitutively expressed in liver and brain, respectively, while CAT-2B is induced in a variety of cell types in response to bacterial endotoxins and pro-inflammatory cytokines [40, 41]. CAT-4 corresponds to a cDNA sequence with 41–42% identity with the other members of the CATs family, but its transport activity has not yet been determined [32, 34, 35]. CAT-1, CAT-2B, and CAT-3 are characterized by high affinity to the substrate ( $K_m = 100\text{--}400\ \mu M$ ) and independency of  $Na^+$ , while CAT-2A has low affinity for cationic amino acids ( $K_m = 2\text{--}5\ mM$ ). Two members of CATs have been reported to be expressed in HUVEC, that is, hCAT-1 and hCAT-2B, while hCAT-2A and hCAT-3 transporters have not been detected in this cell type [34, 36–39] (**Table 1**). Although the hCAT-1 and hCAT-2B transporters have similar kinetic characteristics, it is possible to differentiate them by their different sensitivities to L-lysine trans-stimulation. In *Xenopus laevis* oocytes injected with hCAT-1 and hCAT-2B mRNA, L-lysine increases L-arginine transport by 9.8-fold and 1.8-fold, respectively [42]. Thus, for L-lysine trans-stimulation assays in HUVEC, it has been possible to determine that

Gene	Protein	K <sub>m</sub> (μM)	Distribution
<i>SLC7A1</i>	CAT-1	70–250	All tissues except liver and lacrimal gland
<i>SLC7A2</i>	CAT-2A	2.2–5.2	Liver, skeletal muscle, and pancreas
<i>SLC7A2</i>	CAT-2B	38–380	Endothelium, and inducible in several tissues
<i>SLC7A3</i>	CAT-3	40–120	Thymus, ovary, testes, and brain
<i>SLC7A4</i>	CAT-4	—	Brain, testes, and placenta

*Proteins CATs are coded in different genes (except CAT2A and 2B, same gene), have different kinetic constants for the transport of L-arginine (K<sub>m</sub>) and distribution in tissues.*

**Table 1.**  
 CATs' family members.

the hCAT-1 transporter accounts for 60–80% of the total uptake of L-arginine in physiological conditions [36–38]. The importance of the hCAT-1 transporter in NO synthesis has been confirmed through a transgenic mouse model that overexpresses the protein exclusively in the endothelium. Aortic rings obtained from these transgenic mice have a higher sensitivity to relaxation in response to acetylcholine compared to native mice, while endothelial cell cultures obtained from these animals, that overexpress hCAT-1, exhibit a greater NO synthesis [43].

## 5. Regulation of the expression of hCAT-1

Regarding the gene organization of CAT transporters, it is known that the *SLC7* family is phylogenetically composed of two subfamilies formed by cationic amino acid transporters (CATs) and glycoprotein-associated amino acid transporters (HATs). The cationic amino acid transporter family is encoded by the *SLC7A* (1–4) genes and corresponds to proteins with 14 transmembrane domains [44]. Specifically, the gene that encodes the hCAT-1 protein corresponds to *SLC7A1* whose open reading frame is formed by 11 exons and 10 introns. The gene is located on chromosome 13q12-13q14 [45].

Among the genes encoding CAT-1 in rat, mouse and human have common characteristics: the promoter region lacks TATA box, and they have multiple binding sites for the transcription factor specific protein 1 (Sp1) and they have an extensive 3' non-translatable region (3' UTR) that could perform functions in the regulation of mRNA stability or in translation [46–49]. In rats, stress by amino acids deprivation induces an increase in the rCAT-1 mRNA expression by a mechanism related to increased mRNA stability [46]. This increased mRNA stability would be related to the presence of a regulatory region within the 3' UTR sequence of the gene [47]. Subsequent experiments have shown that the effect of amino acids deprivation on rCAT-1 expression would depend on both transcriptional [48] and posttranscriptional mechanisms [50].

In humans, it is known that insulin increases leg blood flow in healthy subjects via stimulation of endothelial NO synthase (eNOS) [51]. Insulin also increases the synthesis and release of NO and release in primary cultures of HUVEC [38, 52]. Biological effects of insulin involve activation of several transcription factors, including Sp1 in several cell types [53, 54]. Insulin increases Sp1 nuclear protein abundance and its binding to a proximal region (-177 and -105 bp from ATG) of the *SLC7A1* promoter containing four consensus sequences for Sp1 [55]. Interestingly, in patients with essential hypertension, a reduction of *SLC7A1* transcriptional activity due to reduced Sp1 activity in the promoter region has been reported [12]. So, the transcriptional regulation of *SLC7A1* is relevant for cardiovascular physiology,

and the reduction of the promoter activity of this gene could be associated with cardiovascular disease (CVD).

On the other hand, the first intron of *SLC7A1* may play a bifunctional role in regulating the *SLC7A1* transcriptional activity by the binding of the purine-rich element binding protein A (Pur alpha) in physiological conditions and by activating the transcription factor 4 (ATF4) in endoplasmic reticulum stress or by decreasing the *SLC7A1* transcriptional activity by the C/EBP homologous protein 10 (CHOP) binding in C6 rat glioma cells [56].


For the physiological regulation of hCAT-1 activity, both transcriptional regulation of *SLC7A1* and/or posttranscriptional regulation of *SLC7A1* transcript are relevant for the protein expression and L-arginine transport [55]. Insulin increases the expression of *SLC7A1* gene due to an increased transcriptional activity, most likely due to higher Sp1 activity. So, hCAT-1 expression and activity are regulated by insulin in endothelium, suggesting that in insulin resistance there is a reduction of L-arginine transport and NO synthesis that contributes to endothelial dysfunction and cardiovascular diseases.

## **6. High D-glucose and expression and activity of L-arginine/NO pathway**

Hyperglycemia and diabetes mellitus are pathological conditions associated with fetal endothelial dysfunction [55] and type 2 diabetes mellitus (T2DM) [57] or cardiovascular disease (CVD) [58]. CVD in patients with diabetes mellitus is associated with the generation of ROS.

High concentration of D-glucose (25 mM) increases L-arginine transport and cGMP accumulation in endothelium in a similar manner to that observed in HUVEC from pregnancies with gestational diabetes [33, 59]. Increased L-arginine transport in response to incubation with high D-glucose has been related to increased mRNA levels for the hCAT-1 and eNOS activity in HUVEC [60]. In human aortic endothelial cells, prolonged incubation (7 days) with 25 mM D-glucose induces a decrease in eNOS activity (determined by nitrite content), protein abundance, and mRNA level. This effect is associated with a decrease in eNOS promoter activity [61]. In bovine aortic endothelial cells (BAECs), there is a lower production of insulin-induced NO when the cells were incubated with high extracellular concentration of D-glucose, an effect that seems to depend on a signaling pathway that involves the type 1 insulin receptor (IR-1), phosphatidylinositol 3 kinase, and the inhibitor of nuclear factor kappa-B kinase [62]. On the other hand, the increase of cGMP production induced by high D-glucose in HUVEC is blocked by incubating the cells with 1 nM insulin [63]. Incubation with 1 nM insulin (8 h) has been shown in this same cell type to be sufficient to block the effect that D-glucose has on the decreased transport of adenosine [64], an important vasoactive nucleoside [65].

In HUVEC, high extracellular D-glucose increases L-arginine transport, NO synthesis, and  $O_2^-$  generation through eNOS and NADPH oxidase activation. Additionally, high D-glucose increased the contractile response in the human umbilical vein. Insulin reversed these effects of high D-glucose, leading to normal hCAT-1 expression, NO synthesis, ROS generation, and vascular tone. Insulin acts like antioxidant molecules (like tempol, ascorbic acid) to restore high D-glucose-increased oxidative stress in the fetoplacental vascular bed [66]. High D-glucose increases L-arginine transport, likely resulting from higher hCAT-1 expression and protein abundance in the plasma membrane. This mechanism could be an adaptive response of HUVEC to higher ROS generation from high D-glucose-activated




**Figure 3.** Endothelial dysfunction induced by high D-glucose and protection by insulin in HUVEC. Exposure of HUVEC to high D-glucose leads to an increase ( $\uparrow$ ) in the plasma membrane abundance of the human cationic amino acid transporter 1 (hCAT-1) and higher L-arginine uptake. High D-glucose activates NADPH oxidase, leading to higher generation of ROS, including  $O_2^{\cdot -}$ . Insulin restores ROS and  $O_2^{\cdot -}$  generation to values in cells exposed to 5 mM D-glucose (normal), resulting in the restoration of hCAT-1-mediated L-arginine transport and nitric oxide (NO) synthesis. High D-glucose and insulin also activate the SLC7A1 promoter region (coding for hCAT-1) up to -650 bp from the ATG via a mechanism involving ROS and  $O_2^{\cdot -}$  generation. In addition, insulin restores hCAT-1 protein abundance and its distribution in the cells via an NADPH oxidase-independent mechanism (data from González et al. [66]).

NADPH oxidase. In parallel, high D-glucose increased NO synthesis. Insulin reversed the high D-glucose-mediated alterations in L-arginine transport involving the modulation of SLC7A1 gene expression, leading to altered umbilical vein reactivity. Modulation of hCAT-1 expression and activity by insulin is the key to maintaining umbilical vein tone and endothelial function in physiologic and pathophysiological conditions (Figure 3) [66].

## 7. Role of potassium channels in endothelial function

Another important mechanism that regulates the endothelial function is the activity of ion channels that modulate the cell membrane potential. The calcium-activated potassium channels (KCa) have been shown to be relevant to induce the necessary hyperpolarization to stimulate the relaxation of vascular smooth muscle cells (related with EDHF). In systemic circulation, large conductance KCa (BKCa) channels have been shown preferentially expressed in VSMC, meanwhile small (SKCa) and intermediate (IKCa) conductance KCa are preferentially expressed in endothelium [67, 68]. However, potassium currents inhibited by iberiotoxin (BKCa inhibitor) have been described in HUVEC stimulated by sildenafil or insulin [69]. In fact, insulin (10 nM) can directly activate native and recombinant BKCa currents in cell-attached patch-clamping experiments with a rapid effect that is MAPK-dependent when the hormone was added in the pipette [70]. There is evidence that insulin may induce endothelial cell hyperpolarization by modulating K channels activity [38, 71]. The insulin-induced relaxation in human placental veins (~368  $\mu$ m diameter), pre-constricted with U46619, is a mechanism dependent on the BKCa channel activity. The co-incubation of vessels with genistein (tyrosine kinases inhibitor) and wortmannin (PI3K inhibitor) did not block the insulin's relaxation, and by contrast potentiated the insulin-induced






**Figure 4.**

*Proposal of mechanism for KCa activation by insulin. Evidence obtained in endothelial cells (ECs) shows that insulin activates KCa (mainly BKCa) in a mechanism still not fully understood. The activation of KCa by insulin induces hyperpolarization ( $\uparrow\Delta V$ ), leading to activation of eNOS for NO synthesis from L-arginine uptake by hCAT-1 (modified from Rojas et al. [71]).*

vasodilation. Also, insulin decreased perfusion pressure ( $34 \pm 3\%$ ) in the isolated cotyledon of normal placenta with a basal perfusion pressure of  $64 \pm 5$  mmHg (or pre-constricted with U46619) [72]. The effects of insulin on BKCa activity are associated with evidences that show that the constriction induced by U46619 and  $H_2O_2$  in placental vasculature is partially decreased with 10 nM insulin preincubation (10 min) in a mechanism totally dependent of BKCa activity [72]. Recently, it has been determined that insulin-mediated NO synthesis requires the participation of both IKCa/ BKCa channels and eNOS activity in HUVECs [71]. In the same cell type, insulin increased the open probability (NPo) of BKCa, associated with hyperpolarization in single cell analysis [69]. In human placental arteries, the relaxation induced by the NO donor, SNAP, is partially blocked by charybdotoxin (BKCa inhibitor) and almost totally blocked by charybdotoxin and ODQ (sGC inhibitor) [73]. Therefore, an extracellular stimulus that increases the NO availability activates a mechanism that involves sGC and BKCa activities [71]. These findings constitute evidence for postulating a new mechanism induced by insulin in human vasculature related with the physiological regulation of KCa activity for NO synthesis (**Figure 4**).

## 8. Final remarks

The relevance of the endothelium for cardiovascular physiology is well established, mainly by findings related to the capacity of endothelial cells to synthesize NO and regulate the plasma membrane potential of smooth muscle cells. **Figure 5** shows a graphical summary of the L-arginine/NO pathway in the human blood vessels that highlight the capacity of endothelial cells to respond to extracellular stimuli and translate the mechanical forces and endocrine signals to intracellular mechanisms leading to NO synthesis and activation of potassium channels. It is important to note that the subcellular distribution of hCAT-1 and eNOS is also relevant for endothelial cells function. In physiological state, hCAT-1 colocalizes with caveolin-1 in the plasma membrane caveolae in proximity to eNOS.



**Figure 5.**  
 Role of endothelium in the regulation of vascular tone. Endothelial cells, as a part of blood vessels walls, respond to mechanical stress induced by flow (shear stress) by activation of L-arginine/NO pathway to induce the NO release and relaxation of smooth muscle cells (SMCs). Subcellular localization of hCAT-1 in caveolae is relevant for its function, and the role of potassium channels (BKCa, mainly) has been recently described as important for endothelial cells function. The activity of the endothelium is regulated by different agonists like acetylcholine (Ach) through plasma membrane receptor (AchR) and others like insulin or serotonin, etc.

## Acknowledgements

The authors would like to acknowledge the staff at Laboratorio de Investigación Materno-Fetal (LIMaF) and Department of Obstetrics and Gynecology from the Universidad de Concepción. The authors express special thanks to Alexandra Elbakyan for support the open science.

IntechOpen

IntechOpen

### **Author details**

Marcelo González\* and José Carlos Rivas  
Laboratorio de Investigación Materno Fetal (LIMaF), Departamento de Obstetricia  
y Ginecología, Facultad de Medicina, Universidad de Concepción, Concepción,  
Chile

\*Address all correspondence to: [mgonzalezo@udec.cl](mailto:mgonzalezo@udec.cl)

### **IntechOpen**

---

© 2020 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

## References

- [1] Augustin HG, Kozián DH, Johnson RC. Differentiation of endothelial cells: Analysis of the constitutive and activated endothelial cell phenotypes. *BioEssays: News and Reviews in Molecular, Cellular and Developmental Biology*. 1994;**16**(12):901-906
- [2] Fishman AP. Endothelium: A distributed organ of diverse capabilities. *Annals of the New York Academy of Sciences*. 1982;**401**:1-8
- [3] Galley HF, Webster NR. Physiology of the endothelium. *British Journal of Anaesthesia*. 2004;**93**(1):105-113
- [4] Furchgott RF, Zawadzki JV. The obligatory role of endothelial cells in the relaxation of arterial smooth muscle by acetylcholine. *Nature*. 1980;**288**(5789):373-376
- [5] Griffith TM, Edwards DH, Lewis MJ, Newby AC, Henderson AH. The nature of endothelium-derived vascular relaxant factor. *Nature*. 1984;**308**(5960):645-647
- [6] Ignarro LJ, Buga GM, Wood KS, Byrns RE, Chaudhuri G. Endothelium-derived relaxing factor produced and released from artery and vein is nitric oxide. *Proceedings of the National Academy of Sciences of the United States of America*. 1987;**84**(24):9265-9269
- [7] Moncada S, Palmer RM, Higgs EA. The discovery of nitric oxide as the endogenous nitrovasodilator. *Hypertension (Dallas, Tex.: 1979)*. 1988;**12**(4):365-372
- [8] Endemann DH, Schiffrin EL. Endothelial dysfunction. *Journal of the American Society of Nephrology: JASN*. 2004;**15**(8):1983-1992
- [9] Aird WC. Endothelium in health and disease. *Pharmacological Reports*. 2008;**60**(1):139-143
- [10] De Meyer GR, Herman AG. Vascular endothelial dysfunction. *Progress in Cardiovascular Diseases*. 1997;**39**(4):325-342
- [11] Kurowska EM. Nitric oxide therapies in vascular diseases. *Current Pharmaceutical Design*. 2002;**8**(3):155-166
- [12] Yang Z, Kaye DM. Endothelial dysfunction and impaired L-arginine transport in hypertension and genetically predisposed normotensive subjects. *Trends in Cardiovascular Medicine*. 2006;**16**(4):118-124
- [13] González M. Regulation of expression and activity of l-arginine transporters by nutrients and hormones: A focus in transcriptional mechanisms regulated by glucose and insulin. En: Patel VB, Preedy VR, Rajendram R, editores. *L-Arginine in Clinical Nutrition (Nutrition and Health)*. Cham: Springer International Publishing; 2017. pp. 71-83. DOI: 10.1007/978-3-319-26009-9\_6 [Accessed: 12 July 2020]
- [14] Kalinowski L, Malinski T. Endothelial NADH/NADPH-dependent enzymatic sources of superoxide production: Relationship to endothelial dysfunction. *Acta Biochimica Polonica*. 2004;**51**(2):459-469
- [15] Förstermann U. Oxidative stress in vascular disease: Causes, defense mechanisms and potential therapies. *Nature Clinical Practice. Cardiovascular Medicine*. 2008;**5**(6):338-349
- [16] Valko M, Leibfritz D, Moncol J, Cronin MTD, Mazur M, Telser J. Free radicals and antioxidants in normal physiological functions and human disease. *The International Journal of Biochemistry & Cell Biology*. 2007;**39**(1):44-84

- [17] Berry CE, Hare JM. Xanthine oxidoreductase and cardiovascular disease: Molecular mechanisms and pathophysiological implications. *The Journal of Physiology*. 2004;555(Pt 3): 589-606
- [18] Dworakowski R, Alom-Ruiz SP, Shah AM. NADPH oxidase-derived reactive oxygen species in the regulation of endothelial phenotype. *Pharmacological Reports*. 2008;60(1):21-28
- [19] Münzel T, Daiber A, Ullrich V, Mülsch A. Vascular consequences of endothelial nitric oxide synthase uncoupling for the activity and expression of the soluble guanylyl cyclase and the cGMP-dependent protein kinase. *Arteriosclerosis, Thrombosis, and Vascular Biology*. 2005;25(8):1551-1557
- [20] Cadenas E, Sies H. The lag phase. *Free Radical Research*. 1998;28(6):601-609
- [21] Balla J, Vercellotti GM, Nath K, Yachie A, Nagy E, Eaton JW, et al. Haem, haem oxygenase and ferritin in vascular endothelial cell injury. *Nephrology, Dialysis, Transplantation: Official Publication of the European Dialysis and Transplant Association - European Renal Association*. 2003;18(Suppl 5):v8-v12
- [22] Lambeth JD. NOX enzymes and the biology of reactive oxygen. *Nature Reviews. Immunology*. 2004;4(3):181-189
- [23] Antoniadis C, Shirodaria C, Warrick N, Cai S, de Bono J, Lee J, et al. 5-methyltetrahydrofolate rapidly improves endothelial function and decreases superoxide production in human vessels: Effects on vascular tetrahydrobiopterin availability and endothelial nitric oxide synthase coupling. *Circulation*. 2006;114(11):1193-1201
- [24] Li J-M, Shah AM. Endothelial cell superoxide generation: Regulation and relevance for cardiovascular pathophysiology. *American Journal of Physiology. Regulatory, Integrative and Comparative Physiology*. 2004;287(5):R1014-R1030
- [25] Carr AC, McCall MR, Frei B. Oxidation of LDL by myeloperoxidase and reactive nitrogen species: Reaction pathways and antioxidant protection. *Arteriosclerosis, Thrombosis, and Vascular Biology*. 2000;20(7):1716-1723
- [26] Rolo AP, Palmeira CM. Diabetes and mitochondrial function: Role of hyperglycemia and oxidative stress. *Toxicology and Applied Pharmacology*. 2006;212(2):167-178
- [27] Hadi HAR, Suwaidi JA. Endothelial dysfunction in diabetes mellitus. *Vascular Health and Risk Management*. 2007;3(6):853-876
- [28] Rask-Madsen C, King GL. More sugar, less blood vessels: Another piece in the puzzle of increased cardiovascular risk in diabetes. *Arteriosclerosis, Thrombosis, and Vascular Biology*. 2008;28(4):608-610
- [29] Gu Y, Lewis DF, Zhang Y, Groome LJ, Wang Y. Increased superoxide generation and decreased stress protein Hsp90 expression in human umbilical cord vein endothelial cells (HUVECs) from pregnancies complicated by preeclampsia. *Hypertension in Pregnancy*. 2006;25(3):169-182
- [30] Escudero C, Sobrevia L. A hypothesis for preeclampsia: Adenosine and inducible nitric oxide synthase in human placental microvascular endothelium. *Placenta*. 2008;29(6):469-483
- [31] Harrison DG, Cai H, Landmesser U, Griending KK. Interactions of angiotensin II with NAD(P)H oxidase,

oxidant stress and cardiovascular disease. *Journal of the Renin-Angiotensin-Aldosterone System*. 2003;**4**(2):51-61

[32] Mann GE, Yudilevich DL, Sobrevia L. Regulation of amino acid and glucose transporters in endothelial and smooth muscle cells. *Physiological Reviews*. 2003;**83**(1):183-252

[33] Sobrevia L, Mann GE. Dysfunction of the endothelial nitric oxide signalling pathway in diabetes and hyperglycaemia. *Experimental Physiology*. 1997;**82**(3):423-452

[34] Casanello P, Escudero C, Sobrevia L. Equilibrative nucleoside (ENTs) and cationic amino acid (CATs) transporters: Implications in foetal endothelial dysfunction in human pregnancy diseases. *Current Vascular Pharmacology*. 2007;**5**(1):69-84

[35] Sobrevia L, González M. A role for insulin on L-arginine transport in fetal endothelial dysfunction in hyperglycaemia. *Current Vascular Pharmacology*. 2009;**7**(4):467-474

[36] Casanello P, Sobrevia L. Intrauterine growth retardation is associated with reduced activity and expression of the cationic amino acid transport systems  $\gamma^+$ /hCAT-1 and  $\gamma^+$ /hCAT-2B and lower activity of nitric oxide synthase in human umbilical vein endothelial cells. *Circulation Research*. 2002;**91**(2):127-134

[37] Flores C, Rojas S, Aguayo C, Parodi J, Mann G, Pearson JD, et al. Rapid stimulation of L-arginine transport by D-glucose involves p42/44(mapk) and nitric oxide in human umbilical vein endothelium. *Circulation Research*. 2003;**92**(1):64-72

[38] González M, Flores C, Pearson JD, Casanello P, Sobrevia L. Cell signalling-mediated insulin increase of mRNA expression for cationic amino acid transporters-1 and -2 and membrane

hyperpolarization in human umbilical vein endothelial cells. *Pflügers Archiv*. 2004;**448**(4):383-394

[39] Arancibia-Garavilla Y, Toledo F, Casanello P, Sobrevia L. Nitric oxide synthesis requires activity of the cationic and neutral amino acid transport system  $\gamma^+$ L in human umbilical vein endothelium. *Experimental Physiology*. 2003;**88**(6):699-710

[40] Devés R, Boyd CA. Transporters for cationic amino acids in animal cells: Discovery, structure, and function. *Physiological Reviews*. 1998;**78**(2):487-545

[41] Palacín M, Estévez R, Bertran J, Zorzano A. Molecular biology of mammalian plasma membrane amino acid transporters. *Physiological Reviews*. 1998;**78**(4):969-1054

[42] Closs EI, Gräf P, Habermeier A, Cunningham JM, Förstermann U. Human cationic amino acid transporters hCAT-1, hCAT-2A, and hCAT-2B: Three related carriers with distinct transport properties. *Biochemistry*. 1997;**36**(21):6462-6468

[43] Yang Z, Venardos K, Jones E, Morris BJ, Chin-Dusting J, Kaye DM. Identification of a novel polymorphism in the 3'UTR of the L-arginine transporter gene SLC7A1: Contribution to hypertension and endothelial dysfunction. *Circulation*. 2007;**115**(10):1269-1274

[44] Verrey F, Closs EI, Wagner CA, Palacin M, Endou H, Kanai Y. CATs and HATs: The SLC7 family of amino acid transporters. *Pflügers Archiv*. 2004;**447**(5):532-542

[45] Hammermann R, Brunn G, Racké K. Analysis of the genomic organization of the human cationic amino acid transporters CAT-1, CAT-2 and CAT-4. *Amino Acids*. 2001;**21**(2):211-219

- [46] Aulak KS, Liu J, Wu J, Hyatt SL, Puppi M, Henning SJ, et al. Molecular sites of regulation of expression of the rat cationic amino acid transporter gene. *The Journal of Biological Chemistry*. 1996;**271**(47):29799-29806
- [47] Aulak KS, Mishra R, Zhou L, Hyatt SL, de Jonge W, Lamers W, et al. Post-transcriptional regulation of the arginine transporter Cat-1 by amino acid availability. *The Journal of Biological Chemistry*. 1999;**274**(43):30424-30432
- [48] Fernandez J, Lopez AB, Wang C, Mishra R, Zhou L, Yaman I, et al. Transcriptional control of the arginine/lysine transporter, cat-1, by physiological stress. *The Journal of Biological Chemistry*. 2003;**278**(50):50000-50009
- [49] Hatzoglou M, Fernandez J, Yaman I, Closs E. Regulation of cationic amino acid transport: The story of the CAT-1 transporter. *Annual Review of Nutrition*. 2004;**24**:377-399
- [50] Lopez AB, Wang C, Huang CC, Yaman I, Li Y, Chakravarty K, et al. A feedback transcriptional mechanism controls the level of the arginine/lysine transporter cat-1 during amino acid starvation. *The Biochemical Journal*. 2007;**402**(1):163-173
- [51] Steinberg HO, Baron AD. Vascular function, insulin resistance and fatty acids. *Diabetologia*. 2002;**45**(5):623-634
- [52] Scherrer U, Randin D, Vollenweider P, Vollenweider L, Nicod P. Nitric oxide release accounts for insulin's vascular effects in humans. *The Journal of Clinical Investigation*. 1994;**94**(6):2511-2515
- [53] Samson SL-A, Wong NCW. Role of Sp1 in insulin regulation of gene expression. *Journal of Molecular Endocrinology*. 2002;**29**(3):265-279
- [54] Solomon SS, Majumdar G, Martinez-Hernandez A, Raghov R. A critical role of Sp1 transcription factor in regulating gene expression in response to insulin and other hormones. *Life Sciences*. 2008;**83**(9-10):305-312
- [55] González M, Gallardo V, Rodríguez N, Salomón C, Westermeier F, Guzmán-Gutiérrez E, et al. Insulin-stimulated L-arginine transport requires SLC7A1 gene expression and is associated with human umbilical vein relaxation. *Journal of Cellular Physiology*. 2011;**226**(11):2916-2924
- [56] Huang CC, Chiribau C-B, Majumder M, Chiang C-M, Wek RC, Kelm RJ, et al. A bifunctional intronic element regulates the expression of the arginine/lysine transporter Cat-1 via mechanisms involving the purine-rich element binding protein A (Pur alpha). *The Journal of Biological Chemistry*. 2009;**284**(47):32312-32320
- [57] Damm P. Future risk of diabetes in mother and child after gestational diabetes mellitus. *International Journal of Gynaecology and Obstetrics: The Official Organ of the International Federation of Gynaecology and Obstetrics*. 2009;**104**(Suppl 1):S25-S26
- [58] Brewster S, Zinman B, Retnakaran R, Floras JS. Cardiometabolic consequences of gestational dysglycemia. *Journal of the American College of Cardiology*. 2013;**62**(8):677-684
- [59] Sobrevia L, Nadal A, Yudilevich DL, Mann GE. Activation of L-arginine transport (system y<sup>+</sup>) and nitric oxide synthase by elevated glucose and insulin in human endothelial cells. *The Journal of Physiology*. 1996;**490**(Pt 3):775-781
- [60] Vásquez R, Farías M, Vega JL, Martin RS, Vecchiola A, Casanello P, et al. D-glucose stimulation of

L-arginine transport and nitric oxide synthesis results from activation of mitogen-activated protein kinases p42/44 and Smad2 requiring functional type II TGF-beta receptors in human umbilical vein endothelium. *Journal of Cellular Physiology*. 2007;**212**(3):626-632

[61] Srinivasan S, Hatley ME, Bolick DT, Palmer LA, Edelstein D, Brownlee M, et al. Hyperglycaemia-induced superoxide production decreases eNOS expression via AP-1 activation in aortic endothelial cells. *Diabetologia*. 2004;**47**(10):1727-1734

[62] Kim F, Tysseling KA, Rice J, Gallis B, Haji L, Giachelli CM, et al. Activation of IKKbeta by glucose is necessary and sufficient to impair insulin signaling and nitric oxide production in endothelial cells. *Journal of Molecular and Cellular Cardiology*. 2005;**39**(2):327-334

[63] Sobrevia L, Yudilevich DL, Mann GE. Activation of A2-purinoceptors by adenosine stimulates L-arginine transport (system y<sup>+</sup>) and nitric oxide synthesis in human fetal endothelial cells. *The Journal of Physiology*. 1997;**499**(Pt 1):-40. DOI: 135

[64] Muñoz G, San Martín R, Farías M, Cea L, Vecchiola A, Casanello P, et al. Insulin restores glucose inhibition of adenosine transport by increasing the expression and activity of the equilibrative nucleoside transporter 2 in human umbilical vein endothelium. *Journal of Cellular Physiology*. 2006;**209**(3):826-835

[65] San Martín R, Sobrevia L. Gestational diabetes and the adenosine/L-arginine/nitric oxide (ALANO) pathway in human umbilical vein endothelium. *Placenta*. 2006;**27**(1):1-10

[66] González M, Rojas S, Avila P, Cabrera L, Villalobos R, Palma C, et al.

Insulin reverses D-glucose-increased nitric oxide and reactive oxygen species generation in human umbilical vein endothelial cells. *PLoS One*. 2015;**10**(4):e0122398

[67] Kerr PM, Tam R, Narang D, Potts K, McMillan D, McMillan K, et al. Endothelial calcium-activated potassium channels as therapeutic targets to enhance availability of nitric oxide. *Canadian Journal of Physiology and Pharmacology*. 2012;**90**(6):739-752

[68] Sandow SL, Grayson TH. Limits of isolation and culture: Intact vascular endothelium and BKCa. *American Journal of Physiology. Heart and Circulatory Physiology*. 2009;**297**(1):H1-H7

[69] Wiecha J, Reineker K, Reitmayer M, Voisard R, Hannekum A, Mattfeldt T, et al. Modulation of Ca<sup>2+</sup>-activated K<sup>+</sup> channels in human vascular cells by insulin and basic fibroblast growth factor. *Growth Hormone & IGF Research: Official Journal of the Growth Hormone Research Society and the International IGF Research Society*. 1998;**8**(2):175-181

[70] O'Malley D, Harvey J. Insulin activates native and recombinant large conductance Ca(2+)-activated potassium channels via a mitogen-activated protein kinase-dependent process. *Molecular Pharmacology*. 2004;**65**(6):1352-1363

[71] Rojas S, Basualto E, Valdivia L, Vallejos N, Ceballos K, Peña E, et al. The activity of IKCa and BKCa channels contributes to insulin-mediated NO synthesis and vascular tone regulation in human umbilical vein. *Nitric Oxide: Biology and Chemistry*. 2020;**99**:7-16

[72] Cabrera L, Saavedra A, Rojas S, Cid M, Valenzuela C, Gallegos D, et al. Insulin induces relaxation and decreases hydrogen peroxide-induced vasoconstriction in human placental



vascular bed in a mechanism mediated by calcium-activated potassium channels and L-arginine/nitric oxide pathways. *Frontiers in Physiology*. 2016;7:529

[73] Sand A, Andersson E, Fried G. Nitric oxide donors mediate vasodilation in human placental arteries partly through a direct effect on potassium channels. *Placenta*. 2006;27(2-3):181-190

IntechOpen