



SUPPLEMENT ARTICLE

High genetic diversity and differentiation of the *Babesia ovis* population in Turkey

Anabela Mira^{1,2} | Ahmet Hakan Unlu³ | Huseyin Bilgin Bilgic⁴ | Serkan Bakirci⁴ | Selin Hacilarlioglu⁴ | Tulin Karagenc⁴ | Tamara Carletti¹ | William Weir⁵ | Brian Shiels⁵ | Varda Shkap⁶ | Munir Aktas⁷  | Monica Florin-Christensen^{1,2} | Leonhard Schnittger^{1,2} 

¹Instituto de Patobiología Veterinaria, CICVyA, INTA-Castelar, Hurlingham, Argentina

²Consejo Nacional de Ciencia y Tecnología (CONICET), Buenos Aires, Argentina

³Vocational School of Gevas, Van Yuzuncu Yil University, Van, Turkey

⁴Department of Parasitology, Faculty of Veterinary Medicine, Aydin Adnan Menderes University, Aydin, Turkey

⁵Institute of Biodiversity, Animal Health and Comparative Medicine, College of Medical, Veterinary & Life Sciences, University of Glasgow, Glasgow, UK

⁶Division of Parasitology, Kimron Veterinary Institute, Bet Dagan, Israel

⁷Department of Parasitology, Faculty of Veterinary Medicine, Firat University, Elazig, Turkey

Correspondence

Leonhard Schnittger, Instituto de Patobiología Veterinaria, CICVyA, INTA-Castelar, Los Reseros y Nicolas Repetto, s/n, 1686 Hurlingham, Prov. de Buenos Aires, Argentina. Email: schnittger.leonhard@inta.gob.ar

Funding information

European Commission, Grant/Award Number: FP7 INCO245145; Argentine Ministry of Science and Technology, Grant/Award Number: PICT 2013-1249; National Institutes of Agricultural Technology, Grant/Award Number: PNBIO 1131034

Summary

Babesia ovis is a tick-transmitted protozoan haemoparasite causing ovine babesiosis in sheep and goats leading to considerable economic loss in Turkey and neighbouring countries. There are no vaccines available, therapeutic drugs leave toxic residues in meat and milk, and tick vector control entails environmental risks. A panel of eight mini- and micro-satellite marker loci was developed and applied to study genetic diversity and substructuring of *B. ovis* from western, central and eastern Turkey. A high genetic diversity ($H_e = 0.799$) was found for the sample of overall *B. ovis* population ($n = 107$) analyzed. Principle component analysis (PCoA) revealed the existence of three parasite subpopulations: (a) a small subpopulation of isolates from Aydin, western Turkey; (b) a second cluster predominantly generated by isolates from western Turkey; and (c) a third cluster predominantly formed by isolates from central and eastern Turkey. Two *B. ovis* isolates from Israel included in the analysis clustered with isolates from central and eastern Turkey. This finding strongly suggests substructuring of a major Turkish population into western versus central–eastern subpopulations, while the additional smaller *B. ovis* population found in Aydin could have been introduced, more recently, to Turkey. STRUCTURE analysis suggests a limited exchange of parasite strains between the western and the central–eastern regions and *vice versa*, possibly due to limited trading of sheep. Importantly, evidence for recombinant genotypes was obtained in regionally interchanged parasite isolates. Important climatic differences between the western and the central/eastern region, with average yearly temperatures of 21°C versus 15°C, correspond with the identified geographical substructuring. We hypothesize that the different climatic conditions may result in variation in the activity of subpopulations of *Rhipicephalus* spp. tick vectors, which, in turn, could selectively maintain and transmit different parasite populations. These findings may have important implications for vaccine development and the spread of drug resistance.

KEYWORDS

Babesia ovis, genetic diversity, multilocus typing, ovine babesiosis, population structure, satellite marker

1 | INTRODUCTION

Ovine babesiosis is a tick-borne disease principally caused by *B. ovis* and is endemic in southern Europe, Africa, the Middle East and Asia (Ahmed et al., 2002; Mehlhorn & Schein, 1984; Schnittger et al., 2012). The disease has been reported to result in important economic losses in Turkey and Iran (Esmailnejad et al., 2015; Ranjbar-Bahadori et al., 2012; Sevinc et al., 2013, 2014). Clinical signs of the infection include fever, haemolytic anaemia, haemoglobinuria and icterus leading to high morbidity, with a significant number of fatalities observed in some areas (Hurtado et al., 2015; Sevinc et al., 2013). Currently, ovine babesiosis is predominantly controlled by chemotherapeutic treatment of animals during acute disease but, to a limited degree, chemical control of the vector tick is also applied. The drawback of the former approach is that it is expensive, requires early diagnosis of the disease, and drug residues may be introduced into the food chain, whereas the downside of the latter approach is that it poses environmental risks and acaricide resistance is observed.

The transmitting tick vectors of *B. ovis* are species of the genus *Rhipicephalus*, such as *R. bursa*, *R. turanicus*, *R. sanguineus* and *R. annulatus*, which are abundant in western, central and eastern Turkey, and also endemic in Iran (Aydin & Bakirci, 2007; Sayin et al., 1997a,b,c; Shayan et al., 2007). Accordingly, a recent comprehensive molecular epidemiological study demonstrated the highest *B. ovis* infection rates in sheep and goats from the Kutahya and Afyon regions of western, and the Konya and Aksaray regions of eastern Turkey (Bilgic et al., 2017). Notwithstanding the observed high infection rate of and seroprevalence against *B. ovis* in Turkey, endemic instability prevails in most regions, thus an environment-friendly immunoprophylactic approach is required to control the disease (Ekici et al., 2012; Mahoney & Ross, 1972; Sevinc et al., 2014).

Although attenuated live vaccines have been shown to be effective against bovine babesiosis caused by the closely related species, *B. bovis*, and by *B. bigemina*, a similar vaccine is not yet available for ovine babesiosis (Florin-Christensen et al., 2014; Schnittger et al., 2012; Sevinc et al., 2014; Shkap et al., 2007). In a recent preliminary study, 12 successive blood passages in splenectomized lambs mitigated but could not eliminate the virulence of a pathogenic *B. ovis* strain when tested in lambs (Sevinc et al., 2014). Analogous to the situation for *B. bovis* and *B. bigemina*, effective attenuation may be achieved through additional *in vivo* passage or, alternatively, through *in vitro* passage of cultured *B. ovis* strains (Horta et al., 2014; Florin-Christensen et al., 2014).

Population genetic analysis of *Babesia* allows the investigation of parasite genetic diversity and structure, and can estimate the extent of genetic exchange in the field. Estimated parameters are able to inform on the most promising approaches for vaccine development and application, and provide information on the risk of dissemination of drug resistance-encoding genes in the population. Population genetic studies are currently lacking for *B. ovis* but have been carried out for the closely related species *B. bovis* (Flores et al., 2013;

Perez-Llaneza et al., 2010; Schnittger et al., 2012; Simuunza et al., 2011). An extremely high level of genetic differentiation and diversity of the *B. bovis* worldwide metapopulation has been estimated by comparing reference isolates originating from distant countries and continents (Perez Llaneza et al., 2010; Flores et al., 2013). Likewise, a study of the *B. bovis* populations in Turkey and Zambia suggests a high level of genetic diversity and frequent genetic exchange, under preconditions of a relatively low-geographic distance between populations, a high-transmission rate, and exchange of infected animals (Schnittger et al., 2012; Simuunza et al., 2011). In contrast, linkage disequilibrium (LD) is commonly observed when isolates from distant geographic regions are compared.

In this study, a panel of micro- and mini-satellite markers for multilocus typing based on the available *B. ovis* genome was developed and applied, as these types of markers have been shown to be highly suitable for population genetic analysis of related *Babesia* and *Theileria* parasites. Satellite markers were used to assess the genetic diversity, population structure, frequency of recombination and multiplicity of infection of the *B. ovis* population of Turkey. The data obtained indicated separation of major populations of parasites within Turkey, which has implications for both vaccine development and the dissemination of genes that confer drug resistance.

2 | MATERIALS AND METHODS

2.1 | Parasite sampling, genomic DNA isolation and satellite marker development

A total of 107 blood samples of sheep from different areas in Turkey were analysed (Figure 1). These originated from (a) the western



FIGURE 1 Map of Turkey showing the main sampling regions: western Turkey comprising of the Aegean (AR) and Mediterranean Region (MR), central Turkey represented by the Central Anatolian Region (CAR), and eastern Turkey consisting of the Eastern Anatolian Region (EAR) and the Southeastern Anatolia Region (SAR). Western Turkey has a medium temperature of 15.9°C and a medium humidity of 62.4%; central Turkey has a medium temperature of 10.5°C and a medium humidity of 62.6%; EAR and SAR of eastern Turkey have a medium temperature of 9.7°C and 16.5°C and a medium humidity of 60.9% and 53.4%, respectively [Colour figure can be viewed at wileyonlinelibrary.com]

of Afyon ($n = 22$), Aydin ($n = 7$), Burdur ($n = 2$) and Kütahya ($n = 20$), (b) the central [central Anatolian region ($n = 34$), including the provinces of Aksaray-Eskil ($n = 7$), Konya ($n = 19$) and Niğde ($n = 8$)] and (c) the eastern region [(Eastern Anatolia and Southeastern Anatolia region ($n = 22$), including the provinces of Bingöl ($n = 1$), Elazığ ($n = 2$), Erzincan ($n = 4$), Erzurum ($n = 1$), Iğdır ($n = 2$), Malatya ($n = 2$), Mardin ($n = 2$), Muş ($n = 3$) and Şırnak ($n = 5$)] (Figure 1). Genomic DNA was isolated using the DNAeasy blood and tissue kit, according to the recommendation of the manufacturer (Qiagen, Hilden, Germany). In addition, genomic DNA was also isolated from the in vitro cultured *B. ovis* strain, Israel and from the field isolate *B. ovis* strain, Itamar (Kimron Veterinary Institute, Israel). Isolated DNA was quantified using a Nanodrop spectrophotometer, adjusted to 500 pg/μl and stored at -20°C until further use. Selected blood samples were shown to test positive in a PCR assay, using RLB primers generic for *Babesia* spp. infecting small ruminants (Horta et al., 2014; Schnittger et al., 2004). Micro- and mini-satellite markers were developed as described in Perez-Llaneza et al. (2010), and designed based on the as yet publicly unavailable *B. ovis* strain Israel genome sequence (University of Glasgow, Glasgow, UK).

2.2 | PCR amplification of micro- and mini-satellite loci

PCR was performed using primers designed to amplify each of the eight selected marker repeat regions on the *B. ovis* isolates and strains (Table S1). The reverse primer of each pair was labelled at the 5' end with one of the standard fluorescence dyes: 6-FAM (Blue), NED (Yellow), PET (Red), or VIC (Green) (Applied Biosystems), enabling detection of product on an ABI Genetic Analyzer. PCR amplification was performed in a final reaction volume of 10 μl comprising of 500 pg of template genomic DNA, 0.25 U DreamTaq polymerase (ThermoFisher Scientific), 0.2 mM dNTPs, 1 μM of each primer, 2 mM MgCl_2 and DreamTaq buffer (10x) (ThermoFisher Scientific). Nuclease-free water and *B. ovis* genomic DNA were used as negative and positive controls, respectively. The thermocycler program consisted of an initial denaturation step at 95°C for 5 min, followed by 35 cycles comprising denaturation at 95°C for 30 s, annealing at 55 to 60°C (see Table S1) for 30 s, and extension at 72°C for 20 s; and a final extension step at 72°C for 7 min. Five-microliters of each PCR reaction were analysed on an ethidium bromide-containing 1.8% agarose gel to verify amplification success. Genotyping was performed at the Unidad de Genómica, Instituto de Biotecnología, CICVyA, INTA-Castelar. DNA fragment sizes were analysed relative to ROX-labelled GS600 LIZ size standards (Applied Biosystems) using Peak Scanner software (ThermoFisher Scientific), which allowed resolution of 1 bp differences in amplicon size. The presence of multiple amplification products generated in a single PCR reaction of a sample demonstrated the presence of two or more *B. ovis* genotypes. Data output generated by capillary electrophoresis included the area under the peak as a quantitative measurement of allele amplification. The predominant allele of each marker locus of a given sample was identified by the largest peak area, and a multilocus

genotype (MLG) representing the most abundant *B. ovis* strain present in the isolate was generated. An excel file of MLG data was produced to assess genetic diversity and structure of the population. A second file comprised the allelic marker profile of each sample that was used to estimate the multiplicity of infection (MOI).

2.3 | Data analysis

Population genetic analysis for the estimation of F-statistics was performed using Genepop version 4.2 (<http://genepop.curtin.edu.au/index.html>; Raymond & Rousset, 1995; Rousset, 2008). The Microsoft Excel plug-in software, GenAlEx6.5 (<http://biology.anu.edu.au/GenAlEx/>; Peakall & Smouse, 2012), was used to estimate the Analysis of MOlecular VAriance (AMOVA), to construct a similarity matrix, and perform principal component analysis (PCoA) of MLG data. Population structure was also assessed using Bayesian clustering analysis as implemented in STRUCTURE 2.3.4 (Pritchard et al., 2000). The EVANNO Method Delta K and STRUCTURE Harvester was used to derive the best fit for the parameter value K (Earl & von Holdt, 2012). Assuming that the four populations from each sample location are genetically distinct and using the admixture model with correlated allele frequencies, 20 iterations were run for each K ranging from 1 to 10 selecting a 'burn in' of 100,000 and 200,000 MCMC simulations.

The null hypothesis of linkage equilibrium (LE) was tested using LIAN which computes the standardized index of association (I_A^S) and quantifies LE/LD as previously defined by Haubold & Hudson (2000) (<http://guanine.evolbio.mpg.de/cgi-bin/lian/lian.cgi.pl/query>). The I_A^S measures the association between alleles at pairs of loci. I_A^S values close to 0 or negative are indicative of panmixia, while those significantly greater than 0 are indicative of a non-panmictic situation. LE is characterized by the statistical independence of alleles across all loci and is investigated by initially determining the number of loci at which each pair of MLGs differs. From the distribution of mismatch values, a variance (V_D) is calculated which is compared to the variance expected for LE, termed V_e . The null hypothesis that $V_D = V_e$ is tested by both a Monte Carlo simulation and a parametric method in order to estimate a 95% confidence limit (95% CI), which are denoted L_{mc} and L_{para} respectively. When V_D is found to be greater than L , the null hypothesis is rejected and LD is accepted.

The mean number of all identified alleles for all eight loci was calculated to represent the MOI as an estimation of the number of genotypes within an isolate. The average MOI of each population was estimated by calculating the overall mean of the MOI of all representative samples.

3 | RESULTS

3.1 | Satellite marker diversity

In each isolate, the presence of numerous *B. ovis* multilocus genotypes (MLG) was evident through the observation of more than one

allele at one or multiple marker loci. The foremost allele for each marker loci was scored to determine the prominent MLG in each sample. This MLG was used to assess satellite marker and genetic diversity as well as the genetic differentiation within the population. Of the eight employed micro- and mini-satellite markers, five showed a very high level of genetic diversity, ranging from 0.802 (Oms_8) to 0.899 (OMS_47), compared to the remaining three that displayed a somewhat lower genetic diversity, ranging from 0.703 (Oms_11) to 0.755 (OMS_103) (Table 1). This high polymorphism indicated that markers can be assumed to be informative for LD analysis of the *B. ovis* population and also suitable to describe population diversity. The number of alleles per marker identified in all isolates studied was very high, ranging from 14 alleles for marker locus OMS_103 up to 28 alleles for markers OMS_47 and OMS_93. Across all eight markers, an average of 21.6 different alleles was determined. Predominant allele frequencies ranged from 20.6% for marker OMS_47 to 51.4% for marker Oms_11. As an example, the allele frequency distribution of three markers in the three studied geographic regions of Turkey is shown in Figure 2, and for the remaining markers in Figure S1.

3.2 | Population diversity and structure

The genetic diversity (H_e) estimated for the three geographic regions of Turkey was found to be moderate for the population of the western region (0.663), compared to a very high value obtained for both the population of the central (0.770) and eastern region (0.791), which were found to be similar to that of the overall population (0.799) (Figure 3, Table S2). The number of different alleles (N_a) in the western, central and eastern population was 10.8, 10.6 and 8.8 respectively, with a mean number of 19.5 different alleles in the overall population. In contrast, the mean number of effective alleles (N_e) was determined as 4.1, 5.3 and 6.0 and the mean number of private alleles was found to be 5.5, 4.5 and 2.4, with a high mean number of 12.4 private alleles at each marker locus in the overall population. This considerable number of private alleles for each marker can be directly observed in Figure 2 and Figure S1, and this information is detailed in Table S2.

The predominant allele at each marker locus was identified and the MLG for each isolate established. Altogether, 100 different MLG could be identified in the 107 isolates, demonstrating a high level of genomic diversity in the overall *B. ovis* population of Turkey. On the basis of the MLG data, a matrix of Nei's genetic distance was




FIGURE 2 Allele frequency histograms of selected satellite markers of *Babesia ovis* field populations of the western, central and eastern regions of Turkey. (a) Satellite marker OMS_62 shows a high number of private alleles ($n = 8$) in the central population, and low to moderate number of private alleles in the eastern ($n = 4$) and western populations ($n = 2$). (b) A bell-like frequency distribution is observed from allele 211 to 260 bp for marker Oms_8. (c) Marker OMS_47 shows a high number of private alleles ($n = 14$) in the western parasite population. [Colour figure can be viewed at wileyonlinelibrary.com]

calculated to carry out PCoA. PCoA allows visualization of the genetic relationships between MLGs and the comparison of genotypes from the three geographic regions together with *B. ovis* reference isolates from Israel (Figure 4). Two main genetic clusters are observed, one cluster is predominantly composed of samples from western Turkey although it also includes seven (corresponding to six data points in Figure 4) from central Turkey and two from eastern Turkey (cluster 1), while the second cluster is mainly composed of samples from central and eastern Turkey but also includes five isolates (corresponding to four data points in Figure 4) from western Turkey and the two Israel isolates (cluster 2). Clusters 1 and 2 touch to the right of the perpendicular axis. Importantly, a third smaller

TABLE 1 Allelic variation in the *Babesia ovis* population of Turkey

	Oms_8	Oms_10	Oms_11	OMS_47	OMS_58	OMS_62	OMS_93	OMS_103	Mean
Total number of alleles	17	18	15	28	16	20	28	14	21.6
Predominant allele frequency (size in bp)	0.393 (313)	0.346 (181)	0.514 (142)	0.206 (241)	0.421 (130)	0.290 (189)	0.290 (250)	0.393 (282)	0.747
Genetic diversity (H_e)	0.802	0.812	0.703	0.899	0.723	0.823	0.865	0.755	0.799




FIGURE 3 Genetic diversity (H_e), mean number of different (N_a), effective (N_e) and private alleles in the *B. ovnis* populations of western, central and eastern Turkey [Colour figure can be viewed at wileyonlinelibrary.com]

separate grouping (cluster 3) is formed by some samples from western Turkey.

With regard to sample origin, most samples from western Turkey clustered to the right of the perpendicular axis of the plot into cluster 1, with the exception of all isolates originating from the Aydin region ($n = 7$), situated at the Aegean coast, that formed a separate cluster in the lower left quadrant (cluster 3). In contrast, samples from central and eastern Turkey clustered mainly left of the perpendicular axis of the plot including also the two isolates from Israel to produce cluster 2. Seven samples that originate from Aksaray, central Turkey, clustered jointly with the majority of western Turkish isolates to the right of the perpendicular axis forming a subgroup within cluster 1.

The STRUCTURE analysis predicts three different *B. ovnis* genotype clusters ($K = 3$) as determined by the EVANNO delta K method that match the three clusters identified by PCoA. The three genotype clusters are represented in Figure 5 as red genotype 1 (western population including seven isolates from Aksaray, central Turkey, one from Mus and one from Sirnak, eastern Turkey and four isolates of a mixed genotype from Kutahya, western Turkey corresponding to cluster 1), blue/green genotype 2 (central and eastern population including one isolate from Afyon, two from Burdur, and two from Kutahya, western Turkey corresponding to cluster 2), and red genotype 3 (seven isolates from Aydin, corresponding to cluster 3).




FIGURE 4 Principal Component Analysis (PCoA) of the multilocus genotypes (MLGs) of *B. ovnis* isolates of western, central and eastern Turkey. The two principal axes are shown and the portion of variation in the dataset explained by each axis is indicated in parenthesis. The subgroup indicated with broken lines is formed by samples from Aksaray, eastern Turkey and falls within cluster 1, mainly composed by samples from western Turkey. 1: *B. ovnis* strain Israel, 2: *B. ovnis* strain Itamar, Israel [Colour figure can be viewed at wileyonlinelibrary.com]




FIGURE 5 STRUCTURE and EVANNO Method Delta K analysis. The assumption of three genotype clusters ($K = 3$) was found to best fit data which are represented by green (Aydin, western Turkey), green/blue (predominantly central and eastern Turkey) and red stripes (predominantly western Turkey with the exclusion of Aydin) within the bar. The origin of population samples are given below. Genotypes that did not fit into the genotype background of the respective geographic population are given above [Colour figure can be viewed at wileyonlinelibrary.com]

two from Kutahya, western Turkey and both parasite strains from Israel, corresponding to cluster 2) and the green genotype 3 (seven isolates from Aydin, corresponding to cluster 3). AMOVA analysis showed that the molecular variation between the three geographically defined populations of western Turkey (cluster 1), eastern/central Turkey (cluster 2), and Aydin (cluster 3) was 24%, whereas the variation within populations accounted for the remaining 76% of the observed variation.

In order to verify the existence of a separate Aydin population as predicted by the PCoA and STRUCTURE analyses, samples were divided into the four geographic populations Aydin, western Turkey* (excluding Aydin samples), central Turkey and eastern Turkey; and the pairwise genetic differentiation determined by estimation of the unbiased Nei's genetic distance (u_D) and Wright's fixation index (F_{st}) (Table 2). An extremely high level of genetic differentiation was estimated between the Aydin versus the western population ($u_D = 2.464$; $F_{st} = 0.544$) and was also very high between samples from Aydin versus the central ($u_D = 0.421$; $F_{st} = 0.241$) and also versus the eastern population ($u_D = 0.380$; $F_{st} = 0.244$), strengthening the evidence that Aydin isolates should be considered as a discrete population. Furthermore, a high-genetic differentiation was observed between the population of western Turkey* versus that of central ($u_D = 0.889$; $F_{st} = 0.241$) and eastern Turkey ($u_D = 0.830$; $F_{st} = 0.233$). In contrast, the population of central Turkey showed negligible genetic differentiation to that of eastern Turkey ($u_D = 0.049$; $F_{st} = 0.011$), corroborating their joint grouping into cluster 2 and exhibiting a similar blue/green genotype 2 in the PCoA and STRUCTURE analysis, respectively. Consequently, it seems justified to regard *B. ovnis* isolates from central and eastern Turkey as one single population.

3.3 | Linkage analysis

Using the assessed MLG dataset, the standard index of association (I^s_A) was calculated in order to determine whether an association between alleles could be observed when all pairwise combinations of loci were compared. After estimation of I^s_A , the null hypothesis of LE

TABLE 2 Pairwise comparison of Wright's fixation index (F_{st}) and unbiased Nei's genetic distance (uD)

Unbiased Nei's Genetic Distance (uD)	Wright's fixation index (F_{st})			
	Aydin	Western Turk.*	Central Turk.	Eastern Turk.
Aydin	-	0.544	0.241	0.244
Western Turk.*	2.464	-	0.241	0.233
Central Turk.	0.421	0.889	-	0.011
Eastern Turk.	0.380	0.830	0.049	-

Western Turk.*, isolates from western Turkey except Aydin.

for MLG data was tested. Using the MLG data, the LD analysis was carried out for all available data from Turkey, by pairwise combination of the data representing western, central, and eastern regions, for each region separately, and also for individual farms, whenever a sufficient number of samples were available. This approach allows testing hierarchically nested population subdivisions until eventually LE is observed. The observance of LE demonstrates a geographic area, which may correspond to a single farm, within which frequent recombination of MLGs is observed.

The combined population of Turkey resulted in an I_A^s of 0.1615 demonstrating the existence of LD (Table 3). Likewise, pairwise combination of the population representing western/central Turkey and western/eastern Turkey showed even higher values for LD of 0.1992 and 0.1916 respectively. In contrast, the pairwise combination of data representing central/eastern Turkey resulted in a substantially lower I_A^s of 0.0889, indicating a much less pronounced LD. Thus, the central and eastern populations of Turkey are best regarded as a single population as they are substantially closer to equilibrium than the central versus the western or the eastern versus the western population, which is consistent with the population differentiation between the western versus the central/eastern population, evident on the PCoA and F_{ST} analysis. LD was also observed when testing each region, the western population showing a higher I_A^s of 0.2111 followed by the central and eastern populations with 0.1317 and 0.0919

respectively. Further geographic subdivision resulted in the observation of significant LE in samples originating from Konya and Aydin, but in the remaining subpopulations LD was still observed. Notwithstanding that, in the case of Aydin ($n = 7$), the sample size was relatively low; this result indicates that shorter geographic distances facilitate genetic exchange between MLGs, resulting eventually in the observation of LE in populations from local geographic or farm populations as has been reported for related piroplasmid species (Gomes et al., 2016; Simuunza et al., 2011; Weir et al., 2011).

3.4 | Multiplicity of infection

In a single sample, the number of alleles at a given loci ranged from 1 to a maximum of 5 for marker Oms_8 and from 1 to a maximum of 15 for marker Oms_11 and OMS_62. Overall, the mean number of alleles per sample across all markers was found to range from 1.65 for Oms_8 up to 4.36 for OMS_62 (Table S2). For each isolate, the detection of several alleles at one or more markers was indicative of multiple genotypes within all isolates originating from the three studied regions. Correspondingly, the number of genotypes for each sample could be estimated by calculating the average number of alleles across all marker loci for a given sample. In turn, the MOIs determined for each isolate were used to estimate the average MOI in the populations of the western (3.31), central (2.48) and eastern

TABLE 3 Linkage disequilibrium analysis of *Babesia ovis* populations of Turkey

Region	n	H_e	I_A^s	V_D	L	p -value	Linkage
Turkey	107	0.799	0.1615	2.5879	1.2863	<0.01	LD
Western/central Turk.	85	0.784	0.1992	3.1198	1.4101	<0.01	LD
Central/eastern Turk.	56	0.812	0.0889	1.8018	1.2209	<0.01	LD
Western/eastern Turk.	73	0.770	0.1916	3.1558	1.4702	<0.01	LD
Western Turk.	51	0.677	0.2111	3.7629	1.7158	<0.01	LD
Central Turk.	34	0.793	0.1317	2.2986	1.4165	<0.01	LD
Eastern Turk.	22	0.829	0.0919	1.6608	1.2347	<0.01	LD
Konya	20	0.690	0.0331	1.8630	1.9571	0.11	LE
Afyon	22	0.503	0.0660	2.0908	1.7604	<0.01	LD
Kutahya	20	0.643	0.1483	2.6554	1.7877	<0.01	LD
Aydin	7	0.143	0.0196	0.9286	1.9286	0.12	LE
Aksaray/Nigde	15	0.801	0.2120	2.9557	1.5518	<0.01	LD

region (2.62), as well as for the total *B. ovis* population of Turkey (2.91) (Table 4).

4 | DISCUSSION

Population genetic studies using micro- and mini-satellites have been shown to be invaluable for the assessment of parasite diversity, recombination frequency and behaviour of the parasite at the population level. Genome sequence analysis allows the identification of putatively neutral, highly diverse micro- and mini-satellites with relative ease and these markers have been shown to be a useful tool for population genetic studies.

Ovine babesiosis has an important economic impact in Turkey and population genetics should be applied to provide insights into the most promising approaches and strategies for parasite and disease control. Similar to other apicomplexan haemoparasites, *Babesia* spp. are haploid and the only diploid stage is the zygote, which is formed during sexual reproduction in the tick host. Accordingly, only a single allele can be detected in the vertebrate host and thus, the detection of two or more alleles is evidence of dual or multiple infections. Consequently, heterozygosity cannot be directly observed, and is estimated based on the observed allele frequencies in the population.

As measured using expected heterozygosity, the overall genetic diversity of the *B. ovis* population in Turkey was found to be extremely high ($H_e = 0.799$) and similar to that observed for its phylogenetically closest relative, the cattle-infecting *B. bovis* ($H_e = 0.837$) (Simuunza et al., 2011). Likewise, in a recent study comparing reference isolates from diverse countries and continents around the world, a comparably high genetic diversity was observed in the *B. bovis* metapopulation ($H_e = 0.780$, Flores et al., 2013). *Babesia ovis* belongs, together with the closely related species *B. bovis*, to the *Babesia* sensu stricto group of piroplasmids, characterized by transovarial transmission.

In contrast, population genetic studies of *Theileria lestoquardi* (classified within *Theileria* sensu stricto), the causative agent of malignant ovine theileriosis in sheep, have shown a considerably lower genetic diversity in Sudan ($H_e = 0.550$ to 0.572) and Oman ($H_e = 0.528$ to 0.610) as compared to its sister species *T. annulata* in Oman ($H_e = 0.833$). Furthermore, the genetic diversity of *T. lestoquardi* was also found to be considerably lower to that determined in the

present study on *B. ovis* ($H_e = 0.799$) and to *B. bovis* ($H_e = 0.837$) in Turkey (Al-Hamidhi et al., 2016; Ali et al., 2017; Awad et al., 2018; Simuunza et al., 2011). While each of these species was analysed using a different set of markers, these various markers were identified using broadly similar criteria and for this reason, they are broadly comparable. It has been proposed that the generally lower diversity of *T. lestoquardi* compared to that of the closely related *T. annulata* is because present day *T. lestoquardi* and *T. annulata* evolved relatively recently from a common ancestor, following adaptation of *T. lestoquardi* to the small ruminant host, as supported by high 18S rRNA gene identity (Awad et al., 2018; Schnittger et al., 2000, 2003). Importantly, although *B. ovis* and *B. bovis* also represent sister species, it can be predicted that they have undergone a strikingly longer independent evolution, as evidenced by phylogenetic analysis (Schnittger et al., 2012).

A high diversity of the *B. ovis* population in Turkey ($n = 107$) is further substantiated by the high mean number of different alleles identified at each marker loci (19.5), whereas lower mean numbers of 11.3, 10.8 and 9.7 different alleles were observed in the *B. bovis* population of Turkey ($n = 40$), *T. annulata* population in Portugal ($n = 90$), and in the *T. lestoquardi* population of Sudan ($n = 36$) respectively (Ali et al., 2017; Gomes et al., 2016). Noteworthy, a high mean number of 12.4 private alleles in the overall *B. ovis* population suggests that future population admixture and subsequent recombination have the potential to significantly further increase the *B. ovis* genetic diversity in Turkey.

The observation of a lower genetic diversity in the *B. ovis* population of western Turkey ($H_e = 0.663$), as compared to the central ($H_e = 0.770$) and eastern populations ($H_e = 0.791$) suggests the existence of a degree of genetic differentiation between these two populations. Indeed, PCoA demonstrated two larger discrete clusters of *B. ovis* isolates, one comprising predominantly isolates originating from western Turkey (cluster 1) and another composed predominantly of isolates from the central and eastern regions (cluster 2). Strikingly, a third cluster (cluster 3) was observed, indicating that an additional separate *B. ovis* subpopulation exists within the parasite population of western Turkey. A number of assumptions may be drawn from this analysis. Firstly, cluster 1 includes *B. ovis* isolates originating from Aksaray, central Turkey, strongly suggesting that sheep infected with *B. ovis* originating from western Turkey have been introduced into the Aksaray region of central Turkey. This finding strongly suggests that some, though restricted, trading of sheep from western to eastern Turkey has taken place. Secondly, cluster 3 comprises exclusively *B. ovis* isolates from Aydin, a coastal region of western Turkey. Thus, Aydin isolates are distinctly different from the other western and from the eastern/central *B. ovis* populations, which may be due to strain selection by unique sheep breeds such as, *sakız*, *karya* and *çine çaparı* kept in this region. Thirdly, isolates from the *B. ovis* population of central and eastern Turkey cluster together suggesting that they represent a single parasite population. Fourthly, the two *B. ovis* isolates from Israel sit within cluster 2, demonstrating that they possess a MLG similar to those defined for central/eastern Turkey.

TABLE 4 Multiplicity of *Babesia ovis* infection in different regions of Turkey

Region	n	Multiplicity of infection			
		Mean	SD	Min	Max
Turkey	107	2.91	0.93	1.25	6.13
Western Turk.	51	3.31	0.97	1.33	6.13
Central Turk.	34	2.48	0.75	1.25	5
Eastern Turk.	22	2.62	0.69	1.63	4

n, number of samples; SD, standard deviation.

Significant LD was observed in the overall population of Turkey, in pairwise combinations and within each of the regions of western, central and eastern Turkey. A somewhat lower LD ($I_A^S = 0.0889$) was found for the pairwise combination of the population of central and eastern Turkey suggesting that there is a state of panmixia but recombination is not frequent enough to reach LE. Exclusively in two more confined regions, Konya and Aydin, LE could be observed. This finding also suggests that no genotypes have been introduced into these two regions, whereas, as shown by PCoA and STRUCTURE analysis, introduction of genotypes from other regions of Turkey to Afyon, Aksaray and Kutahya can be implied and may be responsible for the observed LD due to a recent admixture of novel genotypes. In order to reach LE, recombination frequency must be high which is underpinned by conditions where transmission intensity is high, the vector tick is abundant, and a high level of sheep infestation is observed (Weir et al., 2011). Due to the restricted mobility of the tick vector off the host, LE is most likely observed at the farm level or a more confined local geographic area under the premise that no novel genotypes are introduced. Thus, LD is generally observed in the overall population and in larger geographical regions, while LE is observed in the two smaller confined regions of Konya and Aydin. This observation suggests that in any scenario where resistance to a drug has arisen in a particular region, dissemination of the resistance gene(s) would occur primarily in a local manner, unless resistant parasites were spread via trading of infected animals across regions.

A high MOI of 3.31 was found in sheep of western Turkey, as compared to that of central (2.48) and eastern Turkey (2.62). In comparison, a considerably lower MOI of 2.04 has been observed for *B. bovis* infection of cattle of Turkey (Simuunza et al., 2011). This high MOI in combination with a high genetic diversity is indicative of a high-infection pressure with ticks carrying distinct parasite genotypes. It may be concluded that pathogen transmission is considerably more intense in western and elevated in eastern Turkey, which coincides with a higher *B. ovis* infection of sheep as reported for these regions by Bilgic et al. (2017).

Babesia ovis is characterized by transovarial transmission which allows direct vertical transfer of the parasite into tick eggs and successive larvae, nymph and adult tick stages in the absence of an infected vertebrate host. Although it has been recently reported that exclusively the adult tick effectively transmits *B. ovis* while larvae and nymph stages have none or inefficient vectorial competence (Erster et al., 2016), it may be assumed that compared to transstadial transmission as seen for *Theileria*, transovarial transmission might have the potential to exert higher infection intensity.

Importantly, two *B. ovis* isolates from Israel show a similar genotype to those from the central/western population of Turkey. A limited exchange of parasite genotypes seems to take place between the western and central/eastern populations, possibly due to mutual trading of sheep. This is supported by the observation that some *B. ovis* MLGs display combinations of western and central/eastern genotypes, suggesting recent recombination events.

Interestingly, the substructuring of the western Turkish parasite population versus the eastern/central parasite population corresponds with important climatic differences in the western region with an average yearly temperature of about 21°C as compared to the central-eastern region with an average yearly temperature of about 15°C. We hypothesize that the different climatic conditions result in activity differences of the respective *Rhipicephalus* spp. subpopulations, which, in turn, possess discrete parasite populations.

The significance of the observed genetic differentiation is that the development of an attenuated live vaccine based on a single strain may possibly not protect against challenges of strains from both populations. Importantly, the in vitro *B. ovis* strain Israel may be suitable for the development of a vaccine against the central/eastern parasite population but would be predicted to less likely protect against challenges from parasites of the population of western Turkey.

ACKNOWLEDGMENTS

The authors are grateful for the financial contribution of the National Research Council of Argentina (CONICET, salaries of AM, MFC and LS) and for the financial support of the European Commission (FP7 INCO245145, PIROVAC), the Argentine Ministry of Science and Technology (PICT 2013-1249), and the National Institutes of Agricultural Technology, Argentina (INTA, PNBIO 1131034).

ETHICS APPROVAL

This study was approved by the Adnan Menderes University Animal Experiment Ethic Committee dated 28 August 2012 in accordance with decision number B.30.2.ADÜ.0.00.00.00/050.04/2012/047.

CONFLICT OF INTEREST

None declared.

ORCID

Munir Aktas  <https://orcid.org/0000-0002-3188-8757>

Leonhard Schnittger  <https://orcid.org/0000-0003-3484-5370>

REFERENCES

- Ahmed, J., Yin, H., Schnittger, L., & Jongejan, F. (2002). Ticks and tick-borne diseases in Asia with special emphasis on China. *Parasitology Research*, 88, S51–S55. <https://doi.org/10.1007/s00436-001-0574-3>
- Al-Hamidhi, S., Weir, W., Kinnaird, J., Tagedin, M., Beja-Pereira, A., Morrison, I., ... Babiker, H. A. (2016). *Theileria lestoquardi* displays reduced genetic diversity relative to sympatric *Theileria annulata* in Oman. *Infection, Genetics and Evolution*, 43, 297–306. <https://doi.org/10.1016/j.meegid.2016.05.007>

- Ali, A. M., Salih, D. A., Njahira, M. N., Hassan, S. K., El Hussein, A. M., Liu, Z., ... Skilton, R. A. (2017). Genotyping of *Theileria lestoquardi* from sheep and goats in Sudan to support control of Malignant Ovine Theileriosis. *Veterinary Parasitology*, 239, 7–14. <https://doi.org/10.1016/j.vetpar.2017.04.005>
- Awad, H., Al-Hamidhi, S., El Hussein, A. M., Yousif, Y. M. Z., Taha, K. M., Salih, D. A., ... Babiker, H. A. (2018). *Theileria lestoquardi* in Sudan is highly diverse and genetically distinct from that in Oman. *Infection, Genetics and Evolution*, 62, 46–52. <https://doi.org/10.1016/j.meegid.2018.04.014>
- Aydin, L., & Bakirci, S. (2007). Geographical distribution of ticks in Turkey. *Parasitology Research*, 101, 163–166. <https://doi.org/10.1007/s00436-007-0694-5>
- Bilgic, H. B., Bakirci, S., Kose, O., Unlu, A. H., Hacilarlioglu, S., Eren, H., ... Karagenc, T. (2017). Prevalence of tick-borne haemoparasites in small ruminants in Turkey and diagnostic sensitivity of single-PCR and RLB. *Parasites & Vectors*, 27, 211. <https://doi.org/10.1186/s13071-017-2151-2153>
- Earl, D. A., & von Holdt, B. M. (2012). STRUCTURE HARVESTER: A website and program for visualizing STRUCTURE output and implementing the Evanno method. *Conservation Genetics Resources*, 4, 359–361. <https://doi.org/10.1007/s12686-011-9548-7>
- Ekici, O. D., Sevinc, F., & Isik, N. (2012). Instability of ovine babesiosis in an endemic area in Turkey. *Veterinary Parasitology*, 188, 372–375. <https://doi.org/10.1016/j.vetpar.2012.04.001>
- Erster, O., Roth, A., Wolkomirsky, R., Leibovich, B., Savitzky, I., & Shkap, V. (2016). Transmission of *Babesia ovis* by different *Rhipicephalus bursa* developmental stages and infected blood injection. *Ticks and Tick-Borne Diseases*, 7, 13–19. <https://doi.org/10.1016/j.ttbdis.2015.07.017>
- Esmailnejad, B., Tavassoli, M., Asri-Rezaei, S., Dalir-Naghadeh, B., Mardani, K., Golabi, M., ... Jalilzadeh, G. (2015). Determination of prevalence and risk factors of infection with *Babesia ovis* in small ruminants from West Azerbaijan Province, Iran by Polymerase Chain Reaction. *Journal of Arthropod-Borne Diseases*, 11, 246–252.
- Flores, D. A., Minichiello, Y., Araujo, F. R., Shkap, V., Benítez, D., Echaide, I., ... Schnittger, L. (2013). Evidence for extensive genetic diversity and substructuring of the *Babesia bovis* metapopulation. *Transboundary and Emerging Diseases*, 60, S131–S136. <https://doi.org/10.1111/tbed.12121>
- Florin-Christensen, M., Suarez, C. E., Rodriguez, A. E., Flores, D. A., & Schnittger, L. (2014). Vaccines against bovine babesiosis: Where we are now and possible roads ahead. *Parasitology*, 28, 1–30.
- Gomes, J., Salgueiro, P., Inácio, J., Amaro, A., Pinto, J., Tait, A., ... Weir, W. (2016). Population diversity of *Theileria annulata* in Portugal. *Infection, Genetics and Evolution*, 42, 14–19. <https://doi.org/10.1016/j.meegid.2016.04.023>
- Haubold, B., & Hudson, R. R. (2000). LIAN 3.0: Detecting linkage disequilibrium in multilocus data. *Linkage Analysis. Bioinformatics*, 16, 847–848. <https://doi.org/10.1093/bioinformatics/16.9.847>
- Horta, S., Barreto, M. C., Pepe, A., Campos, J., & Oliva, A. (2014). Highly sensitive method for diagnosis of subclinical *B. ovis* infection. *Ticks and Tick-borne Diseases*, 5, 902–906. <https://doi.org/10.1016/j.ttbdis.2014.07.005>
- Hurtado, A., Barandika, J. F., Oporto, B., Minguijón, E., Povedano, I., & García-Pérez, A. L. (2015). Risks of suffering tick-borne diseases in sheep translocated to a tick infested area: A laboratory approach for the investigation of an outbreak. *Ticks and Tick-borne Diseases*, 6, 31–37. <https://doi.org/10.1016/j.ttbdis.2014.09.001>
- Mahoney, D. F., & Ross, D. R. (1972). Epizootiological factors in the control of bovine babesiosis. *Australian Veterinary Journal*, 48, 292–298. <https://doi.org/10.1111/j.1751-0813.1972.tb05160.x>
- Mehlhorn, H., & Schein, E. (1984). The piroplasm: Life cycle and sexual stages. *Advances in Parasitology*, 23, 37–103.
- Peakall, R., & Smouse, P. E. (2012). GenA1Ex 6.5: Genetic analysis in excel. Population genetic software for teaching and research—an update. *Bioinformatics*, 28, 2537–2539. <https://doi.org/10.1093/bioinformatics/bts460>
- Perez-Llaneza, A., Caballero, M., Baravalle, E., Mesplet, M., Mosqueda, J., Suarez, C. E., ... Schnittger, L. (2010). Development of a tandem repeat-based multilocus typing system distinguishing *Babesia bovis* geographic isolates. *Veterinary Parasitology*, 167, 196–204. <https://doi.org/10.1016/j.vetpar.2009.09.021>
- Pritchard, J. K., Stephens, M., & Donnelly, P. (2000). Inference of population structure using multilocus genotype data. *Genetics*, 155, 945–959.
- Ranjbar-Bahadori, S., Eckert, B., Omidian, Z., Shirazi, N. S., & Shayan, P. (2012). *Babesia ovis* as the main causative agent of sheep babesiosis in Iran. *Parasitology Research*, 110, 1531–1536. <https://doi.org/10.1007/s00436-011-2658-z>
- Raymond, M., & Rousset, F. (1995). GENEPOP (version 1.2): Population genetics software for exact tests and ecumenicism. *Journal of Heredity*, 86, 248–249. <https://doi.org/10.1093/oxfordjournals.jhered.a111573>
- Rousset, F. (2008). Genepop'007: A complete reimplementation of the Genepop software for Windows and Linux. *Molecular Ecology Resources*, 8, 103–106. <https://doi.org/10.1111/j.1471-8286.2007.01931.x>
- Sayin, F., Dincer, S., Cakmak, A., Inci, A., Yukari, B. A., Vatansever, Z., ... Deniz, A. (1997c). Tick-borne diseases in Turkey. *Tropical Animal Health and Production*, 29, 535. <https://doi.org/10.1007/BF02632925>
- Sayin, F., Dincer, S., Karaer, Z., Cakmak, A., Yukari, B. A., Eren, H., ... Nalbantoglu, S. (1997a). Status of the tick-borne diseases in sheep and goats in Turkey. *Parassitologia*, 39, 153–156.
- Sayin, F., Dincer, S., Karaer, Z., Dumanli, N., Cakmak, A., Inci, A., ... Vatansever, Z. (1997b). Status of tick infestation of sheep and goats in Turkey. *Parassitologia*, 39, 145–152.
- Schnittger, L., Hong, Y., Jianxun, L., Ludwig, W., Shayan, P., Rahbari, S., & Ahmed, J. S. (2000). Phylogenetic analysis by rRNA comparison of the highly pathogenic sheep infecting parasites *Theileria lestoquardi* and a *Theileria* species identified in China. *Annals of the New York Academy of Sciences*, 916, 271–275.
- Schnittger, L., Rodriguez, A. E., Florin-Christensen, M., & Morrison, D. A. (2012). Babesia: A world emerging. *Infection, Genetics and Evolution*, 12, 1788–1809. <https://doi.org/10.1016/j.meegid.2012.07.004>
- Schnittger, L., Yin, H., Gubbels, M. J., Beyer, D., Niemann, S., Jongejan, F., & Ahmed, J. S. (2003). Phylogeny of sheep and goat *Theileria* and *Babesia* parasites. *Parasitology Research*, 91, 398–406. <https://doi.org/10.1007/s00436-003-0979-2>
- Schnittger, L., Yin, H., Qi, B., Gubbels, M. J., Beyer, D., Niemann, S., ... Ahmed, J. S. (2004). Simultaneous detection and differentiation of *Theileria* and *Babesia* parasites infecting small ruminants by reverse line blotting. *Parasitology Research*, 92, 189–196. <https://doi.org/10.1007/s00436-003-0980-9>
- Sevinc, F., Sevinc, M., Ekici, O. D., Yildiz, R., Isik, N., & Aydogdu, U. (2013). *Babesia ovis* infections: Detailed clinical and laboratory observations in the pre- and post-treatment periods of 97 field cases. *Veterinary Parasitology*, 191, 35–43. <https://doi.org/10.1016/j.vetpar.2012.07.025>
- Sevinc, F., Sevinc, M., Koc, Y., Alkan, F., Derinbay, O., Yildiz, R., ... Aydogdu, U. (2014). The effect of 12 successive blood passages on the virulence of *Babesia ovis* in splenectomized lambs: A preliminary study. *Small Ruminant Research*, 116, 66–70. <https://doi.org/10.1016/j.smallrumres.2013.10.010>
- Shayan, P., Hooshmand, E., Rahbari, S., & Nabian, S. (2007). Determination of *Rhipicephalus* spp. as vectors for *Babesia ovis* in Iran. *Parasitology Research*, 101, 1029–1033. <https://doi.org/10.1007/s00436-007-0581-0>

- Shkap, V., de Vos, A. J., Zwegarth, E., & Jongejan, F. (2007). Attenuated vaccines for tropical theileriosis, babesiosis and heartwater: The continuing necessity. *Trends in Parasitology*, 23, 420–426. <https://doi.org/10.1016/j.pt.2007.07.003>
- Simuunza, M., Bilgic, H., Karagenc, T., Syakalima, M., Shiels, B., Tait, A., & Weir, W. (2011). Population genetic analysis and sub-structuring in *Babesia bovis*. *Molecular and Biochemical Parasitology*, 177, 106–115. <https://doi.org/10.1016/j.molbiopara.2011.02.002>
- Weir, W., Karagenc, T., Gharbi, M., Simuunza, M., Aypak, S., Aysul, N., ... Tait, A. (2011). Population diversity and multiplicity of infection in *Theileria annulata*. *International Journal of Parasitology*, 41, 193–203. <https://doi.org/10.1016/j.ijpara.2010.08.004>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: Mira A, Unlu AH, Bilgic HB, et al. High genetic diversity and differentiation of the *Babesia ovis* population in Turkey. *Transbound Emerg Dis*. 2020;67(Suppl. 2):26–35. <https://doi.org/10.1111/tbed.13174>