

## SHORT REPORT

## Open Access



# Genome-wide discovery and development of polymorphic microsatellites from *Leishmania panamensis* parasites circulating in central Panama

Carlos M. Restrepo<sup>1,2</sup>, Alejandro Llanes<sup>1,2</sup>, Carolina De La Guardia<sup>1,2</sup> and Ricardo Lleonart<sup>1\*</sup> **Abstract**

**Background:** The parasite *Leishmania panamensis* is the main cause of leishmaniasis in Panama. The disease is largely uncontrolled, with a rising incidence and no appropriate control measures. While microsatellites are considered some of the best genetic markers to study population genetics and molecular epidemiology in these and other parasites, none has been developed for *L. panamensis*.

**Findings:** Here we have developed and tested a new panel of microsatellites for this species, based on high-throughput genome-wide screening. The new set of microsatellites is composed of seventeen loci, mainly spanning trinucleotide or longer motifs. We have evaluated the sensitivity and specificity of the panel based on a sample of 27 isolates obtained from cutaneous leishmaniasis patients from central Panama and also several reference species from both *L. (Leishmania)* and *L. (Viannia)* subgenera. The genetic equilibrium was assessed both intra- and inter-loci, while the reproductive mode was evaluated using several tests. The new SSR panel shows high polymorphism and sensitivity, as well as good specificity. The preliminary data described here for *L. panamensis* suggest extensive departure from Hardy-Weinberg proportions, significant linkage disequilibrium and strong deficit of heterozygotes. Several recombination tests involving multilocus linkage disequilibrium and a phylogenetic approach allowed rejection of frequent recombination in our dataset.

**Conclusions:** The genome-wide strategy described here proved to be useful to identify and test new polymorphic SSR loci in *Leishmania*. The new panel of polymorphic microsatellites is a valuable contribution to the existing molecular markers for the study of genetic structure and other aspects of this important species.

**Keywords:** *Leishmania panamensis*, Genome-wide screening, Microsatellite, Polymorphisms

**Findings**

In Panama, *Leishmania (Viannia) panamensis* is responsible for most of the reported clinical cases [1], causing cutaneous and mucocutaneous leishmaniasis. Some previous works have explored the genetic composition of local populations of this important parasite using kinetoplast DNA RFLP [2] and AFLP [3]. As these systems have some known limitations, more powerful genetic markers are needed for this species.

Full genome sequencing is increasingly being used to uncover a vast number of DNA polymorphisms in many organisms; most of them being single nucleotide polymorphisms (SNPs). While genome-wide SNP genotyping is becoming more feasible and attractive for population genetics and molecular epidemiology in many organisms, markers such as microsatellites may still have advantages over SNPs due to high polymorphism information content and fast mutation rate [4]. Microsatellites (also simple sequence repeats, SSR) are still considered very useful markers for applications involving short temporal/spatial scales, or slowly evolving or clonal organisms [4–6].

\* Correspondence: rlleonart@indicat.org.pa

<sup>1</sup>Instituto de Investigaciones Científicas y Servicios de Alta Tecnología (INDICASAT AIP), Building 219, Ciudad del Saber, Apartado 0843-01103, Ciudad de Panamá, República de Panamá

Full list of author information is available at the end of the article

There is growing evidence that some SSR may have low transferability among species, limiting the usefulness of these markers for interspecies studies [7, 8]. Additionally, the high intraspecific genetic variability reported for species of the *L. (Viannia)* subgenus further warrants the necessity to develop species-specific microsatellite panels. In this study we describe, develop and characterize a novel panel

of microsatellite markers specific for *L. panamensis* based on genome-wide high-throughput screening (Additional file 1: Extended Methods). We have used for this purpose the sequenced *L. panamensis* chromosomes (Genbank accession numbers CP009370 to CP009404) [9]. Ethical approval for the experiments described here was obtained from INDICASAT AIP Institutional Review Board (Additional

**Table 1** Primer sequences and locus characteristics.

Locus	Primers (5' → 3') <sup>a</sup>	Chr <sup>b</sup>	Loc <sup>c</sup>	Repeat motif	Region <sup>d</sup>	Ta <sup>e</sup> (°C)	Fragment sizes (bp) <sup>f</sup>
M75	TCCCTAATCGTCCGTGCAC CAGTGACGAGCTGGAAATGC	35	2041025	AGG	NC	56	258-270
M78	AACAAGGCACGGAAATTGG CGTCTGTTGCTCCGTCTTTG	35	2220491	AAC	NC	56	333-339
M80	CCGCCGAGAAGCCTGAAG GCAACAGTACGAGAAGTGGC	29	193463	AGC	C	56	237-255
M81	CATCATTAGCAGTGCAGG AAAGGAAGTATGCGTGCAC	29	246586	AGG	NC	56	329-347
M140	GATTGTCGCCCTCATCAACG GAGAAGAGAGGGTCGAGCTC	17	494896	AGG	NC	64	236-263
M147	TCTCCTTGCTTTCCTCTCC ACATATTCGCGCCTAATG	26	483774	AGG	NC	66	192-210
M149	TTCTGCCTTGTGTGCTCC CACATCGGCCTCCATAACAC	13	534713	AGG	NC	56	288-312
M155	ACAAGAATGCCAAGCTGTGC CCTCTGTGCTCCTGTTTC	13	329823	AGC	C	62	314-323
M166	ACTCCTGGATAGTGGCAAC ATGCGGTGCGTGATGTTATG	7	107131	AGC	C	62	194-206
M196	ATCAGCAGCAACTCTCCAG GAGATACCGGAGACAGCGG	12	147953	AGC	C	62	288-300
M202	CAGGGAGAAGAAGGGATGGC TCCTTTGCCTCAGTCCGTAC	5	267314	AGG	C	56	285-318
M207	CTTTGACGGCTCATCAGTG AAGGAGCGAGAAGTAGAGGC	5	251336	AGC	C	56	321-333
M233	TCTGCGGTGAGGATGCTAC GCTACTACCTCTGCTACCC	3	102343	ACC	C	56	195-300
M269	TGCTGTTGACTGAATGCACTG TTTCCTCCACTCCCGTCTTC	33	279723	AGG	NC	62	327-366
M292	GACTTAACGGATGCGCTGAC GGAGCACTGAAGTAAAGCGG	20	158471	ACC	NC	62	240-252
M329	CCAGTCTCTCCACCTACCC GATGGAAGGGTGCACACATC	27	715534	AGGG	NC	56	228-232
M402	GCAGCAGTCGATTGTGAAGG CCCTTTACATGCTGCTTGGG	5	228009	AGAGG	NC	64	130-140

<sup>a</sup>Upper line, forward primer; lower line, reverse primer. Forward primer had tail attached at 5' (GCCTCCCTCGCGCC)

<sup>b</sup>Chromosomal position based on *L. panamensis* full genome assembly

<sup>c</sup>Chromosomal location, considering numbering of the first nucleotide of repeat 1

<sup>d</sup>Location of microsatellite in coding (C) or non-coding (NC) regions

<sup>e</sup>Annealing temperature

<sup>f</sup>Range of fragment sizes found in *L. panamensis* strains tested so far

file 1: Extended Methods). As far as we know, this is the first microsatellite panel developed for *Leishmania panamensis*, using a bioinformatic pipeline at genomic level. We demonstrate that this new microsatellite panel is highly polymorphic and has good sensitivity and specificity, making it very promising for studying the genetic diversity in this parasite.

#### **In silico microsatellite screening and polymorphism assessment**

The SSR mining procedure we used on the reference *L. panamensis* genome allowed the detection of a large number of potential loci, being the AC, AG, AGC, AT, ACC, AGG and GC the most frequent motifs (Additional file 1: Extended Methods; Additional file 2: Table S1). This distribution of abundance of SSR types differ from what was found in *L. braziliensis* [10]. However, results are not directly comparable as different search strategies were used. Out of 19,297 perfect microsatellite loci, 18,921 were considered appropriate for primer design and 12,871 flagged as potentially unique sequences. Four hundred and six microsatellite loci including di-, tri-, tetra- and pentanucleotides were evaluated for consistent amplification and polymorphism using two DNA pools, each one

containing DNA from 10 different *L. panamensis* isolates. After several steps of selection, 104 potentially polymorphic loci were tested for polymorphism and consistent amplification patterns with DNA from individual isolates. All SSR loci were detected by fluorescent capillary electrophoresis using a three primer method (Additional file 1: Extended Methods). Finally, 17 microsatellite loci were selected, showing coherent segregation patterns of all peaks found in the pools, as well as low levels of stuttering (Table 1 and Table 2). The new set of *L. panamensis* microsatellites was validated by genotyping a group of 27 isolates obtained from cutaneous leishmaniasis patients from the central region of Panama (Additional file 1: Extended Methods).

We decided to exclude dinucleotide SSRs from the final panel, to improve allele assignment and stuttering levels. Additionally, we also chose to use perfect repeat loci to avoid errors due to ambiguous allele assignment and undefined marker evolution models [4].

None of the new microsatellite loci described showed more than two clear amplification peaks, suggestive of average diploidy in the corresponding chromosomes of the *L. panamensis* genome. This pattern of only 1–2 alleles per locus at microsatellite loci is consistently observed at each

**Table 2** Summary of population genetics descriptive statistics of *L. panamensis* SSRs.

Locus	Repeat number <sup>a</sup>	Genotype no <sup>b</sup>	Na <sup>c</sup>	He <sup>d</sup>	Ho <sup>e</sup>	PIC <sup>f</sup>	F <sub>IS</sub> <sup>g</sup>	HWE <sup>h</sup>
M75	5-9	2	3	0.072	0.037	0.071	0.500	0.019
M78	6-8	3	3	0.171	0.185	0.161	-0.066	1.000
M80	2-8	3	3	0.172	0.111	0.165	0.371	0.022
M81	7-13	6	6	0.407	0.111	0.387	0.736	0.000
M140	6-15	9	7	0.682	0.185	0.639	0.737	0.000
M147	6-12	7	4	0.693	0.370	0.633	0.481	0.000
M149	5-13	5	5	0.444	0.556	0.409	-0.232	0.723
M155	7-10	4	3	0.612	0.111	0.532	0.825	0.000
M166	5-9	5	4	0.655	0.259	0.584	0.616	0.000
M196	3-7	3	3	0.535	0.000	0.426	1.000	0.000
M202	9-20	7	6	0.339	0.154	0.329	0.560	0.000
M207	7-11	4	3	0.536	0.077	0.469	0.861	0.000
M233	7-42	19	15	0.887	0.519	0.877	0.431	0.000
M269	9-22	5	6	0.385	0.111	0.372	0.720	0.000
M292	5-9	4	3	0.417	0.074	0.379	0.828	0.000
M329	5-6	3	2	0.431	0.111	0.338	0.751	0.000
M402	6-8	3	3	0.426	0.111	0.371	0.748	0.000
Overall		5.4	4.6	0.463	0.181	0.420	0.620	

<sup>a</sup>Number of repeats observed

<sup>b</sup>Number of different genotypes observed

<sup>c</sup>Number of alleles

<sup>d</sup>Expected heterozygosity

<sup>e</sup>Observed heterozygosity

<sup>f</sup>Polymorphism information content

<sup>g</sup>Wright inbreeding coefficient F<sub>IS</sub>

<sup>h</sup>P-values of Hardy-Weinberg equilibrium exact test

*Leishmania* species tested so far, including *L. panamensis* from Ecuador and Peru [11].


#### Sensitivity, specificity and cross-species amplification of the microsatellite panel

Eleven loci could be amplified producing robust amplification signals even when tested with very low amounts of template *Leishmania* DNA (down to 0.01 ng) (Additional file 2: Table S3). These results may open the possibility to genotype parasites directly from infected tissues or insects with this new SSR panel. The described SSR set has high specificity for *Leishmania* DNA, showing no cross-reactivity with human DNA or with DNA from the insect vector, *Lutzomyia* sp. (Additional file 2: Table S2).

In general, the new microsatellite panel showed much better cross-species amplification with other species of the *L. (Viannia)* subgenus than with species of *L. (Leishmania)*. Almost all markers were able to show correct size amplification products when tested with isolates of *L. guyanensis*, *L. braziliensis*, *L. peruviana* and *L. lainsoni* (Additional file 2: Table S5). Several markers did not amplify in some of the species evaluated, indicating potential null alleles.

#### Genetic polymorphism and preliminary population data

Even when all *L. panamensis* isolates were collected from patients from a small geographic area, no repeated multilocus genotypes were detected (Additional file 2: Table S4). This result is congruent with AFLP data obtained with the same group of isolates, ruling out the hypothesis of full clonality in this population [3]. The polymorphism information content ranged from 0.071 (M75) to 0.877 (M233), with a mean value of 0.420 (Table 2). Markers M202 and M207 did not produce amplification products in isolates P7 and P14 respectively, indicating possible null alleles (Additional file 2: Table S4). Plotting genotypic diversity as a function of the number of loci showed that about 8–10 markers were enough to explain most of the genetic diversity in our sample of *L. panamensis* isolates (Fig. 1).




Observed heterozygosity ( $H_o$ ) ranged from 0 to 0.556 with a mean value of 0.181 and was notably lower than the expected heterozygosity ( $H_e$ ) for almost all loci (Table 2).

Within-population inbreeding coefficient ( $F_{IS}$ ) was first estimated by the method of moments, showing that all loci had positive  $F_{IS}$  (range 0.371–1; overall  $F_{IS} = 0.620$ , 95 % CI = 0.418–0.717), except markers M78 and M149. As an important step in the validation of the new microsatellite loci for population analyses, the possible presence of null alleles should be evaluated, particularly if excess homozygosity is observed. Several methods and algorithms have been reported in the literature to test microsatellites for possible null alleles [12]. However, most methods commonly used for that purpose make the assumption of Hardy-Weinberg equilibrium, which we cannot fulfil in the case of *Leishmania* data. Therefore, in order to verify if the observed positive value of  $F_{IS}$  was influenced by the possibility of having null alleles, we used Bayesian estimation of this coefficient and compared models considering or not the presence of null alleles in our data. This analysis revealed a large significant positive value ( $F_{IS} = 0.613$ , 95 % CI: 0.549–0.667). The model considering inbreeding (nfb) had a lower deviance information criterion (DIC = 1403) than the model considering only null alleles (nb, DIC = 1478), suggesting that inbreeding is the significant component of the model, rather than null alleles.

The exact test for Hardy-Weinberg equilibrium revealed a systematic departure from these theoretical proportions, as 13 out of 17 loci had significant  $P$  values (Table 2). Similarly, pairwise linkage disequilibrium (LD) was tested on all possible pairs of loci, showing significant values. Out of 136 comparisons, and expecting only seven significant values by chance, 40 pairs of loci showed  $P$  values below the strict Bonferroni corrected threshold (0.00036) (Additional file 2: Table S6).

Furthermore, multilocus LD was assessed using the new microsatellites on our sample. While the expected mean values for measures of multilocus LD under panmixia are around zero, the values observed were significantly larger ( $I_A = 2.19$  and  $\bar{r}_d = 0.145$ ; both significant at  $P < 0.001$ ) (Fig. 2a and b). The phylogenetic test for recombination shows an observed parsimony tree length of 163, significantly shorter ( $P < 0.001$ ) than those generated from shuffled datasets (mean tree length: 229 steps, Fig. 3). Both approaches allow rejection of the null hypothesis of panmixia. Using a different marker system on the same group of isolates, we have already shown strong signatures of LD, as independent AFLP markers generated highly correlated distance matrices between isolates [3]. Diverse levels of LD have been reported in various *Leishmania* species using several marker systems [13–15]. Due to the limited sample size we cannot rule




out that some additional factors may be influencing the LD estimations reported here, such as Wahlund effect, hidden population subdivision, cryptic species within the sampling units, recombination rate, genetic drift, mutation rates, epistasis or selection.

Although several tests performed on our data allow rejection of the hypothesis of panmixia/frequent recombination, the values of association indexes as well as those of inbreeding coefficient are not consistent with strict clonality either. Predominant clonality would produce strongly negative  $F_{IS}$  values and a higher index of association ( $I_A$  and  $\bar{r}_d$ ). Taken together, these results suggest a mixed mode of reproduction that involves both

clonality and selfing with sporadic recombination. The same mode of reproduction has been suggested for other species of the *L. (Viannia)* subgenus, including *L. braziliensis* and *L. guyanensis* [13, 16] as well as some other parasites such as *Trypanosoma brucei* and *Plasmodium falciparum* [17, 18]. In the closely related *L. guyanensis*, a simulation approach has shown that a reproductive mating scheme involving frequent sexual events may account for the observed levels of linkage disequilibrium and heterozygote deficit [16].

We have developed and tested a new panel of microsatellites using *L. panamensis* genomic sequence data. The bioinformatic pipeline employed allowed us to identify



significantly more potential loci for evaluation than previously reported in *Leishmania* using conventional methodologies [19, 20]. The new panel of microsatellites will contribute to the existing tools for better understanding of the distribution and dynamics of the genetic diversity in this important parasite.

## Additional files

**Additional file 1: Extended Methods.** (DOCX 32 kb)

**Additional file 2: Supplementary Data Table S1.** Perfect microsatellite loci found in *L. panamensis* genome. **Table S2.** Specificity of new *L. panamensis* SSR panel on non-target DNAs. **Table S3.** Characterization of limits of detection for each described *L. panamensis* SSR locus. **Table S4.** Raw genotype data for all tested loci and *Leishmania* species. **Table S5.** Amplification profiles of described panel of microsatellites in species other than *L. panamensis*. **Table S6.** *P* values for exact test of pairwise linkage disequilibrium, as estimated in Arlequin v3.5.1.3. (XLSX 28 kb)

## Competing interests

The authors declare that they have no competing interests.

## Authors' contributions

RL conceived and designed the experiments; CMR, CD and AL performed the experiments; CMR, AL, CD and RL analyzed the data; RL, CMR, CD and AL wrote and reviewed the manuscript. All authors read and approved this manuscript.

## Acknowledgements

We deeply thank Dr. Patricia Llanes, Dr. José Calzada, Dr. Octavio Sousa, Dr. Marcelo Bozza and Dr. Carmenza Spadafora for the kind gift of *Leishmania* strains; and Dr. Gabrielle Britton for reviewing the manuscript. We thank the reviewers for careful examination of the manuscript, leading to its improvement. This project was partially financed by the Panamanian National Secretary of Science, Technology and Innovation (SENACYT; Grants No. FID08-104, CID09-010 and SNI187-2011).

## Author details

<sup>1</sup>Instituto de Investigaciones Científicas y Servicios de Alta Tecnología (INDICASAT AIP), Building 219, Ciudad del Saber, Apartado 0843-01103, Ciudad de Panamá, República de Panamá. <sup>2</sup>Department of Biotechnology, Acharya Nagarjuna University, Guntur, India.

Received: 28 May 2015 Accepted: 8 October 2015

Published online: 12 October 2015

## References

- Vásquez A, Paz H, Alvar J, Pérez D, Hernández C. Informe Final: Estudios Sobre La Epidemiología de La Leishmaniasis En La Parte Occidental de La República de Panamá. 1998.
- Miranda A, Carrasco R, Paz H, Pascale JM, Samudio F, Saldaña A, et al. Molecular epidemiology of American tegumentary leishmaniasis in Panama. *Am J Trop Med Hyg.* 2009;81:565–71.
- Restrepo CM, De La Guardia C, Sousa OE, Calzada JE, Fernández PL, Leonart R. AFLP polymorphisms allow high resolution genetic analysis of American Tegumentary Leishmaniasis agents circulating in Panama and other members of the *Leishmania*. *PLoS One.* 2013;8, e73177.
- Guichoux E, Lagache L, Wagner S, Chaumeil P, Léger P, Lepais O, et al. Current trends in microsatellite genotyping. *Mol Ecol Resour.* 2011;11:591–611.
- Bhargava A, Fuentes FF. Mutational dynamics of microsatellites. *Mol Biotechnol.* 2010;44:250–66.
- Putman AJ, Carbone I. Challenges in analysis and interpretation of microsatellite data for population genetic studies. *Ecol Evol.* 2014;4:4399–428.
- Jamjoom MB, Ashford RW, Bates PA, Kemp SJ, Noyes HA. Polymorphic microsatellite repeats are not conserved between *Leishmania donovani* and *Leishmania major*. *Mol Ecol Notes.* 2002;2:104–6.

- Schwenkenbecher JM, Fröhlich C, Gehre F, Schnur LF, Schönián G. Evolution and conservation of microsatellite markers for *Leishmania tropica*. *Infect Genet Evol.* 2004;4:99–105.
- Llanes A, Restrepo CM, Del Vecchio G, Anguizola FJ, Leonart R: The genome of *Leishmania panamensis*: insights into genomics of the *L. (Viannia)* subgenus. *Sci Rep.* 2015:8550. doi:10.1038/srep08550.
- Fakhar M, Motazedian MH, Daly D, Lowe CD, Kemp SJ, Noyes HA. An integrated pipeline for the development of novel panels of mapped microsatellite markers for *Leishmania donovani* complex, *Leishmania braziliensis* and *Leishmania major*. *Parasitology.* 2008;567–574.
- Rougeron V, De Meeüs T, Hide M, Walecx E, Dereure J, Arevalo J, Llanos-Cuentas A, Bañuls AL: A battery of 12 microsatellite markers for genetic analysis of the *Leishmania (Viannia) guyanensis* complex. *Parasitology* 2010:1879–84. doi:10.1017/S0031182010000776.
- van Oosterhout C, Weetman D, Hutchinson WF. Estimation and adjustment of microsatellite null alleles in nonequilibrium populations. *Mol Ecol Notes.* 2006;255–256.
- Rougeron V, De Meeüs T, Hide M, Walecx E, Bermudez H, Arevalo J, et al. Extreme inbreeding in *Leishmania braziliensis*. *Proc Natl Acad Sci U S A.* 2009;106:10224–9.
- Tibayrenc M, Ayala FJ. How clonal are *Trypanosoma* and *Leishmania*? *Trends Parasitol.* 2013;29:264–9.
- Baleela R, Llewellyn MS, Fitzpatrick S, Kuhls K, Schönián G, Miles MA, et al. *Leishmania donovani* populations in Eastern Sudan: temporal structuring and a link between human and canine transmission. *Parasit Vectors.* 2014;7:496.
- Rougeron V, Bañuls AL, Carme B, Simon S, Couppié P, Nacher M, et al. Reproductive strategies and population structure in *Leishmania*: substantial amount of sex in *Leishmania Viannia guyanensis*. *Mol Ecol.* 2011;3116–3127.
- Gibson W, Peacock L, Ferris V, Williams K, Bailey M. The use of yellow fluorescent hybrids to indicate mating in *Trypanosoma brucei*. *Parasit Vectors.* 2008;1:4.
- Annan Z, Durand P, Ayala FJ, Arnathau C, Awono-Ambene P, Simard F, et al. Population genetic structure of *Plasmodium falciparum* in the two main African vectors, *Anopheles gambiae* and *Anopheles funestus*. *Proc Natl Acad Sci U S A.* 2007;104:7987–92.
- Bulle B, Millon L, Bart JM, Gállego M, Gambarelli F, Portús M, et al. Practical Approach for Typing Strains of *Leishmania infantum* by Microsatellite Analysis. *J Clin Microbiol.* 2002;40(9):3391–7.
- Russell R, Iribar MP, Lambson B, Brewster S, Blackwell JM, Dye C, et al. Intra and inter-specific microsatellite variation in the *Leishmania* subgenus *Viannia*. *Mol Biochem Parasitol.* 1999;103:71–7.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit

