

## 1I. Vitamin D receptor-like receptors (version 2019.4) in the IUPHAR/BPS Guide to Pharmacology Database

Sylvia Christakos<sup>1</sup>

1. Rutgers University, USA

### Abstract

Vitamin D (VDR), Pregnane X (PXR) and Constitutive Androstane (CAR) receptors (**nomenclature as agreed by the NC-IUPHAR Subcommittee on Nuclear Hormone Receptors [47]**) are members of the NR1I family of nuclear receptors, which form heterodimers with members of the retinoid X receptor family. PXR and CAR are activated by a range of exogenous compounds, with no established endogenous physiological agonists, although high concentrations of bile acids and bile pigments activate PXR and CAR [47].

### Contents

This is a citation summary for 1I. Vitamin D receptor-like receptors in the [Guide to Pharmacology](#) database (GtoPdb). It exists purely as an adjunct to the database to facilitate the recognition of citations to and from the database by citation analyzers. Readers will almost certainly want to visit the relevant sections of the database which are given here under database links.

[GtoPdb](#) is an expert-driven guide to pharmacological targets and the substances that act on them. GtoPdb is a reference work which is most usefully represented as an on-line database. As in any publication this work should be appropriately cited, and the papers it cites should also be recognized. This document provides a citation for the relevant parts of the database, and also provides a reference list for the research cited by those parts.

Please note that the database version for the citations given in GtoPdb are to the most recent preceding version in which the family or its subfamilies and targets were substantially changed. The links below are to the current version. If you need to consult the cited version, rather than the most recent version, please contact the GtoPdb curators.

### Database links

View metadata, citation and similar papers at [CORE](#)  <http://www.guidetopharmacology.org/GRAC/FamilyDisplayForward?familyId=90>

[Introduction to 1I. Vitamin D receptor-like receptors](#)

<http://www.guidetopharmacology.org/GRAC/FamilyIntroductionForward?familyId=90>

Receptors

[Vitamin D receptor](#)

<http://www.guidetopharmacology.org/GRAC/ObjectDisplayForward?objectId=605>

[Pregnane X receptor](#)

## References

1. Baes M, Gulick T, Choi HS, Martinoli MG, Simha D and Moore DD. (1994) A new orphan member of the nuclear hormone receptor superfamily that interacts with a subset of retinoic acid response elements. *Mol. Cell. Biol.* **14**: 1544-52 [PMID:8114692]
2. Baker AR, McDonnell DP, Hughes M, Crisp TM, Mangelsdorf DJ, Haussler MR, Pike JW, Shine J and O'Malley BW. (1988) Cloning and expression of full-length cDNA encoding human vitamin D receptor. *Proc. Natl. Acad. Sci. U.S.A.* **85**: 3294-8 [PMID:2835767]
3. Bertilsson G, Heidrich J, Svensson K, Asman M, Jendeborg L, Sydow-Bäckman M, Ohlsson R, Postlind H, Blomquist P and Berkenstam A. (1998) Identification of a human nuclear receptor defines a new signaling pathway for CYP3A induction. *Proc. Natl. Acad. Sci. U.S.A.* **95**: 12208-13 [PMID:9770465]
4. Bhalla S, Ozalp C, Fang S, Xiang L and Kemper JK. (2004) Ligand-activated pregnane X receptor interferes with HNF-4 signaling by targeting a common coactivator PGC-1alpha. Functional implications in hepatic cholesterol and glucose metabolism. *J. Biol. Chem.* **279**: 45139-47 [PMID:15322103]
5. Bishop JE, Collins ED, Okamura WH and Norman AW. (1994) Profile of ligand specificity of the vitamin D binding protein for 1 alpha,25-dihydroxyvitamin D3 and its analogs. *J. Bone Miner. Res.* **9**: 1277-88 [PMID:7976510]
6. Blumberg B, Sabbagh W, Juguilon H, Bolado J, van Meter CM, Ong ES and Evans RM. (1998) SXR, a novel steroid and xenobiotic-sensing nuclear receptor. *Genes Dev.* **12**: 3195-205 [PMID:9784494]
7. Boehm MF, Fitzgerald P, Zou A, Elgort MG, Bischoff ED, Mere L, Mais DE, Bissonnette RP, Heyman RA, Nadzan AM, Reichman M and Allegretto EA. (1999) Novel nonsteroidal vitamin D mimics exert VDR-modulating activities with less calcium mobilization than 1,25-dihydroxyvitamin D3. *Chem. Biol.* **6**: 265-75 [PMID:10322128]
8. Burmester JK, Maeda N and DeLuca HF. (1988) Isolation and expression of rat 1,25-dihydroxyvitamin D3 receptor cDNA. *Proc. Natl. Acad. Sci. U.S.A.* **85**: 1005-9 [PMID:2829212]
9. Cavailles V, Dauvois S, L'Horset F, Lopez G, Hoare S, Kushner PJ and Parker MG. (1995) Nuclear factor RIP140 modulates transcriptional activation by the estrogen receptor. *EMBO J.* **14**: 3741-51 [PMID:7641693]
10. Chen Y, Kissling G, Negishi M and Goldstein JA. (2005) The nuclear receptors constitutive androstane receptor and pregnane X receptor cross-talk with hepatic nuclear factor 4alpha to synergistically activate the human CYP2C9 promoter. *J. Pharmacol. Exp. Ther.* **314**: 1125-33 [PMID:15919766]
11. Choi HS, Chung M, Tzameli I, Simha D, Lee YK, Seol W and Moore DD. (1997) Differential transactivation by two isoforms of the orphan nuclear hormone receptor CAR. *J. Biol. Chem.* **272**: 23565-71 [PMID:9295294]
12. Colston KW, Mackay AG, James SY, Binderup L, Chander S and Coombes RC. (1992) EB1089: a new vitamin D analogue that inhibits the growth of breast cancer cells in vivo and in vitro. *Biochem. Pharmacol.* **44**: 2273-80 [PMID:1472092]
13. Dotzlaw H, Leygue E, Watson P and Murphy LC. (1999) The human orphan receptor PXR messenger RNA is expressed in both normal and neoplastic breast tissue. *Clin. Cancer Res.* **5**: 2103-7 [PMID:10473093]
14. Dussault I, Yoo HD, Lin M, Wang E, Fan M, Batta AK, Salen G, Erickson SK and Forman BM. (2003) Identification of an endogenous ligand that activates pregnane X receptor-mediated sterol clearance. *Proc. Natl. Acad. Sci. U.S.A.* **100**: 833-8 [PMID:12569201]
15. Erben RG, Soegiarto DW, Weber K, Zeitz U, Lieberherr M, Gniadecki R, Möller G, Adamski J and Balling R. (2002) Deletion of deoxyribonucleic acid binding domain of the vitamin D receptor abrogates genomic and nongenomic functions of vitamin D. *Mol. Endocrinol.* **16**: 1524-37 [PMID:12089348]
16. Ferguson SS, Chen Y, LeCluyse EL, Negishi M and Goldstein JA. (2005) Human CYP2C8 is

- transcriptionally regulated by the nuclear receptors constitutive androstane receptor, pregnane X receptor, glucocorticoid receptor, and hepatic nuclear factor 4alpha. *Mol. Pharmacol.* **68**: 747-57 [PMID:15933212]
17. Forman BM, Tzamelis I, Choi HS, Chen J, Simha D, Seol W, Evans RM and Moore DD. (1998) Androstane metabolites bind to and deactivate the nuclear receptor CAR-beta. *Nature* **395**: 612-5 [PMID:9783588]
  18. Frank C, Gonzalez MM, Oinonen C, Dunlop TW and Carlberg C. (2003) Characterization of DNA complexes formed by the nuclear receptor constitutive androstane receptor. *J. Biol. Chem.* **278**: 43299-310 [PMID:12896978]
  19. Fraser DJ, Zumsteg A and Meyer UA. (2003) Nuclear receptors constitutive androstane receptor and pregnane X receptor activate a drug-responsive enhancer of the murine 5-aminolevulinic acid synthase gene. *J. Biol. Chem.* **278**: 39392-401 [PMID:12881517]
  20. Fujishima T, Kojima Y, Azumaya I, Kittaka A and Takayama H. (2003) Design and synthesis of potent vitamin D receptor antagonists with A-ring modifications: remarkable effects of 2alpha-methyl introduction on antagonistic activity. *Bioorg. Med. Chem.* **11**: 3621-31 [PMID:12901907]
  21. Geick A, Eichelbaum M and Burk O. (2001) Nuclear receptor response elements mediate induction of intestinal MDR1 by rifampin. *J. Biol. Chem.* **276**: 14581-7 [PMID:11297522]
  22. Germain P, Staels B, Dacquet C, Spedding M and Laudet V. (2006) Overview of nomenclature of nuclear receptors. *Pharmacol. Rev.* **58**: 685-704 [PMID:17132848]
  23. Gonzalez MM and Carlberg C. (2002) Cross-repression, a functional consequence of the physical interaction of non-liganded nuclear receptors and POU domain transcription factors. *J. Biol. Chem.* **277**: 18501-9 [PMID:11891224]
  24. Goodwin B, Gauthier KC, Umetani M, Watson MA, Lochansky MI, Collins JL, Leitersdorf E, Mangelsdorf DJ, Kliewer SA and Repa JJ. (2003) Identification of bile acid precursors as endogenous ligands for the nuclear xenobiotic pregnane X receptor. *Proc. Natl. Acad. Sci. U.S.A.* **100**: 223-8 [PMID:12509506]
  25. Goodwin B, Moore LB, Stoltz CM, McKee DD and Kliewer SA. (2001) Regulation of the human CYP2B6 gene by the nuclear pregnane X receptor. *Mol. Pharmacol.* **60**: 427-31 [PMID:11502872]
  26. Guo GL, Staudinger J, Ogura K and Klaassen CD. (2002) Induction of rat organic anion transporting polypeptide 2 by pregnenolone-16alpha-carbonitrile is via interaction with pregnane X receptor. *Mol. Pharmacol.* **61**: 832-9 [PMID:11901222]
  27. Herdick M, Steinmeyer A and Carlberg C. (2000) Carboxylic ester antagonists of 1alpha,25-dihydroxyvitamin D(3) show cell-specific actions. *Chem. Biol.* **7**: 885-94 [PMID:11094341]
  28. Honkakoski P, Sueyoshi T and Negishi M. (2003) Drug-activated nuclear receptors CAR and PXR. *Ann. Med.* **35**: 172-82 [PMID:12822739]
  29. Huang W, Zhang J, Washington M, Liu J, Parant JM, Lozano G and Moore DD. (2005) Xenobiotic stress induces hepatomegaly and liver tumors via the nuclear receptor constitutive androstane receptor. *Mol. Endocrinol.* **19**: 1646-53 [PMID:15831521]
  30. Huang W, Zhang J, Wei P, Schrader WT and Moore DD. (2004) Meclizine is an agonist ligand for mouse constitutive androstane receptor (CAR) and an inverse agonist for human CAR. *Mol. Endocrinol.* **18**: 2402-8 [PMID:15272053]
  31. Hughes MR, Malloy PJ, Kieback DG, Kesterson RA, Pike JW, Feldman D and O'Malley BW. (1988) Point mutations in the human vitamin D receptor gene associated with hypocalcemic rickets. *Science* **242**: 1702-5 [PMID:2849209]
  32. Ikeda M, Takahashi K, Dan A, Koyama K, Kubota K, Tanaka T and Hayashi M. (2000) Synthesis and biological evaluations of A-ring isomers of 26,26,26,27,27,27-hexafluoro-1,25-dihydroxyvitamin D3. *Bioorg. Med. Chem.* **8**: 2157-66 [PMID:11003160]
  33. Jia Y, Guo GL, Surapureddi S, Sarkar J, Qi C, Guo D, Xia J, Kashireddi P, Yu S, Cho YW, Rao MS, Kemper B, Ge K, Gonzalez FJ and Reddy JK. (2005) Transcription coactivator peroxisome proliferator-activated receptor-binding protein/mediator 1 deficiency abrogates acetaminophen hepatotoxicity. *Proc. Natl. Acad. Sci. U.S.A.* **102**: 12531-6 [PMID:16109766]
  34. Jones SA, Moore LB, Shenk JL, Wisely GB, Hamilton GA, McKee DD, Tomkinson NC, LeCluyse EL, Lambert MH, Willson TM, Kliewer SA and Moore JT. (2000) The pregnane X receptor: a promiscuous

- xenobiotic receptor that has diverged during evolution. *Mol. Endocrinol.* **14**: 27-39 [PMID:10628745]
35. Kamei Y, Kawada T, Fukuwatari T, Ono T, Kato S and Sugimoto E. (1995) Cloning and sequencing of the gene encoding the mouse vitamin D receptor. *Gene* **152**: 281-2 [PMID:7835717]
  36. Kanno Y, Tanuma N, Takahashi A and Inouye Y. (2013) TO901317, a potent LXR agonist, is an inverse agonist of CAR. *J Toxicol Sci* **38**: 309-15 [PMID:23665929]
  37. Kast HR, Goodwin B, Tarr PT, Jones SA, Anisfeld AM, Stoltz CM, Tontonoz P, Kliewer S, Willson TM and Edwards PA. (2002) Regulation of multidrug resistance-associated protein 2 (ABCC2) by the nuclear receptors pregnane X receptor, farnesoid X-activated receptor, and constitutive androstane receptor. *J. Biol. Chem.* **277**: 2908-15 [PMID:11706036]
  38. Kliewer SA, Moore JT, Wade L, Staudinger JL, Watson MA, Jones SA, McKee DD, Oliver BB, Willson TM, Zetterström RH, Perlmann T and Lehmann JM. (1998) An orphan nuclear receptor activated by pregnanes defines a novel steroid signaling pathway. *Cell* **92**: 73-82 [PMID:9489701]
  39. Kliewer SA, Umesono K, Mangelsdorf DJ and Evans RM. (1992) Retinoid X receptor interacts with nuclear receptors in retinoic acid, thyroid hormone and vitamin D3 signalling. *Nature* **355**: 446-9 [PMID:1310351]
  40. Kodama S, Koike C, Negishi M and Yamamoto Y. (2004) Nuclear receptors CAR and PXR cross talk with FOXO1 to regulate genes that encode drug-metabolizing and gluconeogenic enzymes. *Mol. Cell. Biol.* **24**: 7931-40 [PMID:15340055]
  41. Lehmann JM, McKee DD, Watson MA, Willson TM, Moore JT and Kliewer SA. (1998) The human orphan nuclear receptor PXR is activated by compounds that regulate CYP3A4 gene expression and cause drug interactions. *J. Clin. Invest.* **102**: 1016-23 [PMID:9727070]
  42. Li YC, Pirro AE, Amling M, Delling G, Baron R, Bronson R and Demay MB. (1997) Targeted ablation of the vitamin D receptor: an animal model of vitamin D-dependent rickets type II with alopecia. *Proc. Natl. Acad. Sci. U.S.A.* **94**: 9831-5 [PMID:9275211]
  43. Maglich JM, Parks DJ, Moore LB, Collins JL, Goodwin B, Billin AN, Stoltz CA, Kliewer SA, Lambert MH, Willson TM and Moore JT. (2003) Identification of a novel human constitutive androstane receptor (CAR) agonist and its use in the identification of CAR target genes. *J. Biol. Chem.* **278**: 17277-83 [PMID:12611900]
  44. Makishima M, Lu TT, Xie W, Whitfield GK, Domoto H, Evans RM, Haussler MR and Mangelsdorf DJ. (2002) Vitamin D receptor as an intestinal bile acid sensor. *Science* **296**: 1313-6 [PMID:12016314]
  45. Miki Y, Suzuki T, Kitada K, Yabuki N, Shibuya R, Moriya T, Ishida T, Ohuchi N, Blumberg B and Sasano H. (2006) Expression of the steroid and xenobiotic receptor and its possible target gene, organic anion transporting polypeptide-A, in human breast carcinoma. *Cancer Res.* **66**: 535-42 [PMID:16397270]
  46. Miyamoto K, Murayama E, Ochi K, Watanabe H and Kubodera N. (1993) Synthetic studies of vitamin D analogues. XIV. Synthesis and calcium regulating activity of vitamin D3 analogues bearing a hydroxyalkoxy group at the 2 beta-position. *Chem. Pharm. Bull.* **41**: 1111-3 [PMID:8396500]
  47. Moore DD, Kato S, Xie W, Mangelsdorf DJ, Schmidt DR, Xiao R and Kliewer SA. (2006) International Union of Pharmacology. LXII. The NR1H and NR1I receptors: constitutive androstane receptor, pregnane X receptor, farnesoid X receptor alpha, farnesoid X receptor beta, liver X receptor alpha, liver X receptor beta, and vitamin D receptor. *Pharmacol. Rev.* **58**: 742-59 [PMID:17132852]
  48. Moore LB, Goodwin B, Jones SA, Wisely GB, Serabjit-Singh CJ, Willson TM, Collins JL and Kliewer SA. (2000) St. John's wort induces hepatic drug metabolism through activation of the pregnane X receptor. *Proc. Natl. Acad. Sci. U.S.A.* **97**: 7500-2 [PMID:10852961]
  49. Moore LB, Parks DJ, Jones SA, Bledsoe RK, Consler TG, Stimmel JB, Goodwin B, Liddle C, Blanchard SG, Willson TM, Collins JL and Kliewer SA. (2000) Orphan nuclear receptors constitutive androstane receptor and pregnane X receptor share xenobiotic and steroid ligands. *J. Biol. Chem.* **275**: 15122-7 [PMID:10748001]
  50. Mu Y, Stephenson CR, Kendall C, Saini SP, Toma D, Ren S, Cai H, Strom SC, Day BW, Wipf P and Xie W. (2005) A pregnane X receptor agonist with unique species-dependent stereoselectivity and its implications in drug development. *Mol. Pharmacol.* **68**: 403-13 [PMID:15872116]
  51. Mu Y, Zhang J, Zhang S, Zhou HH, Toma D, Ren S, Huang L, Yaramus M, Baum A, Venkataramanan R

- and Xie W. (2006) Traditional Chinese medicines Wu Wei Zi (*Schisandra chinensis* Baill) and Gan Cao (*Glycyrrhiza uralensis* Fisch) activate pregnane X receptor and increase warfarin clearance in rats. *J. Pharmacol. Exp. Ther.* **316**: 1369-77 [PMID:16267138]
52. Nishimura M, Naito S and Yokoi T. (2004) Tissue-specific mRNA expression profiles of human nuclear receptor subfamilies. *Drug Metab. Pharmacokinet.* **19**: 135-49 [PMID:15499180]
  53. Ourlin JC, Lasserre F, Pineau T, Fabre JM, Sa-Cunha A, Maurel P, Vilarem MJ and Pascussi JM. (2003) The small heterodimer partner interacts with the pregnane X receptor and represses its transcriptional activity. *Mol. Endocrinol.* **17**: 1693-703 [PMID:12805410]
  54. Peleg S, Uskokovic M, Ahene A, Vickery B and Avnur Z. (2002) Cellular and molecular events associated with the bone-protecting activity of the noncalcemic vitamin D analog Ro-26-9228 in osteopenic rats. *Endocrinology* **143**: 1625-36 [PMID:11956143]
  55. Rochel N, Wurtz JM, Mitschler A, Klaholz B and Moras D. (2000) The crystal structure of the nuclear receptor for vitamin D bound to its natural ligand. *Mol. Cell* **5**: 173-9 [PMID:10678179]
  56. Saito N, Matsunaga T, Saito H, Anzai M, Takenouchi K, Miura D, Namekawa J, Ishizuka S and Kittaka A. (2006) Further synthetic and biological studies on vitamin D hormone antagonists based on C24-alkylation and C2alpha-functionalization of 25-dehydro-1alpha-hydroxyvitamin D(3)-26,23-lactones. *J. Med. Chem.* **49**: 7063-75 [PMID:17125259]
  57. Sato K, Nishii Y, Woodiel FN and Raisz LG. (1993) Effects of two new vitamin D3 derivatives, 22-oxa-1 alpha-25-dihydroxyvitamin D3 (OCT) and 2 beta-(3-hydroxypropoxy)-1 alpha, 25-dihydroxyvitamin D3 (ED-71), on bone metabolism in organ culture. *Bone* **14**: 47-51 [PMID:8443002]
  58. Schuetz EG, Brimer C and Schuetz JD. (1998) Environmental xenobiotics and the antihormones cyproterone acetate and spironolactone use the nuclear hormone pregnenolone X receptor to activate the CYP3A23 hormone response element. *Mol. Pharmacol.* **54**: 1113-7 [PMID:9855641]
  59. Shan L, Vincent J, Brunzelle JS, Dussault I, Lin M, Ianculescu I, Sherman MA, Forman BM and Fernandez EJ. (2004) Structure of the murine constitutive androstane receptor complexed to androstenol: a molecular basis for inverse agonism. *Mol. Cell* **16**: 907-17 [PMID:15610734]
  60. Shiina Y, Abe E, Miyaura C, Tanaka H, Yamada S, Ohmori M, Nakayama K, Takayama H, Matsunaga I, Nishii Y, DeLuca HF and Suda T. (1983) Biological activity of 24,24-difluoro-1 alpha, 25-dihydroxyvitamin D3 and 1 alpha, 25-dihydroxyvitamin D3-26,23-lactone in inducing differentiation of human myeloid leukemia cells. *Arch. Biochem. Biophys.* **220**: 90-4 [PMID:6572492]
  61. Shiraki T, Sakai N, Kanaya E and Jingami H. (2003) Activation of orphan nuclear constitutive androstane receptor requires subnuclear targeting by peroxisome proliferator-activated receptor gamma coactivator-1 alpha. A possible link between xenobiotic response and nutritional state. *J. Biol. Chem.* **278**: 11344-50 [PMID:12551939]
  62. Sicinski RR, Prahil JM, Smith CM and DeLuca HF. (1998) New 1alpha,25-dihydroxy-19-norvitamin D3 compounds of high biological activity: synthesis and biological evaluation of 2-hydroxymethyl, 2-methyl, and 2-methylene analogues. *J. Med. Chem.* **41**: 4662-74 [PMID:9804706]
  63. Sonoda J, Xie W, Rosenfeld JM, Barwick JL, Guzelian PS and Evans RM. (2002) Regulation of a xenobiotic sulfonation cascade by nuclear pregnane X receptor (PXR). *Proc. Natl. Acad. Sci. U.S.A.* **99**: 13801-6 [PMID:12370413]
  64. Staudinger JL, Goodwin B, Jones SA, Hawkins-Brown D, MacKenzie KI, LaTour A, Liu Y, Klaassen CD, Brown KK, Reinhard J, Willson TM, Koller BH and Kliewer SA. (2001) The nuclear receptor PXR is a lithocholic acid sensor that protects against liver toxicity. *Proc. Natl. Acad. Sci. U.S.A.* **98**: 3369-74 [PMID:11248085]
  65. Sueyoshi T, Kawamoto T, Zelko I, Honkakoski P and Negishi M. (1999) The repressed nuclear receptor CAR responds to phenobarbital in activating the human CYP2B6 gene. *J. Biol. Chem.* **274**: 6043-6 [PMID:10037683]
  66. Sugatani J, Kojima H, Ueda A, Kakizaki S, Yoshinari K, Gong QH, Owens IS, Negishi M and Sueyoshi T. (2001) The phenobarbital response enhancer module in the human bilirubin UDP-glucuronosyltransferase UGT1A1 gene and regulation by the nuclear receptor CAR. *Hepatology* **33**: 1232-8 [PMID:11343253]

67. Sugatani J, Nishitani S, Yamakawa K, Yoshinari K, Sueyoshi T, Negishi M and Miwa M. (2005) Transcriptional regulation of human UGT1A1 gene expression: activated glucocorticoid receptor enhances constitutive androstane receptor/pregnane X receptor-mediated UDP-glucuronosyltransferase 1A1 regulation with glucocorticoid receptor-interacting protein 1. *Mol. Pharmacol.* **67**: 845-55 [PMID:15557560]
68. Synold TW, Dussault I and Forman BM. (2001) The orphan nuclear receptor SXR coordinately regulates drug metabolism and efflux. *Nat. Med.* **7**: 584-90 [PMID:11329060]
69. Tabb MM, Sun A, Zhou C, Grün F, Errandi J, Romero K, Pham H, Inoue S, Mallick S, Lin M, Forman BM and Blumberg B. (2003) Vitamin K2 regulation of bone homeostasis is mediated by the steroid and xenobiotic receptor SXR. *J. Biol. Chem.* **278**: 43919-27 [PMID:12920130]
70. Takeshita A, Taguchi M, Koibuchi N and Ozawa Y. (2002) Putative role of the orphan nuclear receptor SXR (steroid and xenobiotic receptor) in the mechanism of CYP3A4 inhibition by xenobiotics. *J. Biol. Chem.* **277**: 32453-8 [PMID:12072427]
71. Tzamelis I, Pissios P, Schuetz EG and Moore DD. (2000) The xenobiotic compound 1,4-bis[2-(3,5-dichloropyridyloxy)]benzene is an agonist ligand for the nuclear receptor CAR. *Mol. Cell. Biol.* **20**: 2951-8 [PMID:10757780]
72. Verlinden L, Verstuyf A, Van Camp M, Marcelis S, Sabbe K, Zhao XY, De Clercq P, Vandewalle M and Bouillon R. (2000) Two novel 14-Epi-analogues of 1,25-dihydroxyvitamin D3 inhibit the growth of human breast cancer cells in vitro and in vivo. *Cancer Res.* **60**: 2673-9 [PMID:10825140]
73. Wang W, Prosser WW, Chen J, Taremi SS, Le HV, Madison V, Cui X, Thomas A, Cheng KC and Lesburg CA. (2008) Construction and characterization of a fully active PXR/SRC-1 tethered protein with increased stability. *Protein Eng. Des. Sel.* **21**: 425-33 [PMID:18456871]
74. Wei P, Zhang J, Dowhan DH, Han Y and Moore DD. (2002) Specific and overlapping functions of the nuclear hormone receptors CAR and PXR in xenobiotic response. *Pharmacogenomics J.* **2**: 117-26 [PMID:12049174]
75. Wei P, Zhang J, Egan-Hafley M, Liang S and Moore DD. (2000) The nuclear receptor CAR mediates specific xenobiotic induction of drug metabolism. *Nature* **407**: 920-3 [PMID:11057673]
76. Wentworth JM, Agostini M, Love J, Schwabe JW and Chatterjee VK. (2000) St John's wort, a herbal antidepressant, activates the steroid X receptor. *J. Endocrinol.* **166**: R11-6 [PMID:10974665]
77. Weyts FA, Dhawan P, Zhang X, Bishop JE, Uskokovic MR, Ji Y, Studzinski GP, Norman AW and Christakos S. (2004) Novel Gemini analogs of 1 $\alpha$ ,25-dihydroxyvitamin D(3) with enhanced transcriptional activity. *Biochem. Pharmacol.* **67**: 1327-36 [PMID:15013848]
78. Wiberg K, Ljunghall S, Binderup L and Ljunggren O. (1995) Studies on two new vitamin D analogs, EB 1089 and KH 1060: effects on bone resorption and osteoclast recruitment in vitro. *Bone* **17**: 391-5 [PMID:8573413]
79. Wyde ME, Bartolucci E, Ueda A, Zhang H, Yan B, Negishi M and You L. (2003) The environmental pollutant 1,1-dichloro-2,2-bis (p-chlorophenyl)ethylene induces rat hepatic cytochrome P450 2B and 3A expression through the constitutive androstane receptor and pregnane X receptor. *Mol. Pharmacol.* **64**: 474-81 [PMID:12869653]
80. Xie W, Barwick JL, Downes M, Blumberg B, Simon CM, Nelson MC, Neuschwander-Tetri BA, Brunt EM, Guzelian PS and Evans RM. (2000) Humanized xenobiotic response in mice expressing nuclear receptor SXR. *Nature* **406**: 435-9 [PMID:10935643]
81. Xie W, Radomska-Pandya A, Shi Y, Simon CM, Nelson MC, Ong ES, Waxman DJ and Evans RM. (2001) An essential role for nuclear receptors SXR/PXR in detoxification of cholestatic bile acids. *Proc. Natl. Acad. Sci. U.S.A.* **98**: 3375-80 [PMID:11248086]
82. Xie W, Yeuh MF, Radomska-Pandya A, Saini SP, Negishi Y, Bottroff BS, Cabrera GY, Tukey RH and Evans RM. (2003) Control of steroid, heme, and carcinogen metabolism by nuclear pregnane X receptor and constitutive androstane receptor. *Proc. Natl. Acad. Sci. U.S.A.* **100**: 4150-5 [PMID:12644700]
83. Xu SZ, Zeng F, Boulay G, Grimm C, Harteneck C and Beech DJ. (2005) Block of TRPC5 channels by 2-aminoethoxydiphenyl borate: a differential, extracellular and voltage-dependent effect. *Br. J. Pharmacol.* **145**: 405-14 [PMID:15806115]

84. Yoshinari K, Sueyoshi T, Moore R and Negishi M. (2001) Nuclear receptor CAR as a regulatory factor for the sexually dimorphic induction of CYB2B1 gene by phenobarbital in rat livers. *Mol. Pharmacol.* **59**: 278-84 [[PMID:11160864](#)]
85. Yoshizawa T, Handa Y, Uematsu Y, Takeda S, Sekine K, Yoshihara Y, Kawakami T, Arioka K, Sato H, Uchiyama Y, Masushige S, Fukamizu A, Matsumoto T and Kato S. (1997) Mice lacking the vitamin D receptor exhibit impaired bone formation, uterine hypoplasia and growth retardation after weaning. *Nat. Genet.* **16**: 391-6 [[PMID:9241280](#)]
86. Zeitz U, Weber K, Soegiarto DW, Wolf E, Balling R and Erben RG. (2003) Impaired insulin secretory capacity in mice lacking a functional vitamin D receptor. *FASEB J.* **17**: 509-11 [[PMID:12551842](#)]
87. Zhang H, LeCulyse E, Liu L, Hu M, Matoney L, Zhu W and Yan B. (1999) Rat pregnane X receptor: molecular cloning, tissue distribution, and xenobiotic regulation. *Arch. Biochem. Biophys.* **368**: 14-22 [[PMID:10415106](#)]
88. Zhang J, Huang W, Chua SS, Wei P and Moore DD. (2002) Modulation of acetaminophen-induced hepatotoxicity by the xenobiotic receptor CAR. *Science* **298**: 422-4 [[PMID:12376703](#)]