

See discussions, stats, and author profiles for this publication at:
<http://www.researchgate.net/publication/278025548>

Rare Earth Elements of modern shelf and deep-water articulated brachiopods: evaluation of seawater masses

CONFERENCE PAPER · MAY 2015

DOWNLOADS

13

VIEWS

12

5 AUTHORS, INCLUDING:



Uwe Brand

Brock University

107 PUBLICATIONS 1,960 CITATIONS

SEE PROFILE



Karem Azmy

Memorial University of Newfoundland

52 PUBLICATIONS 1,262 CITATIONS

SEE PROFILE

Available from: Uwe Brand
Retrieved on: 24 June 2015

or carbonate-rock facies) from the Carboniferous to the Permian indicates that the body size changes of *Neochonetes* were not controlled by the substrate. The body size of *Neochonetes* decreased in the Middle Permian and increased in the Wuchiapingian (early Late Permian), which is inferred to respectively link to the temperature rising in the Middle Permian and the transgression in the early Late Permian. Comparison of the occurrence frequency of *Neochonetes* and *Tethyochonetes* in three different facies (shallow-water clastic-rock facies and carbonate-rock facies and deep-water siliceous-rock facies) demonstrates that both genera occurred more commonly in the shallow-water facies, and in the deep-water facies *Tethyochonetes* dominates and has a larger average size, which means that *Tethyochonetes* could perform better in deep-water environments than *Neochonetes*. More interestingly, *Tethyochonetes* occurred more frequently in the Early Triassic than *Neochonetes* did, suggesting *Tethyochonetes* coping with the end-Permian crisis stress better and surviving more during the Early Triassic.

A new sample processing protocol: an important update for procuring reliable ree signatures

Amir ZAKY

Department of Earth Sciences, Memorial University of Newfoundland, St. John's, NL, A1B 3X5 Canada <ahb485@mun.ca>

Uwe BRAND

Department of Earth Sciences, Brock University, St. Catharines, ON, Canada L2S 3A1 Canada

Karem AZMY

Department of Earth Sciences, Memorial University of Newfoundland, St. John's, NL, A1B 3X5 Canada

Cleaning of brachiopod shells is a fundamental step that should be conducted with great care and attention to detail prior to processing their rare earth element (REE) contents. Without cleaning, the concentrations would represent the REE compositions not only of the shells calcitic structure but also of lattice bound oxides, detritus, particulates and organic remnants. Developing a sensitive and precise analytical protocol is of utmost importance required for isolating contaminants on biogenic and abiogenic carbonates in order to acquire a robust REE signal that enhances our confidence in oceanographic applications. To achieve this goal, 26 shell fragments separated from recently dead *Liothyrella neozelanica* shells, recovered from the deep water's of the southwest Pacific, north of New Zealand, were extensively washed and rinsed with distilled water. Five parts did not receive any further cleaning; protocol – 1 (P-1). A set of four parts was physically cleaned using a sharp stainless-steel blade (P-2). Another set of four parts was immersed in 2.5 % hydrogen peroxide (H₂O₂) for three continuous days (P-3). A further set of four parts was physically cleaned then dropped into 10% hydrochloric acid (HCl) until the fragments were deemed clean (P-4). The last nine parts were processed using all cleaning protocols: water washed, physically

cleaned, H₂O₂ immersed, and then HCl leached (P-5).

Attached detritus and nano-particulates adsorbed on the calcitic structure of brachiopod shells besides the organic tissue, proteinaceous periostracum (Carpenter and Lohmann, 1995) may lead to elevated ΣREE, anomalous Ce anomaly (Fig. 1A) and elevated Mn, Fe (Fig. 2A) and U concentrations as documented by the P-1 set. Physical cleaning removes mainly the adsorptive nano-particulates and the periostracum. This protocol leads to depleted ΣREE and lower Mn, Fe, and U concentrations but with a slight Ce anomaly: set P-2. Hydrogen peroxide eliminates primarily the organic tissue; however it requires physical cleaning to remove the periostracum or the nano-particulates, which leads to partial depletion in the Ce anomaly: set P-3. Physical cleaning followed by chemical cleaning removes adsorptive particulates, organic remnants, periostracum and the primary layer. This protocol is reflected in the drastic reductions of the ΣREE, Mn, Fe, U concentrations and 'normal' Ce anomalies in the P-4 and P-5 sets (Figs. 1A, 2A).

Protocol P-4 was applied to Ordovician, Silurian, Pennsylvanian and Permian brachiopods and their enclosing lime matrix collected from Canada, USA and China to test for the reliability of REEs in fossil and Deep-Time carbonates. The immersion in H₂O₂ was omitted since no organic tissue was documented in any fossil samples. The fossil shells display ΣREEs, Mn, Fe, U concentrations and Ce anomalies comparable to their modern counterparts, while the lime matrixes are slightly to extensively enriched (Figs. 1B& 2B). This unequivocally documents the veracity of the processing protocols and if we expect, want reliable REEs, the need for its rigorous application to fossil material such as brachiopods, conodonts, corals, molluscs, foraminiferas and whole rock.

Acknowledgements: We thank the National Institute of Water and Atmospheric research in New Zealand (NIWA) for supplying us with the brachiopod, and NSERC for financial support to UB.

References

- Azmy, K., Brand, U., Sylvester, P., Gleeson, S. A., Logan, A., and Bitner, M. A. 2011. Biogenic and abiogenic low-Mg calcite (bLMC and aLMC): evaluation of seawater-REE composition, water masses and carbonate diagenesis. *Chemical Geology*, 280(1), 180–190.
- Bau, M., and Koschinsky, A. 2009. Oxidative scavenging of cerium on hydrous Fe oxide: evidence from the distribution of rare earth elements and yttrium between Fe oxides and Mn oxides in hydrogenetic ferromanganese crusts. *Geochemical Journal*, 43(1), 37–47.
- Brand, U., Logan, A., Hiller, N., and Richardson, J. 2003. Geochemistry of modern brachiopods: applications and implications for oceanography and paleoceanography. *Chemical Geology*, 198(3), 305–334.
- Brand, U., Logan, A., Bitner, M. A., Griesshaber, E., Azmy, K., and Buhl, D. 2011. What is the ideal proxy of Palaeozoic seawater chemistry?. *Memoirs of the Association of Australasian Palaeontologists*, 41, 9–24.
- Brand, U., Azmy, K., Bitner, M. A., Logan, A., Zuschin, M., Came, R., and Ruggiero, E. 2013. Oxygen isotopes and




Fig. 1: Ce anomaly $(Ce/Ce^*)_{SN}$ - ΣREE evaluation of the different cleaning protocols conducted with a modern brachiopod shell (A) and on Ordovician, Silurian, Pennsylvanian and Permian brachiopods and their whole rock.




Fig. 2: Distribution of Mn and Fe contents with the different cleaning protocols of the modern brachiopod shell (A) and in their Paleozoic counterparts and enclosing whole rock (B).

MgCO₃ in brachiopod calcite and a new paleotemperature equation. *Chemical Geology*, 359, 23–31.

Carpenter, S. J., and Lohmann, K. C. 1995. $\delta^{18}O$ and $\delta^{13}C$ values of modern brachiopod shells. *Geochimica et Cosmochimica Acta*, 59(18), 3749–3764.

de Baar, H. J., German, C. R., Elderfield, H., and Van Gaans, P. 1988. Rare earth element distributions in anoxic waters of the Cariaco Trench. *Geochimica et Cosmochimica Acta*, 52(5), 1203–1219.

German, C. R., and Elderfield, H. 1989. Rare earth elements in Saanich Inlet, British Columbia, a seasonally anoxic basin. *Geochimica et Cosmochimica Acta*, 53(10), 2561–2571.

Haley, B. A., Klinkhammer, G. P., and McManus, J. 2004. Rare earth elements in pore waters of marine sediments. *Geochimica et Cosmochimica Acta*, 68(6), 1265–1279.

James, M. A., Ansell, A. D., Collins, M. J., Curry, G. B., Peck, L. S., and Rhodes, M. C. 1992. Biology of living

brachiopods. *Advances in Marine Biology*, 28, 175–387.

McLennan, S. M. 1989. Rare earth elements in sedimentary rocks; influence of provenance and sedimentary processes. *Reviews in Mineralogy and Geochemistry*, 21(1), 169–200.

Moore, R. C. 1997. Treatise on invertebrate paleontology: part 0, Arthropoda 1, Trilobita. Trilobita. Introduction, order agnostida, order redlichiida/HB Whittington. Geological Society of America.

Rare earth elements of modern shelf and deep-water articulated brachiopods: evaluation of seawater masses

Amir ZAKY




Fig. 1: Geographic distribution of the shelf and deep-water collection stations of the investigated brachiopod archives.

Table 1: Location, setting and depth of the shelf and deep-water stations and brachiopod taxonomy.

	Station	Depth (m)	Bathymetry	Order	Taxon
CARIBBEAN SEA					
VENEZUELA					
Venezuela Basin	1	3986	Deep Sea	<i>Terebratulida</i>	<i>Chlidonophora incerta</i>
	2	3998			
US VIRGIN ISLANDS					
Vigin Islands Basin	3	3940	*	"	"
NORTH ATLANTIC OCEAN					
ICELAND					
Iceland Basin	4	2567.7-2568.5	*	"	<i>Waldheimia cranium</i>
	5	2504.7-2531.8	*	"	"
Irminger Basin	6	698.1-678.5		"	"
	7	724.4-704.9	Continental shelf	"	"
Denmark Strait	8	742.5-603.1	*	"	"
	9	696.9-706.4		"	"
CANARY ISLANDS					
Continental slope of northwest Africa	10	1265	Deep Sea	"	<i>Dallina septigera</i>
	11			"	<i>Hispanirhynchia cornea</i>
SOUTHWEST PACIFIC OCEAN					
TONGA					
Lau Arc (Tonga-1)	12	589	Lau Arc	"	<i>Basilioia beecheri</i>
Colville Arc (Tonga-2)	13	660	Colville Arc	<i>Terebratulida</i>	<i>Dallina</i> sp.
NEW ZEALAND					
	14	1244	Thomson seamount	"	<i>Stenosarina crosnieri</i>
Kermadec Arc (New Zealand-1)	15	1460	Clark Seamount, south cone	"	<i>Dallina triangulari</i> , <i>Stenosarina crosnieri</i>
	16	1583	Clark Seamount south flank	"	<i>Dallina triangularis</i>
	17	1740	Brothers Seamount West	"	<i>Goniobrochus ewingi</i>
Havre Trough (New Zealand-2)	18	1172-1115	Seamount west of Volcano L	"	<i>Liothyrella neozelanic</i>
	19	1520	Seamount west of Rumble II West	"	<i>Goniobrochus ewingi</i>
Northland Plateau (New Zealand-3)	20	1293	Seamount #441 "Ballance Seamount"	<i>Rynchonellida</i>	<i>Basilioia pompholyx</i>
	21	1583	Unnamed volcanic peak	<i>Terebratulida</i>	<i>Stenosarina</i> sp. <i>Abyssothyris wyvillei</i>
SOUTHERN OCEAN					
ANTARCTICA					
Balleny Spreading Corridor	22	547	Northeastern knoll of Admiralty seamount	"	<i>Magellania fragilis</i>
	23	1130	Seamount close to Scott Island	"	<i>Liothyrella</i> sp.

Department of Earth Sciences, Memorial University of Newfoundland, St. John's, NL, A1B 3X5 Canada <ahb485@mun.ca>

Uwe BRAND

Department of Earth Sciences, Brock University, St. Catharines, ON, L2S 3A1 Canada

Karem AZMY

Department of Earth Sciences, Memorial University of Newfoundland, St. John's, NL, A1B 3X5 Canada <ahb485@mun.ca>

Alan LOGAN

Centre of Coastal Studies, University of New Brunswick, Saint John, NB, E2L 4L5 Canada

Jörundur SVAVARSSON

Institute of Biology, University of Iceland, Askja, Sturlugata 7, 101 Reykjavík, Iceland

Modern Rhynchonellids and Terebratulids, obtained from water depths below the neritic zone (>500m) at 23 stations in the Caribbean Sea, North Atlantic, South Pacific and Southern Oceans, were investigated for their rare earth element (REE) contents (Fig. 1, Table 1). The Σ REE of shelf (500-1000 m) or deep-water (>1000 m) brachiopod populations do not vary significantly between oceans/seas irrespective of water mass origins or influences by major currents.




Fig. 2. Mean REE_{SN} patterns of the shelf and deep-water masses revealed by articulated brachiopods, and the REE_{SN} pattern of shallow water articulated brachiopods (<500m) of Azmy et al. (2011).

Four brachiopod populations from the shelves of the Irminger Basin and Denmark Strait (North Atlantic) and from the Lau and Colville Arcs (South Pacific) display a mean REE_{SN} pattern of gradual enrichment with increasing atomic number punctuated by negative Ce excursions (Fig. 2). In contrast to the four above, the population from the Ross Sea (Southern Ocean) is relatively depleted in Σ REE reflecting local sea ice formation and conse-

quent down-flow of saline water. Although similar in pattern to the four shelf populations, the REE_{SN} trend of the five deep-water brachiopod populations from the Venezuela Basin (Caribbean Sea), Iceland Basin (North Atlantic) and from the Kermadec Arc, Havre Trough and Northland Plateau (South Pacific) is more pronounced as well as its Ce excursion is more prominent (Fig. 2). Furthermore two populations from the Irminger Basin and Canary Islands are influenced by local environmental parameters such as deep convection and the remineralization of bottom sediments enhanced by turbidity current flows, respectively. Overall, the REE patterns of the shelf and deep-water brachiopod populations are similar to that of open ocean seawater, but remarkably different to that of shallow water counterparts.

Acknowledgements: We thank the National Institute of Water and Atmospheric research in New Zealand (NIWA) for supplying us with some of the brachiopods, and NSERC for financial support to UB.

References

Azmy, K., Brand, U., Sylvester, P., Gleeson, S.A., Logan, A. and Bitner, M.A. 2011. Biogenic and abiogenic low-Mg calcite (bLMC and aLMC): evaluation of seawater-REE composition, water masses, and carbonate diagenesis. *Chemical Geology*, 280, 180–190.

de Baar, H.J., Brewer, P.G. and Bacon, M.P. 1985. Anomalies in rare earth distributions in seawater: Gd and Tb. *Geochimica et Cosmochimica Acta*, 49,1961–1969.

Elderfield, H. and Greaves, M.J. 1982. The rare earth elements in seawater. *Nature*, 296, 214–218.

German, C.R., Masuzawa, T., Greaves, M.J., Elderfield, H. and Edmond, J.M. 1995. Dissolved rare earth elements in the Southern Ocean: Cerium oxidation and the influence of hydrography. *Geochimica et Cosmochimica Acta*, 59, 1551–1558.

A Katian (Late Ordovician) minute *Foliomena* fauna from the Sibumasu paleoplate

ZHAN Renbin

State Key Laboratory of Palaeobiology and Stratigraphy, Nanjing Institute of Geology and Palaeontology, Chinese Academy of Sciences, Nanjing 210008, China <rbzhan@nigpas.ac.cn>

LI Rongyu

Department of Geology, Brandon University, Brandon Manitoba, R7A 6A9 Canada

WEI Xin

State Key Laboratory of Palaeobiology and Stratigraphy, Nanjing Institute of Geology and Palaeontology, Chinese Academy of Sciences, Nanjing 210008, China

JIN Jisuo

Department of Earth Sciences, Western University, London Ontario, N6A 5B7 Canada