



UNIVERSITI PUTRA MALAYSIA

**CONVERSION OF PADDY STRAW TO BIOETHANOL THROUGH
CONSOLIDATED BIOPROCESSING USING LIGNOCELLULOLYTIC
FUNGI**

MONA FATIN SYAZWANEE BINTI MOHAMED GHAZALI

FS 2019 21



**CONVERSION OF PADDY STRAW TO BIOETHANOL THROUGH
CONSOLIDATED BIOPROCESSING USING LIGNOCELLULOLYTIC FUNGI**

By

MONA FATIN SYAZWANEE BINTI MOHAMED GHAZALI

**Thesis Submitted to the School of Graduate Studies, Universiti Putra
Malaysia, in Fulfilment of the Requirements for the Degree of Doctor of
Philosophy**

April 2019

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Doctor of Philosophy

CONVERSION OF PADDY STRAW TO BIOETHANOL THROUGH CONSOLIDATED BIOPROCESSING USING LIGNOCELLULOLYTIC FUNGI

By

MONA FATIN SYAZWANEE BINTI MOHAMED GHAZALI

April 2019

Chairman : Muskhazli Mustafa, PhD
Faculty : Science

Normally, paddy straw was disposed of via open burning even though it contains valuable lignocellulosic materials which can be readily converted into fermentable sugar for bioethanol production. The second-generation of bioethanol production utilizes useful lignocellulosic substrates especially cellulose for bioconversion process. However, this material is enclosed within hemicellulose and lignin matrix in the cell wall, making the accessibility of cellulose become the major problem in bioethanol production from such sources in consolidate bioprocessing (CBP). The CBP is preferable as it produces faster saccharification result, low risk of contamination and cost-effective. Nevertheless, finding an optimize condition for efficient bioethanol production in CBP is still ambiguous as a different strain of lignocellulolytic fungi has their own environment preferences. Therefore, the main aim of this study is to explore a new approach in converting paddy straw into bioethanol using only filamentous fungi throughout the entire CBP process, thus eliminating the use of yeast as a fermenter organism. In this study, the research objectives involves the pretreatment method of paddy straw, selecting the best lignocellulolytic agent for hydrolyzation, optimizing all factors influencing the bioethanol production via one-factor-at-a-time (OFAT) as well as Response Surface Methodology analysis (RSM) and evaluating the final CBP set-up.

Paddy straw sieved into three different sizes; 2 mm, 5 mm and 8 mm were prepared and underwent several physical pretreatment (autoclave, boil) and chemical pretreatment (HNO_3 and NaOH). Size five millimeter paddy straw showed the highest cellulose content (35.61%) and the percentage of cellulose content went escalated to 72.47% when pretreated with 2% (w/v) sodium hydroxide (NaOH). Pretreatment of 2% (w/v) NaOH also shown the most efficient delignification and desilication process (1.02% lignin; 5.44% ash content) compared to others. All strains of fast-growing fungi were

quantitatively assayed and the results indicate that the highest cellulases enzyme producer were *Trichoderma asperellum* B1581 (3.93 U/mL endoglucanase; 2.37 U/mL exoglucanase; 3.00 U/mL β -glucosidase; 54.87 U/mL xylanase), followed by *Aspergillus niger* B2484 (5.60 U/mL endoglucanase; 1.08 U/mL exoglucanase; 1.57 U/mL β -glucosidase; 56.85 U/mL xylanase). A further test on compatibility test revealed mutual intermingling between both *T. asperellum* B1581 and *A. niger* B2484. Six single factors that are crucial for bioethanol production were tested in one-factor-at-a-time (OFAT) analysis for both selected strains of lignocellulolytic fungi. With all factors combined, *T. asperellum* B1581 prefers 2 days of both saccharification and fermentation process at 30°C with an amount of 3% substrate level and 10% of media level. While *A. niger* B2482 prefers 3 days of saccharification, 1 day of fermentation; at 30°C with an amount of 2% substrate level and 20% of media level. The results produced by OFAT were used as the centre point in the Central Composite Design (CCD) through Response Surface Methodology (RSM) software. However, comparison between the actual and the predicted value of ethanol produced in RSM's recommended CBP set-up for both *T. asperellum* B1581 and *A. niger* B2484 showed no significant difference, thus proving the model's stability to navigate experiment. In order to test effectiveness *T. asperellum* B1581 and *A. niger* B2484 as a fungi consortium, several combination of consortia concentrations (spore/mL) were tested and the amount of ethanol was quantified. However, a single strain of *T. asperellum* B1581 (6:0) was able to match the amount of ethanol produced by consortia of *T. asperellum* B1581 and *A. niger* B2484 (5:1, 4:2, 3:3, 2:4 and 1:5) by producing the highest total amount of ethanol (1.11 g/L). The final amount of ethanol detected by GC-FID was 1.25 g/L; which was not significantly different from the ethanol assayed spectrometrically (1.11 g/L).

As a conclusion, a pretreatment of size 5 mm using 2% (w/v) NaOH had enhanced the breaking of cellulose-lignin complex, delignification, and desilication. Thus making the paddy straw becomes feasible for biofuel production. Both *T. asperellum* B1581 and *A. niger* B2484 were found to produce the highest cellulase enzyme and displayed mutual intermingling relationship suggesting the possibility of fungal consortium formation between these two species. Even though the recommended model for CBP set-up by RSM showed no significant differences between an actual and predicted value of ethanol produced, both species unable to improve the value of ethanol produced as consortia compare to single *T. asperellum* B1581 culture set-up. Thus, indicating that the potential of *T. asperellum* B1581 as single culture for bioethanol production in consolidated bioprocessing (CBP).

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

PENGHASILAN BIOETANOL DARI JERAMI PADI MELALUI PENYATUAN BIOPROSES MENGGUNAKAN KULAT LIGNOSELULOLITIK

Oleh

MONA FATIN SYAZWANEE BINTI MOHAMED GHAZALI

April 2019

Pengerusi : Muskhazli Mustafa, PhD
Fakulti : Sains

Pada kebiasaannya, jerami padi dilupuskan secara pembakaran terbuka walaupun ianya mengandungi bahan lignoselulosa yang dapat diubah menjadi gula fermentasi bagi penghasilan bioetanol. Bioetanol generasi kedua menggunakan substrat lignoselulosa yang penting terutamanya selulosa untuk proses biopenukaran. Namun begitu, bahan tersebut dilindungi rapi oleh hemiselulosa dan matriks lignin dalam dinding sel, lantas telah menjadi masalah utama dalam mengakses selulosa untuk pengeluaran bioetanol dari sumber tersebut menggunakan kaedah penyatuan bioproses (CBP). Keadah CBP adalah lebih sesuai digunakan kerana ia membolehkan proses pensakaridaan berlaku lebih cepat, kurang risiko pencemaran dan kos efektif. Walau bagaimanapun, pencarian keadaan yang optima bagi penghasilan bioetanol melalui CBP masih tidak jelas kerana kulat lignoselulolitik yang berlainan mempunyai pilihan persekitaran yang tersendiri. Oleh itu, tujuan utama kajian ini adalah untuk meneroka pendekatan baru dalam mengubah jerami padi menjadi bioetanol dengan hanya menggunakan kulat filamen sepanjang proses CBP, dan seterusnya mengelak penggunaan yis sebagai organisma penapaian. Oleh itu, matlamat penyelidikan merangkumi kaedah pra-rawatan untuk jerami padi, memilih agen lignoselulolitik yang terbaik untuk hidrolisis, mengoptimakan semua faktor yang mempengaruhi penghasilan bioetanol melalui satu faktor-pada-satu masa (OFAT) serta kaedah analisis tindakbalas permukaan (RSM) dan menilai tetapan akhir CBP.

Jerami padi telah ditapis menggunakan saiz yang berlainan; 2 mm, 5 mm dan 8 mm yang disediakan dan dikenakan pelbagai pra-rawatan fizikal (autoklaf, pendidihan) dan pra-rawatan kimia (HNO_3 dan NaOH). Saiz lima millimeter jerami padi telah menunjukkan kandungan selulosa tertinggi (35.61%) dan peratusan kandungan selulosa meningkat sehingga 72.47% apabila dirawat dengan 2% (w/v) Natrium hidroksida (NaOH). Pra-rawatan 2% (w/v) NaOH juga menunjukkan proses penghapusan lignin dan desilikasi yang paling

berkesan (1.02% lignin; 5.44% kandungan abu) berbanding dengan yang lain. Semua jenis kulat yang mempunyai kadar pertumbuhan yang cepat telah diuji secara kuantitatif dan hasilnya telah menunjukkan bahawa pengeluar enzim selulase yang tertinggi adalah *Trichoderma asperellum* B1581 (3.93 U/mL endoglucanase; 2.37 U/mL exoglucanase; 3.00 U/mL β -glucosidase; 54.87 U/mL xylanase), diikuti oleh *Aspergillus niger* B2484 (5.60 U/mL endoglucanase; 1.08 U/mL exoglucanase; 1.57 U/mL β -glucosidase; 56.85 U/mL xylanase). Ujian lanjut mengenai keserasian antara dua jenis kulat yang berbeza telah menunjukkan tiada persaingan antara kedua-dua pencilan *T. asperellum* B1581 dan *A. niger* B2484. Enam faktor telah diuji dalam analisis satu faktor-pada-satu masa (OFAT) untuk kedua-dua jenis kulat lignoselulolitik yang terpilih. Dengan menggabungkan semua faktor, *T. asperellum* B1581 memberi respon yang optima pada hari kedua bagi kedua-dua proses sakarifikasi dan penapaian pada suhu 30°C dengan jumlah paras substrat 3% dan 10% tahap media. Sementara *A. niger* B2482 memberikan respon yang optimum pada hari ketiga proses sakarifikasi, 1 hari penapaian; pada suhu 30°C dengan jumlah substrat 2% dan 20% tahap media. Keputusan yang dihasilkan oleh OFAT telah digunakan sebagai titik tengah dalam Rekabentuk Komposit Sentral (CCD) melalui kaedah analisis tindakbalas permukaan (RSM). Walau bagaimanapun perbandingan antara nilai sebenar dan nilai ramalan etanol yang diberikan oleh perisian RSM untuk *T. asperellum* B1581 dan *A. niger* B2484 tidak menunjukkan perbezaan yang nyata, dan telah membuktikan kestabilan model untuk digunapakai bagi tujuan eksperimen. Untuk menguji keberkesanan *T. asperellum* B1581 dan *A. niger* B2484 sebagai konsortium kulat, beberapa kombinasi kepekatan konsortia (spora/mL) telah diuji dan jumlah penghasilan etanol ditentukan. Namun, *T. asperellum* B1581 (6:0) dapat menandingi jumlah etanol yang dihasilkan oleh konsortium *T. asperellum* B1581 dan *A. niger* B2484 (5:1, 4:2, 3:3, 2:4 dan 1:5) dengan penghasilan jumlah etanol yang tertinggi (1.11 g/L). Jumlah akhir etanol yang dikesan oleh GC-FID adalah 1.25 g/L; di mana nilainya tidak jauh berbeza dengan nilai etanol yang diuji secara spektrometri (1.11 g/L).

Sebagai kesimpulan, pra-rawatan padi dengan saiz 5 mm menggunakan 2% (w/v) NaOH telah meningkatkan pemecahan kompleks selulosa-lignin, penghakisan lapisan lignin dan desilikasi. Oleh itu menjadikan jerami padi layak dan sesuai digunakan untuk penghasilan bioetanol. Kedua-dua *T. asperellum* B1581 dan *A. niger* B2484 menunjukkan keupayaan dalam menghasilkan enzim selulase tertinggi dan mempunyai hubungan percampuran yang baik untuk pembentukan konsortium kulat antara dua spesies ini. Walaupun model yang disyorkan untuk proses CBP oleh RSM tidak menunjukkan perbezaan yang nyata antara nilai sebenar etanol dan ramalan yang dihasilkan, kedua-dua spesies tidak dapat meningkatkan nilai etanol yang dihasilkan secara konsortia berbanding dengan tetapan kultur tunggal *T. asperellum* B1581. Oleh itu, penggunaan kulat *T. asperellum* B1581 secara tunggal mempunyai potensi untuk digunakan bagi pengeluaran bioetanol menerusi proses CBP.

ACKNOWLEDGEMENTS

All praises to Allah S.W.T for giving me a lifetime opportunity to pursue my dreams, an absolute strength to face all the hardships and guide me throughout the journey. May Allah S.W.T continue to bless us all in this life and hereafter.

I would like to express my gratitude to my parents, Hj. Mohamed Ghazali B Mohamed Yousoff and Hjh. Fawziah Bt Md Noor, whom has been my enthusiastic cheerleader, along with their continuous support, duaa's and endless love that has become pillars to my success. Thank you for offering me comfort and always been there for me when I was feeling down. I am truly blessed to have parents that totally understand my passion and always believe in me.

I also wish to thank my elder brothers, Dr. Farid Ezanee, Mohamed Firdauz Mukhzanee, sister in law, Mazwani Wan Ibrahim and nephews, Daniel Mukhreez and Danish Mikhael for bringing me joy, provided me with moral support, love and courage through high and low. I am so lucky and grateful to have such a wonderful family by my side throughout my journey.

A very special gratitude dedicated to my chairman of supervisory committee, Associate Professor Dr. Muskhazli Mustafa, for his guidance, patience, wisdom, advice and support in assisting me to complete my study and produce a better quality of research. He has helped me to understand and explore more about research field, developed new technical and soft skills, inspire my creativity, encourage my critical thinking, building my confidence and shaped me into a better person. I am so grateful to have a very motivating supervisor like him and could not have imagined having a better supervisor for my PhD project. I would also like to wish my gratitude towards my other committee members; Associate Professor Dr. Nur Ain Izzati Mohd Zainudin for providing me with several identified fungi samples from Mycology Laboratory and guides me in mycology; and Associate Professor Dr. Nor Azwady Abdul Aziz, for providing me support, advice and guidance in statistical analysis. May Allah S.W.T continue to reward all of you for your good deeds.

My immense gratitude to the Ministry of Higher Education Malaysia (MOHE) for providing me with financial support under MyBrain15- MyPhD programme throughout my entire study period. To the Universiti Putra Malaysia, thank you for the Geran Putra – Inisiatif Putra Siswazah (GP-IPS/2016/9485600) which has become an immense help towards minimizing our financial stress during the study. I wish to thank all Plant Systematic and Microbes Laboratory members especially Pn. Azleen Ahmad Adli and Pn. Farah Kamaludin for their assistance throughout the length of my projects.

Special thanks to Pn. Norlinawati Abd Halim from Microbiology Laboratory, Faculty of Food Science and Technology for allowing me to used their equipment throughout my study. Lastly, I would like to thank Mr. Hamezan Muhammad@Ahmad from Chromatography Laboratory, Faculty of Food Science and Technology, for his advice and guidance in operating gas-chromatography (GC-FID) during sample analysis.



I certify that a Thesis Examination Committee has met on 24 April 2019 to conduct the final examination of Mona Fatin Syazwaneer binti Mohamed Ghazali on her thesis entitled "Conversion of Paddy Straw to Bioethanol Through Consolidated Bioprocessing Using Lignocellulolytic Fungi" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

Rusea Go, PhD

Professor
Faculty of Science
Universiti Putra Malaysia
(Chairman)

Syaizwan Zahmir bin Zulkifli, PhD

Senior Lecturer
Faculty of Science
Universiti Putra Malaysia
(Internal Examiner)

Shuhaimi bin Mustafa, PhD

Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Internal Examiner)

Anja Meryandini, PhD

Professor
Faculty of Mathematics and Natural Sciences
Bogor Agricultural University
Indonesia
(External Examiner)



RUSLI HAJI ABDULLAH, PhD
Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 26 June 2019

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Muskhazli Mustafa, PhD

Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Chairman)

Nur Ain Izzati Mohd Zainudin, PhD

Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Member)

Nor Azwady Abd Aziz, PhD

Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No.: Mona Fatin Syazwanee Binti Mohamed Ghazali,
GS43701

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____

Name of Chairman of

Supervisory Committee: Associate Professor Dr. Muskhazli Mustafa

Signature: _____

Name of Member of

Supervisory Committee: Associate Professor Dr. Nur Ain Izzati Mohd Zainudin

Signature: _____

Name of Member of

Supervisory Committee: Associate Professor Dr. Nor Azwady Abdul Aziz

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vii
DECLARATION	ix
LIST OF TABLES	xiv
LIST OF FIGURES	xvi
LIST OF ABBREVIATIONS	xviii
CHAPTER	
1 INTRODUCTION	1
1.1 Background of the study	1
1.2 Problem statement	3
1.3 Objectives of the study	4
2 LITERATURE REVIEW	6
2.1 Biofuel from biomass	6
2.1.1 Bioethanol	6
2.2.1 First generation of bioethanol	7
2.2.3 Second generation of bioethanol	7
2.2 Lignocellulosic material	7
2.2.1 Bioethanol from paddy straw	8
2.2.2 Holocellulose	9
2.2.3 Cellulose	9
2.2.4 Hemicellulose	10
2.2.5 Lignin	10
2.3 Pretreatments	10
2.3.1 Physical pretreatment	11
2.3.1.1 Comminution treatment	11
2.3.1.2 Autoclave treatment	11
2.3.2 Chemical treatment	12
2.3.2.1 Dilute acid treatment	12
2.3.2.2 Alkali pretreatment	12
2.3.3 Physico-chemical/combination treatment	13
2.4 Conversion into ethanol	13
2.4.1 Saccharification of cellulose	14
2.4.1.1 Fungi as source of cellulolytic enzyme	17
2.4.1.2 Microbial consortium or co-culture of two fungi strains	17
2.4.2 Fermentation process	18
2.4.2.1 SSF process	20
2.4.2.2 Consolidated bioprocessing (CBP)	21
2.4.3 Direct ethanol production by filamentous fungi	22

3	OPTIMIZATION OF PRETREATMENT METHOD FOR DESILICATION, DELIGNIFICATION AND ENHANCEMENT OF THE CELLULOSE ACCESSIBILITY IN PADDY STRAW	23
3.1	Introduction	23
3.2	Materials and methods	25
3.2.1	Material preparation	25
3.2.2	Substrates preparation with physical pretreatment	25
3.2.3	Pretreatment of the substrates	25
3.2.4	Determination of holocellulose content in the pretreated paddy straw	26
3.2.5	Determination of hemicellulose	27
3.2.6	Ash content determination	27
3.2.7	Determination of lignin composition	28
3.2.7.1	Acid-detergent fiber (ADF)	28
3.2.7.2	Acid-detergent lignin (ADL)	28
3.2.8	Physical observation using Scanning Electron Microscopes (SEM)	29
3.2.9	Data analysis	29
3.3	Results and Discussions	29
3.3.1	Physical pretreatment of paddy straw	29
3.3.2	Physico-chemical pretreatment	31
3.3.3	Scanning Electron Microscope (SEM) images	35
3.4	Conclusion	37
4	SELECTION OF LIGNOCELLULOLYTIC FUNGI AS AN AGENT FOR HYDROLYZATION OF LIGNOCELLULOSIC MATERIALS IN PADDY STRAW	38
4.1	Introduction	38
4.2	Materials and methods	39
4.2.1	Collection of fungi	39
4.2.2	Qualitative assay for lignocellulolytic activity	41
4.2.2.1	Cellulose degradation ability	41
4.2.2.2	Bavendamm test for laccase detection	41
4.2.3	Quantitative assays	41
4.2.3.1	Endoglucanase, exoglucanase and xylanase assays	41
4.2.3.2	β -glucosidase assay	42
4.2.4	Compatibility test	42
4.2.5	Data Analysis	43
4.3	Results and Discussions	43
4.3.1	Qualitative assays using Congo red and Bavendamm test	43
4.3.2	Determination of best cellulase producer	50
4.3.3	Compatibility test of fungal consortia	52
4.4	Conclusion	54

5	ESTABLISHMENT OF PHYSICO-CHEMICAL OPTIMIZATION FOR BIOETHANOL PRODUCTION VIA CONSOLIDATED BIOPROCESSING (CBP) BASED CONSORTIUM TRIAL BETWEEN <i>TRICHODERMA ASPERELLUM</i> B1581 AND <i>ASPERGILLUS NIGER</i> B2484	55
5.1	Introduction	55
5.2	Materials and methods	57
5.2.1	Preparation of culture conditions	57
5.2.2	One-factor-at-a-time (OFAT)	57
5.2.3	Response Surface Methodology (RSM)	58
5.2.4	Formulation of consortium setting	60
5.2.5	Strategy of consortium development	61
5.2.6	Gas chromatography – flame ionization detector (GC-FID)	61
5.2.6.1	Chemicals	61
5.2.6.2	Calibration curve for reference	62
5.2.6.3	Equipment and operations	62
5.2.7	Data Analysis	63
5.3	Results and Discussions	63
5.3.1	Optimizing significant parameters via one-factor-at-a-time (OFAT)	63
5.3.2	Application of Response Surface Methodology (RSM) for optimization	67
5.3.3	Formation of fungal consortium	84
5.3.4	Validation of ethanol using <i>T. asperellum</i> B1581 via GC-FID	86
5.4	Conclusion	89
6	GENERAL DISCUSSIONS, CONCLUSION AND FUTURE RECOMMENDATIONS	90
6.1	General discussions	90
6.2	Conclusion	91
6.3	Future recommendations	91
	REFERENCES	93
	APPENDICES	128
	BIODATA OF STUDENT	134
	LIST OF PUBLICATIONS	135

LIST OF TABLES

Table		Page
3.1	Types of pretreatment and conditions of all pretreatments on the substrates	25
3.2	Percentage of lignocellulosic composition of untreated paddy straw fibres for several sizes	30
3.3	Percentage of lignocellulosic composition of 5 mm paddy straw fibres after treated with different pretreatment	32
4.1	The list of fungi strain used and the source of origin.	40
4.2	Modes of interaction between filamentous fungi in compatibility test (Mohammad, Alam, Kabashi and Adebayo, 2011).	43
4.3	Qualitative assessment on fungi strains for selection of the best fungi producing hydrolytic enzymes	46
4.4	Quantitative assessment on fungi strains for selection of the best fungi producing hydrolytic enzymes.	51
5.1	The pre-determine ranges for each of the parameters in one-factor-at-a-time (OFAT)	58
5.2	The coded and un-coded variables used for the optimizations in Central Composite Design for ethanol production by <i>T. asperellum</i> B1581 and <i>A. niger</i> B2484	59
5.3	The formulation of new setting for consortium based on the average between the two individual settings of <i>T. asperellum</i> B1581 and <i>A. niger</i> B2484	60
5.4	The strategy of consortium development using combination ratios between <i>T. asperellum</i> B1581 and <i>A. niger</i> B2484 strains	61
5.5	RSM experimental settings for ethanol production using <i>T. asperellum</i> B1581 and <i>A. niger</i> B2484 in consolidated bioprocessing (CBP)	69

5.6	Statistical summary of ethanol production by <i>T. asperellum</i> B1581	73
5.7	Statistical summary of ethanol production by <i>A. niger</i> B2484	74
5.8	The summary of quadratic model statistics of <i>T. asperellum</i> B1581 and <i>A. niger</i> B2484	74
5.9	The optimization settings recommended by RSM with comparison to settings produced by OFAT analysis	82
5.10	The comparison between the amounts of ethanol produced by optimized set-up and central point in CCD	83
5.11	The amount of ethanol produced from different consortium composition between <i>T. asperellum</i> B1581 and <i>A. niger</i> B2484	85

LIST OF FIGURES

Figure		Page
1.1	Summary of research design and attainment of objectives	5
2.1	The schematic image of paddy straw structure (Seo and Sakoda 2014)	9
2.2	The synergistic action of all 3 enzymes; endoglucanase, cellobiohydrolases and β -glucosidase in noncomplex cellulase system in degrading cellulose into glucose (Arantes and Saddler, 2010).	15
2.3	The metabolic pathway of glycolysis in producing ethanol by <i>S. cerevisiae</i> (Paola, Isabella and Patrizia, 2011).	19
2.4	Strategy of different bioconversion process (Hamelinck, Hooijdonk and Faaij, 2005; Devarapalli and Atiyeh 2015).	21
3.1	The scanning electron micrographs (SEM) images of 5 mm paddy straw fibres. (a) Untreated (b) Treated with 2% (w/v) sodium hydroxide (NaOH) (c) Treated with 0.5% (v/v) nitric acid (HNO ₃)	36
4.1	The diameter of halo ring indicates the amount of cellulase enzyme produced by <i>T. asperellum</i> (TA), <i>A. niger</i> (AN) and <i>F. oxysporum</i> (FO) after the plates were stained by Congo red.	44
4.2	The diameter of halo ring indicates the amount of cellulase enzyme produced by <i>F. fujikuroi</i> (FF), <i>C. eragrostidis</i> (CE) and <i>C. lunata</i> (CL) after the plates were stained by Congo red.	45
4.3	The presence of polyphenol oxidase (PPO) in Bavendamm test was measured by the diameter of brown pigmentation formed around <i>C. eragrostidis</i> (CE), <i>T. asperellum</i> (TA), <i>F. fujikuroi</i> (FF) and <i>F. oxysporum</i> (FO) growth colonies.	47
4.4	The presence of polyphenol oxidase (PPO) in Bavendamm test was measured by the diameter of brown pigmentation formed around <i>F. fujikuroi</i> (FF), <i>F. oxysporum</i> (FO), <i>C. lunata</i> (CL) and <i>A. niger</i> (AN) growth colonies.	48

4.5	Mode of interaction between <i>T. asperellum</i> B1581 and <i>A. niger</i> B2484 after 5 days of incubation. (a) Front view (b) Rear view.	53
5.1	Optimization of all significant parameters using OFAT analysis; which were carried out in sequential pattern from parameter (a) temperature of fermentation to parameter (f) temperature of saccharification using <i>T. asperellum</i> B1581 and <i>A. niger</i> B2484.	64
5.2	The development of 3D surface plot based response of ethanol production produced by <i>T. asperellum</i> B1581 with interaction among parameters; A = hours of saccharification, B = temperature of saccharification, C = hours of fermentation, D = temperature of fermentation, E = media level and F = substrate level.	77
5.3	The development of 3D surface plot based response of ethanol production produced by <i>A. niger</i> B2484 with interaction among parameters; A = hours of saccharification, B = temperature of saccharification, C = hours of fermentation, D = temperature of fermentation, E = media level and F = substrate level.	79
5.4	The peak of ethanol from sample at retention time of 2.347 min.	87

LIST OF ABBREVIATIONS

ADF	Acid-detergent fiber
ADL	Acid-detergent lignin
ANOVA	Analysis of variance
ATP	Adenosine triphosphate
CAZymes	Carbohydrate active enzymes
CBH	Cellobiohydrolase
CBP	Consolidated bioprocessing
CCD	Central composite design
CMC	Carboxymethyl-cellulose
DP	Degree of polymerization
EI	Enzymatic index
FAO	Food and Agriculture Organization
g	gram
GC-FID	Gas chromatography – flame ionization detector
GHG	Greenhouse gases
h	Hours
HSD	Honestly significant difference
IU	International Unit
kWh	kilowatt hour
kPa	kilopascal
L	Litre
M	Molar
mm	millimeters
mL	milliliters

min	minutes
OFAT	One-factor-at-a-time
pH	potential of Hydrogen
ppb	Parts per billion
PPO	Polyphenol oxidase
psi	pounds per square inch
RSM	Response surface methodology
RPM	Rotation per minute
SD	Standard deviation
SEM	Scanning electron microscope
U/mL	Units per millilitre
w	weight
v	volume
3D	3-dimensional



© COPYRIGHT UPM

CHAPTER 1

INTRODUCTION

1.1 Background of the study

In December 2017, the Food and Agriculture Organization (FAO) has elevated its estimation of global rice production in the year 2017 by 2.9 million tonnes to 759.6 million tonnes (503.9 million tonnes, milled basis) and China remains the largest paddy producer in the Asia region (FAO, 2018). Rice is one of the most favorite staple food for Malaysian, and with the implementation of National Agrofood Policy 2011-2020 (NAP 4), the local rice production has been increased to ensure country's stock of rice is sufficient to meet future demands as the population of Malaysian increases over the years (Rajamoorthy, Abdul Rahim and Munusamy, 2015). About 23% of overall paddy weight will generate a by-product known as paddy straw, which is frequently disposed off by open burning to clear the fields for the next cycle of rice planting (Kaur and Phutela, 2018). Paddy straw are also referred as an agricultural waste comprising of the dry stalks of crops and usually collected after harvesting period (Bakker, Elbersen, Poppens and Lesschen, 2013). The disposal of paddy straw through open burning was carried out to eliminate the sources of rat infestation, insect pests and to prevent rice diseases (Rosmiza, Davies, Rosniza Aznie, Mazdi and Jabil, 2014). Burning of biomass material releases wide-range of gases such as carbon monoxide (CO), carbon dioxide (CO₂), methane (CH₄), aldehyde, organic acid and inorganic elements, volatile and semi-volatile organic compounds, and particulate matter (PM); affecting people lives nearby (Yadav and Devi, 2018).

Paddy straw is normally being utilised as materials for cattle feed, fuel for residential cooking, thatching for rural houses, gasification, power generation, mulching material, and paper mills (Roy and Kaur, 2015). The compositions of paddy straw are 32.15% cellulose, 28% hemicellulose and 19.6% lignin; which contains major constituents of lignocellulosic materials (Shawky, Mahmoud, Ghazy, Asker and Ibrahim, 2011). Ideally, only cellulose and hemicellulose have the ability to be converted into fermentable sugars (Moiser *et al.*, 2005). These lignocellulosic materials (wood and agricultural crops residue) are promising feedstock for generating variety of great products such as bioethanol (Yoswathana, Phuriphapat, Treyawutthiwat and Eshtiaghi, 2010).

Biofuels are renewable and sustainable energy resources which offer an alternative solution to our conventional fuel sources dilemma as well as an effort to put a halt to climate change (Hidayat, Rochmadi, Wijaya and Budiman, 2016). Up to June 2017, the Malaysia Automotive Association (MAA) has released the total number of Malaysian vehicles on the roads standing at 28,181,203 units (MAA, 2017). With the advancements in the transportation

industry in Malaysia, the demands of energy have increased over the years and urged scientist to develop new renewable energy source to replace the ordinary conventional fuel. Research on renewable energy has gained supports from Malaysian government with the implementation of 5-Fuel Policy (2001) under the 8th Malaysia Plan, in which targeted to achieve 5% mixture of RE, but, only manage to achieve 1.8% in the following year (Loh and Choo, 2013).

The development of first-generation bioethanol based on food crops suffers from the criticism due to competition between food supply and bioethanol development; causing a sudden increase in food prices (Naik, Goud, Rout and Dalai, 2010). The weakness of first-generation bioethanol has highlighted the need to develop second-generation bioethanol based on lignocellulosic agro waste (Sims, Mabee, Saddler and Taylor, 2010). The second generation bioethanol develops from woody biomass are more energy efficient, flexible in terms of their feedstock and not competing with the human food resources (Havlik *et al.*, 2011). Currently, the most progressive technologies in bioethanol production are focusing on transforming lignocellulosic feedstock into transportation energy (Bakker *et al.*, 2013). Theoretically, paddy straw can generate up to 205 billion liter ethanol across the globe, from a single biomass feedstock with only 5% of total consumption (Belal, 2013).

In order to develop bioethanol efficiently and cost-effective from cellulosic feedstock, the degradation of biomass by energetic cellulases and fermentation by dynamic fermentative microorganism are basically important (Takano and Hoshino, 2012). In this case, the lignocellulolytic fungi are the best choice. There are many species of lignocellulolytic fungi such as *Aspergillus* sp., *Trichoderma* sp., *Fusarium* sp. and *Neurospora* sp. (Ferreira, Mahboubi, Lennartsson and Taherzadeh, 2016). These natural occurring fungi may have several benefits with their fermentability in producing bioethanol over the standard baker's yeast *Saccharomyces cerevisiae* (Okamoto, Nitta, Maekawa and Yanase, 2011). Most of the lignocellulolytic fungi secrete extracellular and hydrolytic enzymes that work as a biocatalyst for lignin and cellulosic materials degradations (Mtui, 2012). These enzymes comprise of cellulolytic enzymes (e.g cellulase) and ligninolytic enzymes (e.g lignin peroxidase, manganese peroxidase and laccase) (Manavalan, Manavalan and Heese, 2015). Cellulase is a family of at least 3 groups of enzymes; endo-(1,4)- β -D-glucanase (EC 3.2.1.4), exo-(1,4)- β -D-glucanase (EC 3.2.1.91), and β -glucosidases (EC 3.2.1.21) (Kuhad, Gupta and Singh, 2011a). Researchers have developed strong interests in the production of cellulases as of their utilizations in industries of alcohol fermentation, brewing, pulp and paper industry as well as textile industry (Nasr, Badawi, Mona, Demerdash and Barakat, 2015). Besides, cellulose and hemicellulose content can be enzymatically degraded into simple sugars by cellulases and hemicellulases (Berlowska *et al.*, 2016).

There are 4 significant processes for robust lignocellulosic biomass production which depend on the variations in saccharification or fermentation conditions: separate hydrolysis and fermentation (SHF), simultaneous saccharification and

fermentation (SSF), consolidated bioprocessing (CBP) and simultaneous saccharification and co-fermentation (SSCF) (Parisutham, Kim, and Lee, 2014). The consolidated bioprocessing, or CBP, converts lignocellulosic material into desired products in a single step without adding exogenous enzymes and this process certainly has been a subject of research interest in recent years (Olson, McBride, Shaw and Lynd, 2012). A combination of enzyme secretion, saccharification, and fermentation process in the same bioreactor has been known for economical manufacturing of bioethanol by evading the high expenses on investment, feedstock as well as the equipment for microbial enzyme production (Hasunuma *et al.*, 2013). The principal of cost reduction in CBP derives from either: (1) fermentative organism secretes essential cellulolytic enzymes for degradation of biomass or (2) cellulolytic organism which has the ability to ferment and thus eliminating the need for a separate enzyme production step (Linger and Darzins, 2013).

1.2 Problem statement

Even though bioethanol may induce various benefits that lead towards minimizing the environmental impact, more research is required especially in fermentation technology modification, preparation of raw materials and for economical bioethanol production (Bhatia, Johri and Ahmad, 2012). Paddy straw is a good material for the development of bioethanol but the presence of high ash and silica content in the feedstock has limited the bioconversion process to occur efficiently (Ibrahim, 2012). Unlike the first-generation of bioethanol, the second-generation utilizes the lignocellulosic substrates known as cellulose which is enclosed within hemicellulose and lignin matrix in the cell wall, making the accessibility of cellulose become the major problem in bioethanol production from such sources (Wi, Choi, Kim, Kim and Bae, 2013).

The mechanism of bioconversion process involved the production of sugars from biomass which requires new biotechnological improvements in order to ensure their efficiency enhancement and also economically applicable. Although some fungal strains are exclusively known as lignocellulolytic and thermostable, most of these fungal strains failed to secrete satisfactory amounts of cellulolytic enzymes that are mandatory for a productive cellulose conversion into desired product (Dashtban, Schraft and Qin, 2009). Choosing an appropriate cellulase for saccharification process is extremely challenging as each material have structural difference and difference of enzymatic activities in industrial cellulase reagents (Takano and Hoshino, 2018).

Besides selecting a productive strain, designing a suitable culture condition is also crucial to improve the efficiency of ethanol production systematically by either adding or eliminating components from the formulation, which resulted in a more stabilized and reproducible culture conditions (Dong, Zhao, Ma and Zhang, 2012). The amount of cellulase production seems to rely upon several factors such as incubation period, pH, temperature, carbon, nitrogen sources and cations (Gautam *et al.*, 2011).

The consolidated bioprocessing (CBP), in which enzyme secretion, break down of substrate, and fermentation process is achieved in one-step using either single microorganism or a group of compatible microorganisms and also known as the most economically attractive method for bioconversion of lignocellulosic biomass into bioethanol (Olson *et al.*, 2012; Ho *et al.*, 2012; Wang *et al.*, 2015a). However, the most difficult step in CBP is the selection of an appropriate microorganism or microbial consortium that secretes suitable hydrolytic enzymes corresponding to the lignocellulosic material, and produce ethanol (Paulova, Patakova, Branska, Rychtera and Melzoch, 2015).

1.3 Objectives of the study

The idea of this study is to utilize pretreated paddy straw as bioethanol material using consolidated bioprocessing (CBP) approach with the help from lignocellulolytic fungi. Therefore, the main aim of the study is to explore the use of lignocellulolytic fungi in consolidated bioprocessing (CBP) for an efficient conversion of paddy straw into bioethanol. The outline of the research approach is shown in Figure 1.1 and the design of the study was based on the following specific objectives to address the foregoing issues as stated in 1.2:

- i. To determine the optimized pretreatment method for paddy straw in removing silica, delignification and enhanced the accessibility towards cellulosic content
- ii. To determine the best lignocellulolytic agent for hydrolyzation of lignocellulosic materials of paddy straw.
- iii. To determine the factors that influences the optimization of bioethanol production via one-factor-at-a-time (OFAT) analysis and optimizing the physico-chemical parameters using Response Surface Methodology analysis (RSM).
- iv. To evaluate the efficiency of the final CBP set-up using either single microbe or consortium microbes to improve the amount of ethanol yield.




Figure 1.1: Summary of research design and attainment of objectives

REFERENCES

- Abo-State, M.A., Ragab, A.M.E., El-Gendy, N.S., Farahat, L.A., and Madian, H.R. (2014). Bioethanol production from rice straw enzymatically saccharified by fungal isolates, *Trichoderma viride* F94 and *Aspergillus terreus* F98. *Soft*, 3, 19-29.
- Acharya, S., and Chaudhary, A. (2012). Bioprospecting thermophiles for cellulase production: A review. *Brazilian Journal of Microbiology*, 43, 844-856.
- Agbor, V.B., Cicek, N., Sparling, R., Berlin, A., and Levin, D.B. (2011). Biomass pretreatment: Fundamentals toward application. *Biotechnology Advances*, 29, 675–685.
- Aggarwal, N.K., Goyal, V., Saini, A., Yadav, A., and Gupta, R. (2017). Enzymatic saccharification of pretreated rice straw by cellulases from *Aspergillus niger* BK01. *3 Biotech*, 7, 1-10.
- Argo, B.D., Wardani, A.K., Zubaidah, E., and Winarsih, S. (2014). The increasement of rice straw hydrolysis using blend crude cellulose enzyme from *Trichoderma reesei* and *Aspergillus niger*. *Research and Review in Biosciences*, 9, 41-44.
- Ahmed, A., and Bibi, A. (2018). Fungal cellulase; production and applications: Minireview. *LIFE: International Journal of Health and Life Sciences*, 4, 19-36.
- Akhtar, N., Goyal, D., and Goyal, A. (2017). Characterization of microwave-alkali-acid pre-treated rice straw for optimization of ethanol production via simultaneous saccharification and fermentation (SSF). *Energy Conversion and Management*, 141, 133–144.
- Akram, F., Haq, I.U., Imran, W., and Mukhtar, H. (2018). Insight perspectives of thermostable endoglucanases for bioethanol production: A review. *Renewable Energy*, 122, 225-238.
- Al-Bukhaiti, W.Q., Noman, A., Qasim, A.S., and Al-Farga, A. (2017). Gas chromatography: Principles, advantages and applications in food analysis. *International Journal of Agriculture Innovations and Research*, 6, 123-128.
- Albernas-Carvajala, Y., Corsanob, G., Cortésa, M.G., and Suárez, E.G. (2017). Preliminary design for simultaneous saccharification and fermentation stages for ethanol production from sugar cane bagasse. *Chemical Engineering Research and Design*, 126, 232–240.
- Ali, S.S., Khan, M., Mullins, E., and Doohan, F. (2012). The effect of wheat genotype on ethanol production from straw and the implications for multifunctional crop breeding. *Biomass and Bioenergy*, 42, 1-9.

- Ali, S.S., Nugent, B., Mullins, E., and Doohan, F. (2016). Fungal-mediated consolidated bioprocessing: the potential of *Fusarium oxysporum* for the lignocellulosic ethanol industry. *AMB Express*, 6, 1-13.
- Allen, S.A.A., Ana Godson, R.E.E., Ayodeji, S.A.M., and Deborah, S.A.E. (2016). Lignocelluloses: An economical and ecological resource for bioethanol production – A Review. *International Journal of Natural Resource Ecology and Management*, 1, 128-144.
- Alvira, P., Tomás-Pejó, M., Ballesteros, M., and Negro, M.J. (2009). Pretreatment technologies for an efficient bioethanol production process based on enzymatic hydrolysis: a review. *Bioresource Technology*, 101, 4851- 4861.
- Amarasekara, A.S. (2013). Fermentation 1 – Microorganism. In Amarasekara, A.S. (Ed.), *Handbook of cellulosic ethanol*. United States of America: Scrivener Publishing.
- Amiri, H., Karimi, K., and Zilouei, H. (2014). Organosolv pretreatment of rice straw for efficient acetone, butanol, and ethanol production. *Bioresource Technology*, 152, 450–456.
- Amin, F.R., Khalid, H., Zhang, H., Rahman, S.U., Zhang, R., Liu, G., and Chen, C. (2017). Pretreatment methods of lignocellulosic biomass for anaerobic digestion. *AMB Express*, 7, 1-12.
- Anasontzis, G.E., Kourtoglou, E., Villas-Boas, S.G., Hatzinikolaou, D.G., and Christakopoulos, P. (2016). Metabolic engineering of *Fusarium oxysporum* to improve its ethanol-producing capability. *Frontiers in Microbiology*, 7, 1-10.
- Anasontzis, G.E., and Christakopoulos, P. (2014). Challenges in ethanol production with *Fusarium oxysporum* through consolidated bioprocessing. *Bioengineered*, 5, 393-395.
- Arantes, V. and Saddler, J.N. (2010). Access to cellulose limits the efficiency of enzymatic hydrolysis: the role of amorphogenesis. *Biotechnology for Biofuels* 3, 4 – 15.
- Arifa, T., Madiha, A., and Tasnim, F. (2010). Effect of cultural conditions on ethanol production by locally isolated *Saccharomyces cerevisiae* bio-07. *Journal of Applied Pharmacy*, 3, 72-78.
- Arora, A., Priya, S., Sharma, P., Sharma, S., and Nain, L. (2016). Evaluating biological pretreatment as a feasible methodology for ethanol production from paddy straw. *Biocatalysis and Agricultural Biotechnology*, 8, 66–72.
- Artifon, W., Bonatto, C., Bordin, E.R., Bazoti, S.F., Dervanoski, A., Alves, S.L., and Treichel, H. (2018). Bioethanol production from hydrolyzed lignocellulosic after detoxification via adsorption with activated carbon

and dried air stripping. *Frontiers in Bioengineering and Biotechnology*, 6, 1-6.

- Asadollahzadeh, M.T., Ghasemian, A., Saraeian, A.R., Resalati, H., Lennartsson, P.R., and Taherzadeh, M.J. (2017). Ethanol and biomass production from spent sulfite liquor by filamentous fungi. *International Journal of Biotechnology and Bioengineering*, 11, 718-724.
- Ashgar, A., Abdul Rahman, A.A., and Wan Daud, W.M.A. (2014). A comparison of central composite design and Taguchi method for optimizing fenton process. *The Scientific World Journal*, 2014, 1-14.
- Bader, J., Mast-Gerlach, E., Popovic, M.K., Bajpai, R., and Stahl, U. (2010). Relevance of microbial coculture fermentations in biotechnology. *Journal of Applied Microbiology*, 109, 371–387.
- Bakker, R., Elbersen, W., Poppens, R., and Lesschen, J.P. (2013). Rice straw and wheat straw management. In Bakker, R., Elbersen, W., Poppens, R., and Lesschen, J.P. (Eds.), *Rice straw and wheat straw. Potential feedstocks for the biobased economy* (pp. 11 – 16). Netherlands: NL Agency Ministry of Economic Affairs.
- Banoth, C., Sunkar, B., Tondamanati, P.R., and Bhukya, B. (2017). Improved physicochemical pretreatment and enzymatic hydrolysis of rice straw for bioethanol production by yeast fermentation. *3 Biotech*, 7, 1-11.
- Bansal, N., Tewari, R., Soni, R., and Soni, S.K. (2012). Production of cellulases from *Aspergillus niger* NS-2 in solid state fermentation on agricultural and kitchen waste residues. *Waste Management*, 32, 1341–1346.
- Barakat, A., de Vries, H., and Rouau, X. (2013). Dry fractionation process as an important step in current and future lignocellulose biorefineries: A review. *Bioresource Technology*, 134, 362–373.
- Barakat, A., Mayer-Laigle, C., Solhy, A., Arancon, R.A.D., de Vries, H., and Luque, R. (2014). Mechanical pretreatments of lignocellulosic biomass: towards facile and environmentally sound technologies for biofuels production. *RSC Advances*, 4, 48109–48127.
- Bech, L., Herbst, F.A., Grell, M.N., Hai, Z., and Lange, L. (2015). On-site enzyme production by *Trichoderma asperellum* for the degradation of duckweed. *Fungal Genomics and Biology*, 5, 1-10.
- Belal, E.B. (2013). Bioethanol production from rice straw residues. *Brazilian Journal of Microbiology*, 44, 225-234.
- Benoit-Gelber, I., Gruntjes, T., Vinck, A., Veluw, J.G., Wosten, H.A.B., Boeren, S., Vervoort J.J.M. and Vries, R.P. (2017). Mixed colonies of *Aspergillus niger* and *Aspergillus oryzae* cooperatively degrading wheat bran. *Fungal Genetics and Biology*, 102, 31–37.

- Bensah, E.D., and Mensah, M. (2013). Chemical pretreatment methods for the production of cellulosic ethanol: technologies and innovations. *International Journal of Chemical Engineering*, 2013, 1-21.
- Bensah, E.C., and Mensah, M.Y. (2018). Emerging physico-chemical methods for biomass pretreatment. In Basso, T.P., and Basso, L.C. (Eds.), *Fuel Ethanol Production from Sugarcane*. United Kingdom: INTECH Open Access Publisher.
- Behera, S., Arora, R., Nandhagopal, N., and Kumar, S. (2014). Importance of chemical pretreatment for bioconversion of lignocellulosic biomass. *Renewable and Sustainable Energy Reviews*, 36, 91–106.
- Behera, S.S., and Ray, R.C. (2016). Solid state fermentation for production of microbial cellulases: Recent advances and improvement strategies. *International Journal of Biological Macromolecules*, 86, 656–669.
- Behera, B.C., Sethi, B.K., Mishra, R.R., Dutta, S.K., and Thatoi, H.N. (2017). Microbial cellulases – diversity & biotechnology with reference to mangrove environment: A review. *Journal of Genetic Engineering and Biotechnology*, 15, 197–210.
- Berlowska, J., Pielech-Przybylska, K., Balcerek, M., Dziekońska-Kubczak, U, Patelski, P., Dziugan, P., and Krwgiel, D. (2016). Simultaneous saccharification and fermentation of sugar beet pulp for efficient bioethanol production. *BioMed Research International*, 2016, 1-10.
- Bertrand, E., Vandenbergh, L.P.S., Soccol, C.R., Sigoillot, J.C., and Faulds, C. (2016). First Generation Bioethanol. In: Soccol C., Brar S., Faulds C., Ramos L. (Eds.), *Green Fuels Technology*. Switzerland: Springer, Cham.
- Bezzera, M.A., Santelli, R.E., Oliveira, E.P., Villar, L.S., and Escalera, L.A. (2008). Response surface methodology (RSM) as a tool for optimization in analytical chemistry. *Talanta*, 76, 965–977.
- Bhatnagar, V.S., Sharma, N.R., and Kumar, S. (2015). Pretreatment of paddy straw for enhanced saccharification. *Journal of Chemical and Pharmaceutical Research*, 7, 914-920.
- Binod, P., Sindhu, R., Singhania, R.R., Vikram, S., Devi, L., Nagalakshmi, S., Kurien, N., Sukumaran, R.K., and Pandey, A. (2010). Bioethanol production from rice straw: An overview. *Bioresource Technology*, 101, 4767–4774.
- Binod, P., Kuttiraja, M., Archana, M., Janu, K.U., Sindhu, R., Sukumaran, R.K., and Pandey, A. (2012). High temperature pretreatment and hydrolysis of cotton stalk for producing sugars for bioethanol production. *Fuel*, 92, 340–345.

- Biswas, R., Persad, A., and Bisaria, V.S. (2014). Production of cellulolytic enzymes. In Bisaria, V.S., and Kondo, A. (Eds.), *Bioprocessing of Renewable Resources to Commodity Bioproducts* (pp. 105-132). United States of America: John Wiley & Sons, Inc.
- Bhatia, L., Johri, S., and Ahmad, R. (2012). An economic and ecological perspective of ethanol production from renewable agro waste: A review. *AMB Express*, 2, 1-19.
- Boucher, J., Chirat, C., and Lachenal, D. (2014). Extraction of hemicelluloses from wood in a pulp biorefinery, and subsequent fermentation into ethanol. *Energy Conversion and Management*, 88, 1120–1126.
- Brady, S.K., Sreelatha, S., Feng, Y., Chundawat, S.P.S. and Lang, M.J. (2015). Cellobiohydrolase 1 from *Trichoderma reesei* degrades cellulose in single cellobiose steps. *Nature Communications*, 6, 10149.
- Brethauer, S., and Studer, M.H. (2014). Consolidated bioprocessing of lignocellulose by a microbial consortium. *Energy and Environmental Science*, 7, 1446-1453.
- Brethauer, S., and Studer, M.H. (2015). Biochemical conversion processes of lignocellulosic biomass to fuels and chemicals – A review. *CHIMIA International Journal for Chemistry*, 69, 572-581.
- Brijwani, K., and Vadlani, P.V. (2011). Cellulolytic enzymes production via solid-state fermentation: effect of pretreatment methods on physicochemical characteristics of substrate. *Enzyme Research*, 2011, 1-10.
- Brink, J.V., Maitan-Alfenas, G.P., Zou, G., Wang, C., Zhou, Z., Guimaraes, V.M. and Vries, R.P. (2014). Synergistic effect of *Aspergillus niger* and *Trichoderma reesei* enzyme sets on the saccharification of wheat straw and sugarcane bagasse. *Biotechnology Journal*, 9, 1329–1338.
- Brodeur, G., Yau, E., Badal, K., Collier, J., Ramachandran, K.B., and Ramakrishnan, S. (2011). Chemical and physicochemical pretreatment of lignocellulosic biomass: A review. *Enzyme Research*, 2011, 1-17.
- Brus, C. (2015). Techniques for determining alcohol levels in wine and mulled wine. Örebro Universitet.
- Buckee, G.K., and Mundy, A.P. (1993). Determination of ethanol in beer by gas chromatography (direct injection) - collaborative trial. *Journal of the Institute of Brewing*, 99, 381-384.
- Cabrera, E., Muñoz, M.J., Martín, R., Caro, I., Curbelo, C., and Díaz, A.B. (2014). Alkaline and alkaline peroxide pretreatments at mild temperature to enhance enzymatic hydrolysis of rice hulls and straw. *Bioresource Technology*, 167, 1–7.

- Carrasco F. (1989). Fundamentos del fraccionamiento de la biomasa. *Afinidad*, 46, 425-429.
- Cavka, A., Alriksson, B., Rose, S.H., van Zyl, W.H., and Jönsson, L.J. (2014). Production of cellulosic ethanol and enzyme from waste fiber sludge using SSF, recycling of hydrolytic enzymes and yeast, and recombinant cellulase-producing *Aspergillus niger*. *Journal of Industrial Microbiology and Biotechnology*, 41, 1191–1200.
- Chandra, R.P., Bura, R., Mabee, W.E., Berlin, A., Pan, X., and Saddler, J.N. (2007). Substrate pretreatment: the key to effective enzymatic hydrolysis of lignocellulosics? *Advances. Biochemical Engineering Biotechnology*, 108, 67–93.
- Chang, Y.C., Lee, C.L., and Pan, T.M. (2006). Statistical optimization of media components for the production of *Antrodia cinnamomea* AC0623 in submerged cultures. *Applied Microbiology and Biotechnology*, 72, 654-661.
- Chandel, A.K., Silva, S.S., and Singh, O.V. (2013). Detoxification of lignocellulose hydrolysates: biochemical and metabolic engineering toward white biotechnology. *Bioenergy Resource*, 6, 388–401.
- Chaturvedi, V., and Verma, P. (2013). An overview of key pretreatment processes employed for bioconversion of lignocellulosic biomass into biofuels and value added products. *3 Biotech*, 3, 415–431.
- Chen, Y., Stevens, M.A., Zhu, Y., Holmes, J., and Xu, H. (2013). Understanding of alkaline pretreatment parameters for corn stover enzymatic saccharification. *Biotechnology for Biofuels*, 6, 1-10.
- Chen, X., Zhang, Y.L., Gu, Y., Liu, Z., Shen, Z., Chu, H., and Zhou, X. (2014). Enhancing methane production from rice straw by extrusion pretreatment. *Applied Energy*, 122, 34–41.
- Cheng, Y.S., Zheng, Y., Yu, C.W., Dooley, T.M., Jenkins, B.M., and VanderGheynst, J.S. (2010). Evaluation of high solids alkaline pretreatment of rice straw. *Applied Biochemistry and Biotechnology*, 162, 1768–1784.
- Cheng, J.J., and Timilsina, G.R. (2011). Status and barriers of advanced biofuel technologies: A review. *Renewable Energy*, 36, 3541-3549.
- Cheng, J.R., and Zhu, M.J. (2013). A novel co-culture strategy for lignocellulosic bioenergy production: a systematic review. *International Journal of Modern Biology and Medicine*, 1, 166-193.
- Chin, K.L., and H'ng, P.S. (2013). A real story of bioethanol from biomass: Malaysia perspective. In Matovic, M.D. (Ed.), *Biomass Now: Sustainable Growth and Use*. United Kingdom: INTECH Open Access Publisher.

- Christian, G.D. (2004). Gas chromatography. In Christian, G.D. (Ed.), *Analytical chemistry* (pp. 574-593). United States of America: John Wiley & Sons, Inc.
- Correa, S.J., Jaramillo, A.C., Merino, R.A., and Hormaza, A. (2018). Evaluation of individual fungal species and their co-culture for degrading a binary mixture of dyes under solid-state fermentation. *Biotechnology, Agronomy, Social and Environment*, 22, 242-251.
- Cui, Y., Dong, X., Tong, J., and Liu, S. (2015). Degradation of lignocellulosic components in un-pretreated vinegar residue using an artificially constructed fungal consortium. *BioResources*, 10, 3434-3450.
- Czitrom, V. (1999). One-factor-at-a-time versus designed experiments. *The American Statistician*, 53, 126-131.
- Damaso, M.C.T., Terzi, S.C., Farias, A.X., Oliveira, A.C.P., Fraga, M.E. and Couri, S. (2012). Selection of cellulolytic fungi isolated from diverse substrates. *Brazilian Archives of Biology and Technology*, 55, 513 – 520.
- Das, A., Paul, T., Jana, A., Halder, S.K., Ghosh, K., Maity, C., Mohapatra, P.K.D., Pati, B.R., and Mondal, K.C. (2013). Bioconversion of rice straw to sugar using multizyme complex of fungal origin and subsequent production of bioethanol by mixed fermentation of *Saccharomyces cerevisiae* MTCC 173 and *Zymomonas mobilis* MTCC 2428. *Industrial Crops and Products*, 46, 217– 225.
- Das, A., Mondal, C., and Roy, S. (2015). Pretreatment methods of ligno - cellulosic biomass: A Review. *Journal of Engineering Science and Technology Review*, 8, 141 -165.
- Dash, P.K., Mohapatra, S., Swain, M.R., and Thatoi, H. (2017). Optimization of bioethanol production from saccharified sweet potato root flour by co-fermentation of *Saccharomyces cerevisiae* and *Pichia* sp. using OVAT and response surface methodologies. *Acta Biologica Szegediensis*, 61, 13-23.
- Dashtbant, M. Schraft, H. and Qin, W. (2009). Fungal bioconversion of lignocellulosic residues; opportunities and perspectives. *International Journal of Biological Sciences*, 5, 578–595.
- Dasgupta, D., Suman, S.K., Pandey, D., Ghosh, D., Khan, R., Agrawal, D., Jain, R.K., Vadde, V.T., and Adhikari, D.K. (2013). Design and optimization of ethanol production from bagasse pith hydrolysate by a thermotolerant yeast *Kluyveromyces* sp. IPE453 using response surface methodology. *SpringerPlus*, 2, 1-10.
- Del Pozo, M.V., Fernández-Arrojo, L., Gil-Martínez, J., Montesinos, A., Chernikova, T.N., Nechitaylo, T.Y., Waliszek, A., Tortajada, M., Rojas, A., Huws, S.A., Golyshina, O.V., Newbold, C.J., Polaina, J., Ferrer, M. and Golyshin, P.N. (2012). Microbial β -glucosidases from cow rumen

metagenome enhance the saccharification of lignocellulose in combination with commercial cellulase cocktail. *Biotechnology for Biofuels*, 5, 73.

Derman, E., Abdulla, R., Marbawi, H., and Sabullah, M.K. (2018). Oil palm empty fruit bunches as a promising feedstock for bioethanol production in Malaysia. *Renewable Energy*, 129, 285-298.

Devarapalli, M., and Atiyeh, H.K. (2015). A review of conversion processes for bioethanol production with a focus on syngas fermentation. *Biofuel Research Journal*, 7, 268-280.

Dhap, N., and Singh, H. (2017). Pretreatment of rice straw for bio-ethanol production: A review. *Journal of Chemical and Pharmaceutical Research*, 9, 216-220.

Dhillon, G.S., Oberoi, H.S., Kaur, S., Bansal, S., and Brar, S.K. (2011). Value-addition of agricultural wastes for augmented cellulase and xylanase production through solid-state tray fermentation employing mixed-culture of fungi. *Industrial Crops and Products*, 34, 1160– 1167.

Dong, H.N., Zhao, X.M., Ma, Y.Y., and Zhang, M.H. (2012). Optimization of a synthetic medium for ethanol production by xylose-fermenting *Zymomonas mobilis* using response surface methodology. *Chinese Science Bulletin*, 57, 3782-3789.

Dutta, K., Daverey, A., and Lin, J.G. (2014). Evolution retrospective for alternative fuels: First to fourth generation. *Renewable Energy*, 69, 114-122.

Elkins, E.G., Raman, B., and Keller, M. (2010). Engineered microbial systems for enhanced conversion of lignocellulosic biomass. *Current Opinion in Biotechnology*, 21, 657-662

Ellila, S., Fonseca, L., Uchima, C., Cota, J., Goldman, G.H., Saloheimo, M., Sacon, V., and Siika-aho, M. (2017). Development of a low-cost cellulase production process using *Trichoderma reesei* for Brazilian biorefineries. *Biotechnology Biofuels*, 10, 1-17.

El-Naggar, N.E.A., Deraz, S., and Khalil, A. (2014). Bioethanol production from lignocellulosic feedstocks based on enzymatic hydrolysis: Current status and recent development. *Biotechnology*, 13, 1-21

El-Zawawy, W.K., Ibrahim, M.M., Abdel-Fattah, Y.R., Soliman, N.A., and Mahmoud, M.M. (2011). Acid and enzyme hydrolysis to convert pretreated lignocellulosic materials into glucose for ethanol production. *Carbohydrate Polymers*, 84, 865–871.

EPA (2010): Calibration curves: Program use/needs. *Forum on Environmental Measurements*.

- Fan, G., Wang, M., Liao, C., Fang, T., Li, J., and Zhou, R. (2013). Isolation of cellulose from rice straw and its conversion into cellulose acetate catalyzed by phosphotungstic acid. *Carbohydrate Polymers*, 94, 71– 76.
- Fan, Z. (2014). Consolidated bioprocessing for ethanol production. In Qureshi, N., Hodge, D., and Vertes, A.A. (Eds.), *Biorefineries integrated biochemical processes for liquid biofuels* (pp. 141-160). Amsterdam, Netherlands: Elsevier.
- Fang, H., Zhao, C., and Song, X.Y. (2010). Optimization of enzymatic hydrolysis of steam-exploded corn stover by two approaches: Response surface methodology or using cellulase from mixed cultures of *Trichoderma reesei* RUT-C30 and *Aspergillus niger* NL02. *Bioresource Technology*, 101, 4111–4119.
- Fang, H., Zhao, C., Song, X.Y., Chen, M., Chang, Z. and Chu, J. (2013). Enhanced cellulolytic enzyme production by the synergism between *Trichoderma reesei* RUT-C30 and *Aspergillus niger* NL02 and by the addition of surfactants. *Biotechnology and Bioprocess Engineering*, 18, 390 – 398.
- FAO Rice Market Monitor, April 2018, Volume XXI - Issue No. 1
- Fareeha, R., Nasreen, A.R., Uzma, H., Ikram, U.H., and Mariam, I. (2011). Solid state fermentation for the production of β -glucosidase by co-culture of *Aspergillus niger* and *A. oryzae*. *Pakistan Journal of Botany*, 43, 75-83.
- Fakruddin, M., Abdul Quayum, M., Morshed Ahmed, M., and Choudhury, N. (2012). Analysis of key factors affecting ethanol production by *Saccharomyces cerevisiae* IFST-072011. *Biotechnology*, 11, 248-252.
- Ferreira, J.A., Lennartsson, P.R., and Taherzadeh, M.J. (2014). Production of ethanol and biomass from thin stillage using food-grade zygomycetes and ascomycetes filamentous fungi. *Energies*, 7, 3872-3885.
- Ferreira, A., Mahboubi, A., Lennartsson, P.R., and Taherzadeh, M.J. (2016). Waste biorefineries using filamentous ascomycetes fungi: Present status and future prospects. *Bioresource Technology*, 215, 334-345.
- FitzPatrick, M., Champagne, P., Cunningham, M.F., and Whitney, R.A. (2010). A biorefinery processing perspective: Treatment of lignocellulosic materials for the production of value-added products. *Bioresource Technology*, 101, 8915–8922.
- Florencio, C., Couri, S. and Farinas, C.S. (2012). Correlation between agar plate screening and solid-state fermentation for the prediction of cellulase production by *Trichoderma* strains. *Enzyme Research*, 2012, 1 – 7.
- Galbe, M., and Zacchi, G. (2012). Pretreatment: The key to efficient utilization of lignocellulosic materials. *Biomass and Bioenergy*, 46, 70-78.

- Garcia-Maraver, A., Salvachúa, D., Martínez, M.J., Diaz, L.F., and Zamorano, M. (2013). Analysis of the relation between the cellulose, hemicellulose and lignin content and the thermal behavior of residual biomass from olive trees. *Waste Management*, 33, 2245–2249
- Garvey, M., Klose, H., Fischer, R., Lambertz, C., and Commandeur, U. (2013). Cellulases for biomass degradation: comparing recombinant cellulase expression platforms. *Trends in Biotechnology*, 31, 581-593.
- Gautam, S.P., Bundela, P.S., Pandey, A.K., Khan, K., Awasthi, M.K., and Sarsaiya, S. (2011). Optimization for the production of cellulase enzyme from municipal solid waste residue by two novel cellulolytic fungi. *Biotechnology Research International*, 2011, 1-8.
- Ghaffar, S.H., and Fan, M. (2013). Structural analysis for lignin characteristics in biomass straw. *Biomass Bioenergy*, 57, 264-279.
- Giles, R.L., Zackeru, J.C., Galloway, E.R., Elliot, G.D., and Parrow, M.W. (2015). Single versus simultaneous species treatment of wood with *Ceriporiopsis subvermisporea* and *Postia placenta* for ethanol applications, with observations on interspecific growth inhibition. *International Biodeterioration and Biodegradation*, 99, 66-72.
- Gírio, F.M., Fonseca, C., Carvalheiro, F., Duarte, L.C., Marques, S., and Bogel-Lukasik, R. (2010). Hemicelluloses for fuel ethanol: A review. *Bioresource Technology*, 101, 4775 – 4800.
- Goering, H.K., and Van Soest, P.J. (1970). Forage fiber analysis (apparatus, reagents, procedures, and some applications). In Goering, H.K., and Van Soest, P.J. (Eds.), *Agriculture Handbook No. 379* (pp. 1-20). Washington D.C.: USDA-ARS.
- Goff, B.M., Moore, K.J., Fales, S.L., and Pedersen, J.F. (2011). Comparison of gas chromatography, spectrophotometry and near infrared spectroscopy to quantify prussic acid potential in forages. *Journal of the Science of Food and Agriculture*, 91, 1523-1526.
- Gu, F., Wang, W., Jing, L., and Jin, Y. (2013). Sulfite–formaldehyde pretreatment on rice straw for the improvement of enzymatic saccharification. *Bioresource Technology*, 142, 218–224.
- Guan, R., Li, X., Wachemo, A.C., Yuan, H., Liu, Y., Zou, D., Zuo, X., and Gu, J. (2018). Enhancing anaerobic digestion performance and degradation of lignocellulosic components of rice straw by combined biological and chemical pretreatment. *Science of the Total Environment*, 637–638, 9–17.
- Gude, V.G. (2017). Microwave pretreatment of feedstock for bioethanol production. In Gude, V.G. (Ed.), *Microwave-mediated biofuel production*. United States of America: CRC Press.

- Guiochon, G., and Guillemin, C.L. (1988). Journal of chromatography library. In Guiochon, G., and Guillemin, C.L. (Eds.), *Quantitative gas chromatography for laboratory analyses and on-line control* (pp. 795-797). Amsterdam: Elsevier.
- Guo, M., Song, W., and Buhain, J. (2015). Bioenergy and biofuels: History, status, and perspective. *Renewable and Sustainable Energy Reviews*, 42, 712–725.
- Guray, M.Z. (2009). Partial purification and characterization of polyphenol oxidase from thermophilic *Bacillus* sp., (Master Thesis), İzmir Institute of Technology.
- Gutiérrez-Correa, M., and Villena, G.K. (2012). Batch and repeated batch cellulase production by mixed cultures of *Trichoderma reesei* and *Aspergillus niger* or *Aspergillus phoenicis*. *Journal of Microbiology and Biotechnology Research*, 2, 929-935.
- Hamelinck, C.N., Hooijdonk, G.V., and Faaij, A.P. (2005). Ethanol from lignocellulosic biomass: Techno-economic performance in short, middle and long-term. *Biomass Bioenergy*, 28, 384-410.
- Han, J.S., and Rowell, J.S. (1997). Chemical composition of fibers. In Rowell, R.M., Young, R.A., and Rowell, J.K. (Eds.), *Paper and Composites from Agro-Based Resources* (pp. 92-99). United States of America: CRC Press Inc.
- Harmsen, P.F.H., Huijgen, W.J.J., López, L.M.B., and Bakker, R.R.C. (2010). Literature review of physical and chemical pretreatment processes for lignocellulosic biomass (No. 1184). *Wageningen UR, Food & Biobased Research*, 2010, 1-49.
- Harun, R., and Danquah, M.K. (2011). Influence of acid pre-treatment on microalgal biomass for bioethanol production. *Process Biochemistry*, 46, 304–309.
- Harun, S., and Geok, S.K. (2016). Effect of sodium hydroxide pretreatment on rice straw composition. *Indian Journal of Science and Technology*, 9, 1-9.
- Hasunuma, T., and Kondo, A. (2012). Development of yeast cell factories for consolidated bioprocessing of lignocellulose to bioethanol through cell surface engineering. *Biotechnology Advances*, 30, 1207-1218.
- Hasunuma, T., Okazaki, F., Okai, N., Hara, K.Y., Ishii, J., and Kondo, A. (2013). A review of enzymes and microbes for lignocellulosic biorefinery and the possibility of their application to consolidated bioprocessing technology. *Bioresource Technology*, 135, 513–522.

- Havlik, P., Schneider, U.A., Schmid, E., Bottcher, H., Fritz, S., Skalsky, R., Aoki, K., DeCara, S., Kindermann, G., Kraxner, F., Leduc, S., McCallum, I., Mosnier, A., Sauer, T., and Obersteiner, M. (2011). Global land-use implications of first and second generation biofuel targets. *Energy Policy*, 39, 5690–5702.
- He, Y., Pang, Y., Liu, Y., Li, X., and Wang, K. (2008). Physicochemical characterization of rice straw pretreated with sodium hydroxide in the solid state for enhancing biogas production. *Energy Fuels*, 22, 2775–2781.
- Hendriks, A., and Zeeman, G. (2009). Pretreatments to enhance the digestibility of lignocellulosic biomass. *Bioresource Technology*, 100, 10–18.
- Hennig, S., Wenzel, M., Haas, C., Hoftmann, A., Weber, J., Rödel, G., and Ostermann, K. (2018). New approaches in bioprocess-control: Consortium guidance by synthetic cell-cell communication based on fungal pheromones. *Engineering in Life Sciences*, 18, 387- 400.
- Hidayat, A., Rochmadi, Wijaya, K., and Budiman, A. (2016). Removal of free fatty acid in palm fatty acid distillate using sulfonated carbon catalyst derived from biomass waste for biodiesel production. *IOP Conference Series: Materials Science and Engineering*, 105, 1-8.
- Highina, B.K., Bugaje, I.M., and Umar, B. (2014). A review on second generation biofuel: a comparison of its carbon footprints. *European Journal of Engineering and Technology*, 2, 117-125.
- Hložek, T., Bursová, M., and Čabala, R. (2014). Fast determination of ethylene glycol, 1,2-propylene glycol and glycolic acid in blood serum and urine for emergency and clinical toxicology by GC-FID. *Talanta*, 130, 470–474.
- Ho, C.Y., Chang, J.J., Lee, S.C., Chin, T.Y., Shih, M.C., Li, W.H., and Huang, C.C. (2012). Development of cellulosic ethanol production process via co-culturing of artificial cellulosomal *Bacillus* and kefir yeast. *Applied Energy*, 100, 27-32.
- Ho, D.P., Ngo, H.H., and Guo, W. (2014). A mini review on renewable sources for biofuel. *Bioresource Technology*, 169, 742–749.
- Huang, J., Chen, D., Wei, Y., Wang, Q., Li, Z., Chen, Y., and Huang, R. (2014). Direct ethanol production from lignocellulosic sugars and sugarcane bagasse by a recombinant *Trichoderma reesei* strain HJ48. *The Scientific World Journal*, 2014, 1-9.
- Huberman, L.B., Liu, J., Qin, L., and Glass, N.L. (2016). Regulation of the lignocellulolytic response in filamentous fungi. *Fungal Biology Reviews*, 30, 101-111.

- Humbird, D., and Fei, Q. (2016). Scale-up considerations for biofuels. In Eckert, C.A., and Trinh, C.T. (Eds.), *Bioetchnology for biofuels production and optimization* (pp. 513-537). Amsterdam, Netherlands: Elsevier.
- Ibrahim, H.A. (2012). Pretreatment of straw for bioethanol production. *Energy Procedia*, 14, 542 – 551.
- Inglesby, M.K., Wood, D.F., and Gray, G.M. (2006). Effect of chemical fractionation treatments on silicon dioxide content and distribution in *Oryza sativa*. In Stokke, D.D., and Groom, L.H. (Eds.), *Characterization of the cellulosic cell wall* (pp. 192-212). Ames, IA: Blackwell Publishing.
- Islam, S., and Roy, N. (2018). Screening, purification and characterization of cellulase from cellulase producing bacteria in molasses. *BMC Research Notes*, 11, 1-6.
- Iqbal, A.K.M.A., and Khan, A.A. (2010). Modeling and analysis of MRR, EWR and surface roughness in EDM milling through response surface methodology. *American Journal of Engineering and Applied Sciences*, 3, 611-619.
- Iqbal, H.M.N., Kyazze, G., and Keshavarz, T. (2013). Advance in valorization of lignocellulosic materials by biotechnology: An overview. *BioResources*, 8(2), 3157-3176.
- Ja'afaru, M.I. (2013). Screening of fungi isolated from environmental samples for xylanase and cellulase production. *ISRN Microbiology*, 2013, 1 – 7.
- Jäger, G., Girfoglio, M., Dollo, F., Rinaldi, R., Bongard, H., Commandeur, U., Fischer, R., Spiess, A.C., and Büchs, J. (2011). How recombinant swollenin from *Kluyveromyces lactis* affects cellulosic substrates and accelerates their hydrolysis. *Biotechnology for Biofuels*, 4, 1-16.
- Jena, N., and Satpathy, S. (2017). Production of ethanol by *Trichoderma* spp. in solidstate fermentation of sugarcane molasses. *International Journal of Engineering, Science and Mathematics*, 6, 281-288.
- Jeya, M., Nguyen, N.P.T., Moon, H.J., Kim, S.H. and Lee, J.K. (2010). Conversion of woody biomass into fermentable sugars by cellulase from *Agaricus arvensis*. *Bioresource Technology*, 101, 8742–8749.
- Jia, H., Zhao, P., Wang, B., Tariq, P., Zhao, F., Zhao, M., Wang, Q., Yang, T., and Fang, J. (2016). Overexpression of polyphenol oxidase gene in strawberry fruit delays the fungus infection process. *Plant Molecular Biology Reporter*, 34, 592–606.
- Jiang, L.L., Zhou, J.J., Quan, C.S., and Xiu, Z.L. (2017). Advances in industrial microbiome based on microbial consortium for biorefinery. *Bioresources and Bioprocess*, 4, 1-10.

- Jin, M., Slininger, P.J., Dien, B.S., Waghmode, S., Moser, B.R., Orjuela, A., Sousa, L.C., and Balan, V. (2015). Microbial lipid-based lignocellulosic biorefinery: feasibility and challenges. *Trends in Biotechnology*, 33, 43-54.
- Joshi, B., Bhatt, M.R., Sharma, D., Joshi, J., Malla, R., and Sreerama, L. (2011). Lignocellulosic ethanol production: Current practices and recent developments. *Biotechnology and Molecular Biology Review*, 6, 172-182.
- Jouzani, G.S., and Taherzadeh, M.J. (2015). Advances in consolidated bioprocessing systems for bioethanol and butanol production from biomass: A comprehensive review. *Biofuel Research Journal*, 5, 152-195.
- Junko, U., and Kutsuyoshi, K. (2008). Process optimization to prepare high-purity amorphous silica from rice husks via citric acid leaching treatment. *Transaction of JWRI*, 37, 13-17.
- Kadarmoidheen, M., Saranraj, P., and Stella, D. (2012). Effect of cellulolytic fungi on the degradation of cellulosic agricultural wastes. *International Journal of Applied Microbiology Science*, 1, 13- 23.
- Kahr, H., Helmberger, S., and Jäger, A.G. (2011). Yeast adaptation on the substrate straw. *World Renewable Energy Congress*, 2011, 492-499.
- Kalyani, D., Lee, K.M., Kim, T.S., Li, J., Dhiman, S. S., Kang, Y. C. and Lee, J.K. (2013). Microbial consortia for saccharification of woody biomass and ethanol fermentation. *Fuel*, 107, 815-822.
- Kaneko, S., Mizuno, R., Maehara, T., and Ichinose, H. (2012). Consolidated bioprocessing ethanol production by using a mushroom. In Lima, M.A.P. (Ed.), *Bioethanol*. United Kingdom: INTECH Open Access Publisher.
- Kang, Q., Appels, L., Tan, T., and Dewil, R. (2014). Bioethanol from lignocellulosic biomass: current findings determine research priorities. *The Scientific World Journal*, 2014, 1-13.
- Kanmani, P., Karthik, S., Aravind, J., and Kumaresan, K. (2013). The use of response surface methodology as a statistical tool for media optimization in lipase production from the dairy effluent isolate *Fusarium solani*. *ISRN Biotechnology*, 2013, 1-8.
- Karimi, K., and Taherzadeh, M.J. (2016). A critical review on analysis in pretreatment of lignocelluloses: Degree of polymerization, adsorption/desorption, and accessibility. *Bioresource Technology*, 203, 348 – 356.
- Karunanithy, C., and Muthukumarappan, K. (2011). Optimization of switchgrass and extruder parameters for enzymatic hydrolysis using response surface methodology. *Industrial Crops and Products*, 33, 188-199.

- Kausar, H., Sariah, M., Mohd Saud, H., Zahangir Alam, M. and Ismail, M.R. (2010). Development of compatible lignocellulolytic fungal consortium for rapid composting of rice straw. *International Biodeterioration and Biodegradation*, 64, 594 – 600.
- Kaur, K., and Phutela, U.G. (2016). Enhancement of paddy straw digestibility and biogas production by sodium hydroxide-microwave pretreatment. *Renewable Energy*, 92, 178-184.
- Kaur, K., and Phutela, U.G. (2018). Morphological and structural changes in paddy straw influenced by alkali and microbial pretreatment. *DETRITUS - Multidisciplinary Journal for Waste Resources & Residues*, 3, 30-36.
- Kim, J.H., Block, D.E., and Mills, D.A. (2010). Simultaneous consumption of pentose and hexose sugars: an optimal microbial phenotype for efficient fermentation of lignocellulosic biomass. *Applied Microbiology and Biotechnology*, 88, 1077-1085.
- Kim, S.B., Lee, S.J., Lee, J.H., Jung, Y.R., Thapa, L.P., Kim, J.S., Um, Y., Park, C., and Kim, S.W. (2013). Pretreatment of rice straw with combined process using dilute sulfuric acid and aqueous ammonia. *Biotechnology for Biofuels*, 6, 1-11.
- Kim, D. (2018). Physico-chemical conversion of lignocellulose: inhibitor effects and detoxification strategies: A mini review. *Molecules*, 23, 1-21.
- Ko, J.K., Kim, Y., Ximenes, E., and Ladisch, M.R. (2015). Effect of liquid hot water pretreatment severity on properties of hardwood lignin and enzymatic hydrolysis of cellulose. *Biotechnology Bioengineering*, 112, 252-262.
- Kocar, G., and Civas, N. (2013). An overview of biofuels from energy crops: Current status and future prospects. *Renewable and Sustainable Energy Reviews*, 28, 900–916.
- Kolasa, M., Ahring, B.K., Lübeck, P.S., and Lübeck, M. (2014). Co-cultivation of *Trichoderma reesei* RutC30 with three black *Aspergillus* strains facilitates efficient hydrolysis of pretreated wheat straw and shows promises for on-site enzyme production. *Bioresource Technology*, 169, 143–148.
- Koppram, R., and Olsson, L. (2014). Combined substrate, enzyme and yeast feed in simultaneous saccharification and fermentation allow bioethanol production from pretreated spruce biomass at high solids loadings. *Biotechnology for Biofuels*, 7, 1-9.
- Kshirsagar, S.D., Waghmare, P.R., Loni, P.C., Patil, S.A., and Govindwar, S.P. (2015). Dilute acid pretreatment of rice straw, structural characterization and optimization of enzymatic hydrolysis conditions by response surface methodology. *RSC Advances*, 5, 46525–46533.

- Kuhad, R.C., Gupta, R., and Singh, A. (2011a). Microbial cellulases and their industrial applications. *Enzyme Research*, 2011, 1-10.
- Kuhad, R.C., Gupta, R., Khasa, Y.L., Singh, A., and Zhang, Y.H.P. (2011b). Bioethanol production from pentose sugars: Current status and future prospects. *Renewable and Sustainable Energy Reviews*, 15, 4950–4962.
- Kumagai, A., Kawamura, S., Lee, S.H., Endo, T., Rodriguez, M., and Mielenz, J.R. (2014). Simultaneous saccharification and fermentation and a consolidated bioprocessing for Hinoki cypress and *Eucalyptus* after fibrillation by steam and subsequent wet-disk milling. *Bioresource Technology*, 162, 89–95.
- Kumar, A., Gaiind, S., and Nain, L. (2008). Evaluation of thermophilic fungal consortium for paddy straw composting. *Biodegradation*, 19, 395–402.
- Kumar, P., Barrett, D.M., Delwiche, M.J., and Stroeve, P. (2009). Methods for pretreatment of lignocellulosic biomass for efficient hydrolysis and biofuel production. *Industrial & Engineering Chemistry Research*, 48, 3713–3729.
- Kumar, M., Singh, P., and Sukla, L.B. (2016). Addition of expansin to cellulase enhanced bioethanol production. *Process Biochemistry*, 51, 2097–2103.
- Kumar, A.K., and Sharma, S. (2017). Recent updates on different methods of pretreatment of lignocellulosic feedstocks: A review. *Bioresources and Bioprocessing*, 4, 1-19.
- Kumari, D., and Singh, R. (2018). Pretreatment of lignocellulosic wastes for biofuel production: A critical review. *Renewable and Sustainable Energy Reviews*, 90, 877–891.
- Lee, R.A., and Lavoie, J.M. (2013). From first- to third-generation biofuels: Challenges of producing a commodity from a biomass of increasing complexity. *Animal Frontiers*, 3, 6–11.
- Leiviska, K. (2013). Introduction to Experiment Design, University of Oulu, Control Engineering Laboratory.
- Lennartsson, P.R., Erlandsson, P., and Taherzadeh, M.J. (2014). Integration of the first and second generation bioethanol processes and the importance of by-products. *Bioresource Technology*, 165, 3–8.
- Li, J., Henriksson, G., and Gellerstedt, G. (2006). Lignin depolymerization/repolymerization and its critical role for delignification of aspen wood by steam explosion. *Bioresource Technology*, 98, 3061-3068.

- Li, H., Chai, X.S., Deng, Y., Zhan, H., and Fu, S. (2009). Rapid determination of ethanol in fermentation liquor by full evaporation headspace gas chromatography. *Journal of Chromatography A*, 1216, 169–172.
- Li, D. (2015). Cellulases from thermophilic fungi. In Li, L.F. (Ed.), *Thermophilic microorganisms* (pp. 121-140). China: Caister Academic Press.
- Li, C., Lin, F., Li, Y., Wei, W., Wang, H., Qin, L., Zhou, Z., Li, B., Wu, F., and Chen, Z. (2016). A β -glucosidase hyper-production *Trichoderma reesei* mutant reveals a potential role of cel3D in cellulase production. *Microbial Cell Factory*, 15, 1-13.
- Liang, Y.L., Zhang, Z., Wu, M., Wu, Y. and Feng, J.X. (2014). Isolation, screening, and identification of cellulolytic bacteria from natural reserves in the subtropical region of China and optimization of cellulase production by *Paenibacillus terrae* ME27-1. *BioMed Research International*, 2014, 1-13.
- Limayem, A., and Ricke, S.C. (2012). Lignocellulosic biomass for bioethanol production: Current perspectives, potential issues and future prospects. *Progress in Energy and Combustion Science*, 38, 449-467.
- Lin, H., Wang, B., Zhuang, R., Zhou, Q., and Zhao, Y. (2011). Artificial construction and characterization of a fungal consortium that produces cellulolytic enzyme system with strong wheat straw saccharification. *Bioresource Technology*, 102, 10569–10576.
- Lin, X., Fan, J., Wen, Q., Li, R., Jin, X., Wu, J., Qian, W., Liu, D., Xie, J., Bai, J., and Ying, H. (2014). Optimization and validation of a GC–FID method for the determination of acetone-butanol-ethanol fermentation products. *Journal of Chromatographic Science*, 52, 264–270.
- Linger, J., and Darzins, A. (2013). Consolidated bioprocessing. In Lee, J.W. (Ed.), *Advanced biofuels and bioproducts* (pp. 267-280). New York: Springer.
- Liu, H., Zhu, J., and Fu, S., (2010). Effects of lignin-metal complexation on enzymatic hydrolysis of cellulose. *Journal of Agriculture and Food Chemistry*, 58, 7233–7238.
- Liu, D., Zhang, R., Yang, X., Wu, H., Xu, D., Tang, Z., and Shen, Q. (2011). Thermostable cellulase production of *Aspergillus fumigatus* Z5 under solid-state fermentation and its application in degradation of agricultural wastes. *International Biodeterioration and Biodegradation*, 65, 717-725.
- Liu, Y.K., Chen, W.C., Huang, Y.C., Chang, Y.K., Chu, I.M., Tsai, S.L. and Wei, Y.H. (2017). Production of bioethanol from Napier grass via simultaneous saccharification and co-fermentation in a modified bioreactor. *Journal of Bioscience and Bioengineering*, 124, 184 – 188.

- Lo, Y.C., Lu, W.C., Chen, C.Y., and Chang, J.S. (2010). Dark fermentative hydrogen production from enzymatic hydrolysate of xylan and pretreated rice straw by *Clostridium butyricum* CGS5. *Bioresource Technology*, 101, 5885–5891.
- Loh, S.K., and Choo, Y.M. (2013). Prospect, challenges and opportunities on biofuels in Malaysia. In Pogaku, R., and Sarbatly, R. (Eds.), *Advances in Biofuels*. Boston: Springer.
- Loow, Y.L., Wu, T.Y., Md Jahim, J., Mohammad, A.W., and Teoh, W.H. (2016). Typical conversion of lignocellulosic biomass into reducing sugars using dilute acid hydrolysis and alkaline pretreatment. *Cellulose*, 23, 1491–1520.
- MAA (2017). Summary of new passenger & commercial vehicles registered in Malaysia for the year 1980 to 2017. Malaysian Automotive Association.
- Longoni, P., Rodolfi, M., Pantaleoni, L., Doria, E., Concia, L., Picco, A.M., and Cella, R. (2012). Functional analysis of the degradation of cellulosic substrates by a *Chaetomium globosum* endophytic isolate. *Applied and Environmental Microbiology*, 2012, 3693–3705.
- Madadi, M., and Abbas, A. (2017). Lignin degradation by fungal pretreatment: A review. *Journal of Plant Pathology and Microbiology*, 8, 1-6.
- Madian, H.R., El-Gendy, N.S., Farahat, L.A., Abo-State, M.A., and Ragab, A.M.E. (2012). Fungal hydrolysis and saccharification of rice straw and ethanol production. *Biosciences Biotechnology Research Asia*, 9(2).
- Madrera, R.R., and Valles, B.S. (2007). Determination of volatile compounds in cider spirits by gas chromatography with direct injection. *Journal of Chromatographic Science*, 45, 428-434.
- Manavalan, T., Manavalan, A., and Heese, K. (2015). Characterization of lignocellulolytic enzymes from white-rot fungi. *Current Microbiology*, 70, 485–498.
- Manickam, N.K., Rajarathinam, R., Muthuvelu, K.S., and Senniyappan, T. (2018). New insight into the effect of fungal mycelia present in the bio-pretreated paddy straw on their enzymatic saccharification and optimization of process parameters. *Bioresource Technology*, 267, 291–302.
- Mauch, F., Mauch-Mani, B., and Boller, T. (1988). Antifungal hydrolases in pea tissue and inhibition of fungal growth by combinations of chitinase and β -1,3-glucanase. *Plant Physiology*, 88, 936-942.
- Maurya, D.P., Singla, A., and Negi, S. (2015). An overview of key pretreatment processes for biological conversion of lignocellulosic biomass to bioethanol. *3 Biotech*, 5, 597–609.

- Mayer-Laigle, C., Rajaonarivony, R.K., Blanc, N., and Rouau, X. (2018). Comminution of dry lignocellulosic biomass: Part II. Technologies, improvement of milling performances, and security issues. *Bioengineering*, 5, 1-17.
- Meehnian, H., Jana, A.K., and Jana, M.M. (2016). Effect of particle size, moisture content, and supplements on selective pretreatment of cotton stalks by *Daedalea flavida* and enzymatic saccharification. *3 Biotech*, 6, 1-13.
- Menon, V., and Rao, M. (2012). Trends in bioconversion of lignocellulose: Biofuels, platform chemicals and biorefinery concept. *Progress in Energy and Combustion Science*, 38, 522-550.
- Miller, G.L. (1959). Use of dinitrosalicylic acid reagent for determination of reducing sugar. *Analytical Chemistry*, 31, 426-428.
- Minitab (2017): Getting started with Minitab17
- Miricioiu, M.G., Niculescu, V., Dinca, O.R., Mitu, F., and Craciun, M.E. (2016). Analytical errors in routine gas chromatography analysis. *Revista de Chimie*, 67, 396-400.
- Mohamad Remli, N.A., Md Shah, U.K., Mohamad, R., and Abd-Aziz, S. (2014). Effects of chemical and thermal pretreatments on the enzymatic saccharification of rice straw for sugars production. *BioResources*, 9, 510-522.
- Mohammad, N., Alam, M.Z., Kabashi, N.A., and Adebayo, O.S. (2011). Development of compatible fungal mixed culture for composting process of oil palm industrial waste. *African Journal of Biotechnology*, 10, 18657-18665.
- Mohd Azhar, S.H., Abdulla, R., Jambo, S.A., Marbawi, H., Gansau, J.A., Mohd Faik, A.A., and Rodrigues, K.F. (2017). Yeasts in sustainable bioethanol production: A review. *Biochemistry and Biophysics Reports*, 10, 52-61.
- Momayez, F., Karimi, K., Karimi, S., and Horváth, I.S. (2017). Efficient hydrolysis and ethanol production from rice straw by pretreatment with organic acids and effluent of biogas plant. *RSC Advances*, 7, 50537-50545.
- Mood, S.H., Golfeshan, A.H., Tabatabaei, M., Jouzani, G.S., Najafi, G.H., Gholami, M., and Ardjmand, M. (2013). Lignocellulosic biomass to bioethanol, a comprehensive review with a focus on pretreatment. *Renewable and Sustainable Energy Reviews*, 27, 77-93.
- Mosier, N., Wyman, C., Dale, B., Elander, R., Lee, Y.Y., Holtzapple, M., and Ladisch, M. (2005). Features of promising technologies for pretreatment of lignocellulosic biomass. *Bioresource Technology*, 96, 673-686.

- Mrudula, S., and Murugammal, R. (2011). Production of cellulase by *Aspergillus niger* under submerged and solid state fermentation using coir waste as a substrate. *Brazilian Journal of Microbiology*, 42, 1119-1127.
- Muhaji, and Sutjahjo, D.H. (2018). The characteristics of bioethanol fuel made of vegetable raw materials. *IOP Conference Series: Materials Science and Engineering*, 296, 1-6.
- Mukhlis, Mohd Saud, H., Sariah, M., Razi Ismail, M., Habib, S.H. and Kausar, H. (2013). Potential lignocellulolytic *Trichoderma* for bioconversion of oil palm empty fruit bunches. *Australian Journal of Crop Science*, 7, 425-431.
- Murad, H.A., and Azzaz, H.E.H. (2013). Cellulase production from rice straw by *Aspergillus flavus* NRRL 5521. *Science International*, 1, 103-107.
- Mustafa, A.M., Poulsen, T.G., Xia, Y., and Sheng, K. (2017). Combinations of fungal and milling pretreatments for enhancing rice straw biogas production during solid-state anaerobic digestion. *Bioresource Technology*, 224, 174–182.
- Mtui, G.Y.S. (2012). Lignocellulolytic enzymes from tropical fungi: Types, substrates and applications. *Scientific Research and Essays*, 7, 1544-1555.
- Mutreja, R., Das, D., Goyal, D., and Goyal, A. (2011). Bioconversion of agricultural waste to ethanol by SSF using recombinant cellulase from *Clostridium thermocellum*. *Enzyme Research*, 2011, 1-6.
- Mutturi, S., and Liden, G. (2013). Effect of temperature on simultaneous saccharification and fermentation of pretreated spruce and arundo. *Industrial and Engineering Chemistry Research*, 52, 1244–1251.
- Nadeem, M., Aftab, M.U., Irfan, M., Mushtaq, M., Qadir, A., and Syed, Q. (2015). Production of ethanol from alkali-pretreated sugarcane bagasse under the influence of different process parameters. *Frontiers in Life Science*, 8, 358-362.
- Nagarajan, P., Karunanithi, T., and Deng, Y. (2015). Direct bioconversion of cellulose into ethanol by mixed culture –batch kinetic and modeling. *International Journal of Innovative Research in Science, Engineering and Technology*, 4, 1870-1885.
- Naik, S.N., Goud, V.V., Rout, P.K., and Dalai, A.K. (2010). Production of first and second generation biofuels: A comprehensive review. *Renewable and Sustainable Energy Reviews*, 14, 578–597.
- Nair, R.B., Kabir, M.M., Lennartsson, P.R., Taherzadeh, M.J., and Horváth, I.S. (2017). Integrated process for ethanol, biogas, and edible filamentous

fungi-based animal feed production from dilute phosphoric acid-pretreated wheat straw. *Applied Biochemistry and Biotechnology*, 1-15.

- Nasr, A.M., Badawi, Mona, H., Demerdash, M.A., and Barakat, O.S. (2015). Bioconversion of rice straw into ethanol: fungi and yeasts are the backbone microbiota of the process. *International Journal of Current Microbiology and Applied Sciences*, 4, 382-401.
- Nichols, N.N., Hector, R.E., Saha, B.C., Frazer, S.E., and Kennedy, G.J. (2014). Biological abatement of inhibitors in rice hull hydrolyzate and fermentation to ethanol using conventional and engineered microbes. *Biomass and Bioenergy*, 67, 79-88.
- Nikolic, S., Pejin, J., and Mojovic, L. (2016). Challenges in bioethanol production: utilization of cotton fabrics as a feedstock. *Chemical Industry and Chemical Engineering Quarterly*, 22, 375-390.
- Nizami, A.S., Korres, N.E., and Murphy, J.D. (2009). Review of the integrated process for the production of grass biomethane. *Environmental Science and Technology*, 43, 8496-8508.
- Nlewem, K.C., and Trash, M.E. (2010). Comparison of different pretreatment methods based on residual lignin effect on the enzymatic hydrolysis of switchgrass. *Bioresource Technology*, 101, 5426-5430.
- Norfarina, M.N., Mohd Shamzi, M., Teck, C.L., Hooi, L.F., Raha, A.R., Joo, S.T., and Rosfarizan, M. (2017). Comparative analyses on medium optimization using one-factor-at-a-time, response surface methodology, and artificial neural network for lysine-methionine biosynthesis by *Pediococcus pentosaceus* RF-1. *Biotechnology and Biotechnological Equipment*, 31, 935-947.
- Nur Ain Izzati, M. Z., Nur Adni, M. and Mohd Razik, M. (2017). Isolation and characterisation of ascomycetes isolated from *Eurycoma longifolia* jack and malay traditional vegetables. *Pertanika Journal of Tropical Agricultural Science*, 40, 533 – 542.
- Nur Ain Izzati, M.Z., Madihah, M.Z.A., Nor Azizah, K., Najihah, A. and Muskhazli, M. (2019). First report of *Bipolaris cactivora* causing brown leaf spot in rice in Malaysia. *Plant Disease*. doi: 10.1094/PDIS-08-18-1384-PDN.
- Nur Baiti, A.M., Nor Azizah K. and Nur Ain Izzati, M. Z., (2016). Identification and diversity of *Fusarium* species isolated from tomato fruits. *Journal of Plant Protection Research*, 56, 211-216.
- Oberoi, H.S., Vadlani, P.V., Saida, L., Bansal, S., and Hughes, J.D. (2011). Ethanol production from banana peels using statistically optimized simultaneous saccharification and fermentation process. *Waste Management*, 31, 1576-1584.

- Okamoto, K., Nitta, Y., Maekawa, N. and Yanase, H. (2011). Direct ethanol production from starch, wheat bran and rice straw by the white rot fungus *Trametes hirsute*. *Enzyme and Microbial Technology*, 48, 273–277.
- Oladosu, Y., Rafii, M.Y., Abdullah, N., Magaji, U., Hussin, G., Ramli, A., and Miah, G. (2016). Fermentation quality and additives: A case of rice straw silage. *BioMed Research International*, 2016, 1-14.
- Olofsson, K., Bertilsson, M., and Lidén, G. (2008). A short review on SSF – an interesting process option for ethanol production from lignocellulosic feedstocks. *Biotechnology for Biofuels*, 1, 1-14.
- Olson, D.G., McBride, J.E., Shaw, A.J., and Lynd, L.R. (2012). Recent progress in consolidated bioprocessing. *Current Opinion in Biotechnology*, 23, 396-405.
- Ong, L.G.A., APCBEES, Chan, C.H., and Chew, A.L. (2012). Enzymatic hydrolysis of rice straw: Process optimization. *Journal of Medical and Bioengineering (JOMB)*, 1, 14-16.
- Onuki, S., Koziel, J.A., Leeuwen, J.V., Jenks, W.S., Grewell, D.A. and Cai, L. (2008). Ethanol production, purification, and analysis techniques: A review. *ASABE Annual International Meeting*, Paper No. 085136.
- Panasenko, V.T. (1967). Ecology of microfungi. *The Botanical Review*, 33, 189–215.
- Paola, D.C., Isabella, D.B., and Patrizia, R. (2011). Latest frontiers in the biotechnologies for ethanol production from lignocellulosic biomass. In Bernardes, M.A.D.S. (Ed.), *Biofuel Production-Recent Developments and Prospects*. United Kingdom: INTECH Open Access Publisher.
- Parisutham, V., Kim, T.H., and Lee, S.K. (2014). Feasibilities of consolidated bioprocessing microbes: From pretreatment to biofuel production. *Bioresource Technology*, 161, 431–440
- Park, J.Y., Shiroma, R., Al-Haq, M.I., Zhang, Y., Ike, M., Arai-Sanoh, Y., Ida, A., Kondo, M., and Tokuyasu, K. (2010). A novel lime pretreatment for subsequent bioethanol production from rice straw – Calcium capturing by carbonation (CaCCO) process. *Bioresource Technology*, 101, 6805–6811.
- Paschos, P., Xiros, C., and Christakopoulos, P. (2015). Ethanol effect on metabolic activity of the ethalogenic fungus *Fusarium oxysporum*. *BMC Biotechnology*, 15, 15.
- Passos, D.F., Pereira, N., and de Castro, A.M. (2018). A comparative review of recent advances in cellulases production by *Aspergillus*, *Penicillium* and *Trichoderma* strains and their use for lignocellulose deconstruction. *Current Opinion in Green and Sustainable Chemistry*, 14, 60-66.

- Paulova, L., Patakova, P., Branska, B., Rychtera, M., and Melzoch, K. (2015). Lignocellulosic ethanol: Technology design and its impact on process efficiency. *Biotechnology Advances*, 33, 1091–1107.
- Pellera, F.M., and Gidarakos, E. (2018). Chemical pretreatment of lignocellulosic agroindustrial waste for methane production. *Waste Management*, 71, 689–703.
- Pereira, S.C., Maehara, L., Machado, C.M.M., and Farinas, C.S (2015). 2G ethanol from the whole sugarcane lignocellulosic biomass. *Biotechnology Biofuels*, 8, 1-16.
- Piarpuzán, D., Quintero, J.A., and Cardona, C.A. (2011). Empty fruit bunches from oil palm as a potential raw material for fuel ethanol production. *Biomass and Bioenergy*, 35, 1130-1137.
- Pitt, J.I., and Hocking, A.D. (2009). *Aspergillus* and related teleomorphs. In Pitt, J.I., and Hocking, A.D. (Eds.), *Fungi and Food Spoilage* (pp. 275-337). Boston: Springer.
- Pirzadah, T., Garg, S., Singh, J., Vyas, A., Kumar, M., Gaur, N., Bala, M., Rehman, R., Varma, A., Kumar, V., and Kumar, M. (2014). Characterization of Actinomycetes and *Trichoderma* spp. for cellulase production utilizing crude substrates by response surface methodology. *SpringerPlus* 2014, 3, 1-12.
- Pradeep, P., Reddy, O.V.S., Mohan, P.R., and Ko, S. (2012). Process optimization for ethanol production from very high gravity (VHG) finger millet medium using response surface methodology. *Iranian Journal of Biotechnology*, 10, 168-174.
- Puri, M., Abraham, R.E., and Barrow, C.J. (2012). Biofuel production: Prospects, challenges and feedstock in Australia. *Renewable and Sustainable Energy Reviews*, 16, 6022–6031.
- Rabemanantsoa, H., and Saka, S. (2016). Various pretreatments of lignocellulosics. *Bioresource Technology*, 199, 83–91.
- Rahman, M.M., El-Aty, A.M.A., Choi, J.H., Shin, H.C., Shin, S.C., and Shim, J.H. (2015). Basic Overview on Gas Chromatography Columns. In Anderson, J.L., Berthod, A., Estévez, V.P., and Stalcup, A.M. (Eds.), *Analytical Separation Science, First Edition* (pp. 823-834). New York: Wiley-VCH Verlag GmbH & Co. KGaA.
- Rahnama, N., Foo, H.L., Abdul Rahman, N.A., Ariff, A., and Md Shah, U.K. (2014). Saccharification of rice straw by cellulase from a local *Trichoderma harzianum* SNRS3 for biobutanol production. *BMC Biotechnology*, 14, 103.

- Raissi, S., and Farsani, R.E. (2009). Statistical process optimization through multi-response surface methodology. *World Academy of Science, Engineering and Technology*, 51, 267-271.
- Rajamoorthy, Y., Abdul Rahim, K., and Munusamy, S. (2015). Rice industry in Malaysia: Challenges, policies and implications. *Procedia Economics and Finance*, 31, 861-867.
- Ram, L., Kaur, K. and Sharma, S. (2014). Screening isolation and characterization of cellulase producing micro-organisms from soil. *International Journal of Pharmaceutical Science Invention*, 3, 12 – 18.
- Rani, V., Mohanram, S., Tiwari, R., Nain, L. and Arora, A. (2014). Beta-Glucosidase: Key enzyme in determining efficiency of cellulase and biomass hydrolysis. *Journal of Bioprocessing and Biotechniques*, 5, 197 – 205.
- Rawat, R., Kumbhar, B.K., and Tewari, L. (2013). Optimization of alkali pretreatment for bioconversion of poplar (*Populus deltoides*) biomass into fermentable sugars using response surface methodology. *Industrial Crops and Products*, 44, 220– 226.
- Ray, R.C., and Behera, S.S. (2017). Solid state fermentation for production of microbial cellulases. In Brahmachari, G., Demain, A.L., and Adrio, J.L. (Eds.), *Biotechnology of Microbial Enzymes: Production, Biocatalysis, and Industrial Applications* (pp. 43-69). Amsterdam: Elsevier.
- Ren, N.Q., Zhao, L., Chen, C., Guo, W.Q., and Cao, G.L. (2016). A review on bioconversion of lignocellulosic biomass to H₂: Key challenges and new insights. *Bioresource Technology*, 215, 92–99.
- Resch, M.G., Donohoe, B.S., Baker, J.O., Decker, S.R., Bayer, E.A., Beckham, G.T, and Himmel, M.E. (2013). Fungal cellulases and complexed cellulosomal enzymes exhibit synergistic mechanisms in cellulose deconstruction. *Energy and Environmental Science*, 6, 1858-1867.
- Rheem, S., Rheem, I., and Oh, S. (2017). Response surface methodology using a fullest balanced model: A re-analysis of a dataset in the Korean journal for food science of animal resources. *Korean Journal for Food Science of Animal Resources*, 37, 139-146.
- Robinson, M.T., and Oen, O.S. (1963). The channeling of energetic atoms in crystal lattice. *Applied Physic Letter*, 2, 30-32.
- Rocha, G.J.M., Nascimento, V.M., da Silva, V.F.N., Corso, D.L.S., and Goncalves, A.R. (2014). Contributing to the environmental sustainability of the second generation ethanol production: Delignification of sugarcane bagasse with sodium hydroxide recycling. *Industrial Crops and Products*, 59, 63–68.

- Rosmiza, M.Z., Davies, W.P., Rosniza Aznie, C.R., Mazdi, M. and Jabil, M.J. (2014). Farmers' knowledge on potential uses of rice straw: An assessment in MADA and Sekinchan, Malaysia. *Malaysian Journal of Society and Space*, 10, 30 – 43.
- Roy, P., and Kaur, M. (2015). Status and problems of paddy straw management in west Bengal. *International Journal of Advances in Agricultural & Environmental Engineering*, 2, 44-48.
- Ruriani, E., Sunarti, T.C., and Meryandini, A. (2012). Yeast isolation for bioethanol. *Hayati Journal of Bioscience*, 19, 145-149.
- Saeed, M.A., Ma, H., Yue, S., Wang, Q., and Tu, M. (2018). Concise review on ethanol production from food waste: Development and sustainability. *Environmental Science and Pollution Research*, 25, 28851–28863.
- Saha, B.C, Nichols, N.N., Qureshi, N., Kennedy, G.J., Iten, L.B., and Cotta, M.A. (2015). Pilot scale conversion of wheat straw to ethanol via simultaneous saccharification and fermentation. *Bioresource Technology*, 175, 17–22.
- Sahni, N., and Phutela, U.G. (2013). Comparative profile of paddy straw pretreated with standard and isolated lignocellulolytic fungal cultures. *Journal of Yeast and Fungal Research*, 4, 92-97.
- Sakimoto, K., Kanna, M., and Matsumura, Y. (2017). Kinetic model of cellulose degradation using simultaneous saccharification and fermentation. *Biomass and Bioenergy*, 99, 116-121.
- Sakthi, S.S, Saranraj, P., and Rajasekar, M. (2011). Optimization for cellulase production by *Aspergillus niger* using paddy straw as substrate. *International Journal of Advanced Scientific and Technical Research*, 1, 70-85.
- Santos, V.T.O., Siqueira, G., Milagres, A.M.F., and Ferraz, A. (2018). Role of hemicellulose removal during dilute acid pretreatment on the cellulose accessibility and enzymatic hydrolysis of compositionally diverse sugarcane hybrids. *Industrial Crops and Products*, 111, 722–730.
- Sarabana, S.S.H., El-Gabry, K.I.M., and Eldin, A.M. (2015). Optimizing growth conditions provoked ethanol production by fungi grown on glucose. *Middle East Journal of Applied Sciences*, 5, 1222-1231.
- Saratale, G.D., and Oh, M.K. (2015). Improving alkaline pretreatment method for preparation of whole rice waste biomass feedstock and bioethanol production. *RSC Advances*, 5, 97171–97179.
- Sari, F.P., and Budiyo. (2014). Enhanced biogas production from rice straw with various pretreatment : A review. *Waste Technology*, 2, 17-25.

- Saritha, M., Arora, A., and Lata. (2012a). Biological pretreatment of lignocellulosic substrates for enhanced delignification and enzymatic digestibility. *Indian Journal of Microbiology*, 52, 122-130.
- Saritha, M., Arora, A., and Nain, L. (2012b). Pretreatment of paddy straw with *Trametes hirsuta* for improved enzymatic saccharification. *Bioresource Technology*, 104, 459–465.
- Saritha, M., Arora, A., Singh, S., and Nain, L. (2013). *Streptomyces griseorubens* mediated delignification of paddy straw for improved enzymatic saccharification yields. *Bioresource Technology*, 135, 12-17.
- Sarkar, N.S., Ghosh, S.K., Bannerjee, S., and Aikat, K. (2012). Bioethanol production from agricultural wastes: An overview. *Renewable Energy*, 37, 19-27.
- SAS (1999). Introduction to Response Surface Experiments. Institute Inc., SAS Campus Drive, Cary, North Carolina.
- Sawisit, A., Jampatesh, S., Jantama, S.S., and Jantama, K. (2018). Optimization of sodium hydroxide pretreatment and enzyme loading for efficient hydrolysis of rice straw to improve succinate production by metabolically engineered *Escherichia coli* KJ122 under simultaneous saccharification and fermentation. *Bioresource Technology*, 260, 348–356.
- Schuster, B.G., and Chinn, M.S. (2013). Consolidated bioprocessing of lignocellulosic feedstocks for ethanol fuel production. *Bioenergy Research*, 6, 416–435.
- Segato, F., Damásio, A.R.L., Lucas, R.C., Squina, F.M., and Prade, R.A. (2014). Genomics review of holocellulose deconstruction by *Aspergilli*. *Microbiology and Molecular Biology Reviews*, 78, 588–613.
- Sendich, E., Laser, M., Kim, S., Alizadeh, H., Laureano-Perez, L., Dale, B., and Lynd, L., (2008). Recent process improvements for the ammonia fiber expansion (AFEX) process and resulting reductions in minimum ethanol selling price. *Bioresource Technology*, 99, 8429–8435.
- Seo, D.J., and Sakoda, A. (2014). Assessment of the structural factors controlling the enzymatic saccharification of rice straw cellulose. *Biomass and Bioenergy*, 71, 47-57.
- Shah, S.R., Ishmael, U.C., Palliah, J.V., Asras, M.F.F., and Nik Wan Ahmad, S.S. (2016). Optimization of the enzymatic saccharification process of empty fruit bunch pretreated with laccase enzyme. *Bioresources*, 11, 5138-5154.
- Sharifah Siti Maryam, S.A.R., Nor Azwady, A.A. and Nur Ain Izzati, M.Z. (2016). Isolation and diversity of *Trichoderma* species from cultivated crop soil in Malaysia. *Acta Biologica Malaysiana*, 5, 43-48.

- Sharma, N., Kalra, K.L., Oberoi, H.S., and Bansal, S. (2007). Optimization of fermentation parameters for production of ethanol from kinnow waste and banana peels by simultaneous saccharification and fermentation. *Indian Journal of Microbiology*, 47, 310–316.
- Sharma, R.K., and Arora, D.S. (2011). Solid state degradation of paddy straw by *Phlebia floridensis* in the presence of different supplements for improving its nutritive status. *International Biodeterioration and Biodegradation*, 65, 990-996.
- Sharma, N.K., Behera, S., Arora, R., Kumar, S., and Sani, R.K. (2018). Xylose transport in yeast for lignocellulosic ethanol production: Current status. *Journal of Bioscience and Bioengineering*, 125, 259-267.
- Shaw, R., Festing, M.F.W., Peers, I., and Larry, F. (2002). Use of factorial designs to optimize animal experiments and reduce animal use. *ILAR Journal*, 43, 223–232.
- Shawky, B.T., Mahmoud, M.G., Ghazy, E.A., Asker, M.M.S., and Ibrahim, G.S. (2011). Enzymatic hydrolysis of rice straw and corn stalks for monosugars production. *Journal of Genetic Engineering and Biotechnology*, 9, 59–63.
- Shen, D., Xiao, R., Gu, S., and Zhang, H. (2013). The overview of thermal decomposition of cellulose in lignocellulosic biomass. In Kadla, J., and Van de Ven, T.G.M. (Eds.), *Cellulose Biomass Conversion*. United Kingdom: INTECH Open Access Publisher.
- Sheng, P., Huang, J., Zhang, Z., Wang, D., Tian, X. and Ding, J. (2016). Construction and characterization of a cellulolytic consortium enriched from the hindgut of *Holotrichia parallela* larvae. *International Journal of Molecular Sciences*, 17, 1646 – 1658.
- Shinozaki, Y., and Kitamoto, H.K. (2011). Ethanol production from ensiled rice straw and whole-crop silage by the simultaneous enzymatic saccharification and fermentation process. *Journal of Bioscience and Bioengineering*, 111, 320-325.
- Shong, J., Diaz, M.R.J., and Collins, C.H. (2012). Towards synthetic microbial consortia for bioprocessing. *Current Opinion in Biotechnology*, 23, 798-802.
- Silva, R.N. (2016). *Mycology: Current and future developments: Fungal biotechnology for biofuel production* (pp. 343-344). United Arab of Emirates: Bentham Science Publisher.
- Sims, R.E., Mabee, W., Saddler, J.N., and Taylor, M. (2010). An overview of second generation biofuel technologies. *Bioresource Technology*, 101, 1570-1580.

- Sindhu, R., Binod, P., and Pandey, A. (2016). Biological pretreatment of lignocellulosic biomass – An overview. *Bioresource Technology*, 199, 76–82.
- Singh, A., and Bishnoi, N.R. (2012). Optimization of enzymatic hydrolysis of pretreated rice straw and ethanol production. *Applied Microbiology Biotechnology*, 93, 1785–1793.
- Singh, D.P., and Trivedi, R.K. (2013). Acid and alkaline pretreatment of lignocellulosic biomass to produce ethanol as biofuel. *International Journal of ChemTech Research*, 5, 727-734.
- Singh, R., Tiwari, S., Srivastava, M., and Shukla, A. (2014). Microwave assisted alkali pretreatment of rice straw for enhancing enzymatic digestibility. *Journal of Energy*, 2014, 1-7.
- Singh, N., Mathur, A.S., Tuli, D.K., Gupta, R.P., Barrow, C.J., and Puri, M. (2017). Cellulosic ethanol production via consolidated bioprocessing by a novel thermophilic anaerobic bacterium isolated from a Himalayan hot spring. *Biotechnology and Biofuels*, 10, 1-18.
- Singhania, R.R., Patel, A.K., Sukumaran, R.K., Larroche, C., and Pandey, A. (2013). Role and significance of beta-glucosidases in the hydrolysis of cellulose for bioethanol production. *Bioresource Technology*, 127, 500-507.
- Skory, C.D., Freer, S.N., and Bothast, R.J. (1997). Screening for ethanol-producing filamentous fungi. *Biotechnology Letters*, 19, 203–206.
- Sohail, M., Siddiqi, R., Ahmad, A. and Khan, S.A. (2009). Cellulase production from *Aspergillus niger* MS82: Effect of temperature and pH. *New Biotechnology*, 25, 437-442.
- Soliman, S.A., El-Zahwary, Y.A., and El-Mougith, A.A. (2013). Fungal biodegradation of agro-industrial waste. In Kadla, J., and Van de Ven, T.G.M. (Eds.), *Cellulose Biomass Conversion*. United Kingdom: INTECH Open Access Publisher.
- Sorek, N., Yeats, T.H., Szemenyei, H., Youngs, H., and Somerville, C.R. (2014). The implications of lignocellulosic biomass chemical composition for the production of advanced biofuels. *BioScience*, 64, 192-201.
- Srivastava, N., Srivastava, M., Manikanta, A., Ramteke, P.W., Singh, R.L., Mishra, P.K., and Upadhyay, S.N. (2018). Fungal cellulases production for biodegradation of agriculture waste. In Panpatte, D., Jhala Y., Shelat H., and Vyas R. (Eds.), *Microorganisms for Green Revolution. Microorganisms for Sustainability, vol 7* (pp. 75-89). Singapore: Springer.
- Suhag, M. and Singh, J. (2014). Recent Advances in fermentation of lignocellulosic biomass hydrolysate to ethanol. *Journal of Advances in Science and Technology*, 7, 1-8.

- Sukumaran, R., Singhanian, R., and Pandey, A. (2005). Microbial cellulases production, applications and challenges. *Journal of Scientific and Industrial Research*, 64, 832-844.
- Sukumaran, R.K., Surender, V.J., Sindhu, R., Binod, P., Janu, K.U., Sajna, K.V., Rajasree, K.P., and Pandey, A. (2010). Lignocellulosic ethanol in India: Prospects, challenges and feedstock availability. *Bioresource Technology*, 101, 4826–4833.
- Sun, Y., and Cheng, J. (2002). Hydrolysis of lignocellulosic materials for ethanol production: A review. *Bioresource Technology*, 83, 1–11.
- Suresh, K., Ranjan, A., Singh, S., and Moholkar, V.S. (2014). Mechanistic investigations in sono-hybrid techniques for rice straw pretreatment. *Ultrasonics Sonochemistry*, 21, 200–207.
- Suriyachai, N., Weerasaia, K., Laosiripojana, N., Champreda, V., and Unrean, P. (2013). Optimized simultaneous saccharification and co-fermentation of rice straw for ethanol production by *Saccharomyces cerevisiae* and *Scheffersomyces stipitis* co-culture using design of experiments. *Bioresource Technology*, 142, 171–178.
- Suwannarangsee, S., Bunternngsook, B., Arnthong, J., Paemane, A., Thamchaipenet, A., Eurwilaichitr, L., Laosiripojana, N., and Champreda, V. (2012). Optimisation of synergistic biomass-degrading enzyme systems for efficient rice straw hydrolysis using an experimental mixture design. *Bioresource Technology*, 119, 252–261.
- Taha, M., Shahsavari, E., Al-Hothaly, K., Mouradov, A., Smith, A.T., Ball, A.S. and Adetutu, E.M. (2015). Enhanced biological straw saccharification through coculturing of lignocellulose-degrading microorganisms. *Applied Biochemistry and Biotechnology*, 175, 3709–3728.
- Taher, I.B., Bennour, H., Fickers, P., and Hassouna, M. (2017). Valorization of potato peels residues on cellulase production using a mixed culture of *Aspergillus niger* ATCC 16404 and *Trichoderma reesei* DSMZ 970. *Waste and Biomass Valorization*, 8, 183-192.
- Takano, M., and Hoshino, K. (2012). Direct ethanol production from rice straw by coculture with two high-performing fungi. *Frontiers of Chemical Science and Engineering*, 6, 139–145.
- Takano, M., and Hoshino, K. (2018). Bioethanol production from rice straw by simultaneous saccharification and fermentation with statistical optimized cellulase cocktail and fermenting fungus. *Bioresources and Bioprocessing*, 5, 1-12.
- Tayyab, M., Noman, A., Islam, W., Waheed, S., Arafat, Y., Ali, F., Zaynab, M., Lin, S., Zhang, H., and Lin, W. (2017). Bioethanol production from

lignocellulosic biomass by environment-friendly pretreatment methods: A review. *Applied Ecology and Environmental Research*, 16, 225-249.

- Teeravivattanakit, T., Baramée, S., Phitsuwan, P., Sornyotha, S., Waeonukul, R., Pason, P., Tachaapaikoon, C., Poomputsa, K., Kosugi, A., Sakka, K., and Ratanakhanokchai, K. (2017). Chemical pretreatment-independent saccharifications of xylan and cellulose of rice straw by bacterial weak lignin-binding xylanolytic and cellulolytic enzymes. *Applied and Environmental Microbiology*, 83, 1-15.
- Teghammar, A., Karimi, K., Horváth, I.S., and Taherzadeh, M.J. (2012). Enhanced biogas production from rice straw, triticale straw and softwood spruce by NMMO pretreatment. *Biomass and Bioenergy*, 36, 116-120.
- Tesfaw, A., and Assefa, F. (2014). Co-culture: A great promising method in single cell protein production. *Biotechnology and Molecular Biology Reviews*, 9, 12-20.
- Teter, S.A., Sutton, K.B., and Emme, B. (2014). Enzymatic processes and enzyme development in biorefining. In Waldron, K. (Ed.), *Advances in Biorefineries: Biomass and water supply chain exploitation* (pp. 199-226). United Kingdom: Woodhead Publishing.
- Thomas, L., Larroche, C., and Pandey, A. (2013). Current developments in solid-state fermentation. *Biochemical Engineering Journal*, 81, 146–161.
- Thontowi, A., Perwitasari, U., Kholida, L.N., Fahrurrozi, Yopi, and Prasetya, B. (2017). Optimization of simultaneous saccharification and fermentation in bioethanol production from sugarcane bagasse hydrolyse by *Saccharomyces cerevisiae* BTCC 3 using response surface methodology. *IOP Conference Series: Earth and Environmental Science*, 183, 1-9.
- Thormann, M.N., Currah, R.S. and Bayley, S.E. (2002). The relative ability of fungi from *Sphagnum fuscum* to decompose selected carbon substrates. *Canadian Journal of Microbiology*, 48, 204–211.
- Tiwari, R., Rana, S., Singh, S., Arora, A., Kaushik, R., Agrawal, V.V., Saxena, A.K., and Nain, L. (2013). Biological delignification of paddy straw and *Parthenium sp.* using a novel micromycete *Myrothecium roridum* LG7 for enhanced saccharification. *Bioresource Technology*, 135, 7–11.
- Tye, Y.Y., Lee, K.T., Wan Abdullah, W.N., and Leh, C.P. (2011). Second-generation bioethanol as a sustainable energy source in Malaysia transportation sector: status, potential and future prospects. *Renewable and Sustainable Energy Review*, 15, 4521-4536.
- Umikalsom, M.S., Ariff, A.B., Shamsuddin, Z.H., Tong, C.C., Hassan, M.A., and Karim, M.I.A. (1997). Production of cellulase by a wild strain *Chaetomium globosum* using delignified oil palm empty-fruit-bunch fibre as substrate. *Applied Microbiology and Biotechnology*, 47, 590–595.

- Vanitjinda, G., Nimchua, T., and Sukyai, P. (2019). Effect of xylanase-assisted pretreatment on the properties of cellulose and regenerated cellulose films from sugarcane bagasse. *International Journal of Biological Macromolecules*, 122, 503–516.
- van Zyl, W.H., den Haan, R., and la Grange, D.C. (2011). Developing organisms for consolidated bioprocessing of biomass to ethanol. In Bernardes, M.A.D.S. (Ed.), *Biofuel Production-Recent Developments and Prospects*. United Kingdom: INTECH Open Access Publisher.
- van Zyl, W.H., den Haan, R., and la Grange, D.C. (2013). Developing cellulolytic organisms for consolidated bioprocessing of lignocellulosics. In Gupta V., and Tuohy M. (Eds.), *Biofuel Technologies* (pp. 189-220). Berlin, Heidelberg: Springer.
- Visser, E.M., Leal, T.F., de Almeida, M.N., and Guimarães, V.M. (2015). Increased enzymatic hydrolysis of sugarcane bagasse from enzyme recycling. *Biotechnology for Biofuels*, 8, 1-9.
- Waghmare, P.R., Kadam, A.A., Saratale, G.D., and Govindwar, S.P. (2014). Enzymatic hydrolysis and characterization of waste lignocellulosic biomass produced after dye bioremediation under solid state fermentation. *Bioresource Technology*, 168, 136–141.
- Wagner, A.O., Schwarzenauer, T., and Illmer, P. (2013). Improvement of methane generation capacity by aerobic pre-treatment of organic waste with a cellulolytic *Trichoderma viride* culture. *Journal of Environmental Management*, 129, 357–360.
- Wahid, Z., and Nadir, N. (2013). Improvement of one factor at a time through design of experiments. *World Applied Sciences Journal 21 (Mathematical Applications in Engineering)*, 56-61.
- Walker, G.M., and Stewart, G.G (2016): *Saccharomyces cerevisiae* in the production of fermented beverages. *Beverages*, 2, 30 – 42.
- Wan, C., and Li, Y. (2010a). Microbial delignification of corn stover by *Ceriporiopsis subvermispota* for improving cellulose digestibility. *Enzyme and Microbial Technology*, 47, 31–36.
- Wan, C., and Li, Y. (2010b). Microbial pretreatment of corn stover with *Ceriporiopsis subvermispota* for enzymatic hydrolysis and ethanol production. *Bioresource Technology*, 101, 6398–6403.
- Wan, C., and Li, Y. (2011). Effectiveness of microbial pretreatment by *Ceriporiopsis subvermispota* on different biomass feedstocks. *Bioresource Technology*, 102, 7507–7512.
- Wang, M., Li, Z., Fang, X., Wang, L. and Qu, Y. (2012). Cellulolytic Enzyme Production and Enzymatic Hydrolysis for Second-Generation Bioethanol Production. In Bai, F.W., Liu, C.G., Huang, H. and Tsao, G. (Eds.),

Biotechnology in China III: Biofuels and Bioenergy. Advances in Biochemical Engineering Biotechnology, vol 128 (pp. 1-24). Berlin, Heidelberg: Springer.

- Wang, Z., Cao, G., Zheng, J., Fu, D., Song, J., Zhang, J., Zhao, L., and Yang, Q. (2015a). Developing a mesophilic co-culture for direct conversion of cellulose to butanol in consolidated bioprocess. *Biotechnology for Biofuels*, 8, 1-9.
- Wang, K., Xie, X., Si, Z., Jiang, J., and Wang, J. (2015b). Microwave assisted hydrolysis of holocellulose catalyzed with sulfonated char derived from lignin-rich residue. *Advances in Materials Science and Engineering*, 2015, 1-6.
- Wanmolee, W., Sornlake, W., Rattanaphan, N., Suwannarangsee, S., Laosiripojana, N., and Champreda, V. (2016). Biochemical characterization and synergism of cellulolytic enzyme system from *Chaetomium globosum* on rice straw saccharification. *BMC Biotechnology*, 16, 1-12.
- Wati, L., Kumari, S., and Kundu, B.S. (2007). Paddy straw as substrate for ethanol production. *Indian Journal of Microbiology*, 47, 26-29.
- Wen, Z., Liao, W. and Chen, S. (2005). Production of cellulase/ β -glucosidase by the mixed fungi culture *Trichoderma reesei* and *Aspergillus phoenicis* on dairy manure. *Process Biochemistry*, 40, 3087 - 3094.
- Wen, F., Sun, J., and Zhao, H. (2010). Yeast surface display of trifunctional minicellulosomes for simultaneous saccharification and fermentation of cellulose to ethanol. *Applied and Environmental Microbiology*, 76, 1251–1260.
- Wi, S.G., Choi, I.S., Kim, K.H., Kim, H.M., and Bae, H.J. (2013). Bioethanol production from rice straw by popping pretreatment. *Biotechnology for Biofuels*, 6, 1-7.
- Wieczorek, A.S., Biot-Pelletier, D., and Martin, V.J.J. (2013): Recombinant Cellulase and Cellulosome Systems. In Kadla, J., and Van de Ven, T.G.M. (Ed.), *Cellulose Biomass Conversion*. United Kingdom: INTECH Open Access Publisher.
- Woiciechowski, A.L., Karp, S.G., Sobral, K., De Carvalho, J.C., Letti, L.A.J., Soccol, V.T., and Soccol, C.R. (2014). Pretreatment strategies to enhance value addition of agro-industrial wastes. In Brar, S., Dhillon, G., and Soccol, C. (Eds.), *Biotransformation of waste biomass into high value biochemical* (pp. 29-49). New York: Springer.
- Wongwilaiwalin, S., Rattanachomsri, U., Laothanachareon, T., Eurwilaichitr, L., Igarashi, Y., and Champreda, V. (2010). Analysis of a thermophilic lignocellulose degrading microbial consortium and multi-species

lignocellulolytic enzyme system. *Enzyme and Microbial Technology*, 47, 283–290.

Wyman, C.E., Cai, C.M., and Kumar, R. (2017). Bioethanol from lignocellulosic biomass. In Kaltschmitt M. (Ed.), *Energy from Organic Materials (Biomass). Encyclopedia of Sustainability Science and Technology Series* (pp. 997-1022). New York: Springer.

Xu, C., Qin, Y., Li, Y., Ji, Y., Huang, J., Song, H., and Xu, J. (2010). Factors influencing cellulose activity in consolidated bioprocessing of cellulosic ethanol. *Bioresource Technology*, 101, 9560–9569.

Yadav, I.C., and Devi, N.L. (2018). Biomass burning, regional air quality, and climate change. *Reference Module in Earth Systems and Environmental Sciences*, 2018, 1.

Yamada, R., Taniguchi, N., Tanaka, T., Ogino, C., Fukuda, H., and Kondo, A. (2011). Direct ethanol production from cellulosic materials using a diploid strain of *Saccharomyces cerevisiae* with optimized cellulase expression. *Biotechnology for Biofuels*, 4, 1-8.

Yang, L., Cao, J., Jin, Y., Chang, H.M., Jameel, H., Phillips, R., and Li, Z. (2012). Effects of sodium carbonate pretreatment on the chemical compositions and enzymatic saccharification of rice straw. *Bioresource Technology*, 124, 283–291.

Yao, W., and Nokes, S.E. (2013). Perspective: The use of co-culturing in solid substrate cultivation and possible solutions to scientific challenges. *Biofuels, Bioproducts and Biorefining*, 7, 361-372.

Ylittero, P., Franzén, C.J., and Taherzadeh, M.J. (2013). Impact of furfural on rapid ethanol production using a membrane bioreactor. *Energies*, 6, 1604-1617.

Yoswathana, N., Phuriphipat, P., Treyawutthiwat, P., and Eshtiaghi, M.N. (2010). Bioethanol production from rice straw. *Energy Research Journal*, 1, 26-31.

You, C., Zhang, X.Z., Sathitsuksanoh, N., Lynd, L.R., and Zhang, Y.H.P. (2012). Enhanced microbial utilization of recalcitrant cellulose by an ex vivo cellulosome-microbe complex. *Applied and Environmental Microbiology*, 1437–1444.

Yu, Q., Zhuang, X., Yuan, Z., Wang, Q., Qi, W., Wang, W., Zhang, Y., Xu, J. and Xu, H. (2010). Two-step liquid hot water pretreatment of *Eucalyptus grandis* to enhance sugar recovery and enzymatic digestibility of cellulose. *Bioresource Technology*, 101, 4895–4899.

Zambare, V.P., and Christopher, L.P. (2012). Optimization of enzymatic hydrolysis of corn stover for improved ethanol production. *Energy Exploration and Exploitation*, 30, 193-205.

- Zerva, A., Savvides, A.L., Katsifas, E.A., Karagouni, A.D., and Hatzinikolaou, D.G. (2014). Evaluation of *Paecilomyces variotii* potential in bioethanol production from lignocellulose through consolidated bioprocessing. *Bioresource Technology*, 162, 294–299.
- Zhang, Y.H.P., Himmel, M.E., and Mielenz, J.R. (2006). Outlook for cellulose improvement: Screening and selection strategies. *Biotechnology Advances*, 24, 452-481.
- Zhang, C. (2007). Chromatographic methods for environmental analysis. In Zhang, C. (Ed.), *Fundamentals of environmental sampling and analysis* (pp. 247-285). Hoboken, New Jersey: John Wiley & Sons, Inc.
- Zhang, Q., and Cai, W. (2008). Enzymatic hydrolysis of alkali-pretreated rice straw by *Trichoderma reesei* ZM4-F3. *Biomass and Bioenergy*, 32, 1130-1135.
- Zhang, M., Wang, F., Su, R., Qi, W., and He, Z. (2010). Ethanol production from high dry matter corncob using fed-batch simultaneous saccharification and fermentation after combined pretreatment. *Bioresource Technology*, 101, 4959–4964.
- Zhang, W., Lin, Y., Zhang, Q., Wang, X., Wu, D., and Kong, H. (2013). Optimisation of simultaneous saccharification and fermentation of wheat straw for ethanol production. *Fuel*, 112, 331–337.
- Zhao, L.C., He, Y., Deng, X., Yang, G.L., Li, W., Liang, J., and Tang, Q.L. (2012). Response surface modeling and optimization of accelerated solvent extraction of four lignans from *Fructus schisandrae*. *Molecules*, 17, 3618-3629.
- Zhao, X., Moates, G.K., Elliston, A., Wilson, D.R., Coleman, M.J., and Waldron, K.W. (2015). Simultaneous saccharification and fermentation of steam exploded duckweed: Improvement of the ethanol yield by increasing yeast titre. *Bioresource Technology*, 194, 263–269.
- Zheng, Y., Zhao, J., Xu, F., and Li, Y. (2014). Pretreatment of lignocellulosic biomass for enhanced biogas production. *Progress in Energy and Combustion Science*, 42, 35-53.
- Zhu, J.Y., and Pan, X.J. (2010). Woody biomass pretreatment for cellulosic ethanol production: Technology and energy consumption evaluation. *Bioresource Technology*, 101, 4992–5002.
- Zhu, S., Huang, W., Huang, W., Wang, K., Chen, Q., and Wu, Y. (2015). Pretreatment of rice straw for ethanol production by a two-step process using dilute sulfuric acid and sulfomethylation reagent. *Applied Energy*, 154, 190–196.

Zoglowek, M., Hansen, G.H., Lübeck, P.S., and Lübeck, M. (2016). Fungal consortia for conversion of lignocellulose into bioproducts. In Silva, R.N. (Ed.), *Mycology: Current and future developments (Vol. 1) Fungal biotechnology for biofuel productions* (pp. 329-365). Sharjah, United Arab of Emirates: Betham Science Publisher LTD.

Zuas, O., Mulyana, M.R., and Budiman, H. (2016). Analytical method validation of GC-FID for the simultaneous measurement of hydrocarbons (C2-C4) in their gas mixture. *Revista Colombiana de Química*, 45, 22-27.



BIODATA OF STUDENT

Mona Fatin Syazwane Binti Mohamed Ghazali was born on 27th July 1990 at Penang Adventist Hospital. She acquired her pre-university education from SK Dato Kramat, SMK (P) St. George and Penang Matriculation College. She pursued her studies at Universiti Putra Malaysia where she obtained her Bachelor of Science (Honours) - Biology in 2013 and then Master of Science – Conservation Biology from Universiti Kebangsaan Malaysia in 2015. Owing to a great passion in microbiology, she extended her keen interest in that area by continuing with a PhD research on developing bioethanol from paddy straw using lignocellulolytic fungi that is expected to be successfully completed in 2019.



LIST OF PUBLICATIONS

Mona Fatin Syazwane, M.G., Nurul Shaziera, A.G., Nur Ain Izzati, M.Z., Nor Azwady, A.A. and Muskhazli, M. (2018). Improvement of delignification, desilication and cellulosic content availability in paddy straw via physico-chemical pretreatments. *Annual Research & Review in Biology*, 26 (6), 1-11. doi: 10.9734/ARRB/2018/40947

Mona Fatin Syazwane, M.G., Nur Ain Izzati, M.Z., Nor Azwady, A.A. and Muskhazli, M. Screening of lignocellulolytic fungi for hydrolyzation of lignocellulosic materials in paddy straw for bioethanol production. *Malaysian Journal of Microbiology*, 15 (4), pp. xxx-xxx. doi: <http://dx.doi.org/10.21161/mjm.180250>



© COPYRIGHT



UNIVERSITI PUTRA MALAYSIA

STATUS CONFIRMATION FOR THESIS / PROJECT REPORT AND COPYRIGHT

ACADEMIC SESSION : SECOND SEMESTER 2018/2019

TITLE OF THESIS / PROJECT REPORT :

**CONVERSION OF PADDY STRAW TO BIOETHANOL THROUGH CONSOLIDATED
BIOPROCESSING USING LIGNOCELLULOLYTIC FUNGI**

NAME OF STUDENT : MONA FATIN SYAZWANEE BINTI MOHAMED GHAZALI

I acknowledge that the copyright and other intellectual property in the thesis/project report belonged to Universiti Putra Malaysia and I agree to allow this thesis/project report to be placed at the library under the following terms:

1. This thesis/project report is the property of Universiti Putra Malaysia.
2. The library of Universiti Putra Malaysia has the right to make copies for educational purposes only.
3. The library of Universiti Putra Malaysia is allowed to make copies of this thesis for academic exchange.

I declare that this thesis is classified as :

*Please tick (/)

CONFIDENTIAL

(Contain confidential information under Official Secret Act 1972).

RESTRICTED

(Contains restricted information as specified by the organization/institution where research was done).

OPEN ACCESS

I agree that my thesis/project report to be published as hard copy or online open access.

This thesis is submitted for :

PATENT

Embargo from _____ until _____
(date) (date)

Approved by:

(Signature of Student)
New IC No/ Passport No.:

(Signature of Chairman of Supervisory Committee)
Name:

Date :

Date :

[Note : If the thesis is CONFIDENTIAL or RESTRICTED, please attach with the letter from the organization/institution with period and reasons for confidentiality or restricted.]