Psychiatria Danubina, 2020; Vol. 32, No. 2, pp 311-312 © Medicinska naklada - Zagreb, Croatia Letter to Editor

MENTAL HEALTH CRISIS OF PAKISTANIS POPULATION DURING COVID-19 PANDEMIC

Rooh Ullah¹, Muhammad Suleman Rana¹, Mehmood Qadir¹, Muhammad Usman² & Niaz Ahmed³

¹Department of Microbiology, Quaid-i-Azam University, Islamabad, Pakistan ²Department of Bio Sciences National University of Science and Technology, Islamabad, Pakistan ³Department of Microbiology, Harbin Medical University, Harbin, China

* * * * *

The first country that was affected by pandemic of SARS-CoV-2 was China. Several exclusive features of SARS-CoV-2 pandemic and its managing policies globally provoked a sharp public mental health crisis. Pakistan is one of the endemic countries for various infectious diseases and has experienced many outbreaks and epidemics every year which effects it's economy and social life (WHO 2016). SARS-CoV-2 is considered as more contagious than MERS and SARS, with case fatality rate of (6.92%) (Surveillances 2020). The ambiguous incubation period and likely asymptomatic transmission of the virus have induced anxiety and fear in the public of Pakistan. Second, extraordinary largescale quarantine actions in all main cities, which basically restrict peoples to their families, caused negative psycho-social impact on citizens (Brooks et al. 2020). Third, reports of deficiencies personal protective equipment's, medical staff, hospital beds, isolation wards, medicines and sanitizers in Pakistan provoked panic in public. Last, a surplus of misinformation on NEWS and social media increased the fear in public (WHO 2019) and poses major risk to public psychological healthiness during this health crisis.

Pakistan already experienced the outbreaks and epidemics of many infectious diseases including Dengue, Measles, Chickenpox, HIV, TB and Typhoid. Keeping in mind that SARS-CoV-2 is spreading rapidly, infecting many countries of the world. General fear and fear-induced over-reactive conduct is more common among Pakistani community (Shultz et al. 2016). Furthermore, psychiatric conditions, anxiety, depression and post-traumatic stress disorder, have been developed in high-risk personals, especially front-line healthcare staffs (Mak et al. 2009). In this situation it is suggested that Pakistan must establish emergency mental health crisis interventions for decreasing the psycho-social effects of the SARS-CoV-2 pandemic and should be officially included in National Action Plan for Preparedness & Response to Covid-19 Pakistan (NIH 2020) to tackle mental health crisis of Pakistanis population during current pandemic. The emergency psychological crisis interventions staff includes psychological outreach teams led by mental health professionals, psychiatrists and psychological care hotline teams.

According to opinion, mental health crisis interventions contain three (03) important points: 1. understand psychological health position in various population group influenced by SARS-CoV-2 pandemic; 2. recognizing individuals who are at high risk of aggression and suicide, and 3. giving appropriate mental health interventions for those in need. Targeted individuals are divided in four (04) levels: Level (01) includes those population who are most susceptible to psychological health problems, for example hospitalized individuals with confirmed COVID-19, front-line health experts and managerial team. Level (02) includes isolated individuals with atypical infection sign and symptoms (such as individuals with suspected COVID-19 and close contacts). Level (03) includes those individuals who are in close contacts with level (01) and (02), i.e., friends, family, colleagues and rescuers who contributed in the SARS-CoV-2-pandemic response. Level (04) includes population who are affected by the pandemic preventive and control procedures (Li et al. 2020).

The release of emergency psychological crisis interventions plan acknowledges Pakistan's public psychological health needs during current pandemic. However, plan doesn't specify operationalization of in what way different groups would be assessed or screened to determine the level and type of interventions to give to each. This level of detail is required because Pakistan lacks well established psychological healthcare infrastructure and has no present designated personnel's and national level emergency response system to offer mental health crisis interventions during a national health emergency. Other key challenge in successfully implementing emergency mental health crisis interventions include shortage of mental healthcare providers in Pakistan, 342 psychiatrists (0.20 psychiatrists per 100,000 population) (Jooma et al. 2020).

Challenges reported in Pakistan show that telemedicine/telehealth should be considered to help eliminate barriers and retrieving quality-care for mental-health using smartphones. Task sharing or shifting (i.e., shifting service delivery from experts to general population with less qualifications or making new squad of providers with specific trainings) may help, especially in low resources areas (WHO 2007). Pakistan must also consider guidance and requesting support from global mental-healthcare infrastructures and research groups through international collaborations. Given lessons taken from past epidemics in Pakistan and globally, public psychological health interventions must be recruited in public health preparedness and emergency response plans to efficiently control all epidemics. The World Health Organization's (WHO) strategic preparedness and response plan for COVID-19, however, has not yet definite any approaches to target mental health needs (WHO 2019). As the virus rapidly spreads globally, government of Pakistan and international agencies must discourse public mental health requirements by developing and implementing well-coordinated strategic plans to tackle the COVID-19 pandemic.

Acknowledgements:

We would like to acknowledge the support we received from the medical experts working at Quaid-i-Azam University Islamabad Pakistan, National University of Science and Information Technology Islamabad Pakistan and Harbin Medical University China.

Conflict of interest: None to declare.

References

1. Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N & Rubin GJ: The psychological impact of quarantine and how to reduce it: rapid review of the evidence. The Lancet 2020; 395:912-920

- Jooma R, Minhas FA & Saxena S: WHO-AIMS report on mental health system in Pakistan. World Heal. Organ. http://www.who.int/mental_health/pakistan_who_aims_re port. pdf. 2009
- 3. Li W, Yang Y, Liu ZH, Zhao YJ, Zhang Q, Zhang L, Cheung T & Xiang YT: Progression of Mental Health Services during the COVID-19 Outbreak in China. International Journal of Biological Sciences. 2020; 16:1732
- 4. Mak IW, Chu CM, Pan PC, Yiu MG & Chan VL: Longterm psychiatric morbidities among SARS survivors. General hospital psychiatry 2009; 31:318-26
- 5. National Institute of Health: National Action Plan for Corona virus disease (COVID-19) Pakistan. 2020
- 6. Shultz JM, Cooper JL, Baingana F, Oquendo MA, Espinel Z, Althouse BM, Marcelin LH, Towers S, Espinola M, McCoy CB & Mazurik L: The role of fear-related behaviors in the 2013–2016 West Africa Ebola virus disease outbreak. Current psychiatry reports 2016; 18:104
- 7. Surveillances V: The Epidemiological Characteristics of an Outbreak of 2019 Novel Coronavirus Diseases (COVID-19) - China, 2020. China CDC Weekly 2020; 2:113-122
- 8. WHO: WJ. OGAC Technical Consultation on Task Shifting. Key elements of regulatory framework in support of in-country implementation of 'task shifting''. Geneva: World Health Organization, 2007
- 9. World Health Organization: Pakistan health profile 2015. World Health Organization. Regional Office for the Eastern Mediterranean, 2016
- 10. World Health Organization: Novel Corornavirus (2019nCoV): strategic preparedness and response plan, 2019

Correspondence: Rooh Ullah, M.Phil. Department of Microbiology, Quaid-i-Azam University Islamabad, Pakistan E-mail: roohullah@bs.qau.edu.pk