

Ameliorating Drought Stress Effects on Soybean Physiology and Yield by Hydrogen Peroxide

Oqba BASAL (✉)
András SZABÓ

Summary

Soybean is considered as an important legume because of its high content of protein and oil. However, it is sensitive to drought stress that was increasingly recorded recently. Plants respond to drought by several changes on the physiological and molecular levels and by endogenously changing the concentrations of certain substances, such as reactive oxygen species (ROS). An experiment was conducted in 2018 and 2019 to investigate the effects of drought stress and the possible effects of exogenously sprayed hydrogen peroxide (H_2O_2) at R1 stage on two soybean cultivars. Significant decreases in all studied traits were recorded as a result of drought stress. Noticeable enhancements in stomatal conductance, relative chlorophyll content, relative water content, leaf area index, plant height and the final seed yield of both cultivars when treated with H_2O_2 were also recorded. It could be concluded that H_2O_2 spraying can be a good strategy to alleviate presumable negative influence of drought.

Key words

g_s , height, LAI, RWC, SPAD, yield

Department of Crop Production and Applied Ecology, University of Debrecen, 4032 Boszormenyi road 138/A, Debrecen, Hungary

✉ Corresponding author: qqbabasal@gmail.com

Received: March 6, 2020 | Accepted: May 14, 2020

Introduction

Among seed legumes, soybean (*Glycine max* (L.) Merrill) has the greatest global area harvested representing the main source of relatively-cheap protein and vegetable oil (Mutava et al., 2015). Soybean is highly affected by many abiotic stresses, majorly drought stress (Fan et al., 2013) that intensively increased over the past decades, affecting the world's food security (Vurukonda et al., 2016). Soybean is considered highly-sensitive to drought stress compared to other crops (Maleki et al., 2013) especially during certain periods of plant lifecycle (Liu et al., 2004). Drought stress decreases stomatal conductance, biomass, grain yield and its components (Ruppenthal et al., 2016). Drought also affects chlorophyll pigments and photosynthetic electron transport system. As such, production of reactive oxygen species (ROS) is induced (Zgallai et al., 2005) in higher concentrations (Shigeoka et al., 2002) resulting in cellular damage as a result of gene alteration, protein degradation and enzyme inactivation (Mahajan and Tuteja, 2005). High concentrations of ROS cause damages to the cells, yet low concentrations play the role of signaling molecules that can ease several processes like germination and growth (Dowling and Simmons, 2009). For example, it was reported that ROS play noticeable role in regulating stomatal closure in order to optimize water use efficiency (Wang and Song, 2008; Huang et al., 2009). Hydrogen peroxide (H_2O_2) is a compound that belongs to non-radical ROS (Matilla-Vázquez and Matilla, 2012). It regulates many physiological mechanisms, such as growth and development, under both normal and stressed conditions, playing a major role in activating various signal molecules in plants leading to induction of different mechanisms of tolerance (Wendehenne et al., 2004; Bright et al., 2006; Foyer and Noctor, 2009). Many reports have demonstrated that treating plants with suitable concentrations of H_2O_2 increases tolerance to abiotic stresses (Chao et al., 2009; He et al., 2009; Xu et al., 2010; Gondim et al., 2013; Hossain and Fujita, 2013). It was previously reported that treating seeds before sowing with H_2O_2 or applying it as a foliar spray can enhance abiotic stress tolerance in plants. For example, a stimulation in the germination was reported when seeds of *Pseudotsuga menziesii*, *Sorghum nutum*, *Andropogon gerardii*, *Panicum virgatum* (Sarath et al., 2007) and *Zinnia elegans* (Ogawa and Iwabuchi, 2001) were pre-treated with H_2O_2 . Similarly, enhanced germination rates were recorded in maize seeds pre-soaked in 140 mM of H_2O_2 (Ashraf et al., 2015). Jubany-Marí et al. (2009) reported that H_2O_2 is involved in the acclimation of *Cistus albidus* to summer drought. Ishibashi et al. (2011) concluded that spraying soybean plants with H_2O_2 resulted in better net photosynthesis (P_n) and that this application made the plants more tolerant to drought stress. Similar conclusion was also reported for melon plants (Ozaki et al., 2009) and for cucumber seedlings (Sun et al., 2016). Liu et al. (2010) reported improved osmotic stress resistance when two cucumber varieties were pre-treated with H_2O_2 as a result of the activation of antioxidant system. It was suggested that the mechanism by which plants exogenously-sprayed with H_2O_2 better tolerate stress could be by ROS-detoxification modification or by regulation of multi-pathways that respond to stress (Terzi et al., 2014; Hossain et al., 2015). Other authors reported that H_2O_2 alleviated the negative stress effects by either a regulated stomatal closure (Kolla et al., 2007; Quan et al., 2008; Wang and Song, 2008) or by promoting the biosynthesis of the oligosaccharide and, accordingly, maintaining the leaf water content (Ishibashi et al., 2011). Many efforts were

made to exploit this characteristic in enhancing stress tolerance in plants (Jubany-Marí et al., 2009; Liu et al., 2010).

Although some experiments were conducted to investigate the effects of early exogenous H_2O_2 spraying on some physiological and molecular traits of some plant species, yet, to our knowledge, no report was introduced on the influence of the application of H_2O_2 at reproductive stages on soybean physiology and yield, especially in the study area where the soybean production is newly introduced. Therefore, our study aimed at revealing the probable positive effects of exogenously spraying H_2O_2 at R1 stage on both the physiology and the seed yield of two soybean cultivars.

Materials and Methods

Two soybean cultivars, 'Pannonia Kincse' and 'Boglár', were sown in Debrecen University's experimental site (Látókép) (N. latitude 47° 33', E. longitude 21° 27') on April 23rd and 26th and were harvested on September 15th and 16th in 2018 and 2019, respectively. The soil type is calcareous chernozem. The precipitation and the average temperature during the vegetative period of the soybean plants are presented in Figure 1. The experimental design was split-plot with cultivars representing the main plots and irrigation treatments being the sub-plots. Three treatments were applied in three replications: fully-irrigated (FI) treatment, where irrigation was conveniently applied (100 mm in total) to reach water demands as recommended by the experimental site's management (taking into account the precipitation amounts which were recorded as 271.2 mm and 275.9 mm in 2018 and 2019, respectively); drought-stressed treatment (counting only on precipitation) with the application of 1mM of hydrogen peroxide (H_2O_2) as a foliar spray at R1 stage (Fehr and Caviness, 1977) (HP) and drought-stressed treatment with distilled water foliar spray at R1 stage (DW). Hydrogen peroxide and DW treatments received 1.5 liters of H_2O_2 and distilled water per plot, respectively, ensuring that the whole plant parts were completely covered by the sprayed liquid. The final number of plots was 18 (2 cultivars * 3 treatments * 3 replications), with a plot area of 9 m² (3 * 3 m). Each plot consisted of six rows.

The relative chlorophyll content (SPAD) was measured using SPAD-502Plus (Konica Minolta, Japan).




Figure 1. Precipitation (mm) and average temperature (°C) during the vegetative period of the soybean plants in 2018 and 2019

The stomatal conductance (g_s) was measured using AP4 porometer (Delta- T Devices, UK). Leaf area index (LAI) values were recorded using SS1 – SunScan canopy analysis system (Delta-

T Devices, UK). Plant height was measured using a standard ruler. For SPAD, LAI and g_s measurements, 10 randomly-selected plants from the middle rows of each plot were used for the mentioned traits, and three measurements from the second most developed trifoliolate (one measurement for each leaflet) were taken and then averaged.

To calculate the relative water content (RWC), ten fully-matured leaves (homogenous in size) were collected at 6.30 am and fresh weight (FW) of the leaf was measured immediately. Dry weight (DW) was determined via drying the sample at 80°C to constant weight (after 48 h), and turgid weight (TW) was obtained after floating the leaf in distilled water in a tray at 4°C for 48 h. RWC was calculated as $RWC(\%) = (FW - DW)/(TW - DW) \times 100\%$ (Weatherley, 1950). All traits were measured at R2 stage (1 - 2 weeks after H_2O_2 application).

Yield was estimated by manually harvesting the middle two rows of every plot at full maturity (R8) stage. The analysis of variance (ANOVA) was conducted to compare the means and to indicate the effect size of each treatment, and then Tukey post-hoc test was conducted to indicate the statistically-different means (IBM SPSS ver.25, USA Software).

Results

Stomatal Conductance (g_s)

In both cultivars, g_s was significantly higher when irrigation (FI) was applied. However, H_2O_2 -sprayed plots were significantly higher than drought-stressed counterparts in terms of g_s value. Drought application reduced g_s by 51.6 and 47.7% compared to irrigated counterparts, whereas H_2O_2 spraying decreased the reduction ratio to 21.6 and 19.5% in 'Boglár' and 'Pannonia Kincse', respectively (Table 1). Correlation between g_s and irrigation treatment was highly significant (< 0.01) (Table 3), and the effect size of H_2O_2 application (calculated as partial Eta squared) was 81.6 and 90.5% in 'Boglár' and 'Pannonia Kincse', respectively. In other words, H_2O_2 application was responsible for 81.6 and 90.5% of g_s changes in 'Boglár' and 'Pannonia Kincse', respectively.

Relative Chlorophyll Content (SPAD)

In 'Boglár', drought significantly decreased SPAD value by 26.7% compared to the irrigated counterpart, whereas H_2O_2 spraying resulted in better SPAD than both FI and DW counterparts. In 'Pannonia Kincse', on the other hand, irrigation resulted in the highest SPAD value; it was significantly higher (by 22.4%) than drought-stressed counterpart. However, HP treatment enhanced this value (by 13.1%) compared to DW, without reaching the same level of FI treatment as in 'Boglár' (Table 1). The correlation with irrigation was significant (< 0.05) and highly significant (< 0.01) in 'Boglár' and 'Pannonia Kincse' cultivars, respectively (Table 3), and the effect size of H_2O_2 application was noticeably higher (85.7%) in 'Boglár' compared to 'Pannonia Kincse' (59.1%).

Relative Water Content (RWC)

In 'Boglár', drought significantly decreased RWC by 21.2% compared to the irrigated counterpart. However, H_2O_2 -sprayed treatment significantly increased RWC (by 17.3%) compared

Table 1. Stomatal conductance (g_s), relative chlorophyll content (SPAD) and relative water content (RWC) of soybean cultivars 'Boglár' and 'Pannonia Kincse' under three irrigation treatments; fully-irrigated (FI), drought-stressed with H_2O_2 foliar spray (HP) and drought-stressed (DW)

Trait	Treatment	Boglár	Pannonia Kincse	
g_s ($mmol\ m^{-2}\ s^{-1}$)	DW	190.0 ^c	218.3 ^c	
	HP	307.7 ^b	336.3 ^b	
	FI	392.7 ^a	417.7 ^a	
	DW	26.4 ^b	33.6 ^b	
	SPAD	HP	37.2 ^a	38.0 ^b
	FI	36.0 ^a	43.3 ^a	
RWC (%)	DW	57.333 ^b	61.713 ^c	
	HP	69.331 ^a	79.707 ^b	
	FI	72.701 ^a	86.302 ^a	

In each trait, the same letter indicates no significant differences at $p \leq 0.05$ level among the treatments within the same cultivar

to drought-stressed treatment and had very close value to FI treatment. In 'Pannonia Kincse', applying H_2O_2 significantly increased RWC by 29.2% compared to the drought-stressed treatment; however, irrigation treatment had significantly higher RWC (by 8.3 and 39.9%) compared to HP and DW treatments, respectively (Table 1).

Application of H_2O_2 had a significant effect on RWC (by 95.9 and 97.3% in 'Boglár' and 'Pannonia Kincse', respectively) with a highly significant correlation coefficient (Table 3).

Leaf Area Index (LAI)

Both cultivars followed the same trend; LAI was significantly lower in DW treatment (by 13.0 and 17.5% in 'Boglár' and 'Pannonia Kincse', respectively) compared to FI counterpart. Hydrogen peroxide treatment resulted in the highest LAI in both cultivars; LAI was 21.3 and 28.8% higher compared to DW counterparts in 'Boglár' and 'Pannonia Kincse', respectively (Table 2).

Significant correlation was recorded between irrigation and LAI (Table 3). The effect of H_2O_2 application was also significant in both cultivars as well (with ratios of 92.8 and 95.1% in 'Boglár' and 'Pannonia Kincse', respectively).

Plant Height

Drought significantly reduced plant height in both cultivars compared to irrigated counterpart; the reduction ratio was 13.2 and 7.1% in 'Boglár' and 'Pannonia Kincse', respectively. Applying H_2O_2 significantly increased plant height in both cultivars; plant height was 3.6% less in 'Boglár', whereas it was only 0.2% less in 'Pannonia Kincse' compared to irrigated counterparts (Table 2). Application of H_2O_2 had an effect size of 84.2 and 82.8% in 'Boglár' and 'Pannonia Kincse', respectively.

Table 2. Leaf area index (LAI), plant height and yield of soybean cultivars 'Boglár' and 'Pannonia Kincse' under three irrigation treatments: fully-irrigated (FI), drought-stressed with H₂O₂ foliar spray (HP) and drought-stressed (DW)

Trait	Treatment	Boglár	Pannonia Kincse
LAI	DW	4.7 ^b	5.2 ^b
	HP	5.7 ^a	6.7 ^a
	FI	5.4 ^a	6.3 ^a
Plant Height (cm)	DW	86.9 ^b	94.1 ^b
	HP	96.5 ^a	101.1 ^a
	FI	100.1 ^a	101.3 ^a
Yield (t ha ⁻¹)	DW	3.3 ^b	3.7 ^b
	HP	4.0 ^a	4.2 ^{ab}
		4.2 ^a	4.6 ^a

In each trait, the same letter indicates no significant differences at $p \leq 0.05$ level among the treatments within the same cultivar

Yield

Irrigation resulted in the best yield in both cultivars; the yield significantly increased by 27.3 and 24.3% in irrigated treatment compared to drought-stressed counterpart in 'Boglár' and 'Pannonia Kincse', respectively. Spraying with H₂O₂ also increased the yield of both cultivars compared to drought-stressed treatment; the increase ratio was 21.2 and 13.5% in 'Boglár' and 'Pannonia Kincse', respectively (Table 2). The effect of H₂O₂ application was bigger (78.9%) in 'Pannonia Kincse' compared to 'Boglár' (72.1%). Correlation of irrigation with yield was highly significant in both cultivars (Table 3).

Table 3. Correlation coefficient of irrigation treatments with the studied traits

Trait	Boglár	Pannonia Kincse
g_s	.959**	.975**
SPAD	.758*	.926**
RWC	.929**	.948**
LAI	.668*	.720*
Plant Height	.908**	.755*
Yield	.861**	.912**

* Correlation is significant at $p \leq 0.05$ level (2-tailed).

** Correlation is significant at $p \leq 0.01$ level (2-tailed).

Discussion

Drought reduced g_s in both cultivars as compared to irrigated counterparts, whereas H₂O₂ application could measurably alleviate g_s . Drought stress induces stomatal closure, limits gas exchange and photosynthesis (Yordanov et al., 2000). Ohashi et al. (2006) reported that stomatal conductance of soybean plants significantly decreased under drought stress conditions; similar result was reported by Zhang et al. (2016); a 98.8% decrease in g_s under drought. They concluded that this reduction in g_s was a result of the reduced ratio of open stomata and stomatal aperture size in the plants subjected to drought stress. Hao et al. (2013) reported a significant reduction in stomatal conductance from 0.25 to 0.10 mol H₂O m⁻¹ s⁻¹ as a result of drought applied on soybean plants. Mathobo et al. (2017) justified the reduction in g_s in their experiment on dry beans (*Phaseolus vulgaris*) by the prevention of CO₂ from entering the leaf by stomatal closure. Similarly, Rosales et al. (2012) reported a 70% reduction of g_s after 22 days of drought stress application. Tang et al. (2017) concluded that polyethylene glycol (PEG) induced water stress on soybean significantly reduced g_s by 73%. Ishibashi et al. (2011) compared g_s of two groups of soybean seedlings under drought stress conditions. One group was sprayed with H₂O₂ and the other group with distilled water (DW). They reported that g_s was significantly higher in H₂O₂-treated plants than in DW-treated plants. After two days of spraying, g_s levels in H₂O₂-treated and DW-treated plants were 508 and 323 mmol m⁻² s⁻¹, respectively. They concluded that H₂O₂ spraying reduced stomatal closure caused by drought stress; i.e. H₂O₂ treatment reduced soybean sensitivity to drought stress. In experiment of Terzi et al. (2014) maize leaves pretreated with 10 mM H₂O₂ significantly enhanced g_s (by approximately 50%) as compared to drought-stressed leaves. They concluded that spraying leaves with H₂O₂ can reduce water loss under drought stress conditions by increasing the concentrations of metabolites that are involved in osmotic adjustment (such as proline, polyamines and soluble sugars). Other ROS species were also reported to have a role in alleviating drought stress. Razmi et al. (2017) reported that water stress reduced stomatal conductance of three soybean leaves compared to well-watered counterparts, and foliar spray of 0.4 mM salicylic acid (SA) significantly reversed drought induced stomatal closure and increased it.

Drought resulted in lower SPAD values compared to irrigated counterparts of both cultivars, whereas H₂O₂ application enhanced this value; it even increased SPAD to a level more than did irrigation in 'Boglár', but not in 'Pannonia Kincse'. Similarly, Ergo et al. (2018) reported that SPAD values significantly decreased from 35.5 to 22.4 under drought stress applied 30 days after R5.5 stage. Chlorophylls are the main pigments of absorbing, transporting and converting light energy, and their content is a major parameter that indicates photosynthetic performance (Liu et al., 2007). Subjecting plants to drought stress resulted in a significant decline in chl_{a+b} (from 19.5 to 13.0 mg g⁻¹ DW), indicating a reduced capacity of absorbing and converting light energy (Tang et al., 2017). Similarly, Dong et al. (2015) concluded that light absorption was reduced by drought stress that resulted in changing both leaf area index and leaf chlorophyll content. Both chlorophylls *a* and *b* were reduced under drought stress (Farooq et al., 2010). Other papers also reported that chlorophyll content was decreased because of drought in soybean (Makbul et

al., 2011), chickpea (Mafakheri et al., 2010) and pea (Inaki-Iturbe et al., 1998). That reduction was attributed to induced destruction of the chloroplasts and to the instability of the chlorophyll protein complex (Khan et al., 2015). Sun et al. (2016) reported that the application of H_2O_2 significantly increased the leaf chlorophyll content of cucumber plants exposed to medium drought conditions. An evaluation of the effects of H_2O_2 on leaf chlorophyll content during adventitious rooting under drought conditions showed that drought stress resulted in a decline in chlorophyll content after 72 h of its application, producing a 39.1% decrease in the chlorophyll *a* content compared to control. However, applying exogenous H_2O_2 retarded chlorophyll degradation, especially chlorophyll *a* (Liao et al., 2012). Maize leaves had higher levels of both chlorophylls *a* and *b* when seeds were soaked in 140 mM H_2O_2 before sowing (Ashraf et al., 2015). Enhanced chlorophyll levels induced by hydrogen peroxide treatment were justified by H_2O_2 -stimulated antioxidant enzyme activities (Azevedo Neto et al., 2005; Gao et al., 2010). Razmi et al. (2017) reported that drought significantly reduced both chlorophyll *a* and *b* contents in soybean leaves. However, significant increases (by 15% in chlorophyll *a* and 19% in chlorophyll *b*) resulted from foliar application of 0.4 mM of salicylic acid (SA) compared to control treatment (no SA).

Application of H_2O_2 significantly increased RWC in both cultivars compared to drought-stressed counterparts, and irrigation further enhanced this trait. It was previously reported that drought stress reduced RWC of soybean leaves (Razmi et al., 2017). Ishibashi et al. (2011) reported that RWC in H_2O_2 -treated and DW-treated (control, treated with distilled water only) plants was 60 and 40%, respectively after four days of drought stress application, and RWC was also higher in H_2O_2 -treated plants than in DW-treated plants after six days of drought stress imposition. Authors concluded that H_2O_2 spraying enabled the leaves to maintain high levels of RWC by regulating the osmolality in the leaves, consequently ameliorating the negative effects of drought stress. Similar results on cucumber seedlings were reported later by Sun et al. (2016). Application of SA on common bean improved RWC under drought stress conditions (Sadeghipour and Aghaei, 2012).

Irrigation significantly increased LAI in both cultivars compared to drought-stressed counterparts. Furthermore, H_2O_2 application could further increase LAI in both cultivars. Drought stress decreases LAI (Liu et al., 2008). Ashraf et al. (2015) reported that seeds soaked in 20, 80, 100 and 140 mM of H_2O_2 later formed plants with higher leaf area under drought stress conditions compared to non-treated seeds. Using (SA), other authors concluded that treatments with SA could improve LAI in different plants including soybean (Kuchlan et al., 2017; Razmi et al., 2017), strawberry (Ghaderi et al., 2015) and lemongrass (Idrees et al., 2010). This was attributed to increased accumulation of certain proteins (such as proline) and soluble sugars that, in part, enhances cell turgor pressure (Razmi et al., 2017).

Drought decreased plant height in both cultivars, and H_2O_2 application could ameliorate drought's effects and result in enhanced plant height, reaching a very close level of irrigated counterparts, especially in 'Pannonia Kincse' cultivar. Many previous papers reported a reduction in plant height under drought stress conditions (e.g. Atti et al., 2004; Demirtas et al., 2010; Hao et al., 2013; Mak et al., 2014). Moreover, Garcia et al.

(2010) reported a significant difference in plant height of drought-stressed soybean genotypes compared to control counterparts. They also reported the different examined genotypes to be significantly different in plant height, which was demonstrated later by Hossain et al. (2014) who studied the effect of drought stress on the plant height of three soybean genotypes, one drought-susceptible and two drought-tolerant genotypes; they reported plant height to be shortened as a result of drought stress for the three genotypes. However, the drought-susceptible genotype had a 44.3% of height of the control plants, whereas it was 56.7% and 59.1% for the two drought-tolerant genotypes. The authors attributed this reduction to a drought tolerance mechanism, as cell swells, cell wall and synthesis enzymes reduces, consequently, growth and plant height decreases (Levitt, 1980; Austin, 1989). Banon et al. (2006) justified this decrease by a reduction in cell elongation caused by inhibited growth promoting hormones that, in part, led to decrease of cell turgor, cell volume and eventually cell growth and/or by a restriction of xylem and phloem vessels (Lavisalo and Schuber, 1998). Another possible explanation is that drought results in a decrease in the rate at which the stem nodes are produced (Frederick et al., 1989). Abbas and Mohamed (2011) conducted an experiment on common bean (*Phaseolus vulgaris*) seeds where half of the seeds were soaked in hydrogen peroxide (2%) for four hours and then air dried, and the other half of the seeds were soaked in distilled water for four hours and then air dried. Their results showed an increase by 43.6% in the H_2O_2 -treated seedling height under a drought level of 60% of field capacity. Increasing the drought severity (to reach only 40% of field capacity) decreased the seedling height of both treatments. However, H_2O_2 -treated seedlings showed better height by 38.4%.

Yield was significantly reduced by drought in both cultivars, whereas H_2O_2 application measurably recovered yield. The recovery was significant in 'Boglár' cultivar. Soybean seed yield decreases under drought stress conditions (Karam et al., 2005; Dogan et al., 2007; Bajaj et al., 2008; Sincik et al., 2008; Gercek et al., 2009; Sadeghipour and Abbasi, 2012; Li et al., 2013). It was also reported that genotypes significantly differ in yield production under drought stress conditions (Bellaloui and Mengistu, 2008; Maleki et al., 2013; He et al., 2017). Explanations for under drought stress conditions were that drought stress shortens the seed-filling period what explains yield decrease (Smiciklas et al., 1992). Other authors suggested that it is due to the reduction of seeds number (Dornbos et al., 1989), seeds weight (Samarah et al., 2006; Demirtas et al., 2010) and pod number per plant (Atti et al., 2004; Khatun et al., 2016). However, exogenous application of H_2O_2 had improved plant biomass in wheat under drought stress (He et al., 2009), and SA application improved the grain yield of common bean under drought stress conditions (Sadeghipour and Aghaei, 2012). Horvath et al. (2007) reported that SA can enhance metabolite stream to the developing grains which reduces the abortion rate that, in part, can significantly increase pod number per plant and seed number pod^{-1} in soybean (Khatun et al., 2016). Not only yield, but also yield components (number of grains m^{-2} , pods $plant^{-1}$) were enhanced with the application of SA on soybean leaves under drought stress conditions (Razmi et al., 2017). The authors attributed the increase of grain yield due to SA application, improved RWC, reduced restrictions of stomatal conductance and the enhanced biosynthesis of photosynthetic pigments in the leaves.

Conclusions

Drought stress negatively affects soybean morphology, physiology and, consequently, the final seed yield; it resulted in significant reductions in all studied traits. Treating drought-stressed soybean plants with 1 mM of H₂O₂ could alleviate that negative influence and enhance all studied traits. Its effect was more noticeable on the leaf area index LAI as H₂O₂-sprayed plants had higher LAI values than both drought-stressed and fully-irrigated counterparts. The effect was also noticeable on the relative water content RWC of both cultivars. The correlation of irrigation with the studied traits was also significant.

Acknowledgments

The publication is supported by the EFOP-3.6.3-VEKOP-16-2017-00008 Project. The Project is co-financed by the European Union and the European Social Fund.

References

- Abass S. M., Mohamed H. I. (2011). Alleviation of Adverse Effects of Drought Stress on Common Bean (*Phaseolus vulgaris* L.) by Exogenous Application of Hydrogen Peroxide. *Bangladesh J. Bot.* 41(1): 75-83.
- Ashraf M. A., Rasheed R., Hussain I., Iqbal M., Haider M. Z., Parveen S., Sajid M. A. (2015). Hydrogen peroxide modulates antioxidant system and nutrient relation in maize (*Zea mays* L.) under water-deficit conditions. *Arch. Agron. Soil Sci.* 61 (4): 507-523.
- Atti S., Bonnell R., Smith D., Prasher S. (2004). Response of an Indeterminate Soybean (*Glycine max* (L.) Merrill) to Chronic Water Deficit During Reproductive Development Under Greenhouse Conditions. *Can. Water Resour. J. / Revue Canadienne des ressources hydriques* 29 (4): 209-222.
- Austin R. B. (1989). Maximizing crop production in water limited environments. 13-25. in: F. W. G. Baker. *Drought resistance in cereals*. CAB International, Wallingford, England 2220.
- Azevedo Neto A. D., Prisco J. T., Enéas-Filho J., Medeiros J. V., Gomes-Filho E. (2005). Hydrogen peroxide pre-treatment induces salt-stress acclimation in maize plants. *J. Plant Physiol.* 162: 1114-1122.
- Bajaj S., Chen P., Longer D. E., Shi A., Hou A., Ishibashi T., Brye K. R. (2008). Irrigation and planting date effects on seed yield and agronomic traits of early-maturing Soybean. *J. Crop Improv.* 22 (1): 47-65.
- Banon S. J., Ochoa J., Franco J. A., Alarcon J. J., Sanchez-Blanco M. J. (2006). Hardening of oleander seedlings by deficit irrigation and low air humidity. *Environ. Exp. Bot.* 56: 36-43.
- Bellaloui N., Mengistu A. (2008). Seed composition is influenced by irrigation regimes and cultivar differences in soybean. *Irrig. Sci.* 26: 261-268.
- Bright J., Desikan R., Hancock J. T., Weir I. S., Neill S. J. (2006). ABA-induced NO generation and stomatal closure in Arabidopsis are dependent on H₂O₂ synthesis. *Plant J.* 45: 113-122.
- Chao Y. Y., Hsu Y. T., Kao C. H. (2009). Involvement of glutathione in heat shock-and hydrogen peroxide-induced cadmium tolerance of rice (*Oryza sativa* L.) seedlings. *Plant Soil* 318: 37-45.
- Demirtas Ç. D., Yazgan S., Candogan B. C., Sincik M., Büyükcangaz H., Göksoy A. T. (2010). Quality and yield response of soybean (*Glycine max* (L.) Merrill) to drought stress in sub-humid environment. *Afr. J. Biotechnol.* 9(41): 6873-6881.
- Dogan E., Kirnak H., Copur O. (2007). Deficit irrigations during soybean reproductive stages and CROPGRO-soybean simulations under semi-arid climatic conditions. *Field Crops Res.* 103 (2): 154-159.
- Dong J., Xiao X., Wagle P., Zhang G., Zhou Y., Jin C., Torn M. S., Meyers T. P., Suyker A. E., Wang J., Yan H., Biradar Ch., Moore III B. (2015). Comparison of four EVI-based models for estimating gross primary production of maize and soybean croplands and tallgrass prairie under severe drought. *Remote Sens. Environ.* 162: 154-168.
- Dornbos D. L., Mullen R. E., Shibles R. E. (1989). Drought stress effects during seed fill on soybean seed germination and vigor. *Crop Sci.* 29(2): 476-480.
- Dowling D. K., Simmons L. W. (2009). Reactive oxygen species as universal constraints in life-history evolution. *Proc. R. Soc. B. Biol. Sci.* 276: 1737-1745.
- Ergo V. V., Lascano R., Vegad C. R. C., Parola R., Carrera C. S. (2018). Heat and water stressed field-grown soybean: A multivariate study on the relationship between physiological-biochemical traits and yield. *Environ. Exp. Bot.* 148: 1-11.
- Fan X. D., Wang J. Q., Yang N., Dong Y. Y., Liu L., Wang F. W., Li H. Y. (2013). Gene expression profiling of soybean leaves and roots under salt, saline-alkali and drought stress by high throughput Illumina sequencing. *Gene* 512(2): 392-402.
- Farooq M., Wahid A., Lee D. J., Cheema S. A., Aziz T. (2010). Comparative time course action of the foliar applied glycine betaine, salicylic acid, nitrous oxide, brassinosteroids and spermine in improving drought resistance of rice. *J. Agron. Crop Sci.* 196: 336-345.
- Fehr W. R., Caviness C. E. (1977). Stages of soybean development. Special Report 87. <http://lib.dr.iastate.edu/specialreports/87>, 1977.
- Foyer C. H., Noctor G. (2009). Redox regulation in photosynthetic organisms: signaling, acclimation, and practical implications. *Antioxid. Redox Sign.* 11: 861-905.
- Frederick J. R., Woolley J. T., Hesketh J. D., Peters D. B. (1989). Phenological Responses of Old and Modern Soybean Cultivars to Air Temperature and Soil Moisture Treatment. *Field Crops Res.* 21: 9-18.
- Gao Y., Guo Y. K., Lin S. H., Fang Y. Y., Bai J. G. (2010). Hydrogen peroxide pretreatment alters the activity of antioxidant enzymes and protects chloroplast ultrastructure in heat-stressed cucumber leaves. *Sci. Hort.* 126: 20-26.
- Garcia A. G., Persson T., Guerra L. C., Hoogenboom G. (2010). Response of soybean genotypes to different irrigation regimes in a humid region of the southeastern USA. *Agr. Water Manage.* 97: 981-987.
- Gercek S., Boydak E., Okant M., Dikilitas M. (2009). Water pillow irrigation compared to furrow irrigation for soybean production in a semi-arid area. *Agr. Water Manage.* 96 (1): 87-92.
- Ghaderi N., Normohammadi S., Javadi T. (2015). Morpho-physiological responses of strawberry (*Fragaria ananassa*) to exogenous salicylic acid application under drought stress. *J. Agr. Sci. Tech.* 17: 167-178.
- Gondim F. A., Miranda R. S., Gomes-Filho E., Prisco J. T. (2013). Enhanced salt tolerance in maize plants induced by H₂O₂ leaf spraying is associated with improved gas exchange rather than with non-enzymatic antioxidant system. *Theor. Exp. Plant Physiol.* 25: 251-260.
- Hao L., Wang Y., Zhang J., Xie Y., Zhang M., Duan L., Li Z. (2013). Coronatine enhances drought tolerance via improving antioxidative capacity to maintaining higher photosynthetic performance in soybean. *Plant Sci.* 210: 1-9.
- He J., Du Y. L., Wang T., Turner N. C., Yang R. P., Jin Y., Li F. M. (2017). Conserved water use improves the yield performance of soybean (*Glycine max* (L.) Merrill) under drought. *Agr. Water Manage.* 179: 236-245.
- He, L., Gao, Z., Li, R. (2009). Pretreatment of seed with H₂O₂ enhances drought tolerance of wheat (*Triticum aestivum* L.) seedlings. *Afr. J. Biotechnol.* 8: 6151-6157.
- Horvath E., Szalai G., Janda T. (2007). Induction of abiotic stress tolerance by salicylic acid signaling. *J. Plant Growth Regul.* 26: 290-300.
- Hossain M. A., Bhattacharjee S., Armin S. M., Qian P., Xin W., Li H. Y., Burritt D. J., Fujita M., Tran L. S. P. (2015). Hydrogen peroxide priming modulates abiotic oxidative stress tolerance: insights from ROS detoxification and scavenging. *Front. Plant Sci.* 6: 1-19.
- Hossain M. A., Fujita M. (2013). Hydrogen peroxide priming stimulates drought tolerance in mustard (*Brassica juncea* L.). *Plant Gene and Trait.* 4: 109-123.
- Hossain. Md. M., Liu X., Qi X., Lam H-M., Zhang J. (2014). Differences between soybean genotypes in physiological response to sequential

- soil drying and rewetting. *The Crop Journal* 2: 366-380.
- Huang X. Y., Chao D. Y., Gao J. P., Zhu M. Z., Shi M., Lin H. X. (2009). A previously unknown zinc finger protein, DST, regulates drought and salt tolerance in rice via stomatal aperture control. *Gene Dev.* 23: 1709-1713.
- Idrees M., Masroor M., Khan A., Aftab T., Naeem M., Hashmi N. (2010). Salicylic acid-induced physiological and biochemical changes in lemongrass varieties under water stress. *J. Plant Interact.* 5: 293-303.
- Inaki-Iturbe O., Escuredo P. R., Arrese-Igor C., Becana M. (1998). Oxidative damage in pea plants exposed to water deficit or paraquat. *Plant. Physiol.* 116: 173-181.
- Ishibashi Y., Yamaguchi H., Yuasa T., Iwaya-Inoue M., Arima S., Zheng S. H. (2011). Hydrogen peroxide spraying alleviates drought stress in soybean plants. *J. Plant Physiol.* 168 (13): 1562-1567.
- Jubany-Marí T., Munné-Bosch S., López-Carbonell M., Alegre L. (2009). Hydrogen peroxide is involved in the acclimation of the Mediterranean shrub, *Cistus albidus* L., to summer drought. *J. Exp. Bot.* 60 (1): 107-120.
- Karam F., Masaad R., Sfeir T., Mounzer O., Roupheal Y. (2005). Evapotranspiration and seed yield of field grown soybean under deficit irrigation conditions. *Agr. Water Manage.* 75: 226-244.
- Khan M. I. R., Fatma M., Per T. S., Anjum N. A., Khan N. A. (2015). Salicylic acid-induced abiotic stress tolerance and underlying mechanisms in plants. *Front. Plant Sci.* 6: 1-17.
- Khatun S., Roy T. S., Haque Md. N., Alamgir B. (2016). Effect of plant growth regulators and their time of application on yield attributes and quality of soybean. *IJPSS.* 11: 1-9.
- Kolla V. A., Vavasseur A., Raghavendra A. S. (2007). Hydrogen peroxide production is an early event during bicarbonate induced stomatal closure in abaxial epidermis of *Arabidopsis*. *Planta* 225: 1421-1429.
- Kuchlan P., Kuchlan M. K., Husain S. M. (2017). Effect of foliar application of growth activator, promoter and antioxidant on seed quality of soybean. *Legume Res.* 40(2): 313-318.
- Lavisolo C., Schuber A. (1998). Effects of water stress on vessel size xylem hydraulic conductivity in *Vitis vinifera* L. *J. Exp. Bot.* 49 (321): 693-700.
- Levitt J. (1980). Responses of plants to environmental stresses. Academic Press. New York and London. 697 pp.
- Li D., Liu H., Qiao Y., Wang Y., Cai Z., Dong B., Shi Ch., Liu Y., Li X., Liu M. (2013). Effects of elevated CO₂ on the growth, seed yield, and water use efficiency of soybean (*Glycine max* (L.) Merr.) under drought stress. *Agr. Water Manage.* 129: 105-112.
- Liao W-B., Huang G-B., Yu J-H., Zhang M-L. (2012). Nitric oxide and hydrogen peroxide alleviate drought stress in marigold explants and promote its adventitious root development. *Plant Physiol. Bioch.* 58: 6-15.
- Liu F., Jensen Ch. R., Andersen M. N. (2004). Pod Set Related to Photosynthetic Rate and Endogenous ABA in Soybeans Subjected to Different Water Regimes and Exogenous ABA and BA at Early Reproductive Stages. *Ann. Bot-London* 94: 405-411.
- Liu X., Jin J., Wang G., Herbert S. J. (2008). Soybean yield physiology and development of high-yielding practices in Northeast China. *Field Crops Res.* 105: 157-171.
- Liu Z., Guo Y., Bai J. (2010). Exogenous hydrogen peroxide changes antioxidant enzyme activity and protects ultrastructure in leaves of two cucumber ecotypes under osmotic stress. *J. Plant Growth Regul.* 29 (2): 171-183.
- Liu Z. H., Ran S. L., Bai L. R., Zhao K. F. (2007). Salt stress's effects on the contents of chlorophyll and organic solutes of *Aeluropus littoralis*. *J. Plant Physiology Mol. Biol.* 33: 165-172.
- Mafakheri A., Siosemardeh A., Bahramnejad B., Struik P. C., Sohrabi Y. (2010). Effect of drought stress on yield: proline and chlorophyll contents in three chickpea cultivars. *AJCS* 4: 580-585.
- Mahajan S., Tuteja N. (2005). Cold, salinity and drought stresses: An overview. *Arch. Biochem. Biophys.* 444: 139-158.
- Mak M., Babla M., Xu S. C., O'Carriagan A., Liu X. H., Gong Y. M., Holford P., Chen Z. H. (2014). Leaf mesophyll K⁺, H⁺ and Ca²⁺ fluxes are involved in drought-induced decrease in photosynthesis and stomatal closure in soybean. *Environ. Exp. Bot.* 98: 1-12.
- Makbul S., Saruhan Güller N., Druma N., Güven S. (2011). Changes in anatomical and physiological parameters of soybean under drought stress. *Turk. J. Bot.* 35: 369-377.
- Maleki A., Naderi A., Naseri R., Fathi A., Bahamin S., Maleki R. (2013). Physiological Performance of Soybean Cultivars under Drought Stress. *Bulletin of Environment, Pharmacology and Life Sciences* 2(6): 38-44.
- Mathobo R., Marais D., Steyn J. M. (2017). The effect of drought stress on yield, leaf gaseous exchange and chlorophyll fluorescence of dry beans (*Phaseolus vulgaris* L.). *Agr. Water Manage.* 180: 118-125.
- Matilla-Vázquez M. A., Matilla A. J. (2012). Role of H₂O₂ as Signaling Molecule in Plants/ Environmental Adaptations and Stress Tolerance of Plants in the Era of Climate Change. Springer, New York: 361-380.
- Mutava R. N., Prince S. J. K., Syed N. H., Song L., Valliyodan B., Chen W., Nguyen H. T. (2015). Understanding abiotic stress tolerance mechanisms in soybean: A comparative evaluation of soybean response to drought and flooding stress. *Plant Physiol. Bioch.* 86: 109-120.
- Ogawa K., Iwabuchi M. A. (2001). A mechanism for promoting the germination of *Zinnia elegans* seeds by hydrogen peroxide. *Plant Cell Physiol.* 42: 286-291.
- Ohashi Y., Nakayama N., Saneoka H., Fujita K. (2006). Effects of drought stress on photosynthetic gas exchange, chlorophyll fluorescence and stem diameter of soybean plants. *Biol. Plantarum* 50: 138-141.
- Ozaki K., Uchida A., Takabe T., Shinagawa F., Tanaka Y., Takabe T., Hayashi T., Hattori T., Rai AK., Takabe T. (2009). Enrichment of sugar content in melon fruits by hydrogen peroxide treatment. *J. Plant Physiol.* 166(6): 569-578.
- Quan L. J., Zhang B., Shi W. W., Li H. Y. (2008). Hydrogen peroxide in plants: a versatile molecule of the reactive oxygen species network. *J. Integr. Plant Biol.* 50: 2-18.
- Razmi N., Ebadi A., Daneshian J., Jahanbakhsh S. (2017). Salicylic acid induced changes on antioxidant capacity, pigments and grain yield of soybean genotypes in water deficit condition. *J. Plant Interact.* 12(1): 457-464.
- Rosales M. A., Ocampo E., Rodriguez-Valentin R., Olvera-Carrillo Y., Acosta-Gallegos J., Covarrubias A. A. (2012). Physiological analysis of common bean (*Phaseolus vulgaris*) cultivars uncovers characteristics related to terminal drought. *Plant Physiol. Biochem.* 56: 24-34.
- Ruppenthal V., Zoz T., Steiner F., Lana M. C., Castagnara D. D. (2016). Silicon does not alleviate the adverse effects of drought stress in soybean plants. *Semin Cienc Agrar.* 37: 3941-3954.
- Sadeghipour O., Abbasi S. (2012). Soybean Response to Drought and Seed Inoculation. *World Applied Sciences Journal* 17(1): 55-60.
- Sadeghipour O., Aghaei P. (2012). Biochemical changes of common bean (*Phaseolus vulgaris* L.) to pretreatment with salicylic acid (SA) under water stress conditions. *Adv. Environ. Biol.* 2: 14-22.
- Samarah N. H., Mullen R. E., Cianzio S. R., Scott P. (2006). Dehydrin-like proteins in soybean seeds in response to drought stress during seed filling. *Crop Sci.* 46: 2141-2150.
- Sarath G., Hou G., Baird L. M., Mitchell R. B. (2007). Reactive oxygen species, ABA and nitric oxide interactions on the germination of warm-season C4-grasses. *Planta* 226: 697-708.
- Shigeoka S., Ishikawa T., Tamoi M., Miyagawa Y., Takeda T., Yabuta Y., Yoshimura K. (2002). Regulation and function of ascorbate peroxidase isoenzymes. *J. Exp. Bot.* 53: 1305-1319.
- Sincik M., Candogan B. N., Demirtas C., Büyükacangaz H., Yazgan S., Gksoy A. T. (2008). Deficit irrigation of soybean [*Glycine max* (L.) Merr.] in a sub-humid climate. *J. Agron. Crop Sci.* 194: 200-205.
- Smiciklas K. D., Mullen R. E., Carlson R. E., Knapp A. D. (1992). Soybean seed quality response to drought stress and pod position. *Agron. J.* 84: 166-170.
- Sun Y., Wang H., Liu S., Peng X. (2016). Exogenous application of hydrogen peroxide alleviates drought stress in cucumber seedlings. *S. Afr. J. Bot.* 106: 23-28.

- Tang Y., Sun X., Wen T., Liu M., Yang M., Chen X. (2017). Implications of terminal oxidase function in regulation of salicylic acid on soybean seedling photosynthetic performance under water stress. *Plant Physiol. Bioch.* 112: 19-28.
- Terzi R., Kadioglu A., Kalaycioglu E., Saglam A. (2014). Hydrogen peroxide pretreatment induces osmotic stress tolerance by influencing osmolyte and abscisic acid levels in maize leaves. *J. Plant Interact.* 9: 559-565.
- Vurukonda S. S. K. P., Vardharajula S., Shrivastava M., SkZ A. (2016). Enhancement of drought stress tolerance in crops by plant growth promoting rhizobacteria. *Microbiol. Res.* (Vol. 184). Elsevier GmbH.
- Wang P., Song C. (2008). Guard-cell signaling for hydrogen peroxide and abscisic acid. *New Phytol.* 178: 703-718.
- Weatherley P. (1950). Studies in the water relations of the cotton plant: I. The field measurement of water deficits in leaves. *New Phytol.* 49(1): 81-97.
- Wendehenne D., Durner J., Klessig D. F. (2004). Nitric oxide: a new player in plant signaling and defense responses. *Curr. Opin. Plant Biol.* 7: 449-455.
- Xu F. J., Jin C. W., Liu W. J., Zhang Y. S., Lin X. Y. (2010). Pretreatment with H₂O₂ alleviates aluminum-induced oxidative stress in wheat seedlings. *J. Integr. Plant Biol.* 54: 44-53.
- Yordanov I., Velikova V., Tsonev T. (2000). Plant responses to drought, acclimation, and stress tolerance. *Photosynthetica* 38: 171-186.
- Zgallai H., Steppe K., Lemeur R. (2005). Photosynthetic, physiological and biochemical responses of tomato plants to polyethylene glycol-induced water deficit. *J. Integr. Plant Biol.* 47: 1470-1478.
- Zhang J., Liu J., Yang C., Du S., Yang W. (2016). Photosynthetic performance of soybean plants to water deficit under high and low light intensity. *S. Afr. J. Bot.* 105: 279-287.

aCS85_23