



Schwehm, Carolin and Li, Jin and Song, Hongmei and Hu, Xiao and Kellam, Barrie and Stocks, Michael (2015) Synthesis of new DPP-4 inhibitors based on a novel tricyclic scaffold. *ACS Medicinal Chemistry Letters*, 6 (3). pp. 324-328. ISSN 1948-5875

**Access from the University of Nottingham repository:**

[http://eprints.nottingham.ac.uk/28517/1/M%20Stocks\\_ACS%20med%20chem%20letters\\_2015.pdf](http://eprints.nottingham.ac.uk/28517/1/M%20Stocks_ACS%20med%20chem%20letters_2015.pdf)

**Copyright and reuse:**

The Nottingham ePrints service makes this work by researchers of the University of Nottingham available open access under the following conditions.

- Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners.
- To the extent reasonable and practicable the material made available in Nottingham ePrints has been checked for eligibility before being made available.
- Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.
- Quotations or similar reproductions must be sufficiently acknowledged.

Please see our full end user licence at:

[http://eprints.nottingham.ac.uk/end\\_user\\_agreement.pdf](http://eprints.nottingham.ac.uk/end_user_agreement.pdf)

**A note on versions:**

The version presented here may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the repository url above for details on accessing the published version and note that access may require a subscription.

For more information, please contact [eprints@nottingham.ac.uk](mailto:eprints@nottingham.ac.uk)

# Synthesis of new DPP-4 inhibitors based on a novel tricyclic scaffold.

Carolin Schwehm<sup>1</sup>, Jin Li<sup>2</sup>, Hongmei Song<sup>2</sup>, Xiao Hu<sup>2</sup>, Barrie Kellam<sup>1</sup>, Michael J.Stocks<sup>1\*</sup>

<sup>1</sup> Centre for Biomolecular Sciences, University Park Nottingham, Nottingham, NG7 2RD, UK,

<sup>2</sup> Hitgen Ltd., F7-10, Building B3, Tianfu Life Science Park, 88 South Kayuan Road, Chengdu, Sichuan, China 610041


**KEYWORDS** Diabetes, Scaffold, Crystal Structure, dipeptyl peptidase VI (DPP-4) inhibitor, 1,2,4-triazole

**ABSTRACT:** A novel molecular scaffold has been synthesized and its synthesis and incorporation into new analogues of biologically active molecules will be discussed. A comparison of the inhibitory activity of these compounds to the known type-2 diabetes compound (sitagliptin) against dipeptidyl peptidase-4 (DPP-4) will be shown.

Diabetes remains one of the world's largest health problems with numerous different factors contributing to its pathogenesis. According to the WHO in 2013<sup>1</sup>, 347 million people were diagnosed with type-2 diabetes mellitus, with an alarming growth predicted over the next decade.

Type-2 diabetes is a chronic disease, characterized by elevated blood sugar levels, leading to severe vascular complications and an increased mortality risk. Dipeptidyl peptidase-4 (DPP-4), a widely distributed serine protease found solubilized in blood or anchored into tissue membranes, is involved in glucose metabolism and is now a validated target for anti-diabetic therapy. Inhibition of DPP-4 has been shown to result in indirect stimulation of insulin secretion.<sup>2-3</sup> The mechanism of inhibition<sup>4-5</sup> is through an increase in the release of incretin (GLP-1 and GIP) following food intake, therefore inhibiting glucagon release, which in turn increases insulin secretion and decreases blood glucose levels.<sup>6</sup>

Sitagliptin (Januvia<sup>TM</sup>) was the first approved DPP-4 inhibitor launched by Merck in 2006.<sup>7</sup> It was followed by several, structurally diverse DPP-4 inhibitors namely vildagliptin, saxagliptin, alogliptin, linagliptin and Gempigliptin and recent communications highlighting further compounds such as omarigliptin<sup>8</sup> and imigliptin<sup>9</sup> have been recently communicated (Figure 1).




**Figure 1.** Approved DPP-4 inhibitors: sitagliptin, vildagliptin, saxagliptin, alogliptin, linagliptin and the clinical candidates imigliptin and omarigliptin,

The so called “gliptins” are under investigation for other potential therapeutic uses. For example, it was reported that potential substrates of DPP-4 could have implications in other metabolic disorders and that DPP-4 inhibitors could be utilised in the treatment of diseases associated with the immune/inflammatory response, heart failure, cancer and neurodegenerative disorders. In addition it was stated that a positive role of DPP-4 inhibition was


observed in diseases of the kidney and the cardiovascular system.<sup>10</sup>

In our on-going exploration of novel constrained molecular scaffolds containing substituted ring-fused 1,2,4-triazoles, we were drawn to the possibility of substituting the piperazine-fused 1,2,4-triazolo group present in sitagliptin with a new tricyclic octahydro-[1,2,4]triazolo[4,3-*a*][1,6]naphthyridine molecular scaffold (**1a-b**) to generate sitagliptin hybrid structures of the type shown in compounds **2a** and **2b** (Figure 2).



**Figure 2.** Novel tricyclic scaffold **1a-b** and proposed DPP-4 inhibitors **2a** and **2b**. Compounds **2a** and **2b** would exist as a 1:1 mixture of either *cis*- or *trans*-diastereoisomers.

To test our hypothesis, preliminary docking studies were carried out with the known crystal structure of sitagliptin in DPP-4 (pdb code 1X70).<sup>11-12</sup> The *cis*-fused diastereomer **2a** ( $R_2 = CF_3$ ) showed a good overlay with sitagliptin as well as a good topographical fit into the enzyme pocket when compared to the *trans*-fused diastereomer **2b**. As expected, the 2, 4, 5-trifluorophenyl group fully occupied the  $S_1$  pocket, which was previously reported by the Merck group.<sup>13</sup> From the docking studies it was noted that the key interactions observed in sitagliptin; namely the four hydrogen bond interactions with Tyr662, Glu205 and Glu206 resulting from the (*R*)-amino group were still preserved along with the water molecule bridge present between the amide carbonyl of **2a** with Tyr547. Phe357 provides a  $\pi - \pi$  interaction with the triazole core of compound **2a**. However, it was noticed that the inclusion of the sterically demanding tricyclic portion of **2a** led to a change in the orientation of the  $CF_3$  group, losing the known interaction of this moiety in sitagliptin with Arg358 and Ser209. However, we felt that this potential loss in activity would be compensated by allowing us the exciting potential for discovering new interactions within the DPP-4 protease backbone (Figure 3).




**Figure 3.** Compound **2a** (blue) docked into the DPP-4 active site (pdb code 1X70) overlaid with sitagliptin (magenta). Only one of the two possible diastereoisomers of **2a** with the

optimal docking pose is shown. The image was generated using PyMol.

The novel scaffold **1** (represented as examples **9a** and **9b**) was synthesized in a robust six step racemic sequence starting from commercially available ethyl 4-oxopiperidine-1-carboxylate **3**. The first step in the sequence was the enamine alkylation with ethyl acrylate under Dean Stark conditions that resulted in a high yield of the  $\delta$ -keto ester **4**.<sup>14</sup> Conversion of **4** into the *O*-methyl oxime **5** using methoxyamine hydrochloride in pyridine, delivered the desired compound in a 1:1.5 mixture of imine isomers. These were converted to the bicyclic lactam **6** using Raney Nickel in 7*N* ammonia in methanol under an atmosphere of hydrogen. At this stage the isomers generated in the ring closure were separated by flash chromatography on silica gel. Unfortunately, no assignment of the ring geometry was possible for the two separated lactam diastereoisomers (**7a** and **7b**) from NMR spectroscopy. Therefore single crystal X-ray crystallography was needed to determine the relative configuration of the two isomers.<sup>15</sup> Each of the isomers (**7a** and **7b**) was converted into their corresponding thiolactams (**8a** and **8b**) using Lawesson's reagent.<sup>16-17</sup> These thiolactams were converted into the corresponding tricyclic triazoles (**9a** and **9b**) by refluxing in toluene with 2,2,2-trifluoroacetylhydrazide (Scheme 1).

**Scheme 1.** Synthetic route to compounds **9a** and **9b**.<sup>a</sup>




<sup>a</sup>Reagents and conditions: a) 1. Pyrrolidine, benzene, rf, Dean Stark; 2. Ethyl acrylate, benzene, rf, o.n., 3. water, 2h, rf, 69% (2 steps); b)  $MeONH_2 \cdot HCl$ , pyridine, 84%; c) Raney-Nickel, 7*N* ammonia in MeOH,  $H_2$ , o.n., 86%; d) flash chromatography ethyl acetate/hexane/methanol (10:1:1), ratio 1:4; e) Lawesson's reagent, toluene, rf, o.n., 90-94%. f)  $CF_3$ -hydrazide, toluene, 1-3d, 120°C, 40-84%. Compounds **9a** and **9b** exist as a 1:1 mixture of enantiomers.

The molecular scaffolds (**9a** and **9b**) were inserted in the sitagliptin structural motif by deprotection of the carbamates (**9a** and **9b**) in ethanol/water using potassium hydroxide under refluxing conditions to afford the key molecular scaffolds (Scheme 1: **1a** and **1b**,  $R_1 = H$ ,  $R_2 = CF_3$ ) ready for reaction with the commercially available acids **10** and **11**. This coupling was high yielding and the final

deprotection to the HCl salts (**2a-b** and **12a-b**) was carried out with 4N HCl in dioxane (Scheme 2).

**Scheme 2. Synthetic route to compounds 2a-b and 12a-b<sup>a</sup>.**



<sup>a</sup>Reagents and conditions: a) 1) KOH, water/ethanol, rt, o.n., 2) **10** or **11** + EDCI, HOBt, DMF, rt, o.n., 77-91%, b) 4N HCl in dioxane. Compounds **2a-b** and **12a-b** exist as a 1:1 mixture of either *cis*- or *trans*- diastereoisomers.

As the synthesis of the scaffolds **9a** and **9b** was in racemic form, the separation of the diastereoisomers of compounds **2a** and **2b** was required. Rewardingly, both diastereomers proved separable under chiral HPLC conditions with purity of at least 99% *de*. Therefore **2a** was separated into the single diastereomers (**13** and **14**) and **2b** was separated as single diastereomers (**15** and **16**). Due to the robust nature of the synthetic procedure, a small series of further analogues was also produced (examples **17**, **18**, **19** and **20**, see supplementary information for the synthesis and yields). Inhibitory activity against DPP-4 was then determined (Table 1).<sup>18</sup>

**Table 1. DPP-4 inhibitory activity**

Compound	<i>cis</i> - or <i>trans</i> -	R <sub>3</sub>	R <sub>2</sub>	DPP-4 IC <sub>50</sub> [nM] <sup>a</sup>
<b>12a</b>	<i>cis</i> -	H	CF <sub>3</sub>	695±31 <sup>b</sup>
<b>12b</b>	<i>trans</i> -	H	CF <sub>3</sub>	2801±86 <sup>b</sup>
<b>13</b>	<i>cis</i> -	F	CF <sub>3</sub>	28±1 <sup>c</sup>
<b>14</b>	<i>cis</i> -	F	CF <sub>3</sub>	70±2 <sup>c</sup>
<b>15</b>	<i>trans</i> -	F	CF <sub>3</sub>	145±5 <sup>c</sup>
<b>16</b>	<i>trans</i> -	F	CF <sub>3</sub>	537±23 <sup>c</sup>
<b>17</b>	<i>cis</i> -	F	Ethyl	85±3 <sup>b</sup>
<b>18</b>	<i>trans</i> -	F	Ethyl	108±6 <sup>b</sup>
<b>19</b>	<i>cis</i> -	F	CF <sub>2</sub> CF <sub>3</sub>	67±2 <sup>b</sup>
<b>20</b>	<i>trans</i> -	F	CF <sub>2</sub> CF <sub>3</sub>	258±11 <sup>b</sup>


<sup>a</sup>Sitagliptin IC<sub>50</sub> 22±2 nM (n=20); <sup>b</sup>1:1 mixture of diastereomers; <sup>c</sup>single diastereomer with unknown absolute configuration.

Gratifyingly, the compounds demonstrated very good levels of DPP-4 activity when compared to sitagliptin (IC<sub>50</sub> 22±2 nM). Notably, in all the tested analogues the *cis*-isomer reproducibly showed a better level of biological activity when compared to the *trans*-isomer (for example compare **12a** and **12b** and **17** and **18**). As this is an observation throughout all the synthesized derivatives it demonstrates the effect of the connection of the rings to be important to allow the *cis*-fused compounds to establish optimal interactions within the active site of DPP-4. It was also encouraging to see that the separated diastereoisomers of **2a** and **2b** were shown to have either a 2.5-fold (*cis*-isomers, compare compounds **13** and **14**) or 3.6-fold (*trans*-isomers compare compounds **15** and **16**) difference in activity, demonstrating further molecular recognition for either the novel *cis*-fused or *trans*-fused tricyclic scaffolds within the DPP-4 active site. This was in agreement with our original docking studies (see figure S1 supplementary material) where the **2a cis**-isomers (**13-14**) were observed to have a better topographical fit into the DPP-4 active site, preserving the water molecule bridge present between the amide carbonyl of **2a** with Tyr547. However, from the docking work the **2b trans**-isomers (**15-16**) were shown to lack this key interaction and, as a consequence, we proposed they would possess lower DPP-4 inhibitory activity.

The enlargement of the CF<sub>3</sub> group in **2a** to both the ethyl-substituted analogue **17** and the CF<sub>2</sub>CF<sub>3</sub>-substituted analogue **19** led to a decrease in inhibitory activity. The lack of the three fluorine atoms in the aromatic region (**12a** and **12b**) substantially decreased the activity, which concurred with similar observations previously reported by the Merck group.

In order to show the influence of the rigid tricyclic ring system towards DPP-4 inhibitory activity, the bicyclic compound **24** was synthesized. The commercially available amine **21** was converted to the substituted 1,2,4 triazole **23** in 15% yield.<sup>19</sup> Final deprotection of the carbamate group in **23** was carried out under established standard conditions using KOH in water/ethanol and the coupling to the final bicyclic compound **24** was achieved using acid **11**, followed by deprotection using 4N HCl in dioxane (Scheme 3).


**Scheme 3. Synthesis of the open chain compound 24<sup>a</sup>.**



<sup>a</sup>Reagents and conditions: a) Ac<sub>2</sub>O, CH<sub>2</sub>Cl<sub>2</sub>, NEt<sub>3</sub>, rt, o.n. 84%; b) 1) POCl<sub>3</sub>, CHCl<sub>3</sub>, pyridine, 2) CF<sub>3</sub>CONHNH<sub>2</sub>, CHCl<sub>3</sub>, 3) 2M HCl, 15% (3 steps); c) 1) KOH, water/EtOH, 2) **11**, EDCI, HOBt, DMF, rt, o.n., 93%; d) 4N HCl in dioxane.

In order to gain insight into the impact of ring size for DPP-4 inhibitory activity, the synthesis of the novel *cis*-fused hexahydro-6H-[1,2,4]triazolo[4',3':1,5]pyrrolo[3,2-c]pyridine analogue **31** was carried out. The synthesis to the 5-membered lactams (**28** and **29**) followed a slightly different approach than for the 6, 6-membered lactams **7a-b**. Commercially available ethyl 4-oxopiperidine-1-carboxylate **3** was converted into *tert*-butylester **25** using LDA and *tert*-butyl 2-bromoacetate. The *tert*-butylester **25** was converted through to the substituted benzylamine **26** via a reductive amination reaction using benzylamine and sodium triacetoxyborohydride in 1,2-dichloroethane. Compound **26** was trans-esterified with 0.6M HCl in methanol to yield the methyl ester **27**, which was catalytically-hydrogenated using Pd/C in MeOH under an atmosphere of hydrogen. The final ring closure to the key bicyclic lactams **28** and **29** was carried out with potassium carbonate in methanol and the diastereomers were separated by flash column chromatography. Once more structural determination of the separated diastereomers (**28** and **29**) was performed through X-ray crystallography [see reference 15] (Scheme 4).


**Scheme 4. Synthesis of the 6,5-bicyclic intermediates 28 and 29.**



<sup>a</sup>Reagents and conditions: a) 1) LDA, THF, -78°C, 30 mins 2) *tert*-butyl 2-bromoacetate, -78°C to rt, 50%, b) benzylamine, NaBH(OAc)<sub>3</sub>, 1,2-dichloroethane, 94%, c) 0.6M HCl in MeOH, d) 1) Pd/C, MeOH, 2) K<sub>2</sub>CO<sub>3</sub>, MeOH, 45% (3 steps).

The separated *cis*-fused isomer **29** was converted through to the final product through the standard synthetic procedure (Scheme 5).


**Scheme 5. Synthesis of the 6,5,5-tricyclic compound 33<sup>a</sup>.**



<sup>a</sup>Reagents and conditions: a) Lawesson's reagent, toluene, rf, o.n., 66%, b) CF<sub>3</sub>-hydrazide, toluene, 1-3d, rf, 70%, c) KOH, water/ethanol, rf, o.n., 50 %, d) **11** + EDCI, HOBt, DMF, rt, o.n., 37%, e) 4N HCl in dioxane. Compound **31** exists as a 1:1 mixture of *cis*-diastereoisomers.

The DPP-4 inhibitory activity for compounds **24** (IC<sub>50</sub> 100±4 nM) and **31** (IC<sub>50</sub> 94±4 nM) was established showing a reduced level of activity compared to **13** (separated most active isomer of compound **2a**, IC<sub>50</sub> 28±1 nM). In order to understand the DPP-4 inhibitory activity of compounds **2a**, **24** and **31**, the compounds were energy minimized and docked into the known crystal structure of sitagliptin in DPP-4. As previously mentioned, compound **2a** displayed a good topographical fit into the enzyme pocket, whereas compound **31** docked to allow overlay of the trifluoromethyl group of **31** with that of sitagliptin. However, in achieving this topographical fit, the Phe357 π-π interaction with the triazole core of compound **31** is lost due to a steric clash imposed by the rigid tricyclic ring system and this could be an explanation for the observed loss in DPP-4 inhibitory potency of **31** when compared to **2a**. For the bicyclic compound **24**, once more there is a good topographical fit into the enzyme pocket, however the Phe357 π-π interaction with the triazole core

of compound **24** is not present and this could again be an explanation for the observed loss in DPP-4 inhibitory potency (Figure 4).



**Figure 4.** Compounds **2a**, **24** and **31** (blue) docked into the DPP-4 active site (pdb code 1X70) shown overlaid with sitagliptin (magenta). For compounds **2a** and **31** only the single enantiomer with the optimal docking pose is shown. The image was generated using PyMol.

In summary, we have shown the successful synthesis of promising new inhibitors for DPP-4 along with preliminary docking studies of the active compounds into the active site of the protease. The compounds demonstrated a range of activities with compound **13** (unknown absolute conformation) possessing similar levels of DPP-4 inhibitory activity to that of sitagliptin. We have shown the stereochemical preference for the *cis*- diastereomer of the novel octahydro-[1,2,4]triazolo[4,3-a][1,6]naphthyridine tricyclic ring system and we are currently investigating a chiral synthesis of the key intermediates along with further structure design-based synthesis of analogues to interrogate the SAR within this interesting new tricyclic scaffold which we will report in due course.

## ASSOCIATED CONTENT

**Supporting Information.** Preparation and full characterisation of the compounds is given “This material is available free of charge via the Internet at <http://pubs.acs.org>.”

## AUTHOR INFORMATION

### Corresponding Author

\* michael.stocks@nottingham.ac.uk

### Author Contributions

The manuscript was written through contributions of all authors.

## ACKNOWLEDGMENT


We acknowledge Dr Jinqiao Wan from HitGen for his help in arranging the assaying of the compounds. Dr Simon Hirst and Stephanie Barlow (Sygnature Discovery) for their assistance in the separation of the diastereomers using chiral HPLC

## ABBREVIATIONS

DPP-4 (Dipeptidyl peptidase 4); LDA (lithium diisopropylamide); SAR (structure Activity Relationship); WHO (World Health Organisation)

## REFERENCES





<sup>1</sup> <http://www.who.int/diabetes/en/>

<sup>2</sup> Nauck, M.; Stöckmann, F.; Ebert, R.; Creutzfeldt, W.; Reduced incretin effect in type 2 (non-insulin-dependent) diabetes. *Diabetologia*, **2006**, *29*, 46–52.

<sup>3</sup> Drucker, D.J.; Nauck, M. A.; The incretin system: glucagon-like peptide-1receptor agonists and dipeptidyl peptidase-4 inhibitors in type 2 diabetes, *Lancet*, **2006**, *368*, 1696–705.

<sup>4</sup> Thornberry, N. A.; Gallwitz, B.; Mechanism of action of inhibitors of dipeptidyl-peptidase-4(DPP-4) *Best Practice & Research Clinical Endocrinology & Metabolism*, **2009**, *23*, 479–486.

<sup>5</sup> Drucker, D.J.; Nauck, M. A.; The incretin system: glucagon-like peptide-1receptor agonists and dipeptidyl peptidase-4 inhibitors in type 2 diabetes, *Lancet*, **2006**, *368*, 1696–705.

<sup>6</sup> Orsakov, C.; Glucagon-Like Peptide 1, a New Hormone of theEn-teroinsular Axis. *Diabetologia* **1992**, *35*, 701-711.

<sup>7</sup> Herman, G. A.; Stein, P. P.; Thornberry, N. A.; Wagner, J. A. Dipeptidyl Peptidase-4 Inhibitors for the Treatment of Type 2 Diabetes: Focus On Sitagliptin. *Clin. Pharmacol. Therap.* **2007**, *81*, 761–767.

<sup>8</sup> Biftu, T.; Sinha-Roy, R.; Chen, P.; Qian, X.; Feng, D.; Kuethe, J. T.; Scapin, G.; Gao, Y. D.; Yan, Y.; Krueger, D.; Bak, A.; Eiermann, G.; He, L.; Cox, J.; Hicks, J.; Lyons, K.; He, H.; Salituro, G.; Tong, S.; Patel, S.; Doss, G.; Petrov, A.; Wu, J.; Xu, S. S.; Sewall, C.; Zhang, X.; Zhang, B.; Thornberry, N. A.; Weber, A. E. Omarigliptin (MK-3102): A Novel Long-Acting DPP-4 Inhibitor for Once-Weekly Treatment of Type 2 Diabetes. *J. Med. Chem.*, **2014**, *57*, 3205–3212.

<sup>9</sup> Chutian, S.; Hu, G.; Michael, S.; Chen, J.; Zhou, H.; Qi, Q.; Wang, F.; Ma, X.; Yang, X.; Zhang, G.; Ding, Y.; Zhou, D.; Peng, P.; Shih, C.; Xu, J.; Wu, F. Discovery of Imigliptin, a Novel Selective DPP-4 Inhibitor for the Treatment of Type 2 Diabetes, *ACS Med. Chem. Lett.* **2014**, *5*, 921–926.

<sup>10</sup> Juillerat-Jeanneret, L.; Dipeptidyl Peptidase IV and Its Inhibitors: Therapeutics for Type 2 Diabetes and What Else? *J. Med. Chem.* **2014**, *57*, 2197–2212.

<sup>11</sup> Ex: OMEGA version 2.4.6. OpenEye Scientific Software, Santa Fe, NM. <http://www.eyesopen.com>.

<sup>12</sup> Hawkins, P.C.D.; Nicholls, A. Conformer generation with OMEGA: learning from the data set and the analysis of failures *J. Chem. Inf. Model.* **2012**, *52*, 2919–2936.

<sup>13</sup> Kim, D.; Wang, L.; Beconi, M.; Eiermann, G. J.; Fisher, M. H.; He, H.; Hickey, G. J.; Kowalchick, J. E.; Leiting, B.; Lyons, K.; Marsilio, F.; McCann, M. E.; Patel, R. A.; Petrov, A.; Scapin, G.; Patel, S. B.; Roy, R. S.; Wu, J. K.; Wyvratt, M. J.; Zhang, B. B.; Zhu, L.; Thornberry, N. A.; Weber, A. E. (2R)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine: a potent, orally active dipeptidyl peptidase IV inhibitor for the treatment of type 2 diabetes. *J. Med. Chem.* **2005**, *48*, 141–151.

<sup>14</sup> Borne, R. F.; Fifer, K. E.; Waters, I. W.; Conformationally restrained fentanyl analogues. 2. Synthesis and analgetic evaluation of perhydro-1, 6-naphthyridin-2-ones. *J. Med. Chem.* **1984**, *27*, 1271–1275.

<sup>15</sup> Schwehm, C.; Lewis, W.; Blake, A. J.; Kellam, B.; Stocks, M. Preparation and structural analysis of (±)-cis-ethyl 2-sulfanylidenedecahydro-1, 6-naphthyridine-6-carboxylate and (±)-trans-ethyl 2-oxooctahydro-1H-pyrrolo[3,2-c]pyridine-5-carboxylate *Acta Cryst. C.* **2014**, *70*, 1161–1168.

<sup>16</sup> Thomsen, I.; Clausen, K.; Scheibye, S.; Lawesson, S.-O. Thiation with 2,4-Bis(4-methoxyphenyl)-1,3,2,4-Dithiadiphosphetane 2,4-disulfide: N-Methylthiopyrrolidone, *Org. Synth.; Coll. Vol. 7*, **1990** : 372–376.

<sup>17</sup> Occhiato, E. G.; Ferrali, A.; Menchi, G.; Guarna, A.; Danza, G.; Comerci, A.; Mancina, R.; Serio, M.; Garotta, G.; Cavalli, A.; DeVivo, M.; Recanatini, M.; Synthesis, Biological Activity, and Three-Dimensional Quantitative Structure–Activity Relationship Model for a Series of Benzo[c]quinolizin-3-ones, Nonsteroidal Inhibitors of Human Steroid 5 $\alpha$ -Reductase 1, *J. Med. Chem.* **2004**, *47*, 3546–3560.

<sup>18</sup> The compounds were assayed for their DPP-4 inhibitory activity at HitGen using the following method: DPP-4 recombinant human protein (purchased from Sino Biological Inc.) cleaves a non-fluorescent substrate, H-Gly-Pro-AMC (purchased from Bachem Americas, Inc), to release fluorescent, 7-Amino-4-Methyl Coumarin (AMC) (ex/em=360/460 nm). The initial rate of DPP-4 activity is measured over 15 min by following the fluorescent change at ex/em=360/460 nm, and the fits are inspected to insure that the reactions are linear to a correlation coefficient of 0.9. The resulting IC<sub>50</sub> values are obtained by fitting log (inhibitor concentration) vs. the percentage of remaining activity using four-parameter dose-response model (Sigmaplot Version 11.0).

---

<sup>19</sup> Long, L.Y.; Combs, A. P. Preparation of substituted fused aryl and heteroaryl derivatives as PI3K inhibitors, PT, Int. Appl. WO 2011075630.