

Hall, Charlotte L. and Taylor, John and Newell, Karen and Baldwin, Laurence and Sayal, Kapil and Hollis, Chris (2016) The challenges of implementing ADHD clinical guidelines and research best evidence in routine clinical care settings: a Delphi survey and mixed-methods study. *BJPsych open*, 2 (1). pp. 25-31. ISSN 2056-4724

Access from the University of Nottingham repository:

http://eprints.nottingham.ac.uk/31385/1/Published_2016_BJPOpen_Delphi.pdf

Copyright and reuse:

The Nottingham ePrints service makes this work by researchers of the University of Nottingham available open access under the following conditions.

- Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners.
- To the extent reasonable and practicable the material made available in Nottingham ePrints has been checked for eligibility before being made available.
- Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.
- Quotations or similar reproductions must be sufficiently acknowledged.

Please see our full end user licence at:

http://eprints.nottingham.ac.uk/end_user_agreement.pdf

A note on versions:

The version presented here may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the repository url above for details on accessing the published version and note that access may require a subscription.

For more information, please contact eprints@nottingham.ac.uk

The challenges of implementing ADHD clinical guidelines and research best evidence in routine clinical care settings: Delphi survey and mixed-methods study

Charlotte L. Hall,* John A. Taylor,* Karen Newell, Laurence Baldwin, Kapil Sayal and Chris Hollis

Background

The landmark US Multimodal Treatment of ADHD (MTA) study established the benefits of individualised medication titration and optimisation strategies to improve short- to medium-term outcomes in attention-deficit hyperactivity disorder (ADHD). This individualised medication management approach was subsequently incorporated into the National Institute for Health and Care Excellence (NICE) ADHD Clinical Guidelines (NICE CG78). However, little is known about clinicians' attitudes towards implementing these medication management strategies for ADHD in routine care.

Aims

To examine National Health Service (NHS) healthcare professionals' consensus on ADHD medication management strategies.

Method

Using the Delphi method, we examined perceptions on the importance and feasibility of implementing 103 ADHD treatment statements from sources including the UK NICE ADHD guidelines and US medication management algorithms.

Results

Certain recommendations for ADHD medication management were judged as important and feasible to implement, including a stepwise titration of stimulant medication. Other recommendations were perceived as important but not feasible to implement in routine practice, such as weekly clinic follow-up with the family during titration and collection of follow-up symptom questionnaires.

Conclusions

Many of the key guideline recommendations for ADHD medication management are viewed by clinicians as important and feasible to implement. However, some recommendations present significant implementation challenges within the context of routine NHS clinical care in England.

Declaration of interest

C.L.H. and K.S. were members of the Guideline Development Group for the NICE ADHD Clinical Guideline (NICE CG78).

Copyright and usage

© 2016 The Royal College of Psychiatrists. This is an open access article distributed under the terms of the Creative Commons Non-Commercial, No Derivatives (CC BY-NC-ND) licence.

Attention-deficit hyperactivity disorder (ADHD) is a common neurodevelopmental disorder, which left untreated can have a significant impact on the lives of affected young people, their families and society.^{1,2} In the UK, the National Institute for Health and Care Excellence (NICE) ADHD guidelines³ provide the gold standard for the diagnosis and management of ADHD, advocating medication as the first-line treatment for severe or moderate ADHD. The landmark US Multimodal Treatment of ADHD (MTA) study⁴ investigated the clinical effectiveness of a medication management strategy which involved a carefully monitored individualised titration period. This protocol resulted in significantly improved ADHD outcomes in the short to medium term.^{5,6} The Texas Children's Medication Algorithm Project^{7,8} was developed as a clinical tool to implement the MTA's medication protocol into routine clinical practice. Their findings indicated some deviance from the schedule, with a reluctance to titrate upwards and problems obtaining regular ratings from teachers and parents.⁸ These findings indicate that despite the MTA medication protocol being considered by many as the US 'gold standard' in medication management, the Texas project found that some items were not feasible to implement in routine clinical practice. However, given cross-country differences in clinical practice, including differences in the delivery of 'community

care' for ADHD and healthcare systems between the USA and UK, these findings may not translate to UK practice. Furthermore, since the MTA study which was conducted in the 1990s, prescribing practices have changed, with the introduction of long-acting ADHD medication.

In the UK, there is evidence of variable implementation of NICE clinical guidelines in 'real-world' settings⁹ and the specific implementation challenges of the NICE ADHD guideline have been highlighted.¹⁰ More recently, audit data collected within the East Midlands region of the UK have demonstrated poor adherence to NICE guidelines, particularly with regard to measuring blood pressure/heart rate, formal recording of side-effects and the use of rating scales to aid assessment and management.¹¹ These findings indicate a gap between best practice guidelines and implementation of recommendations in 'real-world' practice that needs to be understood and addressed. Given the difficulties in implementing the Texas Algorithm (based on the MTA protocol) in the USA, variable implementation of NICE guidelines in general in the UK and the evidence of poor adherence to aspects of the NICE ADHD guideline, we decided to investigate how the MTA study-based practice parameters and NICE guideline recommendations are perceived by National Health Service (NHS) clinicians with respect to both importance and feasibility for implementation.

We adopted a multi-method approach to determine UK stakeholder (parent/carer and NHS healthcare professional)

*Joint first authors.

perceptions of current and best practice clinical guideline recommendations for assessment and medical management of ADHD. Data were collected via an online survey, a parent/carer focus group and a healthcare professional's workshop. These findings were used to generate a set of statements on ADHD medication management which were then rated for importance and feasibility using a Delphi survey method.

Method

Delphi survey development

The items for the Delphi survey were developed through four different methodological approaches. Initially a scoping literature review was conducted. Eight healthcare databases were searched (Allied and Complementary Medicine Database (AMED), British Nursing Index, Embase, HMIIC (Health Management Information Consortium), MEDLINE, PsycINFO, CINAHL and Health Business Elite) for papers published in English, using the keywords ADHD and medication prescribing/monitoring/guidelines/titration. Informal searching, including hand-searching of article references and web search engines (including Google and Google Scholar) was also undertaken. In addition, colleagues who have contact with patients with ADHD were also consulted with regard to further literature that might be available. On the basis of this literature search, NICE guidelines were shown as the most prominent and 'gold-standard' approach to inform medication monitoring of ADHD in the UK. The MTA and Texas Algorithm were shown to be one of the most cited and influential sources for ADHD practice parameters in the USA. Many of the MTA principles are incorporated within the NICE guideline recommendations. However, the Texas Algorithm provides more specific details on aspects of titration and optimisation than do NICE guideline recommendations and hence we include statements from both sources. The findings from the literature search were used to inform questions for an online survey aimed at establishing healthcare professionals' perceptions of medication prescribing and monitoring.

Online survey

Eighty-five healthcare professionals responsible for the treatment of children and young people with ADHD across four participating NHS trusts (health provider organisations) within the regional Collaboration for Leadership in Applied Health Research and Care (CLAHRC) footprint were invited to participate in the online survey. Consent to participate was implied by survey participation. Responses were obtained from 26 healthcare professionals (response rate 31%), 12 of whom were community paediatricians (46%) and 14 child and adolescent psychiatrists (54%) working in NHS trusts across three English counties (Nottinghamshire ($n=12$), Derbyshire ($n=9$) and Lincolnshire ($n=5$)). One respondent was excluded from analysis because of low question completion within the survey (1% completion), resulting in a sample of 25 healthcare professionals. Although this response rate was relatively low, it is similar to that found in other online surveys aimed at healthcare professionals working in ADHD.¹²

The survey investigated healthcare professionals' current practice in diagnosis, initiation of medication, titration, assessment of response to medication and maintenance, which were based on strategies derived from NICE guidelines, the MTA group and the Texas Children's Medication Algorithm Project.

The online survey questions and semi-structured focus groups were devised by drawing on the current literature and consultation within the study group, which included three consultant psychiatrists, one child and adolescent mental health nurse and two researchers

with expertise in qualitative methods. Each member reviewed the questions to ascertain their appropriateness to the research question.

Parent/carer focus group

To assess parents/caregivers' perceptions of medication management for children with ADHD, nine parents/carers of children currently receiving medication for ADHD attended a focus group, having read an information sheet and provided written consent prior to their participation.

The focus group was semi-structured and consisted of 12 questions pertaining to the families' experiences of starting medication, how and when they were monitored, how information was fed back to them, the effect of the medication and their perceptions of the utility and feasibility of changes to medication management. Audio recordings from the focus group were anonymised, transcribed verbatim and thematically analysed in accordance with the guidelines of Braun & Clarke.¹³ Themes were derived inductively and verified via interrater reliability measures. The researchers utilised an essentiality/realist paradigm¹³ that sought to understand medication management through the words of the participants, as opposed to the researchers' co-created meaning.

Healthcare professionals' workshop

To further elucidate healthcare professionals' views of medication, a workshop was conducted with 22 healthcare professionals. The sample comprised 11 consultant child and adolescent psychiatrists, 4 nurses/nurse specialists, 5 community paediatricians and 2 NHS managers. Consent to participate was implied by attendance. From this sample, five groups were created to discuss different issues which concerned the decision to medicate and choice of drug, titration, medication maintenance, rating scales/outcomes, treatment adherence and engagement with the aim of identifying current and best practice. Verbal feedback from the healthcare professionals' discussions were collated via written notes and summarised.

Delphi study

The Delphi method is an interactive method utilising an expert panel answering questions in two or more rounds to achieve consensus. Participants are given feedback on question responses after each round; the process terminates once a predefined stopping point is reached.^{14,15} In this study, the aim of the Delphi process was to determine expert consensus on medication management strategies. The combination of multiple methodologies used to inform the Delphi survey ensured that items were comprehensive and relevant to aspects of medication management identified as relevant by multiple stakeholders.

Expert panel

The expert panel comprised 12 healthcare professionals working within the four participating NHS trusts: 7 child and adolescent psychiatrists, 3 consultant community paediatricians and 2 ADHD nurse prescribers, all experienced in the assessment and treatment of ADHD. Consent to participate was implied by acceptance of the invitation to form part of the expert group.

Procedure

Healthcare professionals were invited by email to become a member of the expert group. Statements for the Delphi study were drawn from the findings of the online survey, the parent/carer focus group, the healthcare professional workshop, the NICE guidelines,³ the MTA Cooperative Group^{4,16} and the Texas Children's Medication Algorithm Project.^{7,8} Some additional statements were added by the study team to determine clinical opinion on treatment choices or less directive recommendations. For example, reasons to choose atomoxetine and titration doses

should be increased as required up to British National Formulary (BNF) limits (see Table 1 for statement source). The final Delphi questionnaire comprised 103 statements, 92 of which were to be rated on both their importance and feasibility. A subset of statements ($n=11$) pertaining to clinician choice/preference and where issues of feasibility were not applicable were rated on importance alone. A 9-point Likert scale was used as a response option for each statement, with scores 7–9 denoting the statement is ‘important’ and/or ‘feasible’ and 1–3 denoting the statement is ‘unimportant’ and/or ‘unfeasible’.

The expert group completed round one of the questionnaire online using www.surveymonkey.com and individualised questionnaires were emailed to panelists for round two. In the second round, respondents were given the option to either retain their original score or modify it in light of the group median score. Similar to other studies, the survey was performed exclusively online.^{17,18} In line with the guidance of Jones & Hunter,¹⁹ all feedback was anonymous.

Data analysis methodology

Cut-off points for consensus drew on convention/standards adopted by Morrison & Barratt¹⁵ and Langlands *et al.*²⁰ however a broader percentage gap was used to determine whether a statement should be re-rated, to minimise the influence of a smaller number of participants in this study. The following criteria were employed:

1. Items were considered to have met criteria for consensus if 80% or more of the panel (10/12 members) rated the statement as 7, 8 or 9 (important/feasible) or as 1, 2 or 3 (unimportant/unfeasible).
2. Items were re-presented in the second round if 50–79% of the panel rated an item as 7–9 or 1–3.
3. Items were omitted from the second round if less than 50% of the panel rated an item as 7–9 or 1–3.

Round one resulted in 43 statements meeting criteria for consensus on importance and/or feasibility, four of which were rated unimportant and/or unfeasible. Fifty-five statements were re-rated on importance and/or feasibility during round two.

Ethical approval was granted by the local Research Ethics Committee and Research and Development Departments of Nottinghamshire Healthcare NHS Trust, Nottingham University Hospitals NHS Trust, Lincolnshire Partnership NHS Foundation Trust and Derbyshire Healthcare NHS Foundation Trust.

Results

Four different categories were used to cluster statements:

- Category 1: statements rated as important *and* feasible
- Category 2: statements rated as important but *not* feasible
- Category 3: statements rated as feasible but *not* important
- Category 4: statements rated as *neither* important *nor* feasible

For the purpose of this paper, we only report on statements in category 1 or 2 as these two categories allow an understanding of what clinicians consider to be important in ADHD practice as well as highlighting difficulties in implementing these aspects into routine care. At the end of the Delphi study, 24 statements reached consensus as being both ‘important’ (important/very important) and ‘feasible’ (feasible/very feasible), supporting the inclusion of these in ADHD medication management protocol (Table 1). It should be noted that there are some instances where the final median score for a statement was high, but criteria for consensus

(≥ 80 agreement) were not met. For example, a median score of 8, whereby 9/12 panelists scored 7–9 (statement 60).

ADHD assessment

The Delphi study revealed a consensus on both the importance and feasibility of assessing and describing impairment level alongside the NICE recommendation to document comorbid conditions (Table 1, statements 1 and 5). Whereas 80% (20/25) of survey respondents agreed that it was important to distinguish between the NICE ADHD guideline descriptions of moderate and severe ADHD during assessment, only 12% (3/25) of respondents thought that this was an easy distinction to make in practice. The possible difficulty in assessing ADHD-related impairment may potentially be exacerbated by its poor construct definition,^{21–23} the need to assess impairment across multiple domains and disentangle impairment related to comorbid conditions, as well as the weak association between symptom severity and severity of impairment.^{22–24}

Decision to medicate, drug choice and information provision to parents

The Delphi study revealed support for the NICE guideline that drug treatment should be offered to all children with severe ADHD (Table 1, statement 6) and that methylphenidate should normally be considered as the first line of drug treatment (Table 1, statement 15). However, whereas clinicians supported the clinical importance of the NICE guideline recommendation to offer behavioural advice alongside drug treatment (Table 2, statement 7) or to offer behavioural therapy prior to medication in cases of mild/moderate ADHD (Table 2, statement 9), there was no consensus on the feasibility of this. This may reflect a lack of resources within clinics; many NHS clinics are not commissioned or resourced to provide behavioural management advice or evidence-based parent programmes and so NHS healthcare professionals need to recommend that parents seek out this element of treatment from other agencies. This finding highlights the difficulty the NHS may have in implementing NICE ADHD guidance for access to behavioural interventions, suggesting that the recommendations were perhaps more aspirational than a reflection of what could currently be delivered in practice.

The importance and feasibility of discussing medication benefits and adverse effects with families also obtained consensus within the Delphi study (Table 1, statement 25) which was also highlighted as important by parent/carers in the focus group.

Medication initiation and titration

A consensus was reached in the Delphi study on both the importance and feasibility of NICE recommendations to document physical parameters (blood pressure, heart rate, weight and height; Table 1, statement 43) and cardiac risk factors before starting medication (Table 1, statement 44), administering the lowest available dose when prescribing methylphenidate and titrating up to BNF limits (Table 1, statements 45 and 59). Documenting the titration schedule at initiation, specifying drug, dose, duration of each step and review date (Table 1, statement 56), and titrating with regular stepwise dose increases (Table 1, statement 48) also met consensus on importance and feasibility, suggesting acceptance of more protocol-driven, MTA-style, titration schedules. Consensus also emerged on the importance and feasibility of a maximum duration of 12 weeks for titration, with 2–4 weeks between dose increments (Table 1, statement 61), longer than the 4–6 weeks for titration recommended by NICE. Support in our study for a longer duration of titration may reflect clinician views on the challenges of assessing improvement within 4–6 weeks in the multiple outcomes recommended by NICE (symptom reduction, behaviour change,

Table 1 Category 1 Delphi statements (n=24) reaching consensus on both importance and feasibility							
Delphi statement number	Category 1 statements	Importance median score	Feasibility median score	Source: NICE	Source: MTA/Texas	Source: stakeholder	Source: study team
1	Baseline assessment of ADHD should include a description of impairment (severe/mild-moderate).	9	8	✓			
5	Baseline assessment should routinely document comorbid conditions	9	9	✓			
6	Drug treatment should be offered to all children with severe ADHD	9	8.5	✓			
43	A physical examination, including recording of pulse, blood pressure, height and weight should occur before starting medication	9	9	✓			
44	Cardiovascular risk factors should be asked about and documented	9	9	✓			
45	The starting dose of methylphenidate should be the lowest available dose of the prescribed formulation	8.5	9	✓			
49	Atomoxetine should be titrated on a 'milligram per kilogram' basis	7	7	✓			
50	The dose of atomoxetine should be increased 7 days after initiation	7.5	9	✓			
62	Side-effects should be recorded at each dose change by parents	8.5	9	✓			
86	Compliance with medication should be assessed by discussion with the parent/carer and the child/young person	9	9	✓			
95	Height should be measured every 6 months	9	9	✓			
96	Weight should be measured 3 and 6 months after drug treatment has started and every 6 months thereafter	9	9	✓			
97	Height and weight should be recorded on a growth chart	9	9	✓			
101	The continuing benefit of medication should be formally reviewed and a decision documented on an annual basis with input from the parent/carer, young person and school	9	9	✓			
14	Where drug treatment is considered appropriate, methylphenidate, atomoxetine or dexamfetamine should be used	9	n/a ^a	✓			
16	Reasons to choose atomoxetine should include unresponsive to methylphenidate or intolerable adverse effects	9	n/a ^a	✓			
13	Clinicians should check and document whether or not the child can swallow pills	7	7.3		✓		
48	Stimulant drugs should be titrated with regular step-wise dose increases to achieve the minimum effect dose	9	9		✓		
56	The titration schedule should be documented at initiation specifying drug, dose, duration of each step and review date	9	9		✓		
15	Methylphenidate should normally be considered the first-line drug for treatment of ADHD	9	n/a ^a		✓		
25	Before initiation of medication, different drug options in terms of benefits (e.g. different duration of action) and adverse effects should be discussed with parents and children (age >11)	9	9			✓	
28	Drug information sheets should include details of what parents should do if adverse effects occur	9	9			✓	
29	A medication protocol should include drug information sheets	9	8.5			✓	
61	The duration of the dose titration phase should be a maximum of 12 weeks (2-4 weeks between dose increments)	8	8			✓	
84	After the first 6 months (of medication maintenance) clinic follow-up should be every 6 months	8	9			✓	
24	A medication protocol should include a prescribing algorithm/flow chart	8	n/a ^a			✓	
59	During titration the dose should be increased as required up to BNF limits	8	8				✓
32	Before initiation of medication, parents and young people should be asked to list separately the 3 most desired outcomes of treatment	8	8				✓
	Reasons to choose atomoxetine should include:						
17	Parent/child preference	7	n/a ^a				✓
18	History/risk of drug diversion	7	n/a ^a				✓
19	History/risk of substance misuse	7	n/a ^a				✓

ADHD, attention-deficit hyperactivity disorder; BNF, British National Formulary; MTA, Multimodal Treatment of ADHD; n/a, not applicable; NICE, National Institute for Health and Care Excellence; Texas, Texas Children's Medication Algorithm Project.
a. Additional statements (n=7) rated on importance only.

educational/relationship improvement) within a patient group typified by the presence of comorbidities²¹ and where teacher/parent feedback may conflict.²⁵ Furthermore, healthcare professionals from the workshop reported difficulties with patients not

attending follow-up appointments regularly, thus potentially delaying clinical decision-making.

Our findings suggested some conflict between the frequencies of contact deemed most beneficial as opposed to that deemed most

Table 2 Category 2 Delphi statements (n=19): rated as 'important' but 'not feasible'

Delphi statement number	Category 2 statements	Importance median score	Feasibility median score	Source: NICE	Source: MTA/Texas	Source: stakeholder	Source: study team
7	Drug treatment should only be offered to children together with behavioural management advice	8.5	6.5	✓			
9	Children with mild/moderate ADHD should be offered behaviour therapy/parent training before a trial of medication is considered	7.5	5	✓			
54	Monitoring of improvement/adverse-effects using telephone/SMS or email contact with the parent/young person should occur weekly during the titration phase	7.5	4.5	✓		✓	
83	During the first 6 months (of medication maintenance) clinic follow-up should be every 3 months	8.5	8	✓			
98	Heart rate and blood pressure should be monitored after each dose change	9	9	✓			
100	Heart rate and blood pressure should be recorded on a centile chart	8.5	6	✓			
26	Information sheets describing all available potential ADHD medications should be routinely provided to parents, young people (>11) and teachers	8.5	6.5			✓	
27	An information sheet on the medication to be prescribed should be routinely provided to parents, young people (>11), GPs and teachers	9	7.8			✓	
30	A designated individual (teacher/SENCO) within a school should be identified to provide feedback and liaison	8	6			✓	
31	For children in secondary school, more than one teacher's view should be acquired	7.8	5			✓	
42	A side-effect rating scale should be completed before initiation of medication	7.3	6			✓	
52	During titration, the child should be seen in clinic on a monthly basis	8	7			✓	
10	All pre-school children with ADHD should be offered behaviour therapy/parent training before a trial of medication is considered	9	4				✓
12	In children with comorbid ODD/CD or mood disorders, ADHD medication should be used together with psychological interventions	8	7				✓
34	The Conners short form (26 item) should be completed by parents at baseline (pre-medication)	9	8				✓
37	The Conners short form (26 item) should be completed by teachers at baseline (pre-medication)	8.5	5.5				✓
40	The Conners short form (26 item) self-report should be completed by young people (>11 years) at baseline (pre-medication)	7	5				✓
73	At the end of the titration phase the side-effect scale should be repeated by parents	7	6				✓
76	At the end of the titration phase the Conners short form (26 item) should be repeated by TEACHERS	7	3				✓

ADHD, attention-deficit hyperactivity disorder; CD, conduct disorder; GP, general practitioner; MTA, Multimodal Treatment of ADHD; NICE, National Institute for Health and Care Excellence; ODD, oppositional defiant disorder; SENCO, special educational needs coordinator; Texas, Texas Children's Medication Algorithm Project. The statements are based on information from their ascribed source; they have not been taken verbatim from each source.

feasible during titration. Consensus emerged on the importance of monthly clinic contact and weekly telephone/email contact with the family during titration and these schedules of contact were perceived as beneficial by 91% (20/22) and 82% (18/22) of online survey respondents respectively. However, there was no agreement on the feasibility of this approach (Table 2, statements 52 and 54).

Assessing response to medication: use of rating scales and assessment of side-effects

Support for the importance and feasibility of the NICE recommendation for parents to record side-effects at each dose change emerged from the Delphi study (Table 1, statement 62). However, 27% (6/22) of respondents from the online survey indicated that they did not use side-effect rating scales. Healthcare professionals commented at the workshop that assessment of side-effects was primarily achieved by monitoring physical health and via clinical

enquiry; a lack of easily available and validated side-effect scales may be a barrier to their use.

Clinicians reached a consensus on the importance of the administration of the Conners' short form (26 item) questionnaire²⁶ with parents, teachers and the young person at baseline (Table 2, statements 34, 37 and 40). However, variation among healthcare professionals in the range and number of rating scales used in the UK has been previously reported²⁷ and clinical enquiry may be a preferred source of information: all online survey respondents (22/22) placed value on the parents' view, the child's view and the parents' perception of progress at school when determining response to medication.

Medication maintenance: frequency of clinic contact

NICE recommendations for annual reviews of medication, assessment of concordance with medication via discussion with parents

and patient, and monitoring of height and weight at fixed intervals after starting medication all reached consensus for both importance and feasibility in the Delphi study (Table 1, statements 101, 86, 95 and 96). Delphi study findings indicate consensus on the importance and feasibility of clinic follow-ups every 6 months, after the first 6 months of treatment (Table 1, statement 84). This aligns with results from the online survey, whereby biannual contact during maintenance was rated as beneficial and feasible by 77% (17) and 95% (21) of respondents respectively.

Delphi study responses met consensus on the importance of monitoring blood pressure and pulse after every dose change (Table 2, statement 98) and although the median score was high (9, very important) there was no consensus on the feasibility of this NICE recommendation, which healthcare professionals acknowledged as problematic to achieve in practice. This highlights an implementation challenge regarding obtaining recordings between clinic visits. This may be an area where home or ambulatory recordings sent electronically to the clinic using a telemedicine approach may help to bridge this implementation gap.

The greater frequency of contact observed in the MTA's medication management arm provided opportunity to identify and immediately adjust changes in medication needs during maintenance, indicating a need for long-term close monitoring.²⁸ Additional forms of communication (email, SMS, smart telephone apps) could help to ensure adequate assessment of medication response if intensive face-to-face clinic monitoring is not feasible and may help to maintain the lines of communication desired by parents/carers.

Acquisition of feedback from school and importance of school input

The Delphi study revealed consensus for both the importance of having a nominated individual teacher to provide feedback and in the case of secondary school-age children, gathering information from more than one teacher (Table 2, statements 30 and 31). However, there was no consensus on the feasibility of this. Healthcare professionals from the workshop highlighted difficulties in getting school information, citing it was 'time-consuming'.

Obtaining teacher feedback relating to a child's response to treatment has been deemed 'critical'²⁹ and is a NICE recommendation. However, our findings support previous research indicating that resource limitations are a barrier to guideline implementation.³⁰

Discussion

Summary of main findings

Half (50%) of the NICE ADHD guideline recommendations included in the Delphi study were endorsed as being both clinically important and feasible to implement. These included assessment of ADHD-related impairment, baseline physical examination, recording of height, weight and parental reported side-effects, communicating reasons to use atomoxetine and methylphenidate (see Table 1 for complete list). Meanwhile, a subset of NICE recommendations, although regarded as clinically important, was identified as difficult or challenging to implement in routine UK clinical practice. These included delivery of adjunctive behavioural interventions, weekly monitoring (telephone/SMS/email) during titration, measuring pulse and blood pressure after every dose change during titration and recording pulse and blood pressure on a centile chart (see Table 2 for complete list).

Study strengths, limitations and future research

To the best of our knowledge, this is the first study which has explicitly sought to understand current consensus on medication management strategies for ADHD within the constraints of real-world practice. Whereas the number of Delphi panelists ($n=12$) may be considered relatively small, this is sufficient to demonstrate consensus. There is no consensus in the literature on optimum sample size³¹ and panel sizes ranging from 10 to 1142 participants are apparent in published Delphi studies.³² However, Jones & Hunter¹⁹ recommend a Delphi panel size of between 9 and 12 participants. Although the sample size of the online survey, workshop and focus groups were also small, the combination of opinion of multiple stakeholders to inform the items of the Delphi survey enabled a more complete approach to understanding issues pertinent to clinicians, young people, parent and carers in relation to ADHD medication management. Given that the participant is often over-looked in determining 'best practice', our inclusion of this group is a real strength.

This study only elicited the views of participants located within the East Midlands region of England; thus, it is possible that healthcare professionals working in different healthcare systems may make different judgements about the importance and feasibility of ADHD practice recommendations. Furthermore, the characteristics of the patients being treated by the healthcare professionals completing the Delphi study were not determined: it is likely that these too will have influenced judgements about what is important and feasible. Additionally, the use of self-report to determine current practices may have been subject to biases such as overestimation of adherence to guidelines³³ and triangulation with other objective assessments of patient management (e.g. case-note audits) would be a further method to characterise current practice.

Given the lack of understanding of current consensus in medication management in the UK, it was important to first establish current opinion on what constitutes 'best practice' within the constraints of everyday practice. Further research should establish the extent to which protocols are adhered to and factors that influence non-adherence by combining audit data with survey responses or qualitative interviews. Given the importance of regular contact during titration, future research could investigate whether utilisation of technology, such as Skype or apps, could provide a satisfactory interface for such monitoring. Although not specific to ADHD medication monitoring, a recent study did explore service user opinion of routine monitoring via an electronic questionnaire in mental health settings.³⁰ The findings revealed that parents and children were supportive of the use of technology to report progress to their healthcare professional.

Our study applied an iterative, mixed methods approach (literature review, online survey, stakeholder workshop and Delphi study) to determine previously unexplored perceptions of the implementation of national clinical guidelines and evidence-based recommendations for children and young people receiving medication for ADHD. Our findings indicate areas of agreement and discord with existing NICE³ ADHD guidelines and conflict between the perceived importance of some best practice medication management strategies and the feasibility of their implementation. The study has identified a set of ADHD practice recommendations that are viewed by clinicians as important but currently not feasible to implement. These recommendations should be the focus of health service implementation strategies and research.

Clinician feedback indicated that the intensive monitoring schedules of the MTA protocol are currently regarded as

impractical in routine UK practice and the time needed to titrate medication is likely to extend beyond that recommended by NICE guidelines. The use of technology to automate data collection and provide digital communication may be particularly relevant for aspects of practice viewed as important but currently unfeasible. The Delphi study findings have identified recommendations for practice where implementation challenges remain and clinical consensus is still to be reached. These findings could inform the update of the NICE ADHD clinical guideline and focus implementation strategies on those statements identified as important but not presently feasible to deliver in routine care. One potential area for development is the application of digital technologies to assist remote monitoring and follow-up during medication titration and optimisation.

Charlotte L. Hall, BSc, PhD; **John A. Taylor**, BSc, MSc, PhD; **Karen Newell**, BSc, MSc, CLAHRC, University of Nottingham, Nottingham, UK; **Laurence Baldwin**, BSc, MA, PhD, RN, RNT, CAMHS, Derbyshire Healthcare NHS Foundation Trust, Derbyshire, UK; **Kapil Sayal**, PhD, MRCPsych; **Chris Hollis**, PhD, FRCPsych, Developmental Psychiatry, Queen's Medical Centre, University of Nottingham, Nottingham, UK

Correspondence: Charlotte L. Hall, Institute of Mental Health, University of Nottingham, Innovation Park, Nottingham NG7 2TU, UK. Email: Charlotte.hall@nottingham.ac.uk

First received 29 Oct 2015, accepted 13 Dec 2015

Funding

This research was funded by the National Institute of Health Research (NIHR) Collaboration for Leadership in Applied Health Research and Care East Midlands (CLAHRC EM) and NIHR MindTech Healthcare Technology Co-operative. The work has been funded by the National Institute for Health Research. The views expressed in this publication are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health.

References

- Faraone SV, Biederman J, Spencer T, Wilens T, Seidman LJ, Mick E, et al. Attention-deficit/hyperactivity disorder in adults: an overview. *Biol Psychiatry* 2000; **48**: 9–20.
- Wilens TE, McDermott SP, Biederman J, Abrantes A, Hahsey A, Spencer TJ. Cognitive therapy in the treatment of adults with ADHD: a systematic chart review of 26 cases. *J Cogn Psychother* 1999; **13**: 215–26.
- National Institute for Health and Care Excellence. *Attention Deficit Hyperactivity Disorder: Diagnosis and Management*. Clinical guideline 72. NICE, 2008.
- MTA Cooperative Group. A 14-month randomized clinical trial of treatment strategies for attention-deficit/hyperactivity disorder. The MTA Cooperative Group. Multimodal Treatment Study of Children with ADHD. *Arch Gen Psychiatry* 1999; **56**: 1073–86.
- Swanson JM, Kraemer HC, Hinshaw SP, Arnold LE, Conners CK, Abikoff HB, et al. Clinical relevance of the primary findings of the MTA: success rates based on severity of ADHD and ODD symptoms at the end of treatment. *J Am Acad Child Adolesc Psychiatry* 2001; **40**: 168–79.
- Jensen PS, Garcia JA, Glied S, Crowe M, Foster M, Schlander M, et al. Cost-effectiveness of ADHD treatments: findings from the multimodal treatment study of children with ADHD. *Am J Psychiatry* 2005; **162**: 1628–36.
- Pliszka SR, Greenhill LL, Crismon ML, Sedillo A, Carlson C, Conners CK, et al. The Texas Children's Medication Algorithm Project: report of the Texas Consensus Conference Panel on medication treatment of childhood attention-deficit/hyperactivity disorder. Part I. Attention-deficit/hyperactivity disorder. *J Am Acad Child Adolesc Psychiatry* 2000; **39**: 908–19.
- Pliszka SR, Lopez M, Crismon ML, Toprac MG, Hughes CW, Emslie GJ, et al. A feasibility study of the Children's Medication Algorithm Project (CMAP) algorithm for the treatment of ADHD. *J Am Acad Child Adolesc Psychiatry* 2003; **42**: 279–87.
- Sheldon T, Cullum N, Dawson D, Lankshear A, Lowson K, Watt I, et al. What's the evidence that NICE guidance has been implemented? Results from a national evaluation using time series analysis, audit of patients' notes, and interviews. *BMJ* 2004; **329**: 999.
- Atkinson M, Hollis C. NICE guideline: attention deficit hyperactivity disorder. *Arch Dis Child Educ Pract Ed* 2010; **95**: 24–7.
- Hall CL, Walker GM, Valentine AZ, Guo B, Kaylor-Hughes C, James M, et al. Protocol investigating the clinical utility of an objective measure of activity and attention (QbTest) on diagnostic and treatment decision-making in children and young people with ADHD – 'Assessing QbTest Utility in ADHD' (AQUA): a randomised controlled trial. *BMJ Open* 2014; **4**: e006838.
- Hall CL, Newell K, Taylor J, Sayal K, Swift KD, Hollis C. 'Mind the gap' – mapping services for young people with ADHD transitioning from child to adult mental health services. *BMC Psychiatry* 2013; **13**: 186.
- Braun V, Clarke V. Using thematic analysis in psychology. *Qual Res Psychol* 2000; **3**: 77–101.
- Dalkey N, Helmer O. An experimental application of the DELPHI method to the use of experts. *Manage Sci* 1963; **9**: 458–67.
- Morrison AP, Barratt S. What are the components of CBT for psychosis? A Delphi study. *Schizophr Bull* 2010; **36**: 136–42.
- Greenhill LL, Abikoff HB, Arnold LE, Cantwell DP, Conners CK, Elliott G, et al. Medication treatment strategies in the MTA study: relevance to clinicians and researchers. *J Am Acad Child Adolesc Psychiatry* 1996; **35**: 1304–13.
- Elwyn G, O'Connor A, Stacey D, Volk R, Edwards A, Coulter A, et al. Developing a quality criteria framework for patient decision aids: online international Delphi consensus process. *BMJ* 2006; **333**: 417.
- Herdman M, Rajmil L, Ravens-Sieberer U, Bullinger M, Power M, Alonso J. Expert consensus in the development of a European health-related quality of life measure for children and adolescents: a Delphi study. *Acta Paediatr* 2002; **91**: 1385–90.
- Jones J, Hunter D. Qualitative research: consensus methods for medical and health services research. *BMJ* 1995; **311**: 376–80.
- Langlands RL, Jorm AF, Kelly CM, Kitchener BA. First aid recommendations for psychosis: using the Delphi method to gain consensus between mental health consumers, carers, and clinicians. *Schizophr Bull* 2008; **34**: 435–43.
- Bell AS. A critical review of ADHD diagnostic criteria: what to address in the DSM-V. *J Atten Disord* 2011; **15**: 3–10.
- Gathje RA, Lewandowski LJ, Gordon M. The role of impairment in the diagnosis of ADHD. *J Atten Disord* 2008; **11**: 529–37.
- Gordon M, Antshel K, Faraone S, Barkley R, Lewandowski L, Hudziak JJ. Symptoms versus impairment: the case for respecting DSM-IV's Criterion D. *J Atten Disord* 2006; **9**: 465–75.
- Yoshimasu K, Barbaresi WJ, Colligan RC, Voigt RG, Killian JM, Weaver AL, et al. Childhood ADHD is strongly associated with a broad range of psychiatric disorders during adolescence: a population-based birth cohort study. *J Child Psychol Psychiatry* 2012; **53**: 1036–10.
- Mitsis EM, McKay KE, Schulz KP, Newcorn JH, Halperin JM. Parent-teacher concordance for DSM-IV attention-deficit/hyperactivity disorder in a clinic-referred sample. *J Am Acad Child Adolesc Psychiatry* 2000; **39**: 308–13.
- Conners CK. *Conners Rating Scales-Revised (CRS-R)*. Multi-Health Systems, 1997.
- Tettenborn M, Prasad S, Poole L, Steer C, Coghill D, Harpin V, et al. The provision and nature of ADHD services for children/adolescents in the UK: results from a nationwide survey. *Clin Child Psychol Psychiatry* 2008; **13**: 287–304.
- Vitiello B, Severe JB, Greenhill LL, Arnold LE, Abikoff HB, Bukstein OG, et al. Methylphenidate dosage for children with ADHD over time under controlled conditions: lessons from the MTA. *J Am Acad Child Adolesc Psychiatry* 2001; **40**: 188–96.
- Francke AL, Smit MC, de Veer AJ, Mistiaen P. Factors influencing the implementation of clinical guidelines for health care professionals: a systematic meta-review. *BMC Med Inform Decis Mak* 2008; **8**: 38.
- Hall CL, Taylor J, Moldavsky M, Marriott M, Pass S, Newell K, et al. A qualitative process evaluation of electronic session-by-session outcome measurement in child and adolescent mental health services. *BMC Psychiatry* 2014; **14**: 113.
- Hsu C-C, Sandford BA. The Delphi technique: making sense of consensus. *Practical Assess Res Eval* 2007; **12**: 1–8.
- Akins R, Tolson H, Cole B. Stability of response characteristics of a Delphi panel: application of bootstrap data expansion. *BMC Med Res Methodol* 2005; **5**: 37.
- Adams AS, Soumerai SB, Lomas J, Ross-Degnan D. Evidence of self-report bias in assessing adherence to guidelines. *Int J Qual Health Care* 1999; **11**: 187–92.

