



Morling, Joanne R. and Fallowfield, Jonathan A. and Guha, Indra N. and Williamson, Rachel M. and Glancy, Stephen and Strachan, Mark W.J. and Price, Jackie F. (2015) Clinically significant chronic liver disease in people with type 2 diabetes: the Edinburgh Type 2 Diabetes Study. QJM: An International Journal of Medicine . ISSN 1460-2725

Access from the University of Nottingham repository:

http://eprints.nottingham.ac.uk/32085/1/2015_QJM_eprints.pdf

Copyright and reuse:

The Nottingham ePrints service makes this work by researchers of the University of Nottingham available open access under the following conditions.

- Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners.
- To the extent reasonable and practicable the material made available in Nottingham ePrints has been checked for eligibility before being made available.
- Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.
- Quotations or similar reproductions must be sufficiently acknowledged.

Please see our full end user licence at:

http://eprints.nottingham.ac.uk/end_user_agreement.pdf

A note on versions:

The version presented here may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the repository url above for details on accessing the published version and note that access may require a subscription.

For more information, please contact eprints@nottingham.ac.uk

Clinically significant chronic liver disease in people with type 2 diabetes: the Edinburgh Type 2 Diabetes Study

Joanne R. Morling¹, Jonathan A. Fallowfield², Indra N. Guha³, Rachel M. Williamson⁴, Maria Ali⁴, Stephen Glancy⁵, * Mark W. J. Strachan⁴, Jackie F. Price¹

* Mark W. J. Strachan and Jackie F. Price are joint senior authors.

¹Centre for Population Health Sciences, University of Edinburgh, Old Medical Buildings, Teviot Place, Edinburgh EH8 9AG, UK

²Queen's Medical Research Institute, University of Edinburgh, Edinburgh, UK

³NIHR Nottingham Digestive Diseases Biomedical Research Unit, University of Nottingham, Nottingham, UK

⁴Department of Metabolic Medicine, Western General Hospital, Edinburgh, UK

⁵Department of Radiology, Western General Hospital, Edinburgh, UK

Corresponding author: J. R. Morling

Edinburgh Type 2 Diabetes Study, Centre for Population Health Sciences, University of Edinburgh, Old Medical Buildings, Teviot Place, Edinburgh EH8 9AG, UK

Email: J.morling@ed.ac.uk

Tel. 0131 650 3244

Fax. 0131 650 3245

Word count: 2374 (abstract 244)

Tables and figures: 5

Online supplementary data: 0

Abstract

Background

Type 2 diabetes is an independent risk factor for chronic liver disease, however disease burden estimates and knowledge of prognostic indicators are lacking in community populations.

Aims

To describe the prevalence and incidence of clinically significant chronic liver disease amongst community-based older people with type 2 diabetes and to determine risk factors which might assist in discriminating patients with unknown prevalent or incident disease.

Design

Prospective cohort study

Methods

939 participants in the Edinburgh Type 2 Diabetes Study underwent investigation including liver ultrasound and non-invasive measures of NASH, hepatic fibrosis and systemic inflammation. Over 6-years, cases of cirrhosis and hepatocellular carcinoma were collated from multiple sources.

Results

Eight patients had known prevalent disease with 13 further unknown cases identified (prevalence 2.2%) and 15 incident cases (IR 2.9/1000person-years). Higher levels of systemic inflammation, NASH and hepatic fibrosis markers were associated with both unknown prevalent and incident clinically significant chronic liver disease (all $p < 0.001$).

Conclusions

Our study investigations increased the known prevalence of clinically significant chronic liver disease by over 150%, confirming the suspicion of a large burden of undiagnosed disease. The disease incidence rate was lower than anticipated, but still much higher than the general population rate. The ability to identify patients both with and at risk of developing clinically significant chronic liver disease allows for early intervention and clinical monitoring strategies. Ongoing work, with longer follow-up, including analysis of rates of liver function decline, will be used to define optimal risk prediction tools.

[244]

Introduction

Chronic liver disease (CLD) due to non-alcoholic fatty liver disease (NAFLD) in community populations represents a major challenge for general practitioners and a growing burden for healthcare services¹. Of the 25% of the UK population now categorised as obese, most will have NAFLD² and about 10% of these people have been diagnosed in community studies to have evidence of advanced liver fibrosis that leads to cirrhosis³. Of patients with cirrhosis (all cause), 5–20% will develop hepatocellular carcinoma (HCC) in 5 years⁴. Type 2 diabetes, which is increasing in frequency in parallel with obesity, is strongly associated with NAFLD but data on the progression to cirrhosis and HCC in community-based patients with diabetes is limited.

Three major independent reports have highlighted the need for the early detection of liver disease^{2, 5, 6}. The ability to identify patients with and at risk of developing clinically significant CLD (CS-CLD) would promote early intervention strategies and guide clinical follow-up, ensuring timely detection of cirrhosis where assimilation into surveillance programmes for HCC and varices as well as screening for cardiovascular disease has been shown to improve patient outcomes. Despite this, the existing diagnostic pathways for detection and onward referral of suspected CLD are based on traditional liver enzyme tests which lack accuracy and contribute to late diagnosis.

Using a population-based cohort of approximately 1000 patients with type 2 diabetes (The Edinburgh Type 2 Diabetes Study⁷) we employed an extensive screening programme (including liver ultrasound and non-invasive measures of non-alcoholic steatohepatitis [NASH], hepatic fibrosis and systemic inflammation) for clinically significant liver disease in order to investigate the true burden of disease. Over 6-years, cases of cirrhosis, HCC and gastro-oesophageal varices were collated from multiple sources.

Within the diabetic population annual liver enzyme checks are commonly performed, but there are no clear and consistent national guidelines on their application and interpretation, particularly in the community setting. Despite this, they remain the main trigger for referral to specialist liver clinics. Using standard liver enzyme assessment and liver ultrasound scan (USS) to diagnose hepatic steatosis, we identified patients at potential risk of CS-CLD and followed them up prospectively to collect data on clinical outcomes.

Our aim was to describe the prevalence and incidence of CS-CLD amongst older people with type 2 diabetes and to determine whether the detection of abnormal liver enzyme levels and/or the presence of hepatic steatosis on USS in people with type 2 diabetes might assist in discriminating those patients who subsequently go on to develop CS-CLD. Finally, we investigated a wide range of potential risk factors and biomarkers, including complications of diabetes and markers of advanced CLD, which might be useful in identifying at-risk subjects.

Methods

The Edinburgh Type 2 Diabetes Study (ET2DS)

Full methods of the ET2DS have been published elsewhere⁷. Briefly, in 2006/7 patients aged 60-75 years with type 2 diabetes were randomly selected from the Lothian Diabetes Register (LDR), a comprehensive register of patients with diabetes living in Lothian, Scotland, and invited to participate in the ET2DS.

1066 men and women attended a baseline research clinic and were enrolled into the study. This study population has been shown to be largely representative of all patients randomly selected from the LDR and therefore of the target population of older men and women with type 2 diabetes living in the general population⁸.

All survivors were invited to re-attend for clinical and liver investigation at year 1 (2007/8) and at year 4 (2010/11) with 939 (dead n=15, unable to contact n=19, unwilling/unable n=93) and 831 study participants attending, respectively.

Clinical and laboratory examination

Details of all the research clinics held at the Wellcome Trust Clinical Research Facility, Western General Hospital, Edinburgh, UK have been described previously^{7,9}. In brief, fasting venous blood sampling, physical examination and abdominal USS were undertaken.

A liver screen (including viral hepatitis serology, alpha-feto protein [AFP], liver autoantibodies, serum ferritin) was undertaken if the patient met the criteria for referral to specialist Hepatology services (see below) or if hepatic steatosis was present on USS and the participant had never previously had a liver screen.

All biochemical analytes were measured at the time of clinic attendance using a Vitros Fusion chemistry system (Ortho Clinical Diagnostics, Bucks, UK) at the Western General Hospital (Edinburgh, UK).

Biomarkers of liver injury

A wide range of hepatic biomarkers were assessed including those indicating non-specific liver injury (liver enzymes: alanine aminotransferase [ALT], aspartate aminotransferase [AST] and gamma-glutamyl aminotransferase [GGT]), steatosis (USS-graded liver fat¹⁰), NASH (cytokeratin-18 [CK18]), surrogate measures of advanced portal hypertension (spleen size and platelet count), and liver fibrosis (AST to Platelet Ratio Index [APRI]¹¹, AST:ALT ratio, Enhanced Liver Fibrosis [ELF] score¹², Fibrosis-4 score [FIB-4]¹³, hyaluronic acid [HA] and NAFLD Fibrosis Score [NFS]¹⁴).

CK18 and ELF were measured in stored serum samples (-80°C) as previously described¹⁵.

Abnormal liver enzymes were defined as: i) abnormal - greater than the upper limit of normal (ULN) – ALT >50U/L, AST >45 U/L, GGT >55U/L; ii) highly abnormal - greater than 2x ULN for ALT, AST, GGT; and iii) greater than recently proposed sex specific cut-offs for ALT – males >30U/L, females >19U/L¹⁶.

Serum marker panels for hepatic fibrosis were calculated as in original publications.

Identification of liver disease

Identification of liver disease is described in Figure 1. Definite (prevalent or incident) CS-CLD was a composite outcome defined as a diagnosis of cirrhosis, HCC (with confirmatory radiology) or gastro-oesophageal varices (with confirmatory endoscopy) recorded in the patients' medical records. Research study referral criteria for referral to specialist Hepatology services were designed in conjunction with an experienced consultant Hepatologist (Figure

1A), prevalent and incident CS-CLD was identified using a two-stage process across multiple sources (Figure 1B) and CS-CLD was defined as in Figure 1C.

Data analysis

Data were analysed using SPSS v19.0 (SPSS Inc., Illinois, USA). Variables that were not normally distributed were, where necessary, transformed and analysed on a logarithmic scale.

We describe the prevalence (both known and unknown) at the start of the study and subsequent incident CS-CLD. For initial analyses, all participants without known prevalent CS-CLD were included.

Exploratory analysis of potential risk factors and biomarkers associated with the development of advanced liver disease was undertaken using : patient characteristics, diabetes history and treatment, metabolic variables, established risk factors for CLD, markers of liver injury (non-specific, steatosis, NASH, advanced portal hypertension and liver fibrosis), and markers of systemic inflammation. Cox regression was used to investigate the association of potential risk factors with CS-CLD.

Ethical approval was obtained from the Lothian Research Ethics Committee and all subjects gave written informed consent.

Results

Subject characteristics

Participant attendance is shown in Figure 2. Clinical follow-up data was available for all 939 year 1 clinic attendees. Characteristics of the study population are shown in Table 1.

Prevalent clinically significant chronic liver disease

Eight patients had known prevalent CS-CLD and were excluded from the investigation of unknown prevalent disease (n=931, Figure 2A)¹⁷.

Following the year 1 investigations, 13.4% (125/931) of participants met the research study protocol criteria as high-risk for the presence of liver disease and were referred to Hepatology. Risk factors for referral are shown in Table 1.

Of these, 52 (41.6%) were offered an appointment at Hepatology clinic and the remainder returned to standard care after triage by a consultant Hepatologist. The majority of those not seen had low level titres of positive autoantibodies ($\geq 1:40$;) or isolated mildly elevated liver enzymes only ($< 2 \times \text{ULN}$).

Of those referrals seen, 31 (59.6%) were immediately discharged following initial assessment on the basis of having either simple fatty liver with low risk for the presence of significant liver fibrosis (according to non-invasive marker criteria) and/or false positive indicators of CS-CLD (e.g. platelets $< 150 \times 10^9/\text{L}$ but no portal hypertension) as determined by a consultant Hepatologist. Twenty-one participants (40.4) remain under active Hepatology follow-up.

As a consequence of the year 1 investigations, 13 (1.4%) patients were diagnosed with previously unknown prevalent CS-CLD (including one HCC, the remainder cirrhosis). The reasons for their initial referral were wide-ranging.

Risk factors associated with unknown prevalent cases were elevated liver enzymes (but not ALT), inflammatory markers, markers of NASH and hepatic fibrosis (all $p < 0.01$, Tables 2 and 3).

Overall the prevalence of CS-CLD was 2.2% (known plus unknown).

Development of incident clinically significant CLD

918 patients (Figure 2B) did not have prevalent CS-CLD.

Over a mean follow-up period of 5.6 years (sd 1.0, total 5156 person-years) there were 15 incident cases of CS-CLD, IR 2.9 /1000 person-years. These comprised cirrhosis $n=14$ (2.7 /1000 person-years), HCC $n=4$ (0.8 /1000 person-years), and gastro-oesophageal varices $n=5$ (1.0 /1000 person-years), with some patients having more than one complication.

Patients identified as potentially high risk after the year 1 investigations were more likely to develop CS-CLD than those not thought to have liver disease (IR 10.9 vs 1.8 /1000 person-years, $p=0.001$). However, of the incident 15 patients that developed CS-CLD less than half ($n=7$, 46.7%) were identified as high-risk by the extensive liver-related investigations at the year 1 clinic. Of the incident HCC, only 1 (25.0%) occurred in a participant identified through our year 1 investigations. The majority of cases of varices ($n=3$, 60.0%) were identified through entry into an endoscopic surveillance programme.

Rates of incident CS-CLD were significantly higher in those: seen in the Hepatology clinic (IRR 18.7, 95%CI 6.5-54.0, $p<0.001$), with abnormal liver enzymes (IRR 5.7, 95%CI 2.0-16.0, $p=0.001$) and with very abnormal liver enzymes (IRR 16.3, 95%CI 5.2-50.9, $p<0.001$).

Individuals with hepatic steatosis or elevated liver enzymes as defined by the lower revised laboratory reference ranges had incidence rates of CS-CLD not significantly different to those without (IRR 2.9, $p=0.096$ and IRR 1.2, $p=0.800$ respectively).

Patients with abnormal liver enzymes at the year 1 investigation were more likely to develop CS-CLD, (normal 7/776, 0.9%; abnormal 3/113, 2.7%; very abnormal 5/26, 19.2%; $p < 0.001$). In those developing CS-CLD, mean/median levels of liver enzymes were higher, however they remained broadly within normal laboratory limits for transaminases (ALT 44.4 U/L, AST 46.0 U/L), with median GGT levels above the upper limit of normal (median 62 vs 16 U/L, $p < 0.001$) (Table 2), overall 47% of those with incident CS-CLD had normal liver enzymes (Table 3).

Tables 2 and 3 show the associations between potential risk factors and incident CS-CLD. Known causes of liver disease (including alcohol) were similar in both groups ($p = 0.123$). Higher levels of markers of systemic inflammation were associated with an increased incidence of CS-CLD (ln CRP HR 1.66 $p = 0.026$, ln IL-6 HR 2.86 $p = 0.002$, ln TNF- α HR 2.12 $p = 0.017$)

Nearly all patients (80.0%) with incident CS-CLD had hepatic steatosis at the start of the study, although this was not statistically significant. All of the continuous markers of NASH and hepatic fibrosis (except AST:ALT ratio) were higher in those with incident CS-CLD than in those who did not.

Conclusions

We conducted the only large prospective community-based study of older patients with type 2 diabetes. The prevalence of CS-CLD at baseline was 2.2% (0.9% diagnosed clinically prior to enrolment and 1.4% identified by study investigations). The incidence of CS-CLD was 1.4% over nearly 6 years (IR 2.9 /1000 person-years).

Existing estimates of CS-CLD frequency are limited by reliance on confirmatory liver biopsy in secondary care populations or in community-based cohorts by the use of a single screening modality (e.g. FibroTest¹⁸, transient elastography¹⁹) each with their own challenges; for example liver enzymes have low diagnostic accuracy, non-invasive fibrosis markers (e.g. ELF, NFS) have poor positive predictive values for later stages of CLD^{12, 14}, ICD codes only identify hospitalised cases with decompensation, and biopsy has marked ascertainment bias. Accordingly, in the general population, the prevalence of advanced fibrosis in NAFLD ranged from 0-12% in a single study depending on the markers used²⁰, and there is even greater discordance between different studies on CS-CLD prevalence. Therefore, we used a multimodal approach to comprehensively phenotype our cases and maximize detection CS-CLD.

Our findings support the suggestion that NAFLD is an under-diagnosed chronic disease²¹ despite patients with type 2 diabetes having, as a minimum, annual clinical reviews including liver enzyme tests. Our multimodal diagnostic approach increased the prevalence by more than 150% through the diagnosis of unknown CS-CLD, of which almost 70% was attributable to NAFLD. However, it was a labour intensive process with 14% of the study participants referred to a consultant Hepatologist. In the majority of these patients, simple laboratory tests were abnormal (e.g. liver enzymes above the ULN, low platelet count) although, typically, values were only marginally elevated.

Reassuringly, despite a relatively high prevalence of uncomplicated NAFLD at year 1 there were only a small number of patients who went on to develop incident CS-CLD after 6 years (2.9 /1000 person-years). In 103 NAFLD patients followed by Adams et al²² for a mean of 3.2 years, the fibrosis progression rate was 0.35 stages/year in those with diabetes. This is high compared to our cohort as it equates to a potential 2-stage advancement over the 6 year follow-up of our cohort in those with NAFLD. In NASH populations, 32% of subjects progressed fibrosis score over 3 years²³ and 31% had progressive fibrosis²⁴, although no participant developed cirrhosis during 4 years of follow-up.

The only aspect of diabetes history to be associated with the development of CS-CLD was the use of insulin (HR 9.08). Hyperinsulinemia associated with T2DM is a risk factor for NASH, hepatic fibrosis and HCC. Insulin stimulates hepatic stellate cell proliferation and collagen synthesis in vitro and therefore, by extension, exogenous insulin therapy could promote liver fibrosis in vivo²⁵. Our data reinforces the known association between the presence of diabetes and increased the risk of HCC²⁶. Moreover, Scottish Cancer Registry data showed that within the whole Scottish population aged 60-75 the IR for liver-related cancer (including biliary) in 2012 was 0.3/1000 person-years²⁷, which is substantially lower than the rate for HCC alone in our cohort (0.8/1000 person-years).

Participants in the least deprived SIMD quintile had an incidence rate of CS-CLD 74% lower than those in the most deprived quintile. This unadjusted finding may represent confounding by other factors such as alcohol and obesity. Previous studies into the relationship between chronic liver disease and deprivation have found any associations were lost after adjustment for such risk factors²⁸.

Critically, for clinical practice, the majority of incident CS-CLD had normal liver function tests with no steatosis. Whilst those with abnormal liver enzymes per se were more likely to

develop CS-CLD, the mean levels of liver enzymes in those that did were still within the normal laboratory reference range. It may be that given the high prevalence of steatosis in the cohort that it may add little risk of CS-CLD beyond the risk conferred by type 2 diabetes alone.

The main limitation to this study is the lack of liver biopsy, although this represents an imperfect 'gold standard' test for the diagnosis and staging of NAFLD and is impractical in community studies with a low prevalence of CS-CLD. Additionally, we would argue that our non-invasive liver assessment is more applicable to clinical practice and is unlikely to have missed CS-CLD due to the use of validated cut-offs and robust triage for referral to a consultant Hepatologist.

This work suggests that there is little benefit in performing liver USS of patients similar to the study cohort, where the finding of hepatic steatosis could be predicted and provides no indication of future disease progression.

For the first time, we have evaluated the utility of extensive but simple targeted investigation using routinely available clinical tests in order to identify those at high risk of both immediate and future CS-CLD. We plan to conduct longer term follow-up to capture future incident liver related events. Of particular interest will be how the rate of change of potential biomarkers in earlier stages of CLD can be used to predict future development of CS-CLD.

Funding

The ET2DS was funded by a grant from the UK Medical Research Council [grant number R39788]. The liver substudy was additionally supported by a grant from Pfizer. CK18 testing was supported by Peviva. JRM was supported by a Diabetes UK Clinical Research Fellowship for 3 years [grant number R41481].

Acknowledgements

We thank Lisa D. Nee (Western General Hospital, Edinburgh, UK) for her major contribution to the acquisition of ultrasound data.

Duality of interest

MWJS has received fees for speaking from Novo Nordisk, Eli Lilly and Pfizer. JRM, JAF, ING, RMW, MA, SG, and JFP report no disclosures.

Contribution

ING conception and design of the study, data interpretation, drafting the article, article revision and approval. JAF conception and design of the study, data interpretation, drafting the article, article revision and approval. JFP conception and design of the study, data interpretation, drafting the article, article revision and approval. JRM conception and design of the study, data collection, data analysis, data interpretation, drafting the article, article revision and approval. MA design of the study, data collection, data analysis, article revision and approval. MWJS conception and design of the study, data interpretation, drafting the article, article revision and approval. RMW conception and design of the study, data collection, article revision and approval. SG conception and design of the study, data collection, article revision and approval.

References

1. Ahmed MH, Abu EO, Byrne CD. Non-Alcoholic Fatty Liver Disease (NAFLD): new challenge for general practitioners and important burden for health authorities? Primary care diabetes. 2010;4(3):129-37.
2. Williams R, Aspinall R, Bellis M, Camps-Walsh G, Cramp M, Dhawan A, et al. Addressing liver disease in the UK: a blueprint for attaining excellence in health care and reducing premature mortality from lifestyle issues of excess consumption of alcohol, obesity, and viral hepatitis. *The Lancet*. 2014;384(9958):1953-97.
3. Armstrong MJ, Houlihan DD, Bentham L, Shaw JC, Cramb R, Olliff S, et al. Presence and severity of non-alcoholic fatty liver disease in a large prospective primary care cohort. *Journal of Hepatology*. 2012;56(1):234-40.
4. Fattovich G, Stroffolini T, Zagni I, Donato F. Hepatocellular carcinoma in cirrhosis: Incidence and risk factors. *Gastroenterology*. 2004;127(5, Supplement 1):S35-S50.
5. Davies S. Chief Medical Officer annual report 2011: volume 1, chapter 2c, p 163. Department of Health November 2012, available at https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/141773/CMO_Annual_Report_2011_Chapter_2cpdf [accessed 01/07/2015]. 2012.
6. Liver Disease: Today's complacency, tomorrow's catastrophe. All-Party Parliamentary Hepatology Group (APPHG) Inquiry into Improving Outcomes in Liver Disease. . Available at <http://www.wias.org.uk/uploads/APPHG%20report%20March%202014%20FINAL.pdf> [accessed 01/07/2015]. 2014.
7. Price J, Reynolds R, Mitchell R, Williamson R, Fowkes FG, Deary I, et al. The Edinburgh Type 2 Diabetes Study: study protocol. *BMC Endocrine Disorders*. 2008;8(1):18.
8. Marioni RE, Strachan MWJ, Reynolds RM, Lowe GDO, Mitchell RJ, Fowkes FGR, et al. Association Between Raised Inflammatory Markers and Cognitive Decline in Elderly People With Type 2 Diabetes: The Edinburgh Type 2 Diabetes Study. *Diabetes*. 2010;59(3):710-3.
9. Morling JR, Strachan MWJ, Hayes PC, Butcher I, Frier BM, Reynolds RM, et al. Prevalence of abnormal plasma liver enzymes in older people with Type 2 diabetes. *Diabetic Medicine*. 2012;29(4):488-91.
10. Williamson RM, Perry E, Glancy S, Marshall I, Gray C, Nee LD, et al. The use of ultrasound to diagnose hepatic steatosis in type 2 diabetes: intra- and interobserver variability and comparison with magnetic resonance spectroscopy. *Clinical Radiology*. 2011;66(5):434-9.
11. Wai C-T, Greenson JK, Fontana RJ, Kalbfleisch JD, Marrero JA, Conjeevaram HS, et al. A simple noninvasive index can predict both significant fibrosis and cirrhosis in patients with chronic hepatitis C. *Hepatology*. 2003;38(2):518-26.
12. Guha IN, Parkes J, Roderick P, Chattopadhyay D, Cross R, Harris S, et al. Noninvasive markers of fibrosis in nonalcoholic fatty liver disease: validating the European Liver Fibrosis Panel and exploring simple markers. *Hepatology*. 2008;47(2):455-60.
13. Vallet-Pichard A, Mallet V, Nalpas B, Verkarre V, Nalpas A, Dhalluin-Venier V, et al. FIB-4: An inexpensive and accurate marker of fibrosis in HCV infection. comparison with liver biopsy and fibrotest. *Hepatology*. 2007;46(1):32-6.
14. Angulo P, Hui JM, Marchesini G, Bugianesi E, George J, Farrell GC, et al. The NAFLD fibrosis score: a noninvasive system that identifies liver fibrosis in patients with NAFLD. *Hepatology*. 2007;45(4):846-54.

15. Morling JR, Fallowfield JA, Guha IN, Nee LD, Glancy S, Williamson RM, et al. Using non-invasive biomarkers to identify hepatic fibrosis in people with type 2 diabetes mellitus: The Edinburgh type 2 diabetes study. *Journal of Hepatology*. 2014;60(2):384-91.
16. Prati D, Taioli E, Zanella A, Torre ED, Butelli S, Del Vecchio E, et al. Updated Definitions of Healthy Ranges for Serum Alanine Aminotransferase Levels. *Annals of Internal Medicine*. 2002;137(1):1-10.
17. Williamson RM, Price JF, Glancy S, Perry E, Nee LD, Hayes PC, et al. Prevalence of and risk factors for hepatic steatosis and nonalcoholic fatty liver disease in people with type 2 diabetes: the Edinburgh Type 2 Diabetes Study. *Diabetes Care*. 2011;34(5):1139-44.
18. Jacqueminet S, Lebray P, Morra R, Munteanu M, Devers L, Messous D, et al. Screening for liver fibrosis by using a noninvasive biomarker in patients with diabetes. *Clin Gastroenterol Hepatol*. 2008;6(7):828-31.
19. Roulot D, Costes J-L, Buyck J-F, Warzocha U, Gambier N, Czernichow S, et al. Transient elastography as a screening tool for liver fibrosis and cirrhosis in a community-based population aged over 45 years. *Gut*. 2011;60(7):977-84.
20. Wong VW-S, Chu WC-W, Wong GL-H, Chan RS-M, Chim AM-L, Ong A, et al. Prevalence of non-alcoholic fatty liver disease and advanced fibrosis in Hong Kong Chinese: a population study using proton-magnetic resonance spectroscopy and transient elastography. *Gut*. 2012;61(3):409-15.
21. Dixon JB, Bhathal PS, O'Brien PE. Nonalcoholic fatty liver disease: predictors of nonalcoholic steatohepatitis and liver fibrosis in the severely obese. *Gastroenterology*. 2001;121:91 - 100.
22. Adams LA, Sanderson S, Lindor KD, Angulo P. The histological course of nonalcoholic fatty liver disease: a longitudinal study of 103 patients with sequential liver biopsies. *Journal of Hepatology*. 2005;42(1):132-8.
23. Harrison SA, Torgerson S, Hayashi PH. The natural history of nonalcoholic fatty liver disease: a clinical histopathological study. *Am J Gastroenterol*. 2003;98(9):2042-7.
24. Fassio E, Álvarez E, Domínguez N, Landeira G, Longo C. Natural history of nonalcoholic steatohepatitis: A longitudinal study of repeat liver biopsies. *Hepatology*. 2004;40(4):820-6.
25. Svegliati-Baroni G, Ridolfi F, Di Sario A, Casini A, Marucci L, Gaggiotti G, et al. Insulin and insulin-like growth factor-1 stimulate proliferation and type I collagen accumulation by human hepatic stellate cells: differential effects on signal transduction pathways. *Hepatology*. 1999;29(6):1743-51.
26. El-Serag HB, Hampel H, Javadi F. The Association Between Diabetes and Hepatocellular Carcinoma: A Systematic Review of Epidemiologic Evidence. *Clinical Gastroenterology and Hepatology*. 2006;4(3):369-80.
27. Cancer Registry in Scotland [Internet]. Information Services Division. Available at <http://www.isdscotland.org/Health-Topics/Cancer/Scottish-Cancer-Registry/> [accessed 17/12/2014]. 2013.
28. Major JM, Sargent JD, Graubard BI, Carlos HA, Hollenbeck AR, Altekruse SF, et al. Local geographic variation in chronic liver disease and hepatocellular carcinoma: contributions of socioeconomic deprivation, alcohol retail outlets, and lifestyle. *Annals of Epidemiology*. 2014;24(2):104-10.

Tables

Table 1. Study population. Values are mean (sd), median (IQR) or % (n).

	Study population n=923	Met Hepatology referral criteria n=125	<i>P</i> [vs. not referred]	Seen by Hepatology services n=52	<i>P</i> [vs. not seen]
Age, years	68.9 (4.2)	69.3 (4.3)	0.224	69.3 (4.6)	0.423
Sex, % male	52.3% (483)	40.8 (51)	0.007	40.4 (21)	0.087
SIMD quintile, %	I	11.4 (105)	0.041	9.6 (5)	0.031
	V	34.3 (317)		25.0 (13)	
Random glucose, mmol/L	6.89 (2.3)	7.60 (3.0)	0.003	7.93 (3.3)	0.021
HbA1c, %	7.20 (1.1)	7.37 (1.2)	0.046	7.69 (1.3)	0.008
HbA1c, mmol/mol	55.2 (11.7)	57.1 (12.7)		60.5 (14.6)	
Duration of diabetes, % <5 years	25.7 (235)	20.0 (25)	0.124	17.3 (9)	0.191
Diabetes treatment:	Diet, %	19.5 (180)	1.000	19.2 (10)	1.000
	OAHA, %	64.8 (598)	0.688	59.6 (31)	0.456
	Insulin, %	15.7 (145)	0.511	21.2 (11)	0.245
Retinopathy, %	Mild	27.9 (254)	0.182	25.0 (13)	0.035
	Moderate/severe	4.4 (40)		11.5 (6)	
Chronic kidney disease ^a , %	19.6 (179)	22.8 (28)	0.393	27.5 (14)	0.149
Cardiovascular disease ^b , %	36.9 (338)	37.6 (47)	0.842	38.5 (20)	0.769
BMI, kg/m ²	31.3 (5.7)	31.7 (6.1)	0.411	32.0 (6.6)	0.414
Total cholesterol, mmol/L	4.4 (0.8)	4.3 (0.8)	0.021	4.1 (0.8)	0.690
Triglycerides, mmol/L	1.66 (0.9)	1.74 (1.0)	0.305	1.50 (0.6)	0.073
sBP, mmHg	138.2 (18.5)	140.6 (18.6)	0.115	138.7 (17.1)	0.845
Known risk factor for liver disease ^c , %	20.9 (193)	31.2 (39)	0.004	34.6 (18)	0.021
Alcohol excess ^d , %	12.8 (118)	20.8 (26)	0.006	21.2 (11)	0.084
Hepatotoxic medication ^e , %	9.2 (6)	10.4 (13)	0.618	11.5 (6)	0.467
Positive autoantibodies ^f , %	0.7 (6)	3.2 (4)	0.005	3.8 (2)	0.046

BMI body mass index; **IQR** inter-quartile range; **OAHA** oral anti-hyperglycaemic agent; **sBP** systolic blood pressure; **sd** standard deviation; **SIMD** Scottish Index of Multiple Deprivation

^a Defined as estimated glomerular filtration rate <60ml/min.

^b Defined as of myocardial infarction, angina, coronary intervention, intermittent claudication, peripheral artery intervention, stroke, transient ischaemic attack or carotid endarterectomy.

^c Defined as any of d-f below.

^d Defined as females >14 units/week, males >21 units/week or patient disclosed history of a current or prior alcohol problem.

^e Defined as the use of (non-topical) glucocorticoids for >1 week, isoniazid, methotrexate, amiodarone or tamoxifen within the 6 months prior to the liver assessment.

^f Defined as ASMA titre >1:160 or AMA titre >1:40

Table 2. Potential demographic, diabetes and metabolic risk factors for the development of CS-CLD. Values are % (n), odds ratios and hazard ratios.

	No CS-CLD n=903	Unknown prevalent CS-CLD n=13	OR (95% CI) for unknown prevalent CS-CLD	<i>p</i>	Incident CS-CLD n=15	HR (95% CI) for incident CS-CLD	<i>p</i>	
Age, years	68.9 (4.2)	69.5 (4.9)	1.04 (0.91,1.18)	0.600	69.8 (4.1)	1.07 (0.94,1.21)	0.306	
Sex, % male	52.5 (474)	30.8 (4)	2.49 (0.76,8.13)	0.132	40.0 (6)	1.93 (0.69,5.45)	0.213	
SIMD quintile:	I, %	11.2 (101)	-	-	33.3 (5)	Ref		
	V, %	34.9 (315)	0 (0)	-	26.7 (4)	0.26 (0.07,0.97)	0.044	
Duration of diabetes >5 years, %	73.7 (660)	76.9 (10)	1.19 (0.32,4.35)	0.796	93.3 (14)	5.30 (0.70,40.34)	0.107	
Fasting glucose, mmol/L	6.87 (2.3)	7.53(3.2)	1.11 (0.91,1.35)	0.301	6.98 (2.8)	1.21 (0.78,1.88)	0.405	
HbA1c,	%	7.19 (1.1)	7.34 (0.8)	1.13 (0.70,1.84)	0.615	7.53 (1.5)	1.02 (0.98,1.06)	0.405
	mmol/mol	55.1 (11.6)	56.7 (8.3)	1.01 (0.97,1.06)				
Diabetes treatment:	Diet, %	19.7 (178)	15.4 (2)	Ref	6.7 (1)	Ref		
	OAHA, %	65.1 (588)	46.2 (6)	1.06 (0.22,5.15)	0.943	53.3 (8)	2.50 (0.31,20.0)	0.388
	Insulin, %	15.2 (137)	69.2 (9)	2.60 (0.47,14.4)	0.274	40.0 (6)	9.08 (1.09,75.5)	0.041
BMI, kg/m ²	31.2 (5.6)	33.4 (6.7)	1.06 (0.98,1.16)	0.161	34.8 (7.4)	1.09 (1.02,1.17)	0.016	
Cholesterol, mmol/L	4.2 (0.8)	3.9 (0.5)	0.68 (0.32,1.43)	0.310	3.9 (0.8)	0.66 (0.33,1.34)	0.248	
Triglycerides, mmol/L	1.67 (0.9)	1.25 (0.3)	0.39 (0.14,1.11)	0.078	1.46 (0.4)	0.70 (0.32,1.54)	0.371	
CRP ^a , mg/L	1.69 (0.8-3.9)	2.72 (1.4-11.6)	1.83 (1.14,2.94)	0.013	3.98 (1.9-13.6)	1.66 (1.06,2.60)	0.026	

CRP C-reactive protein; CS-CLD clinically significant chronic liver disease; HR hazard ratio; OR odds ratio; sd standard deviation; SIMD Scottish Index of Multiple Deprivation.

CS-CLD defined as clinically significant disease: incident cirrhosis, hepatocellular carcinoma or gastro-oesophageal varices.

^a Analysed on the Ln scale: odds and hazard ratios for a one unit increase in the Ln of the risk factor

Table 3. Potential liver injury related risk factors for the development of CS-CLD. Values are % (n), odds ratios and hazard ratios.

	No CS-CLD n=903	Unknown prevalent CS-CLD n=13	OR (95% CI) for unknown prevalent CS-CLD	<i>p</i>	Incident CS-CLD n=15	HR (95% CI) for incident CS-CLD	<i>p</i>
ALT, U/L	33.3 (12.7)	39.0 (12.6)	1.03 (1.00-1.06)	0.108	44.4 (23.6)	1.04 (1.01,1.06)	0.002
ALT >50 U/L, %	7.8 (70)	23.1 (3)	3.56 (0.96,13.2)	0.058	40.0 (6)	6.69 (2.38,18.8)	<0.001
AST, U/L	30.0 (9.5)	49.7 (12.6)	1.09 (1.06-1.13)	<0.001	46.0 (23.7)	1.07 (1.04,1.09)	<0.001
AST >45 U/L, %	6.8 (61)	69.2 (9)	30.8 (9.22,102.9)	<0.001	40.0 (6)	8.30 (2.94,23.43)	<0.001
GGT ^a , U/L	16.0 (10.0-26.0)	55.0 (34.0-103.0)	5.35 (2.91,9.83)	<0.001	62.0 (21.5-185.0)	3.56 (2.17,5.83)	<0.001
GGT >55 U/L, %	7.9 (71)	69.2 (9)	26.2 (7.87,87.1)	<0.001	53.3 (8)	13.2 (4.79,36.6)	<0.001
CK18 ^a , U/L	100.6 (76.2-135.5)	152.7 (143.1-207.9)	3.42 (1.59,7.34)	0.002	127.6 (83.4-586.5)	4.10 (2.08,8.06)	<0.001
Hepatic steatosis, %	56.8 (513)	46.2 (6)	0.65 (0.22,1.95)	0.445	80.0 (12)	2.93 (0.83,10.38)	0.096
APRI*	0.24 (0.19-0.32)	0.67 (0.35-0.92)	87.7 (20.8,369.6)	<0.001	0.39 (0.29-0.88)	20.4 (6.81,61.0)	<0.001
AST:ALT	0.95 (0.3)	1.33 (0.3)	2.55 (1.22,5.34)	0.013	1.10 (0.3)	1.56 (0.93,2.64)	0.094
AST:ALT ratio >1, %	35.7 (320)	76.9 (10)	6.00 (1.64,22.0)	0.007	60.0 (9)	3.67 (1.29,10.44)	0.015
ELF score	8.9 (0.8)	10.9 (1.0)	4.38 (2.20,8.70)	<0.001	10.2 (1.0)	1.64 (1.30,2.06)	<0.001
FIB4 score	1.50 (0.6)	3.96 (2.0)	7.81 (3.93,15.5)	<0.001	2.56 (1.3)	4.08 (2.71,6.15)	<0.001
HA ^a , micrg/L	50.6 (34.8-81.4)	220.3 (177.9-318.9)	16.4 (5.35,50.1)	<0.001	183.2 (78.9-230.6)	5.80 (2.60,13.0)	<0.001
NFS	-0.40 (1.1)	1.38 (1.5)	3.99 (2.36,6.73)	<0.001	0.74 (1.0)	2.18 (1.54,3.09)	<0.001
Spleen >13 cm, %	4.1 (37)	69.2 (9)	52.5 (15.5,178.5)	<0.001	0	-	
Platelets <150x10 ⁹ /L, %	3.0 (27)	46.2 (6)	27.3 (8.59,86.6)	<0.001	14.3 (2)	4.36 (0.97,19.6)	0.055
Known risk factor for liver disease ^b , %	20.7 (187)	30.8 (4)	1.70 (0.52,5.59)	0.488	40.0 (6)	2.44 (0.87,6.85)	0.091
Alcohol excess ^c , %	12.6 (113)	23.1 (3)	2.09 (0.57,7.71)	0.268	20.0 (3)	1.63 (0.46,5.77)	0.451

ALT alanine aminotransferase; APRI Aspartate to Platelet Ration Index; AST aspartate aminotransferase; CK18 cytokeratin-18; CS-CLD clinically significant chronic liver disease; ELF Enhanced Liver Fibrosis panel; FIB4 Fibrosis-4 Score; GGT gamma glutamyltransferase; HA hyaluronic acid; HR hazard ratio; NFS NAFLD Fibrosis Score; OR odds ratio.

CS-CLD defined as clinically significant disease: incident cirrhosis, hepatocellular carcinoma or gastro-oesophageal varices.

^a Analysed on the Ln scale: odds and hazard ratios for a one unit increase in the ln of the risk factor

^b Defined as any of: alcohol excess (females >14 units/week, males >21 units/week or patient disclosed history of a current or prior alcohol problem), hepatotoxic medication use (use of (non-topical) glucocorticoids for >1 week, isoniazid, methotrexate, amiodarone or tamoxifen within the 6 months prior to the liver assessment) or positive autoantibodies (ASMA titre >1:160 or AMA titre >1:40).

^c Defined as females >14 units/week, males >21 units/week or patient disclosed history of a current or prior alcohol problem

Figure legend

Figure 1. Identification of liver disease

ALP alkaline phosphatase; **ALT** alanine aminotransferase; **AST** aspartate aminotransferase; **CLD** chronic liver disease; **CS-CLD** clinically significant chronic liver disease; **GGT** gamma glutamyl transferase; **HCC** hepatocellular carcinoma; **USS** ultrasound scan.

Alcohol excess was defined as females >14 units/week, males >21 units/week or patient disclosed history of a current or prior alcohol problem.

Use of potentially hepatotoxic medication was defined as the use of non-topical glucocorticoids for >1 week, isoniazid, methotrexate, amiodarone or tamoxifen within the 6 months prior to the year 1 clinic. Strongly positive autoantibodies were defined as ASMA titre >1:160 or AMA titre >1:40.

Figure 2. Patient flowcharts

AFP alpha feto protein; **ALD** alcoholic liver disease; **ALT** alanine aminotransferase; **AST** aspartate aminotransferase; **CS-CLD** clinically significant chronic liver disease; **HA** hyaluronic acid; **HBV** hepatitis B virus; **LFTs** liver function tests; **NAFLD** non-alcoholic fatty liver disease; **PBC** primary biliary cirrhosis

Figures

<p>A. Specialist Hepatology referral criteria, any of:</p> <ul style="list-style-type: none"> • routine liver enzyme tests above the laboratory ULN (ALT >50 U/L, AST >45 U/L, GGT >55 U/L, ALP >125 U/L) • AST:ALT ratio >1 • hyaluronic acid >100 microg/L (in the absence of known joint disease) • positive autoantibodies (any of anti-nuclear antibody, anti-smooth muscle antibody, anti-mitochondrial antibody) • ferritin >1000 ng/mL • alpha-feto protein >6 ng/mL • positive hepatitis B or C serology) • spleen >13cm (in the absence of known haematological cause) • platelets <150 x10⁹/L (in the absence of known haematological cause) • suspected cirrhosis on USS
<p>B. Health record and individual case note verification:</p> <ul style="list-style-type: none"> • <i>possible</i> cases were identified from year 1 and year 4 clinic patient self-reported questionnaires; data linkage to ICD-9/10 and OPCS coding within SMR01 general and acute inpatient discharge records (at NHS National Services Scotland, Information Services Division) in July 2008 and July 2011; non-attenders at the year 4 clinic were sent a postal questionnaire and if no response was received a modified questionnaire was sent to the General Practitioner. • <i>definite</i> cases were confirmed through the review of all patients' electronic secondary care medical records (TrakCare, InterSystems Corp., Cambridge, USA) until 31 December 2013 in order to verify all possible diagnoses and to identify any additional CLD.
<p>C. Clinically significant chronic liver disease definition:</p> <ul style="list-style-type: none"> • Definite (prevalent or incident) CS-CLD was a composite outcome defined as a diagnosis of cirrhosis, HCC (with confirmatory radiology) or gastro-oesophageal varices (with confirmatory endoscopy) recorded in the patients' medical records.

Figure 1. Identification of liver disease

Figure 2. Patient flowcharts

Figure 2A Research clinic outcomes




Figure 2B Development of CS-CLD

