

1 **Perspectives on Synoptic Climate Classification and its Role in Interdisciplinary Research**

2

3 Authors: P. Grady Dixon<sup>1</sup>, Michael Allen<sup>2</sup>, Simon N. Gosling<sup>3</sup>, David M. Hondula<sup>4,5</sup>, Vijendra Ingole<sup>6,7</sup>, Rebekah

4 Lucas<sup>8</sup>, Jennifer Vanos<sup>9</sup>

5 <sup>1</sup>Department of Geosciences, Fort Hays State University, Hays, Kansas, USA

6 <sup>2</sup>Department of Political Science and Geography, Old Dominion University, Norfolk, Virginia, USA

7 <sup>3</sup>School of Geography, University of Nottingham, Nottingham, United Kingdom

8 <sup>4</sup>Center for Policy Informatics, Arizona State University, Phoenix, Arizona, USA

9 <sup>5</sup>School of Geographical Sciences and Urban Planning, Arizona State University, Tempe, Arizona, USA

10 <sup>6</sup>Umeå Centre for Global Health Research, Umeå University, Umeå, Sweden

11 <sup>7</sup>Vadu Rural Health Program, KEM Hospital Research Centre, Pune, India

12 <sup>8</sup>School of Sport, Exercise, and Rehabilitation Sciences, University of Birmingham, Birmingham, United Kingdom

13 <sup>9</sup>Department of Geosciences, Texas Tech University, Lubbock, Texas, USA

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28 **Abstract**

29 Synoptic climatology has a long history of research where weather data are aggregated and composited to gain a  
30 better understanding of atmospheric effects on non-atmospheric variables. This has resulted in an applied  
31 scientific discipline that yields methods and tools designed for applications across disciplinary boundaries. The  
32 spatial synoptic classification (SSC) is an example of such a tool that helps researcher bridge methodological gaps  
33 between disciplines, especially those studying weather effects on human health. The SSC has been applied in  
34 several multi-discipline projects, and it appears that there is ample opportunity for growth into new topical areas.  
35 Likewise, there is opportunity for the SSC network to be expanded across the globe, especially into mid-latitude  
36 locations in the southern hemisphere. There is some question of the utility of the SSC in tropical locations, but such  
37 decisions must be based on the actual weather data from individual locations. Despite all of the strengths and  
38 potential uses of the SSC, there are some research problems, some locations, and some datasets for which it is not  
39 suitable. Nevertheless, the success of the SSC as a cross-disciplinary method is noteworthy because it has become  
40 a catalyst for collaboration.

41

42

43

44

45

46

47

48

49

50

51

52

53

54

## 55 1. Introduction

56 One of the most comprehensive methods of air-mass categorization is the spatial synoptic classification  
57 (SSC) system (Sheridan, 2002, Sheridan and Dolney, 2003). The current SSC was developed by Sheridan (2002) and  
58 was referred to as “SSC2” because it stemmed from an extensive line of research initiated by Muller, Kalkstein, and  
59 others in the late 1970s (Kalkstein et al., 1996, Lamb, 1972, Muller, 1977) that eventually led to an initial version  
60 that is sometimes referred to as “SSC1” (see Hondula et al., 2014 for an in-depth history). A combination of  
61 weather variables (air temperature, dew-point depression, wind speed, mean cloud cover, mean sea-level  
62 pressure, diurnal temperature range, and diurnal dew-point range), is used to numerically characterize the state of  
63 the atmosphere; these quantities are subsequently differentiated into weather-type categories, encompassing  
64 variables that synergistically affect human health (Greene et al., 2011, Davis et al., 2003) and various ecological  
65 systems (e.g., Frank et al., 2008a, Frank et al., 2008b).

66 The relative nature of the SSC daily weather-type classification scheme (i.e., weather-type definitions vary  
67 across space and time) is a strength cited in many studies. The SSC has become one of the key analytical tools  
68 implemented in a diverse range of climate and health research investigations that are location- and time-specific  
69 (Hondula et al., 2014). Other areas of study that have benefited from analyses of SSC data include air-quality  
70 variability (Davis et al., 2010, Pope and Kalkstein, 1996, Power et al., 2006, Rainham et al., 2005, Vanos et al.,  
71 2014b), human health (Hajat et al., 2010, Vanos et al., 2014b, Vanos et al., 2015), the urban heat island (Dixon and  
72 Mote, 2003), and climatological trend analyses (Hondula and Davis, 2011, Knight et al., 2008, Vanos and Cakmak,  
73 2014). Through these studies, we see the SSC is applicable to various topics in cross-cutting disciplines and has a  
74 large geographical range, which includes approximately 400 stations (Figure 1) spanning the United States, Canada,  
75 and Europe, and select cities in Asia with data covering several decades (Bower et al., 2007, Hondula et al., 2014,  
76 Sheridan, 2002, Tan et al., 2004).

77 There are numerous opportunities to expand the application of synoptic-scale impact analyses to new  
78 locations, contexts, and disciplines. In this article, we discuss the identified gaps in both the spatial nature of the  
79 system and the disciplinary applications, providing critical information to researchers outside of the area of  
80 climatology on where and how the SSC can be successfully applied. This review highlights synoptic climatology as a  
81 catalyst for cross-discipline research.

82

## 83 2. Synoptic Climatology

### 84 a. Discipline Review

85 A goal of synoptic climatology is to understand the relationships between the surface environment and  
86 overlying atmospheric circulation (Yarnal, 1993). With a horizontal scale of ~1,000 km and a lifespan of ~5–7 days,  
87 cyclones and anticyclones, which are the main synoptic-scale features of the atmosphere, influence a wide range  
88 of environmental processes including water resources, severe-weather outbreaks, and health. Accordingly, local-  
89 scale analysis of weather often begins with a characterization of the synoptic-scale forcing processes. Such  
90 atmospheric “snapshots” provide simple, useful descriptions that are designed to aid understanding of our physical  
91 world.

92 In synoptic climatology, the classification scheme has been a primary focus of research efforts for many  
93 decades. Multiple variables have been used to classify atmospheric patterns including temperature, pressure,  
94 airflow, and derived properties such as vorticity (Barry, 2005, LeDrew, 1984). Additionally, these features are  
95 classified at multiple spatial (e.g. global or regional) and temporal (e.g. annual or daily) scales. Discrete  
96 classification of synoptic patterns allow synoptic climatologists to communicate with other disciplines so that  
97 environmental relationships may be analyzed (Carleton, 1999). Only during the last two decades has the use of  
98 synoptic climatology accelerated significantly as a tool for applications rather than pure classification (Sheridan  
99 and Lee, 2013, Yarnal et al., 2001).

100 Synoptic climatological classifications often involve one of two approaches. The circulation-to-  
101 environment approach emphasizes the atmospheric patterns. In this case, the overlying atmospheric scenario is  
102 classified *a priori* and then related to the surface variable of interest (e.g., air temperature). In contrast, the  
103 environment-to-circulation approach initially determines the environmental variable of study and then compares  
104 its condition to the circulation pattern(s) (Yarnal, 1993).

105 Within the field of synoptic climatology, multiple classification approaches exist and may be subjective  
106 (manual), objective/computer-automated, or hybrid. Manual map comparisons began very early (Abercromby,  
107 1883, Lamb, 1950, van Bebber and Köppen, 1895), yet this method was subjective and labor-intensive (Frakes and  
108 Yarnal, 1997). In manual approaches, the analysis relies on professional expertise to define *a priori* classifications.

109 While the majority of subjective catalogs (Baur et al., 1944, Lamb, 1972) focus on regional analysis, some have  
110 been developed for larger-scale considerations (Girs, 1948). Recently, automated and hybrid classification  
111 methods have been developed, and the discipline continues to evolve with the increased availability of weather  
112 data and more complex climate models. There is no standard classification scheme, but rather, synoptic  
113 climatology highlights the importance of interpreting map patterns and evaluating surface relationships. Huth et al.  
114 (2008) provide further discussion on synoptic climatological approaches.

115 Along with increased computing ability, more sophisticated, statistically robust techniques for  
116 classification have become increasingly common in synoptic climatology (Yarnal et al., 2001). In addition to  
117 understanding basic circulation controls, statistical and dynamic modeling techniques are used to uncover the  
118 patterns and near-surface processes related to a variety of environmental issues. Techniques such as cluster  
119 analysis (e.g., Esteban et al., 2005) and self-organizing maps (Hewitson and Crane, 2002, Kohonen et al., 2001)  
120 have helped re-shape the discipline. Globally-gridded reanalysis datasets (e.g., Dee et al., 2011, Ebita et al., 2011,  
121 Kalnay et al., 1996) have led to the inclusion of more complex, derived variables such as vorticity and moisture  
122 characteristics. Regional and global climate modeling now offer new approaches to examine the physical  
123 mechanisms linking surface conditions with atmospheric circulation (Giorgi and Mearns, 1999).

124

#### 125 b. Spatial Synoptic Classification

126 The Spatial Synoptic Classification (SSC) is a weather type classification based solely on surface  
127 observations. To determine the SSC weather types for a given time and place, a hybrid system is employed using  
128 both manual and automated processes. First, 'typical' meteorological conditions are chosen for each of the  
129 weather types (Dry Polar [DP], Dry Moderate [DM], Dry Tropical [DT], Moist Polar [MP], Moist Moderate [MM],  
130 Moist Tropical [MT], or Transition [TR]) at each weather station based on climatological knowledge. There is also  
131 the MT+ subset of the MT weather type, which is common in the summer across the mid-latitudes, to differentiate  
132 the days with the greatest potential for heat stress. The MT+ conditions occur when both morning and afternoon  
133 apparent temperatures are above the MT weather-type means for the location (Sheridan and Kalkstein, 2004).  
134 Sliding "seed days" representing each of the weather types are created for four two-week windows during each  
135 season of the year to correspond with the hottest and coldest two weeks annually and the midway points in spring

136 and autumn for the given location (Sheridan, 2002). The sliding seed-day method permits an improved temporal  
137 continuity across various climate types and throughout the entire year, encapsulating the temporally relative  
138 nature of the SSC.

139 Actual conditions are then compared to the seed days and each day is classified as the weather type it  
140 most closely resembles (lowest error score based on equal-weighted z-scoring). The groups of days identified as  
141 certain SSC types are not completely homogeneous, as the synoptic-scale circulation is a complex process not  
142 perfectly described by seven distinct groups. Meteorological variability is also identified within an SSC weather  
143 type at various scales of interest dependent on the research (e.g., division of MT and DT days into categories of  
144 higher or lower severity for heat stress (Sheridan and Kalkstein, 2004), division of TR days into categories  
145 representing various frontal types (Hondula and Davis, 2011)). Complete details of the classification procedure can  
146 be found in Sheridan (2002). SSC data are freely available online at <http://sheridan.geog.kent.edu/ssc.html>.

147

### 148 **3. Spatial Synoptic Classification Uses**

#### 149 a. Temperature and Human Health

150 Among the wide range of potential applications for synoptic classification schemes, the SSC has gained  
151 greatest traction in studies of the relationships between heat and human-health outcomes. SSC-based studies of  
152 heat impacts on morbidity and mortality focus largely on the DT and MT+ weather types, often referred to as the  
153 ‘oppressive’ types (e.g., Isaksen et al., 2015, Saha et al., 2015). These oppressive days have been applied in the  
154 development of several of the initial outcomes-based heat-health watch-warning systems deployed in the USA as  
155 well as in Toronto (Canada), South Korea, Shanghai (China), and select Italian cities (Kalkstein et al., 2011, Kalkstein  
156 et al., 2008, Kirchmayer et al., 2004, Sheridan and Kalkstein, 2004, Tan et al., 2004). More recently, the SSC and  
157 related techniques have been applied to the study of additional health outcomes including respiratory-related  
158 hospital admissions (Hondula et al., 2013, Lee et al., 2012), influenza and pneumonia mortality (Davis et al., 2012),  
159 and cold-season cardiovascular deaths (Lee, 2015).

160

#### 161 b. Air Pollution

162 The SSC has been used to help characterize the relationship between air quality and meteorology in

163 research studies set in Canada, Korea, and the United States. To date, the main pollutants addressed have been  
164 nitrogen dioxide (NO<sub>2</sub>), sulfur dioxide (SO<sub>2</sub>), carbon monoxide (CO), ozone (O<sub>3</sub>), and particulate matter < 2.5 µm  
165 (PM<sub>2.5</sub>). Standard analyses segregate each day into a select weather type, and the individual mean air-pollution  
166 levels are then calculated and statistically compared by weather type. Prior to the current SSC, Cheng et al. (1992)  
167 completed the first SSC air pollution study using the SSC1 to assess concentrations of O<sub>3</sub> and PM in the city of  
168 Philadelphia. Following this, Pope & Kalkstein (1996) used the SSC1 to confirm associations between respirable  
169 particles and mortality in the Utah Valley, and Smoyer et al. (2000) described relationships between weather, air  
170 pollution, and mortality in Birmingham and Philadelphia (USA), also using the SSC1. Over the last 15 years, ambient  
171 air pollution has been shown in over a dozen studies to be closely related to the SSC weather type (e.g., Davis et  
172 al., 2010, Greene et al., 1999, Hanna et al., 2011, Kim et al., 2014, Rainham et al., 2005, Vanos et al., 2014a, Vanos  
173 et al., 2013, Vanos et al., 2014b). The most commonly cited findings show a close association between higher  
174 concentrations of O<sub>3</sub> on DT days, specifically in the summer season (e.g., Davis et al., 2010, Hanna et al., 2011, Kim  
175 et al., 2014, Rainham et al., 2005, Rainham et al., 2001, Smoyer et al., 2000, Vanos et al., 2013). Further, Vanos et  
176 al. (2014b) found that when DT air is present in Canada, other pollutants, such as NO<sub>2</sub> and SO<sub>2</sub>, are significantly  
177 higher than the mean for all weather types. The stagnant, dry, sunny, and hot conditions found within the DT  
178 weather type result in the greatest pollution build up for many pollutants and aid in the photochemical creation of  
179 ozone (Davis and Kalkstein, 1990, Smoyer et al., 2000). Low concentrations of pollutants have been generally  
180 found in moist, cool weather types (e.g., Greene et al., 1999), as well as the TR weather type (e.g., Rainham et al.,  
181 2005, Vanos et al., 2013). TR days are indicated by shifts in synoptic conditions and are commonly associated with  
182 frontal activity (increased wind and precipitation chances), thus resulting in lower air-pollution levels. Newer  
183 research also links higher aeroallergen levels to the presence of MT and DT weather types in 10 Canadian cities  
184 (Hebbern and Cakmak, 2015).

185

### 186 c. Climate Change

187 The potential impacts of climate change on human health have been assessed by applying weather-type-  
188 mortality relationships derived from the present climate to SSC types projected by global climate models (GCMs).  
189 This analysis was first completed using projections of weather types into the 2020s and 2050s for 44 cities in the

190 USA, with subsequent analysis of each city's mortality risks (Kalkstein and Greene, 1997). This analysis was later  
191 updated by Greene et al. (2011) to estimate mortality during excessive heat events (EHEs) for the 2020s, 2050s,  
192 and 2090s across 40 cities in the USA. An application of the SSC by Hayhoe et al. (2010) showed that a 2003  
193 European Heatwave-type event could occur in Chicago by 2050, with a high likelihood of 10 times the city's current  
194 annual average number of heat-related deaths occurring in only a few weeks. In a rare application of synoptic-  
195 weather typing to assess climate-change impacts outside of the US, Cheng et al. (2008) showed that heat-related  
196 mortality could more than double by the 2050s and triple by the 2080s in south-central Canada. The most recent  
197 application of the SSC in climate-change impacts assessment projected future weather types for California for the  
198 2090s and estimated that heat-related mortality among those over 65 could increase by tenfold in major urban  
199 centers (Sheridan et al., 2012a, Sheridan et al., 2012b).

200

#### 201 d. Other SSC Uses

202 The utility of the SSC has not been limited to topics related to human health and the associated impacts of  
203 climate change. Researchers have applied the SSC types to discriminate days that are hot vs cold, arid vs humid, or  
204 synoptically active vs inactive. Almost immediately following Sheridan's (2002) release of the updated SSC, a few  
205 researchers employed the system as an efficient proxy for air-mass types, which were not historically easy to  
206 quantify for most locations (Dixon and Mote, 2003, Grundstein, 2003, Kalkstein and Balling, 2004, Leathers et al.,  
207 2004, Leathers et al., 2002). While some of these projects were focused on how SSC types affect snow cover and  
208 characteristics (Grundstein, 2003, Leathers et al., 2004, Leathers et al., 2002), one paper showed that SSC types  
209 could be used to understand summer thunderstorms initiated by the urban heat island (Dixon and Mote, 2003).  
210 Kalkstein and Balling (2004) then used the SSC to analyze diurnal temperature range following the attack on the  
211 World Trade Center in New York on 11 September 2001. Hence, very early in the life of the SSC, it was becoming  
212 apparent that the system would have widespread applicability in weather and climate research.

213 Following the initial burst of authors using SSC for applied climatology research, subsequent papers were  
214 largely related to weather and health, with further studies addressing urban effects on weather (Brazel et al., 2007,  
215 Chow and Svoma, 2011, Ellis et al., 2015) and diurnal temperature ranges (Scheitlin, 2013, Scheitlin and Dixon,  
216 2010). Further growth was seen as climatologists began to use SSC as a way to define "synoptically weak" or



217 “benign” days, which is important when studying convection, lightning, and other meteorological phenomena that  
218 are driven by thermal instability rather than dynamic forcing (Ashley et al., 2012, Bentley et al., 2010, Bentley et  
219 al., 2012, Owen and Dixon, 2015, Stallins et al., 2013, Haberlie et al., 2015, Mote et al., 2007, Shem and Shepherd,  
220 2009). Similarly, some researchers have discovered the utility of the SSC to efficiently analyze weather conditions  
221 as they relate to tree growth (Huang et al., 2010, Senkbeil et al., 2007) and wildlife behavior (Esslinger et al., 2015,  
222 Palumbo et al., 2015). Our discussion of articles using the SSC is not exhaustive, but it is clear that SSC is continuing  
223 to grow in popularity among researchers studying weather-health interactions as well as several other  
224 applications, mostly within applied climatology.

225

#### 226 **4. Limitations of SSC Methods**

227 The previous sections demonstrate many opportunities to apply the SSC, and it appears that such  
228 opportunities will continue to grow. Therefore, we propose a goal for the SSC of being accessible and applicable for  
229 all possible uses where it has been shown to function well. This could mean establishing an SSC for all regions of  
230 the world, but that is not currently feasible due to a lack of reliable weather data (Hondula et al., 2014). There are  
231 many locations with reliable weather data but no SSC, and there is also a question of whether all climate types are  
232 conducive to daily classification by the SSC. Likewise, not all research topics involving synoptic weather variables  
233 can benefit from the SSC or synoptic classification systems in general. Here, we address some known limitations  
234 and challenges so that researchers from various disciplines can better understand and effectively apply the SSC to  
235 benefit their research goals.

236

##### 237 a. Limitations in Temperature-Health Research

238 With its synoptic-scale resolution, the SSC is not designed to describe human exposure to thermal stress  
239 at microscale levels. This is a limitation from the physiological perspective as behavioral factors, metabolic rate,  
240 and clothing properties are not currently considered. In this sense, it could be argued that the SSC system is not  
241 yet applicable as a heat-stress index for estimating thermal strain in individuals (NIOSH, 1986, Parsons, 2003).  
242 There are, however, many pre-existing heat-stress indices that have been designed for the workplace to establish  
243 safe practices and safe limits for work (Parsons, 2003).

244           With respect to environmental epidemiology, the SSC offers a considerable shift from many of the  
245 traditional and emerging techniques applied to investigate the association between temperature and mortality, in  
246 which continuous variables (e.g., temperature, heat index, Universal Thermal Comfort Index (UTCI)) tend to be  
247 used in statistical models (e.g., McMichael et al., 2008, Anderson and Bell, 2009, Urban and Kyselý, 2014, Petitti et  
248 al., 2015). The association between exposure variables and health outcomes in these models has been shown in  
249 many places to be a smooth, non-linear function. Mapping discrete variables like the SSC weather types into this  
250 continuous exposure-response space would seem to be a challenge (Barnett et al., 2010, Huang et al., 2011).  
251 Operational heat-health warning systems designed around the SSC, however, include linear regression functions  
252 within the subset of days associated with each weather type that allow for continuous prediction of anomalous  
253 mortality (Sheridan and Kalkstein, 2004). Whether the current algorithmic approach utilized by these warning  
254 systems most effectively accounts for within SSC-type variability is an outstanding research question that we  
255 recommend investigating in the years ahead.

256           Evaluation of trigger indicators for heat-health early warning systems is recommended by the World  
257 Health Organization and World Meteorological Organization and should take into account system complexity, error  
258 in weather forecast data, and acceptability among user groups (Åström et al., 2014, McGregor et al., 2010). In an  
259 evaluation of the predictive capacity of four different triggering criteria for heat warning systems (including an SSC-  
260 based approach) in four different cities worldwide, Hajat et al. (2010) found that no system was recommended to  
261 be universally preferable. Other studies from Detroit and New York City in the USA suggest that relatively simple  
262 metrics like minimum temperature and maximum heat index perform comparably to more complex models,  
263 including the SSC, therefore, the simpler triggering criteria were deemed preferable for their locations (Metzger et  
264 al. 2010, Zhang et al. 2012). Urban and Kyselý (2015) also encouraged continued comparison of the current SSC  
265 framework to other approaches for triggering operational heat warning systems, including different methods  
266 based on sequences of SSC types.

267           These comparative studies are of interest because they represent the incorporation of different  
268 perspectives into the design of heat-health warning systems. For example, Hajat et al. (2010) connected research  
269 groups from academic institutions and government research offices across five different countries. The SSC and its  
270 operational extension for heat-health warning systems helped to push the conversation regarding what should be

271 included in the design of effective triggering criteria. Whether or not the SSC is ultimately used as the basis for  
272 triggering a public health alert is, for us, less interesting than the idea that its consideration, along with alternatives  
273 ranging from simple environmental variables (e.g., temperature) to complex, biophysical indices (e.g., UTCI), can  
274 expand how researchers and practitioners think about designs of heat-health warning systems.

275

#### 276 b. Limitations in Air-Pollution Studies

277 Air-pollution and health studies conducted in the 20<sup>th</sup> century supported the development of public  
278 warning systems when potentially harmful pollution was likely due to synoptic conditions (e.g., Smoyer et al.,  
279 2000). Yet, even with technological advancements and numerous studies showing connections between SSC  
280 weather types and air pollution, few studies have attempted to produce such SSC-based forecast models.  
281 Investigations of spatiotemporal connections between air pollution and synoptic weather generally stop short of  
282 providing a physical explanation. Rather, most research yields mean levels of air pollution for each SSC weather  
283 type before proceeding with health-outcomes-based approaches.

284 A potential reason for difficulty in using the SSC for air pollution forecasting is the complexity in  
285 determining the origin of air pollution. Weather types alone cannot be used to identify source regions of pollutants  
286 (Hondula et al., 2010); different circulation regimes can result in the same SSC designation at a given location.  
287 Certain DM days, for example, could advect pollutants from a problematic source region or be more conducive  
288 (e.g., warmer, sunnier) to the formation of secondary pollutants than other days, but such variability would be lost  
289 by simply examining overall differences between SSC types. Indeed using the SSC to supplement back-trajectory  
290 analysis has revealed interactive relationships that are not evident from using only the back-trajectory or synoptic  
291 analytical method (Davis et al., 2010, Hondula et al., 2010).

292 Changing concentrations of ground-level pollution is driven by the variables often used to characterize air  
293 masses and weather types (e.g., temperature, pressure, wind, sunlight), which provides the physical underpinning  
294 to explain why studies examining SSC-air pollution linkages often report strong associations. These results are  
295 quite intuitive, yet highly generalized as they differ by pollutant of interest, location, and time of season. Further,  
296 the SSC is of greater utility for examining air pollution variability primarily in locations that are more susceptible to

297 high concentrations and variability of air pollution (Smoyer et al., 2000). Hence, careful consideration and analysis  
298 is still required when using SSC to assess and/or predict air pollution.

299

### 300 c. The Challenges of SSC outside the Mid-Latitudes

301 A map of SSC locations (Figure 1) highlights the absence of SSC locations in tropical, desert, and  
302 developing locations, with a distinct lack of stations in the southern hemisphere. Access to reliable weather data is  
303 challenging in many developing countries, so there is little that can be done to remedy that in the near term. There  
304 is still a question of whether the SSC provides as much value in tropical and/or desert locations that are less likely  
305 to experience synoptic-scale frontal passages and the associated sudden air-mass changes. Such locations often  
306 experience the same synoptic weather types for months at a time. For example, Miami, Florida (USA) experiences  
307 the MT weather type on 65% of days annually and 80% of summer days (Figure 2). It is certainly feasible to break  
308 down those climates into SSC types that are relative to specific locations, but it may not be very useful if the air  
309 temperature differences between DT and MM SSC types are only a few degrees. Further, some current SSC  
310 locations along the southern tier of the USA never experience as many as three of the seven possible categories  
311 during long periods. Frequency distributions of SSC types throughout the year for select SSC locations (Figure 2)  
312 illustrate that mid-latitude locations tend to experience all SSC categories in every season while sub-tropical  
313 locations are unable to fully take advantage of the seven SSC categories. We encourage continued investigation of  
314 the relationship between SSC weather types and synoptic-scale circulation regimes in these locations to determine  
315 if there is within-SSC-type heterogeneity that may be valuable to capture in new tools that aid the fields of  
316 climatology and applied climatology.

317 A noteworthy example of a tropical SSC location that is also in a developing country is Pune, India (the  
318 only location in India; Figure 1). Previous research has shown associations between temperature and human health  
319 in rural parts of Pune District (Ingole et al., 2012), therefore, the authors of this manuscript collaborated (along  
320 with the help of others, including Scott Sheridan) to develop the SSC for the city of Pune to work toward improved  
321 weather-health research in India and an expanded network of SSC stations. One concern among developers was a  
322 lack of the usual four seasons as Pune is dominated by the Asian monsoon, resulting in just three discernible  
323 seasons: summer, monsoon, and winter (Figure 3). Moreover, due to the altitude and overall aridity of Pune,

324 diurnal temperature ranges can often exceed 20 °C during summer and winter. However, interseasonal differences  
325 are much less dramatic with mean monthly temperatures all within 10 °C of each other, and it is debatable  
326 whether Pune ever experiences weather types that are truly Polar (e.g., Pune has never officially recorded a  
327 temperature below freezing). There is the possibility that the SSC can ultimately prove useful in a location even if  
328 some of the categories are never experienced, but only if it helps to understand and/or predict weather-related  
329 effects on non-atmospheric variables, such as health and ecology. Researchers are currently working to test  
330 associations between SSC and health outcomes in locations like Pune.

331 While confirming the lack of synoptic frontal activity across much of the land located within the tropics,  
332 Berry *et al.* (2011) show that some tropical regions do regularly experience fronts (Figure 4). It is probably not  
333 prudent to describe large regions of the planet as being “good” or “bad” candidates for SSC stations without a  
334 thorough review of the climatology of the locations in question, but it does appear that some locations would fail  
335 to make enough use of the SSC categories to justify creating them.

336

## 337 **5. Advantages and Opportunities**

338 The SSC has been relatively under-utilized in the assessment of climate-change impacts, both in terms of  
339 the region of application (most studies have been focused on the USA) and with respect to impacts being assessed  
340 (most focus on human health). Thus a unique opportunity exists to explore numerous climate-change impacts  
341 around the world using the SSC. GCMs output climate variables required to develop SSCs on a grid covering the  
342 globe at resolutions as fine as 25 km (Roberts et al., 2014), so there is potential for applying the SSC to assess  
343 impacts in many regions of the globe. There is also potential to use the SSC to assess impacts such as (but not  
344 limited to): water stress, food security, energy demand, wildfires, and crop yields. These impacts typically occur  
345 across spatial domains similar to that of the SSC. These vital outcomes are similar to human health as their statuses  
346 also depend on multiple, often simultaneous, weather variables in addition to human decisions. Given the success  
347 of using SSC to study human-health outcomes, the issues listed above are likely to benefit from SSC analyses as  
348 well. In any case, comparison and evaluation between techniques should provide a framework for new  
349 applications of synoptic climatology (Huth, 1996, Huth et al., 2008).

350 Human-weather interactions are dynamic and complex because an individual's response, both  
351 physiological and behavioral, can alter the level of exposure, which determines their well-being, health, or even  
352 survival. This interaction and any resulting physiological strain can be defined by six factors or agents (Fanger,  
353 1970):

- 354 1) ambient air temperature
- 355 2) air motion or wind velocity
- 356 3) relative humidity
- 357 4) mean radiant temperature
- 358 5) metabolic heat production of the body
- 359 6) the clothing worn and its insulation and moisture permeability

360 The first four of these agents are environmental and they should all be considered when assessing the thermal  
361 influence of the environment on the human body (Höppe, 1999), while the remaining two are behavioral. There is  
362 much debate and research surrounding which human thermal index is superior at predicting the human  
363 experience in a given environment. Difficulties arise when accounting for the complexity and interactions of all six  
364 factors. It has even been suggested that there cannot be a universal system for rating thermal stress (Belding,  
365 1970, Epstein and Moran, 2006). In this sense, there may be an advantage to using a system, such as the SSC, that  
366 describes well the four climate variables and does not attempt to assume how humans may behave. Future work  
367 on the SSC system might advantageously consider behavioral factors (particularly varying levels of metabolic heat  
368 production) with respect to heat-health warning systems to determine how a weather type impacts humans  
369 performing varying levels of physical activity.

370 Air temperature alone is frequently used to assess the impact of the climatic environment on human  
371 health (Hondula et al., 2014, Parsons, 2003) even though air temperature is seldom the lone cause of heat stress  
372 (Goldman, 2001). Such a reductionist approach can limit our understanding of human-weather interactions. High  
373 humidity significantly increases heat stress by lowering the efficacy of evaporative heat loss (achieved via  
374 sweating), which is the primary human mechanism for heat loss under warm-hot conditions (Havenith, 1999,  
375 Parsons, 2003). Similarly, increased air velocity (wind) enhances both convective and evaporative heat loss  
376 (Havenith, 1999) in most situations. The radiant temperature is directly related to the heat exchange between the

377 environment and the human body, and can significantly contribute to heat stress, matching the heating effect of  
378 air temperature when air velocity is minimal (Höppe, 1999). Thus, consideration of all four environmental factors  
379 and their interactions is essential in accurately describing the relationship between human health and the climatic  
380 environment. In this respect, the SSC and its comprehensive integration of meteorological parameters (air  
381 temperature, dew point, wind velocity, pressure, and cloud cover as a proxy for radiation) provides a meaningful  
382 and insightful description of climatic variables while combining them into one index, which is more manageable for  
383 subsequent epidemiological analyses.

384           A likely advantage of using the SSC for epidemiological and physiological research is its location and time  
385 specificity because weather-health interactions vary seasonally and geographically due to thermal acclimatization  
386 and adaptation strategies. For example, at the end of winter, a population may be more vulnerable to a sudden  
387 hot day. Further, populations in extreme climates are more resilient to weather variability than those in temperate  
388 regions due to adaptation strategies (behavioral responses, clothing, housing, technology, etc.). The spatial  
389 resolution of the SSC is suitable to characterize the climate sensitivity or vulnerability of different socioeconomic  
390 groups (Kalkstein and Davis, 1989). Such characterization is important in understanding key modifiers that affect  
391 the interaction between human health and climate.

392

## 393 **6. Conclusions**

394           Ultimately, there are three closely related goals in this area of research: increase cross-discipline  
395 research, increase knowledge and awareness of SSC, and increase geographical locations with available SSC data.  
396 The success of any of these three goals seems to depend heavily on the progress of the other two, so working  
397 toward one is indirectly equivalent to working on all of them. There will be challenges in expanding the SSC  
398 network and the demand for SSC data in many parts of the world that have been underserved thus far. However,  
399 history suggests that there will be “tipping points” where it becomes quite efficient to increase the number of SSC  
400 stations in a country after the first few are established, and these bursts of new data will likely be accompanied by  
401 newfound interest in those data by regional researchers. It also seems quite likely that the SSC is simply not  
402 suitable for the climates of some locations. Determining which locations fall into this category will not be easy, but

403 this is an area of potential future research that could lead to improved synoptic classification methods and/or  
404 weather-health assessment tools.

405           Application of the SSC, or any synoptic weather analysis tool, in other disciplines often involves the  
406 introduction of an analytical approach (i.e., synoptic classification) that will be unfamiliar to subject experts. This  
407 situation can potentially create confusion, disagreement, and competition among researchers who ultimately have  
408 shared questions and goals. We suggest, however, that such blending of ideas can lead to a productive scientific  
409 advancement. The application of the SSC to temperature-related mortality is a fertile ground for such cross-  
410 perspective discussions that has only recently begun to appear in the scientific literature. There have been several  
411 conference sessions, workshops, and collaborative research projects available in recent years for researchers to  
412 learn more about the SSC and its potential applications. Such opportunities should be less about learning a specific  
413 tool (i.e., the SSC) and more about learning to embrace the methods, perspectives, and goals of other disciplines.  
414 The simplicity of the SSC categories makes it a great catalyst for crossing disciplinary boundaries and making  
415 meaningful progress toward solving real environmental problems, but it cannot be applied in all scenarios. It would  
416 be a great compliment to those who developed the SSC over the years if cross-discipline researchers beyond  
417 climatology find common ground in their past use of the SSC. In the past several years, the SSC has been applied to  
418 numerous research topics including human health, urban heat islands, tree growth, wildlife behavior, and climate  
419 change, and there are some obvious areas of overlap between these study topics that might lead to future  
420 collaborations. It is conceivable that the SSC could become a potential gateway to interdisciplinary efforts  
421 connecting weather, climate, human health and ecology.

422

423

424

425

426

427

428

429



430 **References**

- 431 ABERCROMBY, R. 1883. On certain types of British weather. *Quarterly Journal of the Royal*  
432 *Meteorological Society*, 9, 1-25.
- 433 ANDERSON, B. G. & BELL, M. L. 2009. Weather-related mortality: how heat, cold, and heat waves affect  
434 mortality in the United States. *Epidemiology*, 20, 205.
- 435 ASHLEY, W. S., BENTLEY, M. L. & STALLINS, J. A. 2012. Urban-induced thunderstorm modification in the  
436 Southeast United States. *Climatic Change*, 113, 481-498.
- 437 ÅSTRÖM, C., EBI, K. L., LANGNER, J. & FORSBERG, B. 2014. Developing a heatwave early warning system  
438 for Sweden: evaluating sensitivity of different epidemiological modelling approaches to forecast  
439 temperatures. *International Journal of Environmental Research and Public Health*, 12, 254-267.
- 440 BARNETT, A. G., TONG, S. & CLEMENTS, A. C. A. 2010. What measure of temperature is the best  
441 predictor of mortality? *Environmental Research*, 110, 604-611.
- 442 BARRY, R. G. 2005. Synoptic Climatology. In: OLIVER, J. E. (ed.) *Encyclopedia of World Climatology*.  
443 Springer.
- 444 BAUR, F., HESS, P. & NAGEL, H. 1944. Kalender der Großwetterlagen Europas 1881–1939. *Tech. Rep.*,  
445 *Forschungsinstitut für langfristige Wettervorhersage*. Bad Homburg.
- 446 BELDING, H. S. 1970. The search for a universal heat stress index. In: HARDY, J., GAGGE, A. & STOLWIJK,  
447 J. (eds.) *Physiological and behavioral temperature regulation*. Springfield, Illinois: Charles C.  
448 Thomas.
- 449 BENTLEY, M. L., ASHLEY, W. S. & STALLINS, J. A. 2010. Climatological radar delineation of urban  
450 convection for Atlanta, Georgia. *International Journal of Climatology*, 30, 1589-1594.
- 451 BENTLEY, M. L., STALLINS, J. A. & ASHLEY, W. S. 2012. Synoptic environments favourable for urban  
452 convection in Atlanta, Georgia. *International Journal of Climatology*, 32, 1287-1294.

- 453 BERRY, G., REEDER, M. J. & JAKOB, C. 2011. A global climatology of atmospheric fronts. *Geophysical*  
454 *Research Letters*, 38, n/a-n/a.
- 455 BOWER, D., MCGREGOR, G. R., HANNAH, D. M. & SHERIDAN, S. C. 2007. Development of a spatial  
456 synoptic classification scheme for western Europe. *International Journal of Climatology*, 27,  
457 2017-2040.
- 458 BRAZEL, A., GOBER, P., LEE, S.-J., GROSSMAN-CLARKE, S., ZEHNDER, J., HEDQUIST, B. & COMPARRI, E.  
459 2007. Determinants of changes in the regional urban heat island in metropolitan Phoenix  
460 (Arizona, USA) between 1990 and 2004. *Climate Research*, 33, 171.
- 461 CARLETON, A. M. 1999. Methodology in Climatology. *Annals of the Association of American*  
462 *Geographers*, 89, 713-735.
- 463 CHENG, C., CAMPBELL, M., LI, Q., LI, G., AULD, H., DAY, N., PENGELLY, D., GINGRICH, S., KLAASSEN, J.,  
464 MACIVER, D., COMER, N., MAO, Y., THOMPSON, W. & LIN, H. 2008. Differential and combined  
465 impacts of extreme temperatures and air pollution on human mortality in south–central Canada.  
466 Part II: future estimates. *Air Quality, Atmosphere & Health*, 1, 223-235.
- 467 CHENG, S., YE, H. & KALKSTEIN, L. S. 1992. An evaluation of pollution concentrations in Philadelphia  
468 using an automated synoptic approach. *Middle States Geographer*, 25, 45-51.
- 469 CHOW, W. T. & SVOMA, B. M. 2011. Analyses of nocturnal temperature cooling-rate response to  
470 historical local-scale urban land-use/land cover change. *Journal of Applied Meteorology and*  
471 *Climatology*, 50, 1872-1883.
- 472 DAVIS, R. E. & KALKSTEIN, L. S. 1990. Using a spatial synoptic climatological classification to assess  
473 changes in atmospheric pollution concentrations. *Physical Geography*, 11, 320-342.
- 474 DAVIS, R. E., KNAPPENBERGER, P. C., MICHAELS, P. J. & NOVICOFF, W. M. 2003. Changing Heat-Related  
475 Mortality in the United States. *Environmental Health Perspectives*, 111, 1712-1718.

- 476 DAVIS, R. E., NORMILE, C. P., SITKA, L., HONDULA, D. M., KNIGHT, D. B., GAWTRY, S. P. & STENGER, P. J.  
477 2010. A comparison of trajectory and air mass approaches to examine ozone variability.  
478 *Atmospheric Environment*, 44, 64-74.
- 479 DAVIS, R. E., ROSSIER, C. E. & ENFIELD, K. B. 2012. The Impact of Weather on Influenza and Pneumonia  
480 Mortality in New York City, 1975–2002: A Retrospective Study. *PLOS One*, 7, e34091.
- 481 DEE, D., UPPALA, S., SIMMONS, A., BERRISFORD, P., POLI, P., KOBAYASHI, S., ANDRAE, U., BALMASEDA,  
482 M., BALSAMO, G. & BAUER, P. 2011. The ERA-Interim reanalysis: Configuration and performance  
483 of the data assimilation system. *Quarterly Journal of the Royal Meteorological Society*, 137, 553-  
484 597.
- 485 DIXON, P. G. & MOTE, T. L. 2003. Patterns and Causes of Atlanta's Urban Heat Island–Initiated  
486 Precipitation. *Journal of Applied Meteorology*, 42, 1273-1284.
- 487 EBITA, A., KOBAYASHI, S., OTA, Y., MORIYA, M., KUMABE, R., ONOGI, K., HARADA, Y., YASUI, S.,  
488 MIYAOKA, K. & TAKAHASHI, K. 2011. The Japanese 55-year Reanalysis" JRA-55": an interim  
489 report. *Sola*, 7, 149-152.
- 490 ELLIS, K. N., HATHAWAY, J. M., MASON, L. R., HOWE, D. A., EPPS, T. H. & BROWN, V. M. 2015. Summer  
491 temperature variability across four urban neighborhoods in Knoxville, Tennessee, USA.  
492 *Theoretical and Applied Climatology*, in press.
- 493 EPSTEIN, Y. & MORAN, D. S. 2006. Thermal comfort and the heat stress indices. *Industrial Health*, 44,  
494 388-398.
- 495 ESSLINGER, Z. A., GIERSCH, A., LANKUTIS, J. & DIXON, P. G. Observing Effects of Weather on White-tailed  
496 Deer. 2015 Annual Meeting, Association of American Geographers, 2015 Chicago, Illinois.
- 497 ESTEBAN, P., JONES, P. D., MARTÍN-VIDE, J. & MASES, M. 2005. Atmospheric Circulation Patterns Related  
498 to Heavy Snowfall Days in Andorra, Pyrenees. *International Journal of Climatology*, 25, 319 - 329.

- 499 FANGER, P. 1970. *Thermal Comfort: Analysis and applications in environmental engineering*, New York,  
500 McGraw Hill.
- 501 FRAKES, B. & YARNAL, B. 1997. A procedure for blending manual and correlation-based synoptic  
502 classifications. *International Journal of Climatology*, 17, 1381-1396.
- 503 FRANK, K., GEILS, B., KALKSTEIN, L. & THISTLE JR, H. 2008a. Synoptic climatology of the long-distance  
504 dispersal of white pine blister rust II. Combination of surface and upper-level conditions.  
505 *International Journal of Biometeorology*, 52, 653-666.
- 506 FRANK, K., KALKSTEIN, L., GEILS, B. & THISTLE JR, H. 2008b. Synoptic climatology of the long-distance  
507 dispersal of white pine blister rust. I. Development of an upper level synoptic classification.  
508 *International Journal of Biometeorology*, 52, 641-652.
- 509 GIORGI, F. & MEARNES, L. O. 1999. Introduction to special section: Regional climate modeling revisited.  
510 *Journal of Geophysical Research: Atmospheres (1984–2012)*, 104, 6335-6352.
- 511 GIRS, A. 1948. Some aspects concerning basic forms of atmospheric circulation. *Meteorol. Gidrol.*, 3, 9-  
512 11.
- 513 GOLDMAN, R. 2001. Introduction to heat-related problems in military operations. *In*: PANDOLF, K. B.,  
514 BURR, R. E., WENGER, C. B. & POZOS, R. S. (eds.) *Medical Aspects of Harsh Environments*.  
515 Washington, D.C.: Department of the Army, Office of the Surgeon General, and Borden Institute.
- 516 GREENE, J., KALKSTEIN, L., YE, H. & SMOYER, K. 1999. Relationships between synoptic climatology and  
517 atmospheric pollution at 4 US cities. *Theoretical and Applied Climatology*, 62, 163-174.
- 518 GREENE, S., KALKSTEIN, L. S., MILLS, D. M. & SAMENOW, J. 2011. An Examination of Climate Change on  
519 Extreme Heat Events and Climate-Related Mortality Relationships in Large U.S. Cities. *Weather,*  
520 *Climate, and Society*, 3, 281-292.
- 521 GRUNDSTEIN, A. 2003. A synoptic-scale climate analysis of anomalous snow water equivalent over the  
522 Northern Great Plains of the USA. *International Journal of Climatology*, 23, 871-886.

- 523 HABERLIE, A. M., ASHLEY, W. S. & PINGEL, T. J. 2015. The effect of urbanisation on the climatology of  
524 thunderstorm initiation. *Quarterly Journal of the Royal Meteorological Society*, 141, 663-675.
- 525 HAJAT, S., SHERIDAN, S. C., ALLEN, M. J., PASCAL, M., LAAIDI, K., YAGOUTI, A., BICKIS, U., TOBIAS, A.,  
526 BOURQUE, D. & ARMSTRONG, B. G. 2010. Heat-health warning systems: a comparison of the  
527 predictive capacity of different approaches to identifying dangerously hot days. *American*  
528 *journal of public health*, 100, 1137.
- 529 HANNA, A. F., YEATTS, K. B., XIU, A., ZHU, Z., SMITH, R. L., DAVIS, N. N., TALGO, K. D., ARORA, G.,  
530 ROBINSON, P. J. & MENG, Q. 2011. Associations between ozone and morbidity using the Spatial  
531 Synoptic Classification system. *Environmental Health*, 10, 15.
- 532 HAVENITH, G. 1999. Heat balance when wearing protective clothing. *Annals of Occupational Hygiene*,  
533 43, 289-296.
- 534 HAYHOE, K., SHERIDAN, S., KALKSTEIN, L. & GREENE, S. 2010. Climate change, heat waves, and mortality  
535 projections for Chicago. *Journal of Great Lakes Research*, 36, 65-73.
- 536 HEBBERN, C. & CAKMAK, S. 2015. Synoptic weather types and aeroallergens modify the effect of air  
537 pollution on hospitalisations for asthma hospitalisations in Canadian cities. *Environmental*  
538 *Pollution*, 204, 9-16.
- 539 HEWITSON, B. C. & CRANE, R. G. 2002. Self-Organizing Maps: Applications to Synoptic Climatology.  
540 *Climate Research*, 22, 13 - 26.
- 541 HONDULA, D., VANOS, J. & GOSLING, S. 2014. The SSC: a decade of climate–health research and future  
542 directions. *International Journal of Biometeorology*, 58, 109-120.
- 543 HONDULA, D. M. & DAVIS, R. E. 2011. Climatology of winter transition days for the contiguous USA,  
544 1951–2007. *Theoretical and Applied Climatology*, 103, 27-37.

- 545 HONDULA, D. M., DAVIS, R. E., KNIGHT, D. B., SITKA, L. J., ENFIELD, K., GAWTRY, S. B., STENGER, P. J.,  
546 DEATON, M. L., NORMILE, C. P. & LEE, T. R. 2013. A respiratory alert model for the Shenandoah  
547 Valley, Virginia, USA. *International Journal of Biometeorology*, 57, 91-105.
- 548 HONDULA, D. M., SITKA, L., DAVIS, R. E., KNIGHT, D. B., GAWTRY, S. D., DEATON, M. L., LEE, T. R.,  
549 NORMILE, C. P. & STENGER, P. J. 2010. A back-trajectory and air mass climatology for the  
550 Northern Shenandoah Valley, USA. *International Journal of Climatology*, 30, 569-581.
- 551 HÖPPE, P. 1999. The physiological equivalent temperature—a universal index for the biometeorological  
552 assessment of the thermal environment. *International Journal of Biometeorology*, 43, 71-75.
- 553 HUANG, C., BARNETT, A. G., WANG, X., VANECKOVA, P., FITZGERALD, G. & TONG, S. 2011. Projecting  
554 Future Heat-related Mortality under Climate Change Scenarios: A Systematic Review.  
555 *Environmental Health Perspectives*, 119, 1681-1690.
- 556 HUANG, J., TARDIF, J. C., BERGERON, Y., DENNELER, B., BERNINGER, F. & GIRARDIN, M. P. 2010. Radial  
557 growth response of four dominant boreal tree species to climate along a latitudinal gradient in  
558 the eastern Canadian boreal forest. *Global Change Biology*, 16, 711-731.
- 559 HUTH, R. 1996. An intercomparison of computer-assisted circulation classification methods.  
560 *International Journal of Climatology*, 16, 893-922.
- 561 HUTH, R., BECK, C., PHILIPP, A., DEMUZERE, M., USTRNUL, Z., CAHYNOVÁ, M., KYSELÝ, J. & TVEITO, O. E.  
562 2008. Classifications of atmospheric circulation patterns. *Annals of the New York Academy of*  
563 *Sciences*, 1146, 105-152.
- 564 INGOLE, V., JUVEKAR, S., MURALIDHARAN, V., SAMBHUDAS, S. & ROCKLOV, J. 2012. The short-term  
565 association of temperature and rainfall with mortality in Vadu Health and Demographic  
566 Surveillance System: a population level time series analysis. *Global Health Action*, 5, 44-52.

- 567 ISAKSEN, T. B., FENSKE, R. A., HOM, E. K., REN, Y., LYONS, H. & YOST, M. G. 2015. Increased mortality  
568 associated with extreme-heat exposure in King County, Washington, 1980–2010. *International*  
569 *Journal of biometeorology*, 1-14.
- 570 KALKSTEIN, A. J. & BALLING, R. C., JR. 2004. Impact of unusually clear weather on United States daily  
571 temperature range following 9/11/2001. *Climate Research*, 26, 1-4.
- 572 KALKSTEIN, L. S. & DAVIS, R. E. 1989. Weather and Human Mortality: An evaluation of demographic and  
573 interregional responses in the United States. *Annals of the Association of American*  
574 *Geographers*, 79, 44-64.
- 575 KALKSTEIN, L. S. & GREENE, J. S. 1997. An evaluation of climate/mortality relationships in large U.S. cities  
576 and the possible impacts of a climate change. *Environmental Health Perspectives*, 105, 84-93.
- 577 KALKSTEIN, L. S., GREENE, J. S., MILLS, D. & SAMENOW, J. 2011. An evaluation of the progress in  
578 reducing heat-related human mortality in major U.S. cities. *Natural Hazards*, 56, 113-129.
- 579 KALKSTEIN, L. S., GREENE, J. S., MILLS, D. M., PERRIN, A. D., SAMENOW, J. P. & COHEN, J.-C. 2008. Analog  
580 European Heat Waves for U.S. Cities to Analyze Impacts on Heat-Related Mortality. *Bulletin of*  
581 *the American Meteorological Society*, 89, 75-85.
- 582 KALKSTEIN, L. S., NICHOLS, M. C., BARTHEL, C. D. & GREENE, J. S. 1996. A New Spatial Synoptic  
583 Classification: Application to Air-Mass Analysis. *International Journal of Climatology*, 16, 983-  
584 1004.
- 585 KALNAY, E., KANAMITSU, M., KISTLER, R., COLLINS, W., DEAVEN, D., GANDIN, L., IREDELL, M., SAHA, S.,  
586 WHITE, G. & WOOLLEN, J. 1996. The NCEP/NCAR 40-year reanalysis project. *Bulletin of the*  
587 *American Meteorological Society*, 77, 437-471.
- 588 KIM, H. C., CHOI, H., NGAN, F. & LEE, P. 2014. Surface Ozone Variability in Synoptic Pattern Perspectives.  
589 *Air Pollution Modeling and its Application XXIII*. Springer.

- 590 KIRCHMAYER, U., MICHELOZZI, P., DE'DONATO, F., KALKSTEIN, L. S. & PERUCCI, C. A. 2004. A national  
591 system for the prevention of health effects of heat in Italy. *Epidemiology*, 15, S100-S101.
- 592 KNIGHT, D. B., DAVIS, R. E., SHERIDAN, S. C., HONDULA, D. M., SITKA, L. J., DEATON, M., LEE, T. R.,  
593 GAWTRY, S. D., STENGER, P. J. & MAZZEI, F. 2008. Increasing frequencies of warm and humid air  
594 masses over the conterminous United States from 1948 to 2005. *Geophysical Research Letters*,  
595 35.
- 596 KOHONEN, T., SCHROEDER, M. & HUANG, T. 2001. *Self-Organizing Maps*, Secaucus, NJ, Springer-Verlag  
597 New York, Inc.
- 598 LAMB, H. 1950. Types and spells of weather around the year in the British Isles: annual trends, seasonal  
599 structure of the year, singularities. *Quarterly Journal of the Royal Meteorological Society*, 76,  
600 393-429.
- 601 LAMB, H. H. 1972. *British Isles weather types and a register of the daily sequence of circulation patterns*,  
602 1861-1971, London, Her Majesty's Stationery Office.
- 603 LEATHERS, D. J., GRAYBEAL, D., MOTE, T., GRUNDSTEIN, A. & ROBINSON, D. 2004. The Role of Airmass  
604 Types and Surface Energy Fluxes in Snow Cover Ablation in the Central Appalachians. *Journal of*  
605 *Applied Meteorology*, 43, 1887-1899.
- 606 LEATHERS, D. J., MOTE, T. L., GRUNDSTEIN, A. J., ROBINSON, D. A., FELTER, K., CONRAD, K. & SEDYWITZ,  
607 L. 2002. Associations between continental-scale snow cover anomalies and air mass frequencies  
608 across eastern North America. *International Journal of Climatology*, 22, 1473-1494.
- 609 LEDREW, E. F. 1984. The role of local heat sources in synoptic activity within the Polar Basin.  
610 *Atmosphere-Ocean*, 22, 309-327.
- 611 LEE, C., SHERIDAN, S. & LIN, S. 2012. Relating Weather Types to Asthma-Related Hospital Admissions in  
612 New York State. *EcoHealth*, 9, 427-439.




- 613 LEE, C. C. 2015. A systematic evaluation of the lagged effects of spatiotemporally relative surface  
614 weather types on wintertime cardiovascular-related mortality across 19 US cities. *International*  
615 *Journal of Biometeorology*, in press.
- 616 MCGREGOR, G., EBI, K., BESSEMOULIN, P. & MENNE, B. 2010. Heat waves and health: Guidance on  
617 warning system development. *Report to the World Meteorological Organization and World*  
618 *Health Organization*.
- 619 MCMICHAEL, A. J., WILKINSON, P., KOVATS, R. S., PATTENDEN, S., HAJAT, S., ARMSTRONG, B.,  
620 VAJANAPOOM, N., NICIU, E. M., MAHOMED, H. & KINGKEOW, C. 2008. International study of  
621 temperature, heat and urban mortality: the 'ISOTHURM' project. *International Journal of*  
622 *Epidemiology*, 37, 1121-1131.
- 623 MOTE, T. L., LACKE, M. C. & SHEPHERD, J. M. 2007. Radar signatures of the urban effect on precipitation  
624 distribution: a case study for Atlanta, Georgia. *Geophysical Research Letters*, 34.
- 625 MULLER, R. A. 1977. A Synoptic Climatology for Environmental Baseline Analysis: New Orleans. *Journal*  
626 *of Applied Meteorology*, 16, 20-33.
- 627 NIOSH 1986. Criteria for a Recommended Standard: Occupational exposure to hot environment.  
628 Washington, D.C.: National Institute for Occupational Safety and Health.
- 629 OWEN, N. O. & DIXON, P. G. 2015. An Investigation into the Impacts of Land-Use/Land-Cover and  
630 Urbanization on Cloud-to-Ground Lightning Activity. *International Journal of Biometeorology*, in  
631 review.
- 632 PALUMBO, M. D., VILELLA, F. J., WANG, G., STRICKLAND, B. K., GODWIN, D. & DIXON, P. G. 2015.  
633 Regional Differences on Gobbling Activity of the Wild Turkey in Mississippi. *Wildlife Society*  
634 *Bulletin*, in press.
- 635 PARSONS, K. C. 2003. *Human Thermal Environments. The effects of hot, moderate and cold temperatures*  
636 *on human health, comfort and performance*, London, Taylor & Francis.

- 637 PETITTI, D. B., HONDULA, D. M., YANG, S., HARLAN, S. L. & CHOWELL, G. 2015. Multiple Trigger Points  
638 for Quantifying Heat-Health Impacts: New Evidence from a Hot Climate. *Environmental Health*  
639 *Perspectives*, in press.
- 640 POPE, C. A. & KALKSTEIN, L. S. 1996. Synoptic weather modeling and estimates of the exposure-response  
641 relationship between daily mortality and particulate air pollution. *Environmental Health*  
642 *Perspectives*, 104, 414-420.
- 643 POWER, H. C., SHERIDAN, S. C. & SENKBEIL, J. C. 2006. Synoptic climatological influences on the spatial  
644 and temporal variability of aerosols over North America. *International Journal of Climatology*,  
645 26, 723-742.
- 646 RAINHAM, D. G., SMOYER-TOMIC, K. E., SHERIDAN, S. C. & BURNETT, R. T. 2005. Synoptic weather  
647 patterns and modification of the association between air pollution and human mortality.  
648 *International Journal of Environmental Health Research*, 15, 347-360.
- 649 RAINHAM, D. G. C., SMOYER, K. & BURNETT, R. 2001. Spatial synoptic classification of air pollution and  
650 human mortality associations in Toronto, Canada: Past relationships and policy implications.  
651 *American Journal of Epidemiology*.
- 652 ROBERTS, M. J., VIDALE, P. L., MIZIELINSKI, M. S., DEMORY, M.-E., SCHIEMANN, R., STRACHAN, J.,  
653 HODGES, K., BELL, R. & CAMP, J. 2014. Tropical Cyclones in the UPSCALE Ensemble of High-  
654 Resolution Global Climate Models. *Journal of Climate*, 28, 574-596.
- 655 SAHA, S., BROCK, J. W., VAIDYANATHAN, A., EASTERLING, D. R. & LUBER, G. 2015. Spatial variation in  
656 hyperthermia emergency department visits among those with employer-based insurance in the  
657 United States—a case-crossover analysis. *Environmental Health*, 14, 20.
- 658 SCHEITLIN, K. 2013. The Maritime Influence on Diurnal Temperature Range in the Chesapeake Bay Area.  
659 *Earth Interactions*, 17, 1-14.

- 660 SCHEITLIN, K. N. & DIXON, P. G. 2010. Variations in diurnal temperature range in the Southeast due to  
661 land-use/land-cover classifications, 1995–2004. *Journal of Applied Meteorology and*  
662 *Climatology*, 49, 879-888.
- 663 SENKBEIL, J. C., RODGERS, J. C., III & SHERIDAN, S. C. 2007. The sensitivity of tree growth to air mass  
664 variability and the Pacific Decadal Oscillation in coastal Alabama. *International Journal of*  
665 *Biometeorology*, 51, 483-491.
- 666 SHEM, W. & SHEPHERD, M. 2009. On the impact of urbanization on summertime thunderstorms in  
667 Atlanta: two numerical model case studies. *Atmospheric Research*, 92, 172-189.
- 668 SHERIDAN, S. C. 2002. The redevelopment of a weather-type classification scheme for North America.  
669 *International Journal of Climatology*, 22, 51-68.
- 670 SHERIDAN, S. C., ALLEN, M. J., LEE, C. C. & KALKSTEIN, L. S. 2012a. Future heat vulnerability in California,  
671 Part II: projecting future heat-related mortality. *Climatic Change*, 115, 311-326.
- 672 SHERIDAN, S. C. & DOLNEY, T. J. 2003. Heat, mortality, and level of urbanization: measuring vulnerability  
673 across Ohio, USA. *Climate Research*, 24, 255-265.
- 674 SHERIDAN, S. C. & KALKSTEIN, L. S. 2004. Progress in Heat Watch-Warning System Technology. *Bulletin*  
675 *of the American Meteorological Society*, 85, 1931-1941.
- 676 SHERIDAN, S. C. & LEE, C. C. 2013. Synoptic Climatology. *Oxford Bibliographies in Geography*.
- 677 SHERIDAN, S. C., LEE, C. C., ALLEN, M. J. & KALKSTEIN, L. S. 2012b. Future heat vulnerability in California,  
678 Part I: projecting future weather types and heat events. *Climatic Change*, 115, 291-309.
- 679 SMOYER, K. E., RAINHAM, D. G. C. & HEWKO, J. N. 2000. Heat-stress-related mortality in five cities in  
680 Southern Ontario: 1980–1996. *International Journal of Biometeorology*, 44, 190-197.
- 681 STALLINS, J. A., CARPENTER, J., BENTLEY, M. L., ASHLEY, W. S. & MULHOLLAND, J. A. 2013. Weekend–  
682 weekday aerosols and geographic variability in cloud-to-ground lightning for the urban region of  
683 Atlanta, Georgia, USA. *Regional Environmental Change*, 13, 137-151.


- 684 TAN, J., KALKSTEIN, L., HUANG, J., LIN, S., YIN, H. & SHAO, D. 2004. An operational heat/health warning  
685 system in Shanghai. *International Journal of Biometeorology*, 48, 157-162.
- 686 URBAN, A. & KYSELÝ, J. 2014. Comparison of UTCI with other thermal indices in the assessment of heat  
687 and cold effects on cardiovascular mortality in the Czech Republic. *International Journal of*  
688 *Environmental Research and Public Health*, 11, 952-967.
- 689 URBAN, A. & KYSELÝ, J. 2015. Application of spatial synoptic classification in evaluating links between  
690 heat stress and cardiovascular mortality and morbidity in Prague, Czech Republic. *International*  
691 *Journal of Biometeorology*.
- 692 VAN BEBBER, W. J. & KÖPPEN, W. 1895. Die Isobarentypen des Nordatlantischen Ozeans und  
693 Westeuropas: ihre Beziehungen zur Lage und Bewegung der barometrischen Maxima und  
694 Minima. *Archiv der Deutschen Seewarte*, 18, 27.
- 695 VANOS, J. & CAKMAK, S. 2014. Changing air mass frequencies in Canada: potential links and implications  
696 for human health. *International Journal of Biometeorology*, 58, 121-135.
- 697 VANOS, J., CAKMAK, S., KALKSTEIN, L. & YAGOUTI, A. 2014a. Association of weather and air pollution  
698 interactions on daily mortality in 12 Canadian cities. *Air Quality, Atmosphere & Health*, 1-14.
- 699 VANOS, J. K., CAKMAK, S., BRISTOW, C., BRION, V., TREMBLAY, N., MARTIN, S. L. & SHERIDAN, S. S. 2013.  
700 Synoptic weather typing applied to air pollution mortality among the elderly in 10 Canadian  
701 cities. *Environmental research*, 126, 66-75.
- 702 VANOS, J. K., CAKMAK, S., KALKSTEIN, L. S. & YAGOUTI, A. 2015. Association of weather and air pollution  
703 interactions on daily mortality in 12 Canadian cities. *Air Quality, Atmosphere & Health*, 8, 307-  
704 320.
- 705 VANOS, J. K., HEBBERN, C. & CAKMAK, S. 2014b. Risk assessment for cardiovascular and respiratory  
706 mortality due to air pollution and synoptic meteorology in 10 Canadian cities. *Environmental*  
707 *Pollution*, 185, 322-332.

- 708 YARNAL, B. 1993. *Synoptic Climatology in Environmental Analysis: A primer*, London, Belhaven Press.
- 709 YARNAL, B., COMRIE, A. C., FRAKES, B. & BROWN, D. P. 2001. Developments and Prospects in Synoptic  
710 Climatology. *International Journal of Climatology*, 21, 1923-1950.
- 711
- 712
- 713
- 714
- 715
- 716
- 717
- 718
- 719
- 720
- 721
- 722
- 723
- 724
- 725
- 726




727


728 **Figure 1:** Map of locations with SSC data available.



729



730



731

732 **Figure 2:** Frequency distributions of each SSC category throughout the year in 10-day periods for (a) Chicago

733 (1946–2014), (b) Las Vegas (1948–2014), and (c) Miami (1948–2014).

734

735

736

737

738

739

740


741

742

743

744





745

746 **Figure 3:** Frequency distributions of each SSC category throughout the year in 10-day periods for Pune, India

747 (1973–2014).

748

749

750

751

752

753

754


755

756

757

758

759



760

761

**Figure 4:** Frequency (percentage of 6-hr intervals) of fronts, 1958–2001 (Berry et al., 2011).