



The University of  
**Nottingham**

UNITED KINGDOM · CHINA · MALAYSIA

Tibby, John and Barr, Cameron and McInerney, Francesca A. and Henderson, Andrew C.G. and Leng, Melanie J. and Greenway, Margaret and Marshall, Jonathan C. and McGregor, Glenn B. and Tyler, Jonathan J. and McNeil, Vivienne (2016) Carbon isotope discrimination in leaves of the common paperbark tree, *Melaleuca quinquenervia*, as a tool for quantifying past tropical and subtropical rainfall. *Global Change Biology* . ISSN 1365-2486

**Access from the University of Nottingham repository:**

[http://eprints.nottingham.ac.uk/32423/1/Tibby%20et%20al%20C%20isotopes%20in%20Melaleuca%20quinquenervia\\_final%20submission.pdf](http://eprints.nottingham.ac.uk/32423/1/Tibby%20et%20al%20C%20isotopes%20in%20Melaleuca%20quinquenervia_final%20submission.pdf)

**Copyright and reuse:**

The Nottingham ePrints service makes this work by researchers of the University of Nottingham available open access under the following conditions.

This article is made available under the University of Nottingham End User licence and may be reused according to the conditions of the licence. For more details see: [http://eprints.nottingham.ac.uk/end\\_user\\_agreement.pdf](http://eprints.nottingham.ac.uk/end_user_agreement.pdf)

**A note on versions:**

The version presented here may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the repository url above for details on accessing the published version and note that access may require a subscription.

For more information, please contact [eprints@nottingham.ac.uk](mailto:eprints@nottingham.ac.uk)

**Title: Carbon isotope discrimination in leaves of the common paperbark tree, *Melaleuca quinquenervia*, as a tool for quantifying past tropical and subtropical rainfall**

Running head: *Melaleuca quinquenervia* leaf carbon isotopes

**Authors**

Tibby, J.<sup>1,2\*</sup>

Barr, C.<sup>1,2</sup>

McInerney, F.A.<sup>2,3</sup>

Henderson, A.C.G.<sup>4</sup>

Leng, M.J.<sup>5,6</sup>

Greenway, M.<sup>7</sup>

Marshall, J.C.<sup>8</sup>

McGregor, G.B.<sup>8</sup>

Tyler, J.J.<sup>2,3</sup>

McNeil, V.<sup>8</sup>

1. Geography, Environment and Population, University of Adelaide, Adelaide, Australia;
2. Sprigg Geobiology Centre, University of Adelaide, Adelaide, Australia;
3. Department of Earth Sciences, University of Adelaide, Adelaide, Australia;
4. School of Geography, Politics & Sociology, Newcastle University, Newcastle upon Tyne, NE1 7RU, United Kingdom;
5. NERC Isotope Geosciences Facilities, British Geological Survey, Keyworth Nottingham, NG12 5GG, United Kingdom;
6. Centre for Environmental Geochemistry, University of Nottingham, Nottingham, NG7 2RD, United Kingdom;
7. Griffith School of Engineering, Environmental Engineering, Griffith University, Brisbane, Australia;
8. Queensland Department of Science, Information Technology, Innovation

**\*Corresponding author**

Telephone: +61 8 8313 5136; email: [john.tibby@adelaide.edu.au](mailto:john.tibby@adelaide.edu.au);

Keywords: carbon isotope ratios, palaeoclimate, CO<sub>2</sub>, discrimination, climate reconstruction, wetlands, Holocene

Paper type: Primary Research Article

## Abstract

Quantitative reconstructions of terrestrial climate are highly sought after but rare, particularly in Australia. Carbon isotope discrimination in plant leaves ( $\Delta_{\text{leaf}}$ ) is an established indicator of past hydroclimate because the fractionation of carbon isotopes during photosynthesis is strongly influenced by water stress. Leaves of the evergreen tree *Melaleuca quinquenervia* have been recovered from the sediments of some perched lakes on North Stradbroke and Fraser Islands, south-east Queensland, eastern Australia. Here, we examine the potential for using *M. quinquenervia*  $\Delta_{\text{leaf}}$  as a tracer of past rainfall by analysing carbon isotope ratios ( $\delta^{13}\text{C}$ ) of modern leaves. We firstly assess  $\Delta_{\text{leaf}}$  variation at the leaf and stand scale and find no systematic pattern within leaves or between leaves due to their position on the tree. We then examine the relationships between climate and  $\Delta_{\text{leaf}}$  for an 11 year timeseries of leaves collected in a litter tray. *M. quinquenervia* retains its leaves for 1-4 years; thus cumulative average climate data are used. There is a significant relationship between annual mean  $\Delta_{\text{leaf}}$  and mean annual rainfall of the hydrological year for 1-4 years (i.e. 365-1460 days) prior to leaf fall ( $r^2=0.64$ ,  $p=0.003$ ,  $n=11$ ). This relationship is marginally improved by accounting for the effect of  $p\text{CO}_2$  on discrimination ( $r^2=0.67$ ,  $p=0.002$ ,  $n=11$ ). The correlation between rainfall and  $\Delta_{\text{leaf}}$ , and the natural distribution of *Melaleuca quinquenervia* around wetlands of eastern Australia, Papua New Guinea and New Caledonia offers significant potential to infer past rainfall on a wide range of spatial and temporal scales.

## Introduction

Quantitative reconstructions of past climate parameters, such as temperature and precipitation, are considerably more useful than dimensionless, relative assessments of climate trends such as “wetter” or “drier”. Quantitative inferences enable direct comparison of the rate and magnitude of climate change and variability between different locations and time periods (e.g. Shakun & Carlson, 2010) and they can be used to explore fundamental questions in environmental and Earth science, such as the relationship between global atmospheric carbon dioxide concentration ( $p\text{CO}_2$ ) and temperature (Rohling *et al.*, 2012, Shakun *et al.*, 2012) and the sensitivity of rainfall to temperature change (Li *et al.* 2013). Furthermore, numerical climate estimates can be compared with climate models to assess their veracity (Phipps *et al.*, 2013) and used as inputs into hydrological (Henley *et al.*, 2011) and/or ecological (Prowse *et al.*, 2013, Verschuren *et al.*, 2000) models. Importantly, quantitative reconstructions can be calibrated with (Emile-Geay *et al.*, 2013b), and tested against (Verschuren *et al.*, 2000), instrumental records, thereby facilitating an estimate of the precision and accuracy of proxy-climate relationships.

Numerical estimates of prehistoric climate change from terrestrial archives are limited in time and space. Many records span less than 1000 years or are spatially restricted (e.g. ice core or coral records). The Australian continent, in particular, has a very limited number of quantitative palaeoclimate reconstructions extending more than 1000 years. The majority of Australian proxy rainfall records, such as those derived from corals and tree rings, are relatively short (< 500 years, e.g. Lough, 2007, O'Donnell *et al.*, 2015), and even though individual records can be merged to lengthen the temporal coverage, this process reduces the proxy's ability to detect low frequency variability (Ault *et al.*, 2013, Emile-Geay *et al.*, 2013a). As a result of the

limited number of archives, key quantitative reconstructions of Australian climate rely on archives that are distant from the Australian mainland (Neukom & Gergis, 2012). These reconstructions are, therefore, based on teleconnections that may not be stable through time (Gallant *et al.*, 2013). Quantitative climate estimates >1000 years in length include temperature reconstructions from pollen (Fletcher & Thomas, 2010, Lloyd & Kershaw, 1997) and branched glycerol dialkyl glycerol tetraethers (Woltering *et al.*, 2014). Although some (pollen-based) rainfall reconstructions exist (Kershaw *et al.*, 2004a, Kershaw & Nix, 1988, Kershaw *et al.*, 2004b), the majority of Australian hydrological reconstructions consist of non-quantitative lake water balance and speleothem oxygen isotope records (e.g. Barr *et al.*, 2014, Denniston *et al.*, 2013, Gouramanis *et al.*, 2013).

Carbon isotope discrimination recorded in leaf tissue ( $\Delta_{\text{leaf}}$ ) reflects the degree of moisture stress experienced by plants, along with factors including photosynthetic rate (Farquhar *et al.*, 1989). Importantly,  $\Delta_{\text{leaf}}$  can be quantitatively related to variables driving moisture stress including vapour pressure deficit (Brett *et al.*, 2014, Turney *et al.*, 1999) and rainfall (Diefendorf *et al.*, 2010, Kohn, 2010). The relationship between leaf carbon isotope ratios ( $\delta^{13}\text{C}$ ) and rainfall has been predominantly used to infer rainfall in pre-Quaternary settings since lengthy (i.e. > 1000 year) preservation of leaves in late Quaternary deposits is rare (Birks, 2001, Spicer, 1989). However, the potential of this proxy was demonstrated by Liu *et al.* (2011) who used  $\delta^{13}\text{C}$  in plant remains from a mixture of species to infer relative changes in rainfall during the Little Ice Age in north-western China.

We have discovered a number of perched acid lakes on North Stradbroke and Fraser Islands, south-east Queensland, Australia, that preserve leaves of the tree species *Meleleuca quinquenervia* (Cav.) S.T. Blake (the broad-leaved paperbark;

family Myrtaceae) in sediments deposited throughout much of the Holocene. In this study we assess the nature and strength of the rainfall- $\Delta_{\text{leaf}}$  relationship in *M. quinquenervia* to evaluate its potential as a rainfall proxy. Firstly, we characterize within-leaf and within-stand  $\Delta_{\text{leaf}}$  variability and then evaluate the relationship between rainfall and  $\Delta_{\text{leaf}}$  in *M. quinquenervia* by determining  $\Delta_{\text{leaf}}$  from an 11-year long monthly collection of leaves. Finally, we identify factors that lead to leaf preservation and map wetlands that have the potential to preserve sub-fossil leaves of *M. quinquenervia*.

### Carbon isotope discrimination in leaves

$\Delta_{\text{leaf}}$  reflects the net photosynthetic carbon isotope fractionation relative to the  $\delta^{13}\text{C}$  of atmospheric  $\text{CO}_2$ .  $\Delta_{\text{leaf}}$  varies among plants with different photosynthetic pathways; i.e.  $\text{C}_3$ ,  $\text{C}_4$  and CAM. *M. quinquenervia* utilises the  $\text{C}_3$  photosynthetic pathway (Kattge *et al.*, 2011). Within  $\text{C}_3$  plants, environmental variables such as the availability of water,  $p\text{CO}_2$  and irradiance can further modify  $\Delta_{\text{leaf}}$  (Cernusak *et al.*, 2013; Shubert and Jahren, 2012). The effects of irradiance are particularly evident in closed canopy environments (Graham *et al.*, 2014).

#### *Relationship between $\Delta_{\text{leaf}}$ and precipitation in $\text{C}_3$ plants*

Plants preferentially fix  $^{12}\text{C}$  over  $^{13}\text{C}$  during photosynthesis. This preference is expressed as carbon isotope discrimination and is defined as:

$$\Delta_{\text{leaf}} = \frac{(\delta_{\text{atm}} - \delta_{\text{leaf}})}{1 + (\delta_{\text{leaf}}/1000)} \quad (1)$$

Where  $\delta_{\text{atm}}$  and  $\delta_{\text{leaf}}$  represent the  $\delta^{13}\text{C}$  of atmospheric  $\text{CO}_2$ , and the leaf, respectively (Farquhar et al., 1982).

The most widely used model for carbon isotope discrimination during  $\text{C}_3$  plant photosynthesis, developed in Farquhar *et al.* (1982), explains carbon isotope discrimination as the result of the isotopic effects of diffusion of  $\text{CO}_2$  through stomata and carboxylation by RuBisCo during photosynthesis, such that:

$$\Delta_{\text{leaf}} = a + (b - a) \left( \frac{c_i}{c_a} \right) \quad (2)$$

where  $a$  and  $b$  are constants that represent fractionation due to diffusion (4.4‰) and carboxylation (27-29‰), respectively;  $c_i$  and  $c_a$ , represent the partial pressure of  $\text{CO}_2$  ( $p\text{CO}_2$ ) inside the leaf, and the ambient atmosphere, respectively (Cernusak *et al.*, 2013, Farquhar *et al.*, 1989). Although more complicated models exist, this model sufficiently describes photosynthetic carbon isotope discrimination in many different settings (Cernusak, *et al.* 2013). In this model carbon isotope discrimination is entirely a function of the  $c_i/c_a$  ratio.

Key factors determining  $c_i/c_a$  are the supply of  $\text{CO}_2$  to the leaf and the photosynthetic demand for  $\text{CO}_2$ . Supply of  $\text{CO}_2$  is a function of stomatal conductance (i.e. the rate of passage of gases through the stomata). Stomatal conductance also controls the flux of water out of the leaf via transpiration, creating a trade-off between water loss and  $\text{CO}_2$  uptake. As moisture availability increases, plants can afford to increase stomatal conductance, increasing  $c_i/c_a$  and increasing carbon isotope discrimination (Cernusak *et al.*, 2013; Diendendorf *et al.*, 2010). On the demand side of the equation, light limitation can lower the photosynthetic fixation of  $\text{CO}_2$  leading to an increase in  $c_i$ , thereby increasing  $c_i/c_a$  and  $\Delta_{\text{leaf}}$ . Our study species, *M.*

*quinquenervia*, lives in relatively open wetland environments and forms the uppermost canopy layer. As a result, canopy shading is unlikely to have a large influence on  $\Delta_{leaf}$ , but self-shading could have an effect and is considered in this study.

Furthermore,  $\Delta_{leaf}$  can be altered by the concentration of carbon dioxide in the atmosphere ( $pCO_2$ ) (Schubert & Jahren, 2012). This relationship is described by the hyperbolic equation (Schubert and Jahren, 2012):

$$\Delta_{leaf} = \frac{[(28.26)(0.21)(pCO_2+25)]}{[28.26 + (0.21) \times (pCO_2+25)]} \quad (3)$$

In this equation, the value of 28.26 represents the asymptote or maximum carbon isotope discrimination as determined empirically through experiments on a number of plants grown under a wide range of  $pCO_2$  and high nutrient and water availability (Schubert and Jahren, 2012).

The relationship between  $pCO_2$  and  $\Delta_{leaf}$  is steepest at low  $pCO_2$  concentrations. Hence, the  $pCO_2$ -induced change in  $\Delta_{leaf}$  through the last glacial-interglacial cycle is argued to exceed 2‰ (Schubert and Jahren, 2012). In the Holocene, the change in  $\Delta_{leaf}$  resulting from the shift in  $pCO_2$  from 251 ppm at the start of the Holocene to 281 ppm at 600 year BP (Monnin *et al.*, 2004) would result in a shift of approximately 0.6‰ in  $\Delta_{leaf}$ . This change is noteworthy compared to those resulting from moisture stress, with a 0.6‰ shift in  $\Delta_{leaf}$  equivalent to that induced by an increase in mean annual rainfall from 1000 to 1285 mm based on the relationship observed for evergreen angiosperms in Diefendorf *et al.* (2010).

*Global rainfall-carbon isotope relationships*



Global syntheses of rainfall- $\Delta_{\text{leaf}}$  relationships demonstrate the potential to quantitatively infer past rainfall from (sub)fossil leaves (Diefendorf *et al.*, 2010, Kohn, 2010). Diefendorf *et al.* (2010) found a statistically significant positive relationship between mean annual rainfall ( $\log_{10}$  transformed) and  $\Delta_{\text{leaf}}$  ( $r^2=0.55$ ,  $p<0.0001$ ). Notably, this global relationship masks variability at the continental scale. For example, there is no statistically significant relationship exhibited in Europe ( $r^2=0.025$ ,  $p=0.125$ ). Kohn (2010) developed a more complex model that includes the influence of altitude and latitude. It showed similarly strong relationships between rainfall and  $\Delta_{\text{leaf}}$ , particularly below  $\sim 1700 \text{ mm a}^{-1}$ , with a largely constant isotopic discrimination at higher rainfall amounts.

While these meta-analyses point to the potential to reconstruct rainfall from  $\Delta_{\text{leaf}}$ , only  $\sim 50\%$  of the variance (on log scales) in isotope discrimination is explained by mean annual precipitation (Diefendorf *et al.*, 2010, Kohn, 2010). Substantial  $\Delta_{\text{leaf}}$  variation between different species has been observed under the same water availability regime in natural (Cernusak *et al.*, 2013) and experimental (Cernusak *et al.*, 2009) settings. The between-species variability in rainfall- $\Delta_{\text{leaf}}$  relationships is so large that it has been argued  $\Delta_{\text{leaf}}$  cannot be used to infer past precipitation from a mixed leaf fossil assemblage (Freeman *et al.*, 2011).

### *Rainfall-carbon isotope relationships in Australia*

Statistically significant relationships between rainfall and  $\delta^{13}\text{C}_{\text{leaf}}$  of plant communities (Stewart *et al.*, 1995),  $\delta^{13}\text{C}_{\text{leaf}}$  of  $\text{C}_3$  grasses (Murphy & Bowman, 2009) and  $\delta^{13}\text{C}$  in charcoal (Turney, 2012) have been demonstrated in Australia across broad spatial gradients. By contrast, Schulze *et al.* (1998) noted that  $\Delta_{\text{leaf}}$  did not increase above 475 mm in mixed species samples from Northern Australia, and only seven out

of thirteen *Eucalyptus* species from the same study region showed significant (positive) relationships between  $\Delta_{\text{leaf}}$  and rainfall (Miller *et al.*, 2001). These studies were undertaken across broad spatial gradients, mirroring research in other locations (e.g. Ehleringer, 1993, Weiguo *et al.*, 2005). Spatial sampling regimes have the potential to overestimate the explanatory power of rainfall due to co-variation with confounding factors (e.g. changes in soil type and moisture, elevation and aspect) and spatial autocorrelation (see discussion in Telford & Birks, 2009). In contrast to previous mixed species and spatial study designs, this study circumvents these uncertainties by sampling leaves of *M. quinquenervia* over a period of 11 years at a single site.

#### *Spatial distribution of Melaleuca quinquenervia*

*M. quinquenervia* is native to the eastern coastal margin of Australia (between 8–34°S) and to Papua New Guinea (PNG) and New Caledonia (Fig. 1a). It is generally found proximal to wetlands as its seeds have an obligate need for moist substrates for propagation, although mature trees will survive in locations with low soil moisture (Doran & Turnbull, 1997). *M. quinquenervia* has a shallow root system, with over 80% of the root biomass occurring in the top 15 cm of the soil in our Carbrook Wetlands study site (Bradley, 1996). In Australia and PNG, *M. quinquenervia* mostly grows at altitudes of less than 100 m above sea level, while in New Caledonia, where *M. quinquenervia* is the dominant savannah tree (Ibanez *et al.*, 2013), it grows at altitudes up to 1000 m. The mean maximum temperature of the hottest month over *M. quinquenervia*'s range in Australia is 26–34°C (Doran & Turnbull, 1997), while the mean minimum temperature of the coolest month is between 4 and 20°C (Doran & Turnbull, 1997). The species occurs across a wide

rainfall gradient, ranging from 400 to 4000 mm per annum (Fig. 1b). *M.*

*quinquenervia* is an invasive species in the Florida Everglades, USA and a substantial literature discusses the reasons for, and possible solutions to, this problem (Dray Jr *et al.*, 2006).

## **Regional Setting**

Our study analyses *M. quinquenervia* from two locations in south-east Queensland, Australia: Carbrook Wetlands (Fig. 1c), where a long-term leaf collection was undertaken, and Swallow Lagoon, North Stradbroke Island, where we analysed leaf and stand-scale variation in  $\Delta_{\text{leaf}}$ . Carbrook Wetlands are located in the catchment of Native Dog Creek and the Logan River Floodplain (27.690 S, 153.276 E) (Fig. 1c). *M. quinquenervia* forest occupies an area of 164 ha in the wetland, which is described by Greenway (1994). Mean annual rainfall at nearby Mount Cotton Farm (Australian Bureau of Meteorology station number 040460), 7.2 km north of Carbrook Wetlands, (Fig. 1d) is 1289 mm (annual rainfall range: 634–2499 mm,  $n=34$ ). Mean annual maximum temperature at the Logan City Water Treatment Plant, approximately 5 km west of Carbook Wetland (Bureau of Meteorology station number 40854) is 26°C, with the highest mean monthly maximum temperature in January of 29°C and the lowest mean monthly maximum of 21.4°C in July.

Swallow Lagoon (27.499 S, 153.455 E) is a small (<1.5 ha) relatively deep (5.8 m maximum depth) perched freshwater lake (Marshall & McGregor, 2011) (maximum recorded elevation of the lake surface: 154 m AHD, April 2011) on North Stradbroke Island, a large (285 km<sup>2</sup>) sand island in Moreton Bay, Queensland (Barr *et al.*, 2013). Swallow Lagoon is fringed by *M. quinquenervia*, which forms a narrow

ring around the lake at a distance of less than 10 m from the water's edge and in water up to 0.5 m deep (in March 2014).

## Material and Methods

The weight of *M. quinquenervia* litterfall and its carbon isotope discrimination was analysed from a leaf litter tray on the Carbrook Wetlands floodplain, 100 m from Native Dog Creek. The litter tray was located in a position which had 2175 trees ha<sup>-1</sup>, with a mean height of 18.6 m ( $\pm 4$  m, 1 s.d.) and a mean diameter at breast height of 17.8 cm ( $\pm 9$  cm 1 s.d.) (Greenway, 1994). The density of the trees, and collection of windblown leaves of species which did not overhang the tray, indicates the tray integrates leaf fall from a number of canopy trees, rather than a single individual (Greenway, 1994). Litterfall was collected from a raised 0.25 m<sup>2</sup> (0.5 m  $\times$  0.5 m) leaf litter tray at *ca.* four-weekly intervals between April 1992 and July 2003. We thereby extend the record of Greenway (1994) through provision of more recent data (April 1994–July 2003) and additional analyses. Hereafter, this data set is referred to as the “temporal” data set, with the sampling interval described herein as “monthly”. Litterfall was sorted into a number of different components (including leaves divided into “young” – identified by their covering of silky white hairs which disappear as the leaf matures – and “total” categories), oven dried at 40°C and weighed. These data provide an understanding of the phenology of *M. quinquenervia* and, in particular, identify the period of new leaf formation and highest leaf production. Leaves were stored in paper bags until 2011. From each monthly collection, a 1 cm slice from the leaf tip was sampled from a mean of nine leaves for  $\delta^{13}\text{C}$  analysis.

We examined variation in *M. quinquenervia*  $\Delta_{\text{leaf}}$  at the leaf, tree and stand scale from trees fringing Swallow Lagoon. Discrimination within a leaf may vary

with distance from the vein due to increased diffusive pathway length (Seibt *et al.*, 2008). The leaves of *M. quinquenervia* have five longitudinal veins (as indicated by the species epithet), which may reduce the effects of diffusive length. We nevertheless undertook experiments to assess within leaf variation in  $\Delta_{\text{leaf}}$ . In these experiments, leaves were cut into 12 sections (Fig. 2) and  $\Delta_{\text{leaf}}$  determined on each subsample. At the stand scale,  $\Delta_{\text{leaf}}$  may be influenced by factors including the leaf's position in the canopy (Marshall *et al.*, 2007), orientation on the tree (as has been observed by Lockheart *et al.*, 1998) and edaphic factors such as temperature and water availability (Diefendorf *et al.*, 2010, Kaplan *et al.*, 2002). Hence, we examined the difference between north facing (more sun exposed) and south facing (more shaded) leaves on four trees and the variation between these trees growing at positions 1 m outside the lake, at the lake edge, and at 3 m and 6 m inside the lake.

All leaf samples were washed in deionised water, freeze dried and milled to a fine powder prior to  $\delta^{13}\text{C}$  analysis.  $\delta^{13}\text{C}$  analyses were performed on a Costech Elemental Analyser coupled to a VG TripleTrap and Optima dual-inlet mass spectrometer, with  $\delta^{13}\text{C}$  values reported on the VPDB scale using a within-run laboratory standard calibrated against NBS-18, NBS-19 and NBS-22. Replicate analysis of well-mixed samples indicated a precision of  $\pm <0.1\%$  VPDB (1 s.d.).

$\Delta_{\text{leaf}}$  was calculated with equation 1 using  $\delta^{13}\text{C}_{\text{atm}}$  derived from an average of monthly values for the two years prior to leaf sample collection as measured at the Mauna Loa Observatory, Hawaii, by the United States of America's NOAA Earth System Research Laboratory, Global Monitoring Division ([www.esrl.noaa.gov/gmd](http://www.esrl.noaa.gov/gmd)). We also derived a relationship that removes the effect of  $p\text{CO}_2$  on  $\Delta_{\text{leaf}}$  by calculating the difference between  $\Delta_{\text{leaf}}$  predicted from  $p\text{CO}_2$  (equation 3) and the  $\Delta_{\text{leaf}}$  from the carbon isotope values measured in our leaves (equation 1). This calculation used

$p\text{CO}_2$  data from Mauna Loa Observatory for 2 years prior to dates of leaf collection. This difference between  $p\text{CO}_2$ -predicted and measurement-based discrimination is referred to as the discrimination anomaly.

While the primary aim of our study is to investigate the relationship between *M. quinquenervia*  $\Delta_{\text{leaf}}$  values and rainfall, some studies (e.g. Diefendorf *et al.*, 2010) have shown that other climate variables (e.g. temperature) may be of secondary importance. Hence, we investigated the degree to which variability in *M. quinquenervia*  $\Delta_{\text{leaf}}$  was explained by a range of potential environmental variables, namely: rainfall, minimum and maximum daily temperature, pan evaporation, precipitation divided by evaporation (hereafter P/E), solar radiation - hereafter termed “radiation”, vapour pressure and relative humidity at both maximum and minimum temperature. Precipitation data were taken from the Mount Cotton Farm rainfall station (Fig. 1). Where no rainfall data were available for a particular month ( $n=4$ ), we estimated rainfall amount from a linear relationship, developed from the remaining study period, between that month’s rainfall and that of the adjacent month most correlated to the month with a single missing value. Other climate data were derived from the Queensland Department of Science, Information Technology and Innovation’s Scientific Information for Land Owners dataset which derives synthetic climate data interpolated across space from variables measured at climate stations (Jeffrey *et al.*, 2001).

Our analyses of the relationships between climate and  $\Delta_{\text{leaf}}$  were undertaken using two temporal data sets. The first determines the relationship between the individual  $\Delta_{\text{leaf}}$  values collected at monthly intervals and climate data for which composite means were calculated for a range of time windows prior to the month of

leaf collection. Climate data in the year immediately prior to leaf fall was excluded because very few young (<1 year old) leaves were collected in the tray (Fig. 3). Thus, the time composites encompassed periods of 1 to 2 years, 1 to 3 and 1 to 4 years prior to leaf fall. For the second set of analyses, annual mean  $\Delta_{\text{leaf}}$  was calculated for each hydrological year (November 1st to October 31st). Annual mean values were compared to both mean rainfall for the hydrological year, and individual month's rainfall, again composited for periods of between 1-2 years and 1-4 years before leaf collection.

In order to identify sites where *M. quinquenervia* leaves may be preserved we identified where this species grows close to wetlands that may favour preservation. Vegetation associations (known as regional ecosystems; Neldner *et al.*, 2012) with *M. quinquenervia* as the dominant or sub-dominant canopy tree were mapped where they fell within 30 m of lacustrine or permanent palustrine wetlands using the Queensland Government's Regional Ecosystem Description Database ([www.qld.gov.au/environment/plants-animals/plants/ecosystems/about/#redd](http://www.qld.gov.au/environment/plants-animals/plants/ecosystems/about/#redd)) vegetation associations. The location of the wetlands was identified from satellite imagery, with all wetlands < 8 ha classified as palustrine and the distinction between lacustrine or palustrine for wetlands > 8 ha based on whether open water (lacustrine) or vegetation (palustrine) dominated the site (Neldner *et al.*, 2012). To further identify sites of, and conditions favouring preservation, we collected sediment cores from 10 lakes and swamps on North Stradbroke and Fraser Islands using piston (Livingstone, 1955) or side sampling (Jowsey, 1966) corers. Continuous sediment sections (between 1 and 5 cm in width) from these cores were sieved with a 250  $\mu\text{m}$  sieve, with the retained material inspected for *M. quinquenervia* leaves. Lastly, at Swallow Lagoon we used a YSI 6920 multi-parameter sonde to determine the lake water dissolved

oxygen profile approximately monthly between June 2011 and May 2012, while historical dissolved oxygen data from Brown Lake, North Stradbroke Island (see supporting information S3) are also discussed.

## Results

### *Phenology of Melaleuca quinquenervia*

*M. quinquenervia* produces new leaves throughout the year but with marked seasonality in production. Peak leaf production (young leaf fall) commences in June and is greatest in August–November (Fig. 3). It is unlikely that young leaf fall is related to wind stress since the period of maximum abscission of older leaves frequently occurs after the abscission of young leaves in the seasonal cycle (Fig. 3). The small number of young leaves collected (<1% of leaf weight over the sampling period) indicates that most leaves remain on the tree for over one year.

### *Intra-leaf $\Delta_{\text{leaf}}$ variation and variation within a stand*

In the three leaves analysed, the within-leaf variability of  $\Delta_{\text{leaf}}$  was less than 1‰. The mean  $\Delta_{\text{leaf}}$  for the 12 sub-samples from the three leaves was 23.82‰  $\pm$  0.29, 24.48‰  $\pm$  0.18 and 23.85‰  $\pm$  0.22 ( $\pm$  1 standard deviation [s.d], Fig. 4). Moreover, the leaf tips – equivalent to the leaf fraction sampled in our temporal sampling – have  $\Delta_{\text{leaf}}$  values within the range measured for the other parts of the leaf (Fig. 4a). We also found no systematic differences between central and marginal leaf  $\Delta_{\text{leaf}}$  for individual leaves (see Fig. 4b). The  $\Delta_{\text{leaf}}$  of north (mean=24.40‰, s.d.=1.16‰, n=4) and south (mean=24.44‰, s.d.=0.83‰, n=4) facing samples of four trees at Swallow Lagoon exhibited no systematic difference (Fig. 5).

### *Temporal relationship between rainfall and $\Delta_{\text{leaf}}$ of *M. quinquenervia**



The  $\Delta_{\text{leaf}}$  of *M. quinquenervia* collected from April 1992 to July 2003 in Carbrook Wetlands ranged from 20.46‰ to 24.90‰, (mean: 22.63‰, s.d.: 0.9‰,  $n=137$ ) (supporting information S1). Carbon isotope discrimination systematically increased in 1996 (Figure 6), with values before August 1996 rarely exceeding 22.50‰ (mean: 21.89‰) and after this time mostly exceeding 22.50‰ (mean: 23.09‰). Importantly, the shift in  $\Delta_{\text{leaf}}$ , corresponds to an increase in rainfall at nearby Mount Cotton Farm in the 24 months prior to leaf fall (Fig. 6).

The relationship between *M. quinquenervia*  $\Delta_{\text{leaf}}$  and climate was first examined using the monthly  $\Delta_{\text{leaf}}$  data. All the climate variables analysed, with the exception of maximum temperature, were significantly, but weakly, correlated with monthly  $\Delta_{\text{leaf}}$  data for at least one of the time periods considered (1-2, 1-3 and 1-4 years before leaf fall) (Table 1). However, spurious correlations can emerge from correlation of biological data with multiple climate variables over multiple timescales. This is particularly problematic in this study as *M. quinquenervia* leaves remain on the tree for a variable period of time (between 1-4 years). Therefore, each monthly collection provides a time-averaged sample spanning up to four years prior to leaf collection, resulting in a need to examine relationships to climate over varied time intervals. In addition, unlike the litter collection, natural accumulations of leaves in sediments are mixed over multiple years, calling for a calibration that examines climate relationships relevant to that coarser sampling resolution. As a result of these considerations, we focus the remaining analysis on the relationship between climate and  $\Delta_{\text{leaf}}$  pooled into yearly means.

When the relationship between climate and annual mean  $\Delta_{\text{leaf}}$  is derived, only two climate variables exhibited significant relationships to  $\Delta_{\text{leaf}}$ : rainfall 1-4 years before leaf collection and radiation 1-4 years before leaf collection (Table 2). Rainfall and radiation are significantly correlated ( $r^2=0.54$ ,  $p=0.011$ ). Based on this outcome, we developed rainfall calibrations that can be applied to sub-fossil records (Table 3). One relationship is based directly on  $\Delta_{\text{leaf}}$  and one takes into account the effect of  $p\text{CO}_2$  on  $\Delta_{\text{leaf}}$  (Schubert & Jahren, 2012) (see methods). There was little difference in the strength of the isotope-rainfall relationship based on  $\Delta_{\text{leaf}}$  and discrimination anomaly (Fig. 7), due to the linear increase in atmospheric  $p\text{CO}_2$  through the study period ([www.esrl.noaa.gov/gmd](http://www.esrl.noaa.gov/gmd)). However, given the influence of  $p\text{CO}_2$  variation on  $\Delta_{\text{leaf}}$  through time, the application of the model that accounts for this effect is preferable. The relationship between rainfall 1–4 years before leaf collection and the anomaly (which accounts for the influence  $p\text{CO}_2$  on  $\Delta_{\text{leaf}}$ ) is slightly stronger ( $r^2=0.67$   $p=0.002$ ,  $n=11$ ) (Fig. 7) than that between  $\Delta_{\text{leaf}}$  and rainfall over the same period ( $r^2=0.64$ ,  $p=0.003$ ,  $n=11$ ).

## Discussion

### *Isotope systematics of leaves*

Analysis of intra-leaf variability shows no discernible difference in  $\Delta_{\text{leaf}}$  values between the leaf tip and other parts of the leaf (Fig. 4). This suggests the leaf tip  $\Delta_{\text{leaf}}$  used in our temporal study represents the whole leaf  $\Delta_{\text{leaf}}$ . Additionally, it appears any given leaf fragment (e.g. those occurring in sub-fossil deposits) can provide a representative whole leaf  $\Delta_{\text{leaf}}$  value for *M. quinquenervia*. Furthermore, our analysis shows leaf position on individual trees (north vs south facing), and the position of the trees relative to the water's edge do not bias the isotopic analysis as there are no

systematic differences between leaves or trees in different positions. Therefore, leaf position on the tree, and the tree's location relative to the water's edge, should have a minimal impact on  $\Delta_{\text{leaf}}$ .

#### *Relationship between *M. quinquenervia* $\Delta_{\text{leaf}}$ and climate*

Our analysis of monthly  $\Delta_{\text{leaf}}$  for *M. quinquenervia* for an 11 year period reveals significant but weak correlations with all of the climate variables examined on at least some of the time intervals prior to leaf collection. The numerous weak correlations reflect the complexity of the relationship between climate and  $\Delta_{\text{leaf}}$  on short timescales. However, when the climate and leaf data are pooled into yearly means, a process akin to filtering the climate data, then only rainfall and radiation (in the 1-4 years preceding leaf fall) are significant. The significant relationship ( $r^2=0.64$ ,  $p<0.003$ ,  $n=11$ ) between  $\Delta_{\text{leaf}}$  and rainfall (Table 2; Fig. 7) mirrors observations made at the scale of individual plant species (Van de Water *et al.*, 2002), communities (Prentice *et al.*, 2011, Stewart *et al.*, 1995) and the globe (Diefendorf *et al.*, 2010, Kohn, 2010). This study, however, assesses the relationship between  $\Delta_{\text{leaf}}$  and rainfall at a single location, rather than across a mean annual precipitation gradient as used in most previous studies. Hence, our analysis represents a robust assessment of rainfall-isotope relationships as it negates effects associated with (spatial) autocorrelation frequently encountered in calibration data sets (Telford & Birks, 2009). The strong relationship between this species and rainfall is in contrast to that observed in other studies (Miller *et al.*, 2001, Schulze *et al.*, 1998, Van de Water *et al.*, 2002). *M. quinquenervia*'s adaptation to a wide range of environments with varying moisture availability (see Fig. 1b) may contribute to the strong relationship observed.

#### *The potential to quantify past tropical and subtropical rainfall*

A review of the locations in eastern Australia where *M. quinquenervia* leaves may preserve highlights enormous potential to establish quantitative rainfall reconstructions from the  $\Delta_{\text{leaf}}$ -rainfall relationships described herein. Our analysis shows there are over 200 permanent wetlands in the Australian state of Queensland where *M. quinquenervia* grows within 30 m of the wetland (Fig. 8).

To investigate whether *M. quinquenervia* leaves are preserved in some of these sites, we recovered sediments from ten lakes and swamps on the sand islands North Stradbroke Island and Fraser Island (Table 3). On these islands, three out of the ten wetlands surveyed preserved leaves of *M. quinquenervia* and two also preserved leaves of *Eucalyptus* (Table 3). *M. quinquenervia* has highly resistant sclerophyllous leaves (Li *et al.*, 2009), which favours their preservation, however sclerophylly does not ensure preservation since there was no preserved material found at a number of other sites we sampled (Table 3). In addition, no leaves were preserved in sediments of Lachlan Nature Reserve swamp, Sydney, Australia (R. Hamilton pers. comm.), a wetland which has no permanent standing water but has *M. quinquenervia* as the current canopy dominant (Hamilton & Penny, 2015). As a result, we hypothesise the preservation of leaves is dependent on anoxia, which reduces chemical and biological breakdown (e.g. by macroinvertebrate “shredders” Li *et al.*, 2009). Anoxia in these low nutrient lake systems is likely to result from their physical characteristics, in particular the ratio of lake surface area to both depth and water volume, and the degree of protection afforded from wind driven overturning by the surrounding terrain (including vegetation) (Hondzo & Stefan, 1996). As an example, monitoring the deepest area of Swallow Lagoon at approximately monthly intervals between June 2011 and May 2012 found that it experiences prolonged anoxia (near bed dissolved

oxygen typically < 1 ppm; supporting information S3) for most of the year due to its sheltered position and small surface area (< 1.5 ha). By contrast, larger more exposed lakes on North Stradbroke Island typically have higher oxygen concentrations at depth. Blue Lake never experienced anoxia for the period 1996–2002 (Barr et al. 2013) and lakebed anoxia was only recorded once in Brown Lake for the period 1992–2001 ( $n=47$ ; supporting information S3). Neither of these sites preserved leaves (Table 3). Beyond our investigations, there are large numbers of potential sites for preservation of *M. quinquenervia* in Queensland, Australia (see Fig. 8). Outside Australia *M. quinquenervia* has been observed on the margins of palaeoecological sites in New Caledonia (Stevenson *et al.*, 2001).

### *Conclusions*

The annual mean carbon isotope discrimination of *M. quinquenervia* leaves collected over an 11-year period exhibits a statistically significant relationship to rainfall. In the modern day, the effect of  $p\text{CO}_2$  is largely negligible for our samples; however, for application to the fossil record, changes in  $p\text{CO}_2$  even during the Holocene has an important effect (Schubert & Jahren, 2015). Variation within leaves and within the canopy is not systematically offset and is small relative to the overall rainfall-isotope relationship. Therefore the utility of  $\Delta_{\text{leaf}}$  in *M. quinquenervia* as a proxy for past rainfall variability has great potential in locations where leaves are preserved.

An analysis of the distribution of *M. quinquenervia* reveals a large number of wetlands across the Australian eastern seaboard that may contain sub-fossil leaves within their sediments. A survey of ten lakes on North Stradbroke Island and Fraser Island found three lakes with abundant preservation within the sediments. The

rainfall- $\Delta_{\text{leaf}}$  relationship, widespread distribution and demonstrated preservation of *M. quinquenervia* leaves, indicate substantial potential for reconstructing past variability in tropical and subtropical precipitation.

### **Acknowledgements**

We thank the Minjerribah Moorgumpin elders, in particular Aunty Marg, for permission to undertake research on North Stradbroke Island. Joanna Blessing, Sarah Lindermann, Norbert Menke, Patrick Moss, Tim Page, Lynda Petherick, Peter Negus, Penny Rogers, Melanie Shaw, Bill Senior and Craig Sloss provided assistance in the field. We would also like to thank Paul Smith for all his assistance with our research on North Stradbroke Island and John Ferris for Brown Lake dissolved oxygen data. Christine Crothers drew Figures 1a and b. Ross (Stan) Mitchell undertook the spatial analysis presented in Figure 8. Chris Kendrick undertook the isotope analyses. This manuscript was improved by the comments of two anonymous reviewers. This project was funded by Australian Research Council (ARC) grants LP34106364 (with co-investment from The Queensland Government and Sibelco Australia) and DP150103875, an ARC Future Fellowship (FT110100793) and Discovery Project (DP130104314) held by FAM and a University of Adelaide interdisciplinary research fund grant to FAM, JT, JJT and CB.

**Supporting information captions**

S1: *Melaleuca quinquenervia*  $\Delta_{\text{leaf}}$  from Carbook Wetland April 1992-July 2003.

S2: Monthly and annual rainfall from Mt Cotton Farm

S3: Swallow Lagoon and Brown Lake, North Stradbroke Island dissolved oxygen concentrations

## References

- Ault TR, Cole JE, Overpeck JT *et al.* (2013) The continuum of hydroclimate variability in western North America during the last millennium. *Journal of Climate*, **26**, 5863-5878.
- Barr C, Tibby J, Gell P, Tyler J, Zawadzki A, Jacobsen GE (2014) Climate variability in south-eastern Australia over the last 1500 years inferred from the high-resolution diatom records of two crater lakes. *Quaternary Science Reviews*, **95**, 115-131.
- Barr C, Tibby J, Marshall JC, Mcgregor GB, Moss PT, Halverson GP, Fluin J (2013) Combining monitoring, models and palaeolimnology to assess ecosystem response to environmental change at monthly to millennial timescales: The stability of Blue Lake, North Stradbroke Island, Australia. *Freshwater Biology*, **58**, 1614-1630.
- Bayly I, Ebsworth E, Wan H (1975) Studies on the lakes of Fraser Island, Queensland. *Australian Journal of Marine and Freshwater Research*, **26**, 1-13.
- Birks HH (2001) Plant macrofossils. In: *Tracking environmental change using lake sediments. Volume 3: terrestrial, algal and siliceous indicators.* (ed Smol JP, Birks, H.J.B., Last, W.M.) pp 49-74. Dordrecht, The Netherlands, Kluwer Academic Publishers.
- Bradley RS (1996) Plant-water relations of a subtropical riparian wetland. Unpublished B.Eng. Griffith University, Brisbane, 96 pp.
- Brett MJ, Baldini JUL, Gröcke DR (2014) Environmental controls on stable isotope ratios in New Zealand Podocarpaceae: Implications for palaeoclimate reconstruction. *Global and Planetary Change*, **120**, 38-45.
- Cernusak LA, Ubierna N, Winter K, Holtum JAM, Marshall JD, Farquhar GD (2013) Environmental and physiological determinants of carbon isotope discrimination in terrestrial plants. *New Phytologist*, **200**, 950-965.
- Cernusak LA, Winter K, Turner BL (2009) Physiological and isotopic ( $\delta^{13}\text{C}$  and  $\delta^{18}\text{O}$ ) responses of three tropical tree species to water and nutrient availability. *Plant, Cell and Environment*, **32**, 1441-1455.
- Denniston RF, Wyrwoll KH, Polyak VJ *et al.* (2013) A stalagmite record of Holocene Indonesian-Australian summer monsoon variability from the Australian tropics. *Quaternary Science Reviews*, **78**, 155-168.
- Diefendorf AF, Mueller KE, Wing SL, Koch PL, Freeman KH (2010) Global patterns in leaf  $^{13}\text{C}$  discrimination and implications for studies of past and future climate. *Proceedings of the National Academy of Sciences*, **107**, 5738-5743.
- Doran JC, Turnbull JW (eds) (1997) *Australian trees and shrubs: Species for land rehabilitation and farm planting in the tropics*, Canberra, Australian Centre for International Agricultural Research.
- Dray Jr FA, Bennett BC, Center TD (2006) Invasion history of *Melaleuca quinquenervia* (Cav.) S.T. Blake in Florida. *Castanea*, **71**, 210-225.
- Ehleringer JR (1993) Variation in leaf carbon isotope discrimination in *Encelia farinosa*: implications for growth, competition, and drought survival. *Oecologia*, **95**, 340-346.
- Emile-Geay J, Cobb KM, Mann ME, Wittenberg AT (2013a) Estimating central equatorial Pacific SST variability over the past millennium. Part I: methodology and validation. *Journal of Climate*, **26**, 2302-2328.



- Emile-Geay J, Cobb KM, Mann ME, Wittenberg AT (2013b) Estimating central equatorial Pacific SST variability over the past millennium. Part II: Reconstructions and Implications. *Journal of Climate*, **26**, 2329-2352.
- Farquhar G, O'Leary M, Berry J (1982) On the relationship between carbon isotope discrimination and the intercellular carbon dioxide concentration in leaves. *Functional Plant Biology*, **9**, 121-137.
- Farquhar GD, Ehleringer JR, Hubick KT (1989) Carbon isotope discrimination and photosynthesis. *Annual Review of Plant Physiology and Plant Molecular Biology*, **40**, 503-537.
- Fletcher MS, Thomas I (2010) A quantitative Late Quaternary temperature reconstruction from western Tasmania, Australia. *Quaternary Science Reviews*, **29**, 2351-2361.
- Freeman KH, Mueller KE, Diefendorf AF, Wing SL, Koch PL (2011) Clarifying the influence of water availability and plant types on carbon isotope discrimination by C3 plants. *Proceedings of the National Academy of Sciences of the United States of America*, **108**, E59-E60.
- Gallant AJE, Phipps SJ, Karoly DJ, Mullan AB, Lorrey AM (2013) Nonstationary Australasian teleconnections and implications for paleoclimate reconstructions. *Journal of Climate*, **26**, 8827-8849.
- Gouramanis C, De Deckker P, Switzer AD, Wilkins D (2013) Cross-continent comparison of high-resolution Holocene climate records from southern Australia — Deciphering the impacts of far-field teleconnections. *Earth-Science Reviews*, **121**, 55-72.
- Greenway M (1994) Litter accession and accumulation in a *Melaleuca quinquenervia* (Cav.) S.T.Blake wetland in south-eastern Queensland. *Australian Journal of Marine and Freshwater Research*, **45**, 1509-1519.
- Hamilton R, Penny D (2015) Ecological history of Lachlan Nature Reserve, Centennial Park, Sydney, Australia: a palaeoecological approach to conservation. *Environmental Conservation*, **42**, 84-94.
- Henley BJ, Thyer MA, Kuczera G, Franks SW (2011) Climate-informed stochastic hydrological modeling: Incorporating decadal-scale variability using paleo data. *Water Resources Research*, **47**.
- Hondzo M, Stefan HG (1996) Long-term lake water quality predictors. *Water Research*, **30**, 2835-2852.
- Ibanez T, Curt T, Hely C (2013) Low tolerance of New Caledonian secondary forest species to savanna fires. *Journal of Vegetation Science*, **24**, 177-188.
- Jeffrey SJ, Carter JO, Moodie KB, Beswick AR (2001) Using spatial interpolation to construct a comprehensive archive of Australian climate data. *Environmental Modelling and Software*, **16**, 309-330.
- Jowsey PC (1966) An improved peat sampler. *New Phytologist*, **65**, 245-248.
- Kaplan JO, Prentice IC, Buchmann N (2002) The stable carbon isotope composition of the terrestrial biosphere: Modeling at scales from the leaf to the globe. *Global Biogeochemical Cycles*, **16**, 8-1 - 8-11.
- Kattge J, Díaz S, Lavorel S *et al.* (2011) TRY - a global database of plant traits. *Global Change Biology*, **17**, 2905-2935.
- Kershaw AP, D'Costa D, Tibby J, Wagstaff B, O'Sullivan P, Heijnis H (2004a) The last million years around Lake Keilambete, Western Victoria. *Proceedings of the Royal Society of Victoria*, **116**, 95-106.

- Kershaw AP, Nix HA (1988) Quantitative palaeoclimatic estimates from pollen data using bioclimatic profiles of extant taxa. *Journal of Biogeography*, **15**, 589-602.
- Kershaw AP, Tibby J, Penny D, Yezdani H, Walkley R, Cook EJ, Johnston R (2004b) Latest Pleistocene and Holocene vegetation and environmental history of the Western Plains of Victoria. *Proceedings of the Royal Society of Victoria*, **116**, 143-163.
- Kohn MJ (2010) Carbon isotope compositions of terrestrial C3 plants as indicators of (paleo)ecology and (paleo)climate. *Proceedings of the National Academy of Sciences of the United States of America*, **107**, 19691-19695.
- Li AOY, Ng LCY, Dudgeon D (2009) Effects of leaf toughness and nitrogen content on litter breakdown and macroinvertebrates in a tropical stream. *Aquatic Sciences*, **71**, 80-93.
- Liu W, Liu Z, An Z, Wang X, Chang H (2011) Wet climate during the 'Little Ice Age' in the arid Tarim Basin, Northwestern China. *Holocene*, **21**, 409-416.
- Livingstone DA (1955) A lightweight piston sampler for lake deposits. *Ecology*, **36**, 137-139.
- Lloyd PJ, Kershaw AP (1997) Late quaternary vegetation and early Holocene quantitative climate estimates from Morwell Swamp, Latrobe Valley, south-eastern Australia. *Australian Journal of Botany*, **45**, 549-563.
- Lockheart MJ, Poole I, Van Bergen PF, Evershed RP (1998) Leaf carbon isotope compositions and stomatal characters: important considerations for palaeoclimate reconstructions. *Organic Geochemistry*, **29**, 1003-1008.
- Lough JM (2007) Tropical river flow and rainfall reconstructions from coral luminescence: Great Barrier Reef, Australia. *Paleoceanography*, **22**, PA2218.
- Marshall JC, McGregor GB (2011) The Influence of water depth on the distribution of the emergent sedge *Lepironia articulata* (Cyperaceae) in two dune lakes of southern Queensland Wallum wetlands. *Proceedings of the Royal Society of Queensland*, **117**, 193-200.
- Marshall JC, Negus P, Steward AL, McGregor GB (2011) Distributions of the freshwater fish and aquatic macroinvertebrates of North Stradbroke Island are differentially influenced by landscape history, marine connectivity and habitat preference. *Proceedings of the Royal Society of Queensland*, **117**, 239-260.
- Marshall JD, Brooks JR, Lajtha K (2007) Sources of variation in the stable isotope composition of plants. In: *Stable Isotopes in Ecology and Environmental Science*. (eds Michener R, Lajtha K) pp Page. Australia, Blackwell Publishing.
- Mettam P, Tibby J, Barr C, Marshall JC (2011) Development of Eighteen Mile Swamp, North Stradbroke: A palaeolimnological approach. *Proceedings of the Royal Society of Queensland*, **117**, 119-131.
- Miller JM, Williams RJ, Farquhar GD (2001) Carbon isotope discrimination by a sequence of *Eucalyptus* species along a subcontinental rainfall gradient in Australia. *Functional Ecology*, **15**, 222-232.
- Monnin E, Steig EJ, Siegenthaler U *et al.* (2004) Evidence for substantial accumulation rate variability in Antarctica during the Holocene, through synchronization of CO<sub>2</sub> in the Taylor Dome, Dome C and DML ice cores. *Earth and Planetary Science Letters*, **224**, 45-54.
- Mosisch TD, Arthington AH (2001) Polycyclic aromatic hydrocarbon residues in the sediments of a dune lake as a result of power boating. *Lakes and Reservoirs: Research and Management*, **6**, 21-32.

- Moss PT, Tibby J, Petherick L, McGowan H, Barr C (2013) Late Quaternary vegetation history of North Stradbroke Island, Queensland, eastern Australia. *Quaternary Science Reviews*, **74**, 257-272.
- Murphy BP, Bowman DMJS (2009) The carbon and nitrogen isotope composition of Australian grasses in relation to climate. *Functional Ecology*, **23**, 1040-1049.
- Neldner VJ, Wilson BA, Thompson EJ, Dillewaard HA (2012) *Methodology for survey and mapping of regional ecosystems and vegetation communities in Queensland. Version 3.2*, Brisbane (Queensland), Department of Science, Information Technology, Innovation and the Arts.
- Neukom R, Gergis J (2012) Southern Hemisphere high-resolution palaeoclimate records of the last 2000 years. *The Holocene*, **22**, 501-524.
- O'Donnell AJ, Cook ER, Palmer JG, Turney CSM, Page GFM, Grierson PF (2015) Tree rings show recent high summer-autumn precipitation in northwest Australia is unprecedented within the last two centuries. *Plos One*, **10**, DOI: 10.1371/journal.pone.0128533.
- Phipps SJ, McGregor HV, Gergis J *et al.* (2013) Paleoclimate data–model comparison and the role of climate forcings over the past 1500 years. *Journal of Climate*, **26**, 6915-6936.
- Prentice IC, Meng T, Wang H, Harrison SP, Ni J, Wang G (2011) Evidence of a universal scaling relationship for leaf CO<sub>2</sub> drawdown along an aridity gradient. *New Phytol*, **190**, 169-180.
- Prowse TAA, Johnson CN, Bradshaw CJA, Brook BW (2013) An ecological regime shift resulting from disrupted predator–prey interactions in Holocene Australia. *Ecology*, **95**, 693-702.
- Rohling EJ, Sluijs A, Dijkstra HA *et al.* (2012) Making sense of palaeoclimate sensitivity. *Nature*, **491**, 683-691.
- Schubert BA, Jahren AH (2012) The effect of atmospheric CO<sub>2</sub> concentration on carbon isotope fractionation in C<sub>3</sub> land plants. *Geochimica et Cosmochimica Acta*, **96**, 29-43.
- Schubert BA, Jahren AH (2015) Global increase in plant carbon isotope fractionation following the Last Glacial Maximum caused by increase in atmospheric pCO<sub>2</sub>. *Geology*, **43**, 435-438.
- Schulze ED, Williams RJ, Farquhar GD, Schulze W, Langridge J, Miller JM, Walker BH (1998) Carbon and nitrogen isotope discrimination and nitrogen nutrition of trees along a rainfall gradient in northern Australia. *Australian Journal of Plant Physiology*, **25**, 413-425.
- Seibt U, Rajabi A, Griffiths H, Berry JA (2008) Carbon isotopes and water use efficiency: sense and sensitivity. *Oecologia*, **155**, 441-454.
- Shakun JD, Carlson AE (2010) A global perspective on Last Glacial Maximum to Holocene climate change. *Quaternary Science Reviews*, **29**, 1801-1816.
- Shakun JD, Clark PU, He F *et al.* (2012) Global warming preceded by increasing carbon dioxide concentrations during the last deglaciation. *Nature*, **484**, 49-54.
- Spicer RA (1989) The formation and interpretation of plant fossil assemblages. In: *Advances in Botanical Research*. (ed Callow JA) pp Page., Academic Press.
- Stevenson J, Dodson JR, Prosser IP (2001) A late Quaternary record of environmental change and human impact from New Caledonia. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **168**, 97-123.
- Stewart GR, Turnbull MH, Schmidt S, Erskine PD (1995) <sup>13</sup>C natural abundance in plant communities along a rainfall gradient: a biological integrator of water availability. *Australian Journal of Plant Physiology*, **22**, 51-55.

- Telford RJ, Birks HJB (2009) Evaluation of transfer functions in spatially structured environments. *Quaternary Science Reviews*, **28**, 1309-1316.
- Turney CSM (2012) Surface  $\delta^{13}\text{C}$  in Australia: A quantified measure of annual precipitation? *Terra Australis*, **34**, 435-443.
- Turney CSM, Barringer J, Hunt JE, McGlone MS (1999) Estimating past leaf-to-air vapour pressure deficit from terrestrial plant  $\delta^{13}\text{C}$ . *Journal of Quaternary Science*, **14**, 437-442.
- Van de Water PK, Leavitt SW, Betancourt JL (2002) Leaf  $\delta^{13}\text{C}$  variability with elevation, slope aspect, and precipitation in the southwest United States. *Oecologia*, **132**, 332-343.
- Verschuren D, Tibby J, Sabbe K, Roberts N (2000) Effects of lake level, salinity and substrate on the benthic invertebrate community of a shallow tropical lake system. *Ecology*, **81**, 164-182.
- Weiguo L, Feng X, Ning Y, Zhang Q, Cao Y, An Z (2005)  $\delta^{13}\text{C}$  variation of C3 and C4 plants across an Asian monsoon rainfall gradient in arid northwestern China. *Global Change Biology*, **11**, 1094-1100.
- Woltering M, Atahan P, Grice K, Heijnis H, Taffs K, Dodson J (2014) Glacial and Holocene terrestrial temperature variability in subtropical east Australia as inferred from branched GDGT distributions in a sediment core from Lake McKenzie. *Quaternary Research*, **82**, 132-145.

**Table 1.** Coefficient of determination for significant ( $p < 0.01$ ) relationships between monthly  $\Delta_{\text{leaf}}$  and climate variables over different time intervals before leaf collection (blc). RHmaxT and RHminT=relative humidity at maximum and minimum temperature, respectively. n/s=not significant.

Climate variables	Coefficient of determination of relationship with monthly $\Delta_{\text{leaf}}$		
	1 - 2 years blc	1 - 3 years blc	1 - 4 years blc
Evaporation	0.13	0.10	n/s
Minimum temperature	0.30	0.28	0.15
P/E	0.19	0.28	0.18
Radiation	0.18	0.23	0.26
Rainfall	0.17	0.28	0.19
RHmaxT	0.15	0.16	0.16
RHminT	n/s	n/s	0.06
Vapour Pressure	0.19	0.18	0.14

**Table 2.** Relationship between  $\Delta_{\text{leaf}}$  and climate variables for 1–2, 1–3 and 1–4 hydrologic (Nov-Oct) years before leaf collection (blc). Only significant ( $p < 0.01$ )

variables are displayed. Rainfall data are expressed as yearly mean (mm) while radiation data are daily mean (MJ/m<sup>2</sup>).

Variable	Relationship	Linear regression
Annual average rainfall 1–4 years (365–1460 days) blc	$r^2=0.64, p=0.003$	$y = 0.004x + 17.936$
Radiation 1–4 years (365–1460 days) blc	$r^2=0.57, p=0.008$	$y = -2.1557x + 62.815$

**Table 3.** Wetlands with *M. quinquenervia* growing on the wetland surface or around the margin that we investigated for leaf preservation. For additional background information about North Stradbroke Island (NSI) wetlands see Marshall *et al.* (2011).

Wetland	Site type	Location (and island)	Reference (for lake characteristics)	Water depth at time of sampling	Wetland area	Leaves recovered?
18 Mile Swamp (pools)	Freshwater coastal swamp	27.526444 S 153.496778 E (NSI)	Mettam <i>et al.</i> (2011)	2.3 m	0.1 ha	No
Brown Lake	Perched lake	27.488291 S 153.434740 E (NSI)	Mosisch and Arthington (2001); Marshall <i>et al.</i> (2011)	6.1 m	46 ha	No
Blue Lake	Groundwater window lake	27.518056 S 153.477417 E (NSI)	Barr <i>et al.</i> (2013); Marshall and McGregor (2011); Marshall <i>et al.</i> (2011)	7.2 m	10.3 ha	No
Fern Gully Lagoon	Perched palustrine wetland	27.25053 S 153.27663 E (NSI)	Marshall <i>et al.</i> (2011)	1 m	7.3 ha	No
Swallow Lagoon	Perched lake	27°.41772 S 153°.46118 E (NSI)	Mosisch and Arthington (2001); Marshall and McGregor (2011); Marshall <i>et al.</i> (2011)	5.7 m	<1.5 ha	Yes. <i>Eucalyptus</i> leaves also preserved
Welsby Lagoon	Perched palustrine wetland	27.43717 S 153.45010 E (NSI)	Moss <i>et al.</i> (2013); Marshall <i>et al.</i> (2011)	1.2 m	19.25 ha	No
Barga Lake	Perched lake	25.5 S 153.05 E (Fraser Island)	Bayly <i>et al.</i> (1975)	0.8 m*	7.0 ha	No
Basin Lake	Perched lake	25.466667 S 153.05 E	Bayly <i>et al.</i> (1975)	7.5 m	3.8 ha	Yes

		(Fraser Island)				
Benaroon Lake	Perched lake	25.516667 S 153.05 E (Fraser Island)	Bayly <i>et al.</i> (1975)	5.7 m	83 ha	No
Jennings Lake	Perched lake	25.494 S 153.055 E (Fraser Island)	Bayly <i>et al.</i> (1975)	4.1 m	12.2 ha (open water area)	Yes. <i>Eucalyptus</i> leaves also preserved

\*deepest part of the lake not sampled.