



The University of
Nottingham

UNITED KINGDOM • CHINA • MALAYSIA

Lee, Darren S. and Durán-Peña, Maria Jesus and Burroughs, Laurence and Woodward, Simon (2016) Efficient preparation of TMSCl_2Br and its use in dichlorocyclopropanation of electron deficient alkenes. *Chemistry - a European Journal*, 2016 (22). ISSN 1521-3765

Access from the University of Nottingham repository:

<http://eprints.nottingham.ac.uk/32995/3/Lee-2016-Chem%20Eur%20J-Final-Archive%20Version-26.04.2016.pdf>

Copyright and reuse:

The Nottingham ePrints service makes this work by researchers of the University of Nottingham available open access under the following conditions.

This article is made available under the University of Nottingham End User licence and may be reused according to the conditions of the licence. For more details see:
http://eprints.nottingham.ac.uk/end_user_agreement.pdf

A note on versions:

The version presented here may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the repository url above for details on accessing the published version and note that access may require a subscription.

For more information, please contact eprints@nottingham.ac.uk

Efficient preparation of TMSCCl_2Br and its use in dichlorocyclopropanation of electron deficient alkenes

Darren S. Lee,* María Jesús Durán-Peña, Laurence Burroughs and Simon Woodward*


Dr. D. S. Lee, Dr. M. J. Durán-Peña, Dr. L. Burroughs, Prof. Dr. S. Woodward

School of Chemistry, University of Nottingham, University Park, Nottingham NG7 2RD (United Kingdom)
darren.lee@nottingham.ac.uk; simon.woodward@nottingham.ac.uk

Abstract: The reaction of excess TMSCl and LiCCl_2Br at low temperature is a technically simple high yield route to TMSCCl_2Br . The latter is a stable source of the dichlorobromomethide carbanion, which undergoes 1,4-addition with cyclic nitroalkenes and (*E*)-fumarates leading to dichlorocyclopropanes after bromide expulsion. For nitrostyrenes the reaction arrests at the 1,4-addition product. Low temperature NMR studies and DFT calculations suggest the formation of an 'ate' species [(nitronate) SiFMe_3] which, upon boil off of TMSF at 10 - 20 °C, yields the cyclopropane. DFT calculations also support the experimental differences between fluoride and acetate as promoters.

Introduction


Dihalocyclopropane motifs are attractive targets frequently associated with biological activity, especially in insects.^[1] Additionally, they are potent starting points in their own right for ring modification and expansion chemistry.^[2] While electron-rich alkenes react readily with a wide variety of electrophilic $:\text{CCl}_2$ sources [e.g. from diazocompounds^[3] or via CHX_3 ($\text{X} = \text{Cl}, \text{Br}$) deprotonations^[4]] providing the cyclopropanes directly. Electron deficient alkenes are insufficiently nucleophilic^[5] for such strategies. While some difluorocyclopropanes have been attained from electron-poor alkenes, high temperature reactions of TMSCF_3 or related analogues^[6] were necessary. For the dichloro- and dibromocyclopropanes typically the only generic viable methodology employs stoichiometric quantities of the highly toxic Seyferth reagents PhHgCX_3 ($\text{X} = \text{Cl}, \text{Br}$).^[7] Clearly, the latter are also undesirable on environmental grounds. In seeking for a user-friendly approach to dichlorocyclopropanation we had settled on use of TMSCCl_3 and recently could demonstrate both its high yield synthesis and its use in the conjugate addition of trichloromethyl units (**A** Scheme 1).^[8] However, under the published conditions^[8a,c] base-induced closure of **A** to the dichlorocyclopropanes **B** could lead to poor yields or the formation of alternative (elimination) by-products.



Scheme 1. Approaches to dichlorocyclopropanes based on TMSCCl_2X reagents ($\text{X} = \text{Cl}, \text{Br}$).

Results and Discussion

A potential route to overcoming the difficulties outlined above would involve the use of TMSCCl_2Br ^[9] (**1**). The greater leaving group ability of bromide (vs. chloride) is expected to facilitate direct closure to the cyclopropane. Existing literature routes to TMSCCl_2Br **1** involving bromination of TMSCCl_2Li with Br_2 ^[9a] are undesirable for preparation of large quantities of reagent. By switching from chloroform to HCCl_2Br we could adapt our previous^[8] TMSCCl_3 synthesis allowing TMSCCl_2Br **1** to be synthesised in excellent yield (87%) in a simple one-pot reaction from low cost commercially available materials. With TMSCCl_2Br **1** now accessible cleanly and in large quantities, its reactivity and potential as a cyclopropanation reagent was explored. Cyclic nitroolefin (**2a**) was picked as the model alkene for the reaction, as nitroolefins have been reported as potent Michael acceptors for TMSCCl_3 .^[8a,c] Table 1 outlines the effects of different promoters on its cyclopropanation reaction with TMSCCl_2Br **1**. Initial trials were carried out at room temperature and catalytic TBAF (5 mol-%), however, no conversion of **2a** was observed (Table 1, entry 1). When a stoichiometric quantity of TBAF was used poor conversion to mixtures of **3a-5a** also resulted (Table 1, entry 2). It is worth noting that the use of TBAF at room temperature was also accompanied by the formation of a significant quantity of decomposition product which arises from rapid fluoride-promoted activation of TMSCCl_2Br **1**. We believe this is due to the formation of $(\text{CCl}_2)_n$ oligomeric species, but the lack of a convenient spectroscopic or other handles has made their characterisation untenable thus far. This behaviour prevents practical use of the high temperature conditions developed by Hu^[6] with **1**. Dosing room temperature reactions with extra **1** slightly improves the conversion of **2a**. The milder fluoride source TBAT ($\text{NBu}_4[\text{Ph}_3\text{SiF}_2]$) provided good selectivity in favour of **3a** (Table 1, entry 3); however, removal of Ph_3SiF by-product from **3a** proved difficult so subsequently TBAT was avoided. Using less silylophilic promoters (Cl^- and Br^- , 1.1 equiv.) limited excessive decomposition of the silane reagent **1** (Table 1, entries 4-5); however, with bromide little or no conversion of **2a** was attained and chloride resulted in poor chemoselectivity favouring **5a**. The acetate NBu_4OAc (1.1 equiv.) was found to be an efficient and mild promoter, with no rapid decomposition of **1** observed. In this case starting material **2a** was consumed rapidly (<2 min) at room temperature; however, a mixture of products **3a-5a** was isolated again, with the 1,4-addition product **4a** as the major component (Table 1, entry 6). Using fluoride promoters at lower temperatures avoided the decomposition of **1** and favoured the formation of cyclopropane **3a** (Table 1, entries 7-9). Optimal conditions were found to be dropwise addition of TBAF (1.1 equiv.) at -78°C with complete consumption of **2a** in <15 min followed by conversion to **3a** exclusively after warming to room temperature (Table 1, entry 9). The analogous reaction with TMSCCl_3 led to no cyclopropane formation and only the 1,4-addition product was observed. Interestingly, repeating the optimum conditions with a slow addition of NBu_4OAc at -78°C favoured the elimination product **5a** and only 11% **3a** was observed (Table 1, entry 10). Cyclopropane formation is favoured in more polar solvents (such as DMF, MeCN, THF), less polar solvents (e.g. toluene, Et_2O) lead to more observed elimination (**5a**) and 1,4-addition (**4a**) products being observed.^[10] It is likely that in less polar solvents intermolecular aggregation is favoured leading to more observed **4a** and **5a** (see computational studies later). All cyclopropanes were assigned on the basis of their highly characteristic ^1H NMR spectra: compounds **3a**, **3d** – **3g** give singlets at 3.8 – 4.0 ppm, compounds **3b** – **3c** give double doublets at 2.7 – 3.0, and compounds **3h** – **3j**^[11] give singlets at 3.0 – 3.2. Further support for cyclopropane formation arises from the characteristic quaternary CCl_2 at 71 – 79 ppm exhibited by all dihalocyclopropanes in their ^{13}C NMR spectra. Additionally, quaternary C- NO_2 peaks at 62 – 65 ppm are seen for cyclopropanes from nitroalkenes. The presence of the NO_2 groups is also supported by characteristic IR stretches between 1550-1475 and 1360-1290 cm^{-1} . Parent ions were observed in the mass spectra of most of our cyclopropanes but fragile compound **3c** had to be characterised on the first daughter peaks observed. The possibility of alternative connectivity, e.g. 2,3-dichloroprop-1-enes, could be discounted on the basis of our previous work.^[8b]

Table 1. Effects of different promoters on the dichlorocyclopropanation reaction.^[a]

Run	NBu ₄ X (X =) ^[b]	Loading (equiv.)	Conc. (M)	Temp (°C)	Time (h)	2a:3a:4a:5a (%) ^[c]
1	F	0.05	0.2	RT	16	0:0:0:0
2	F	1.1	0.2	RT	16	92:5:0:3
3	Ph ₃ SiF ₂	1.1	0.2	RT	16	0:80:20:0
4	Br	1.1	0.2	RT	16	0:0:0:0
5	Cl	1.1	0.2	50	16	16:29:12:43
6	OAc	1.1	0.2	RT	<1	0:28:50:22
7	F	1.1	0.1	-40	<1	0:76:0:24
8	F	1.1	0.1	-50	<1	0:89:0:11
9 ^[d]	F	1.1	0.1	-78	<1	0:100:0:0
10	OAc	1.1	0.1	-78	<1	0:11:38:51

^[a] Runs 1-6 carried out on: **2a** (0.143 mmol), **1** (1.5 equiv., 0.215 mmol) in THF (0.72 mL, 0.2 M). Runs 7-10 carried out on: **2a** (0.143 mmol), **1** (2 equiv., 0.286 mmol) in THF (1.43 mL, 0.1 M). ^[b] The TBAF used was a 1 M solution in THF. ^[c] Yields were determined using ¹H NMR spectroscopy. ^[d] When repeated on 1 mmol scale, 2.2 equiv. TBAF and 2.5 equiv. of TMSCCl₂Br **1** were used to achieve complete conversion to **3a**.

The difference in reactivity between the promoters in Table 1 prompted a mechanistic investigation of the transformation. Multi-nuclear NMR studies of a mixture of TMSCCl₂Br **1** and **2a** (2.5:1) in d₈-toluene were carried out. Toluene was chosen as the NMR solvent as it allows observation of all 3 major reaction products. After **1** was treated with TBAF (2.2 equiv.) at -70 °C, ¹H and ¹³C NMR spectra were recorded as the mixture was raised in 10 °C steps to 20 °C. Initial 1,4-addition began at -70 °C leading to complete consumption of **2a** by -50 °C (30 min). The new species observed (Figure 1) was assigned to the free nitronate anion (**6**) on the basis of its similarity to the previously characterised CCl₃-analogue.^[8a] Nitronate **6** was stable to 10 °C (See Supporting Information). However, when the reaction was warmed from 10 to 20 °C, apparent free nitronate **6** was completely consumed and peaks due to the cyclopropane (**3a**), proton quench (**4a**) and elimination product (**5a**) were detected. In fluoride promoted reactions the chemoselectivity to the cyclopropane was always higher than reactions promoted by NBu₄OAc, where the conjugate addition (**4a**) and elimination (**5a**) products were formed in approximately equimolar amounts (Figure 1).




Figure 1. Representative variable temperature partial ¹H NMR spectra of the reaction mixture showing the free nitronate **6** *in situ* at 10 °C (top, O = **6**). The unlabelled peaks are due to toluene and THF. Decomposition products of **6** at 20 °C are also shown (bottom: cyclopropane Δ = **3a**, 1,4-addition □ = **4a**, elimination ☆ = **5a**), these are formed between 10-20 °C.

To support these results DFT computational studies were carried out on the cyclopropane formation (e.g. modelling Table 1, run 9) from TBAF (Chart 1). [All calculations were carried out at the B3LYP/6-31+G(d,p)^[12] level of theory using the polarizable continuum model with THF as the solvent, as higher levels of theory proved computationally very time demanding]. In these complex systems we simplified the amine salt to NMe₄F, again to improve computational throughput. The conjugate addition of **1** to **2a** was calculated to proceed over a barrier of 28.4 kcal mol⁻¹ via transition state I (Chart 1 and Figure 2a). This subsequently forms intermediate **6**, which is 23.5 kcal mol⁻¹ lower in energy than the starting structure. From **6**, ring-closure via transition state II (Figure 2b) to **3a** is hindered due to induced 3-ring strain and a barrier of 22.3 kcal mol⁻¹ was calculated, potentially allowing other reaction pathways to compete.




Chart 1. Reaction Pathway of Conjugate addition of TMSBr to **2a** and closure of intermediate **6** to **3a**.




Figure 2. (a) Transition state I with highlighted distances; (b) transition state II with highlighted distances.

We propose that alkene **5a** is formed by rearrangement of **6** to spectroscopically undetected **7** (Scheme 2). However, computationally at least [with B3LYP/6-31+G(d,p)], unimolecular formation of **7** from **6** was not viable. Using **6'** as a simple model for a second equivalent of nitronate **6** (Chart 2) we were able to obtain a sensible transition state (III, Figure 3) that supports viable intermolecular proton transfer between two molecules of nitronate **6**. The validity of this proposal is further supported by the observations from the low temperature NMR experiments described previously (Figure 1): the formation of **5a** is observed with concomitant formation of an equimolar quantity of **4a** without the introduction of an external proton source. The NMR experiment was carried out in toluene-*d*₈ so increased aggregation compared to THF is expected, favouring bimolecular elimination over cyclopropane formation. In THF we cannot completely

rule out deprotonation by bases other than **6** as in some cases **5a** is observed experimentally in the absence of **4a**.




Figure 3. Calculated transition state (**III**) in the formation of **5a**, with the distance between the forming and breaking C-H bond and the breaking C-Br bond highlighted.

If we directly compare the cyclopropane formation (Chart 1) and elimination (Chart 2) pathways, we can see that the $\Delta E_{\text{act}}(\text{calc})$ for formation of **3a** from **6** (via transition state **II**) is 16.7 kcal mol⁻¹ lower than the $\Delta E_{\text{act}}(\text{calc})$ for the formation of **5a** from **6** (via transition state **III**).






Chart 2. The reaction pathway from intermediate **6** to **5a** versus **6** to **3a**.

The improved chemoselectivity observed for use of TBAF over NBu_4OAc in THF deserves separate mechanistic comment. The by-products from the formation of free **6** in these two cases are TMSF and

TMSOAc respectively; the latter poorly intercepts **6** leading to significant formation of **4a/5a** via the routes already discussed (Scheme 2). The TMSF case is more interesting: the strength of the Si-F bond (135 kcal mol⁻¹) precludes formation of a neutral TMS-nitronate ether from **6** + TMSF. The formation of the hypervalent ‘ate’ species **8** from **6** and TMSF (Scheme 2) appears much more viable ($\Delta G_r^\circ(\text{calc}) = 21.2$ kcal mol⁻¹, see Supporting Information). It is likely that **8** acts as a stable reservoir of **6** that is released slowly at ~20 °C due to loss of TMSF (b.p. 16 °C) favouring annulation. Variable temperature ¹⁹F and ²⁹Si NMR spectra of the reaction mixture at low temperature (-30 °C to 20 °C) provide some evidence to support this idea. The signals for TMSF in the reaction mixture display a shift that could be interpreted as supporting rapid equilibration with unseen **8**. At 20 °C the TMSF signals become very attenuated supporting ‘boil off’ of TMSF. In the absence of TMSF, the reactive nitronate **6** is consumed non-selectively via **7**.



Scheme 2. Proposed behaviour for the kinetic nitronate (**6**), derived from the reaction of **2a** with **1** and NBu₄X (X = OAc, F) sources.

Given the proposals of Scheme 2 we sought a computational comparison of the energy barriers of the cyclopropanation that results from the free nitronate (**6b**) versus the TMS-nitronate ether (**6b-TMS**) proposed in the presence of NBu₄OAc (Chart 3). These simple models showed us $\Delta E_{\text{act}}(\text{calc}) = 25.5$ kcal mol⁻¹ for formation of the cyclopropane from the free nitronate and $\Delta G_r^\circ(\text{calc}) = -15.8$ kcal mol⁻¹ (Chart 3). The $\Delta E_{\text{act}}(\text{calc})$ for the TMS-nitronate closure was significantly higher at 40.1 kcal mol⁻¹ (Chart 3); however, the cyclopropane product from this system is 58.7 kcal mol⁻¹ lower in energy than transition state **V** resulting in a similar Gibbs Free Energy of reaction ($\Delta G_r^\circ(\text{calc}) = -18.6$ kcal mol⁻¹). The higher energy barrier for cyclopropane closure from the TMS-nitronate indicates that for this system, elimination is likely to be highly competitive with cyclopropane formation. This is backed up by our experimental observations, which show that the NBu₄OAc system [from which our proposed TMS-nitronate (**6-TMS**) may arise] suffers from poor chemoselectivity.




Chart 3. Comparison of cyclopropanation pathways from free nitronate model and TMS-nitronate ether.






Figure 4. (a) Transition State IV showing select distances; (b) transition state V showing selected distances.

By applications of the conditions of Table 1, or slight modifications thereof, a range of dichlorocyclopropanes could be isolated (Scheme 3). Fumarates also participated in the reaction yielding the desired cyclopropanes (in this case NBu_4Cl at 50 °C gave cyclopropane exclusively, whereas TBAF at -78 °C gave 1:1 cyclopropane:elimination). Other common Michael acceptors (e.g. enals, chalcones, alkylidene malonates, cyclic enones and acyclic enones) screened favoured 1,2-addition over the thermodynamic 1,4-addition and were not pursued further.^[8c] All of the cyclic nitroalkenes we tried formed the corresponding cyclopropanes, however, substrates containing halogens (**3e-3f**) were found to require the addition to 20 mol-% TEMPO to provide acceptable yields. This behaviour is attributed to aerobic triggered competing radical reactions, which we have noted before.^[8a, 13] Control reactions (absence of anionic promoters and use of other radical traps) indicated that the TEMPO does act as an inhibitor in this case.^[14]




Scheme 3. Dichlorocyclopropanes resulting from the 1,4-addition of TMSCCl_2Br . **3a-3h**: Optimal Conditions: Substrate in THF (0.1 M) with TMSCCl_2Br **1** (2.5 equiv.), then addition of TBAF (1 M in THF, 2.2 equiv.) at -78 °C. Monitored by tlc. **3i-3j**: Optimal Conditions: Substrate in THF (0.1 M) with TMSCCl_2Br **1** (1 equiv.) NBu_4Cl added and heated to 50 °C, after 1 h a second portion of **1** was added (1 equiv.) and monitored by tlc until complete. Footnote [a]: **3h** was isolated as an inseparable mixture of cyclopropane: elimination product (1:1).

Acyclic nitroalkenes were also subjected to the conditions of Scheme 3 but favoured the 1,4-addition product exclusively (**9a-h**, Scheme 4). We assume that the 1,4-addition produces a suitable nitronate intermediate, however, this fails to shut to desired cyclopropane due to lack of reactivity and/or induced ring strain. Treatment of the reaction mixtures with halophilic Ag^+ , Fe^{3+} or Al^{3+} salts did not aid in the ring closure. Heating the intermediate to 50 °C did lead to consumption of the intermediate nitronate but only undefined polymeric materials were obtained. Clearly cyclisation is not favoured. We note that in his seminal paper Cunico^[8d] attained related products and, in only one case (with a nitrostyrene predisposed to close due to the presence of a Thorpe-Ingold effect inducing CR_2 unit) were traces of a cyclopropane observed. We believe similar issues operate in our own system. The motifs **9a-h** are also accessible using our previously reported anionic promotor pool methodology where using 5 mol-% NBu_4OAc typically furnishes the 1,4-addition products in <30 min at room temperature.^[8a,c] When NBu_4OAc was used at 5 mol-% with cyclic nitroalkene **2a**, in an attempt to obtain the 1,4-addition product **4a** exclusively, only traces of cyclopropane product **3a** (<10%) were detected by ^1H NMR spectroscopy along with significant degradation.



Scheme 4. 1,4-addition of TMSOAc to acyclic nitroalkenes. Optimal Conditions: Substrate in THF (0.1 M) with TMSOAc **1** (1.1 equiv.), then NBu₄OAc (5 mol-%) added at room temp., monitored by tlc.

In the search for a selective catalytic process we found that metal-based salts (i.e. MF, MOAc where M = Na, K, Cs) had potential as stoichiometric promoters for the reactions of Schemes 1-4 as in the absence of suitable catalysts only minimal substrate conversion rates result. Extensive screening of a very wide range of diverse promoters (library of >200 entities) revealed that dibenzo-18-crown-6 (DB18-C-6) resulted in significant ligand accelerated catalysis^[15] when in combination with an excess of MF or MOAc. For example, while **2a** is converted to a mixture of **3-5a** by KOAc over 36 h this reaction is complete within 20 minutes in the presence of 5 mol-% DB18-C-6. In these catalytic systems the chemoselectivity to the cyclopropane **3a** was favoured by use of fluoride. However, this also promotes degradation of the TMSOAc **1** reagent. Potassium acetate was the most active stoichiometric promoter in the presence of catalytic DB18-C-6 (10 mol-%) but these reactions showed poorer chemoselectivity to the cyclopropane (**3a:4a:5a** ≈ 3:4:18) for the reasons indicated in Scheme 2. Significant attempts were made to extend the catalytic promotion seen for DB18-C-6 to chiral analogues but in all cases (crowns **10**^[16]-**11**^[17] are representative) negligible enantioselectivity and at best only modest catalysis were observed.



Due to the lack of stereoselectivity in these reactions we sought information on the putative transition states via B3LYP-DFT calculation. To the best of our knowledge no prior attempts have been made to understand the transition state associated with any 1,4-addition of any TMSOAc species to a Michael acceptor. This step is critical as it defines which stereo face of the acceptor is attacked by a potential asymmetric catalyst. As a simple model we studied the reaction of **2a**, TMSOAc and NMe₄F *in silico* at the [B3LYP/6-31+G(d,p)] level of theory (Supporting Information). Regardless of where we initialised the reaction coordinate search from, we found that the 1,4-addition step transition state has quite specific requirements with the nitroalkene **2a**, reagent, fluoride and quaternary amine all being co-linear (Figure 5). Based on this simple picture it is clear why attaining efficient chirality transfer is challenging – the triggering promoter (F in NBu₄F) is >9 Å away from the forming stereocenter and the supporting (potentially chiral)

catalytic cation even further away, and with high conformational freedom. While our model uses a simple quaternary amine $[\text{NMe}_4]^+$ and a simple promoter (F^-) it is likely that this transition state applies to all the runs with chiral quaternary amines and chiral crown ether complexes, providing some explanation as to why asymmetric induction is so challenging.




Figure 5. Transition state for the 1,4-addition of TMSBr **1** to **2a** in THF, showing the large distance between the substrate, quaternary amine and promoter [B3LYP/6-31+G(d,p)].

Conclusion

By a mixture of synthetic, NMR and basic computational experiments the underlying features of the catalytic cyclopropanation of nitroalkenes have been built up. The kinetic product of the reaction of such alkenes with TMSBr is the nitronate **6** which can either close to the cyclopropane, disproportionate to the 1,4-addition and elimination products, or suffer other related events. Conditions controlling the behaviour of **6** were identified by the mechanistic studies and an efficient synthesis of dichlorocyclopropanes from electron deficient alkenes demonstrated. Conditions for the use of largely inert promoters that are activated in the presence of crown ethers have been identified. The reasons for the extreme difficulty in attaining asymmetric dichlorocyclopropanation of electron deficient alkenes have also been identified. It is clear from this study that in order to obtain an efficient enantioselective process the reagent that delivers the haloform anion must be redesigned, allowing two point coordination of both the substrate and the cyclopropanation reagent. Given that and appropriate face-blocking strategies, asymmetric catalysis should be possible given that ligand accelerated catalysis has already been attained.

Experimental

A representative example is given, see supporting information for full details.

To a flame dried Schlenk tube under argon was added the cyclic nitroalkene (50 mg, 0.285 mmol), TMSBr (169 mg, 0.714 mmol) and THF (3 mL) [N.B. TEMPO 20 mol-% was added to reactions containing aromatic halogens]. The reaction vessel was cooled to -78°C in an acetone/dry ice bath and allowed to equilibrate for 15 min. TBAF (0.63 mL, 1 M in THF) was added dropwise to the reaction mixture. The mixture was stirred for 10 min and monitored by TLC. Upon consumption of the starting material the reaction vessel was removed from the cold bath and allowed to warm to room temperature (ca. 20 min). Upon warming, the reaction was examined by TLC and consumption of the intermediate nitronate was monitored. Once the reaction is complete it is quenched immediately with 1 M HCl (2 mL) and extracted with EtOAc (10 mL). After further washing with water (10 mL) the organic fraction was dried over MgSO_4 ,

filtered and the solvent evaporated. The cyclopropanes were isolated by flash chromatography through a small silica plug using 95:5 pentane:EtOAc. Prolonged exposure to silica gel proved detrimental to overall yield and sample quality.

Acknowledgements

We are grateful to the Engineering and Physical Sciences Research Council (EPSRC) for support of this work through grant EP/K000578/1. The authors would like to acknowledge the use of the EPSRC UK National Service for Computational Chemistry Software (NSCCS) at Imperial College London in carrying out this work. M. J. Durán-Peña acknowledges award of a Contrato Puente para Doctores supported by University of Cádiz.


References

- [1] Representative biological activity in dichlorocyclopropanes: a) T. Kudou, D. Tanimu, Y. Masuzawa, T. Yano, *PCT Int. Appl.* **2010**, WO2010026989, pp. 343 [*Chem. Abs.* 2010:306382]. b) D.G. Kuhn, J. A. Furch, D. A. Hunt, M. Asselin, S. P. Baffic, R. E. Diehl, T. P. Miller, Y. L. Palmer, M. F. Treacy, S. H. Trotto, in *Synthesis and Chemistry of Agrochemicals V*, **1998**, 686, 185-193. c) J. Mann, P. D. Kane, *Tetrahedron Lett.* **1985**, 26, 4677-4680. d) G. Holan, R. Walser, *Ger. Offen*, DE2653189 A1, **1977**, pp. 43 [*Chem. Abs.* 1977:534658]. e) G. Holan, *US Patent* US3962356 A, **1976**, pp. 5 [*Chem. Abs.* 1976:494027]. f) G. Holan, *Pat. Specif. (Aust.)* **1968**, AU283356, pp. 17 [*Chem. Abs.* 1969:412815]. For an overview of the insecticide area see: g) K. Naumann, *Synthetic Pyrethroid Insecticides: Chemistry and Patents*, Springer-Verlag Berlin, **1990**, pp. 390.
- [2] a) G. Keglevich, J. Dvorszki, V. Ujj, K. Ludányi, *Heteroat. Chem.* **2010**, 21, 271–277. b) M. G. Banwell, *Pure Appl. Chem.* **2008**, 80, 669–679. c) P. C. Stanislawski, A. C. Willis, M. G. Banwell, *Chem. Asian. J.* **2007**, 2, 1127–1136. d) A. C. Bissember, A. T. Phillis, M. G. Banwell, A. C. Willis, *Org. Lett.* **2007**, 9, 5421–5424. e) M. G. Banwell, M. E. Reum in *Advances in Strain in Organic Chemistry*, Vol 1 (Ed.: B. Halton) JAI Press Ltd, London, **1991**, pp 19–64. f) S. Kagabu, S. Naruse, Y. Tagami, Y. Watanabe, *J. Org. Chem.* **1989**, 54, 4275-4277. g) K. J. Shea, W. M. Fruscella, W. P. England, *Tetrahedron Lett.* **1987**, 28, 5623-5626. h) P. Mueller, J. Pfyffer, *Chimia* **1984**, 38, 79–80. i) D. C. Remy, P. S. Anderson, M. E. Christy, B. E. Evans, *J. Org. Chem.* **1978**, 43, 4311–4315. j) L. Skattebøl, *J. Org. Chem.* **1970**, 35, 3200–3201. k) S. R. Sandler, *J. Org. Chem.* **1967**, 32, 3876–3881. l) W. E. Parham, R. J. Sperley, *J. Org. Chem.* **1967**, 32, 926–931. m) R. B. Woodward, R. Hoffmann, *J. Am. Chem. Soc.* **1965**, 87, 395–397. n) C. H. DePuy, L. G. Schack, J. W. Hausser, W. Wiedemann, *J. Am. Chem. Soc.* **1965**, 87, 4006. o) S. J. Cristol, R. M. Sequeira, C. H. DePuy, *J. Am. Chem. Soc.* **1965**, 87, 4007–4008. p) W. R. Moore, W. R. Moser, J. E. LaPrade, *J. Org. Chem.* **1963**, 28, 2200–2205.

- [3] a) R. A. Moss, *J. Org. Chem.* **2010**, *75*, 5773–5783. b) R. A. Moss, *J. Phys. Org. Chem.* **2010**, *23*, 293–299. c) R. A. Moss, J. Tian, R. R. Sauers, D. H. Ess, K. N. Houk, K. Krogh-Jespersen, *J. Am. Chem. Soc.* **2007**, *129*, 5167–5174. d) R. A. Moss, J. Tian, G. Chu, R. R. Sauers, K. Krogh-Jespersen, *Pure Appl. Chem.* **2007**, *79*, 993–1001. e) R. A. Moss, J. Tian, R. R. Sauers, C. Skalit, K. Krogh-Jespersen, *Org. Lett.* **2007**, *9*, 4053–4056. f) A. P. Molchanov, V. S. Korotkov, R. R. Kostikov, *Russ. J. Org. Chem.* **2006**, *42*, 1146–1150. g) G. Chu, R. A. Moss, R. R. Sauers, *J. Am. Chem. Soc.* **2005**, *127*, 14206–14207.
- [4] For a comprehensive reviews on the synthesis of dichlorocarbenes and *gem*-dihalocyclopropanes: a) M. Fedoryński, *Chem. Rev.* **2003**, *103*, 1099–1032. b) M. Mąkosza, M. Fedoryński, In *Encyclopedia of Catalysis* Vol. V (Ed.: I. T. Horváth) Wiley, New York, **2003**, pp. 511–564. c) M. Mąkosza, M. Fedoryński, In *Interfacial Catalysis*; (Ed.: A. G. Volkov) Marcel Dekker, New York, **2003**, pp. 159. d) R. A. Jones, *Quaternary Ammonium Salts: Their Use in Phase-Transfer Catalysis*; Academic Press: London, **2001**. e) M. Mąkosza, M. Fedoryński, *Handbook of Phase Transfer Catalysis* (Eds.: Y. Sasson, R. Neumann) Blackie Academic & Professional: London, **1997**; pp. 135. f) M. Mąkosza, M. Fedoryński, *Pol. J. Chem.* **1996**, *70*, 1093. g) C. M. Starks, C. L. Liotta, M. Halpern, *Phase-Transfer Catalysis, Fundamentals, Applications and Industrial Perspectives*; Chapman & Hall: London, **1994**. h) E.V. Dehmlow, S. S. Dehmlow, *Phase Transfer Catalysis, 3rd ed.*, VCH: Weinheim, Germany, **1993**. i) M. Mąkosza, M. Fedoryński, *Adv. Catal.* **1987**, *35*, 375–422.
- [5] a) H. Cang, R. A. Moss, K. Krogh-Jespersen, *J. Am. Chem. Soc.* **2015** *137*, 2730–2737. b) R. A. Moss, L. Wang, C. M. Odorisio, M. Zhang, K. Krogh-Jespersen, *J. Phys. Chem. A* **2010**, *114*, 209–217. c) R. A. Moss, M. Zhang, K. Krogh-Jespersen, *Org. Lett.* **2009**, *11*, 5702–5704. d) R. A. Moss, M. Zhang, K. Krogh-Jespersen, *Org. Lett.* **2009**, *11*, 1947–1950. e) R. A. Moss, L. Wang, M. Zing, C. Skalit, K. Krogh-Jespersen, *J. Am. Chem. Soc.* **2008**, *130*, 5634–5635. f) R. A. Moss, J. Tian, *J. Am. Chem. Soc.* **2005**, *127*, 8960–8961.
- [6] a) L. Li, F. Wang, C. Ni, J. Hu, *Angew. Chem. Int. Ed.* **2013**, *52*, 12390–12394. b) F. Wang, T. Luo, J. Hu, Y. Wang, H. S. Krishnan, P. V. Jog, S. K. Ganesh, G. K. S. Prakash, G. A. Olah, *Angew. Chem. Int. Ed.* **2011**, *50*, 7153–7157. c) F. Wang, W. Zhang, J. Zhu, H. Li, K. Huang, J. Hu, *Chem. Comm.* **2011**, *47*, 2411–2413.
- [7] a) J. B. Lambert, B. T. Ziemnicka-Merchant, *J. Org. Chem.* **1990**, *55*, 3460–3464. b) L. W. Jenneskens, W. H. de Wolf, F. Bickelhaupt, *Angew. Chem. Int. Ed.* **1985**, *24*, 585–586. c) H. Mayr, U. W. Heigl, *Angew. Chem.* **1985**, *97*, 567–568. d) J. B. Lambert, E. G. Larson, R. J. Bosch, M. L. E. TeVrucht, *J. Am. Chem. Soc.* **1985**, *107*, 5443–5447. e) D. Seyferth, H. Shih, *J. Am. Chem. Soc.* **1973**, *95*, 8464–8465. f) D. Seyferth, J. M. Burlitch, R. J. Minas, J. Yick-Pui Mui, H. D. Simmons, A. J. H. Treiber, S. R. Dowd, *J. Am. Chem. Soc.* **1965**, *87*, 4259–4270.
- [8] a) N. Wu, B. Wahl, S. Woodward, W. Lewis, *Chem. Eur. J.* **2014**, *20*, 7718–7724. b) B. Wahl, A. Cabré, S. Woodward, W. Lewis, *Tetrahedron Lett.* **2014**, *55*, 5829–5831. c) B. Wahl, D. S. Lee, S. Woodward, *Eur. J. Org. Chem.* **2015**, 6033–6039. d) R. F. Cunico, C. P. Zhang, *Synth. Commun.* **1991**, *21*, 2189–2195.

- [9] a) D. Seyferth, F. M. Jr. Armbrecht, E. M. Hanson, *J. Organomet. Chem.* **1967**, *10*, P25–P27. b) D. Seyferth, E. M. Hanson, F. M. Jr. Armbrecht, *J. Organomet. Chem.* **1970**, *23*, 361–371. c) D. Seyferth, R. L. Jr. Lambert, E. M. Hanson, *J. Organomet. Chem.* **1970**, *24*, 647–661. d) G. Fritz, K. H. Schmid, *Z. Anorg. Allg. Chem.* **1978**, *441*, 125–164.
- [10] The conditions of table 1, entry 9 were repeated in different solvents; DMF and MeCN gave the same results as THF, i.e. exclusively **3a**. MeOH, EtOH, acetone, and DMSO resulted in significant degradation of **1**, no conversion of **2a** was observed. For toluene and Et₂O complete conversion of **2a** was observed and yielded a mixture of **3a**:**4a**:**5a** in a 4:5:4 ratio.
- [11] D. Seyferth, H. Shih, *J. Org. Chem.* **1974**, *39*, 2336–2341.
- [12] a) C. Lee, W. Yang, R. G. Parr, *Phys. Rev. B* **1988**, *37*, 785–789. b) A. D. Becke, *Phys. Rev. A* **1988**, *38*, 3098–3100. c) A. D. Becke, *J. Chem. Phys.* **1993**, *98*, 1372. d) A. D. Becke, *J. Chem. Phys.* **1993**, *98*, 5648–5652. e) P. J. Stephens, F. J. Devlin, C. F. Chabalowski, M. J. Frisch, *J. Phys. Chem.* **1994**, *98*, 11623–11627.
- [13] M. S. Kharasch, B. M. Kuderna, W. Urry, *J. Org. Chem.* **1948**, *13*, 895–902; for an alternative procedure of high synthetic utility and substrate scope discussions, see: b) D. Vofsi, M. Asscher, *Org. Synth.* **1965**, *45*, 104.
- [14] Examples of TEMPO acting as a radical inhibitor: a) S. Zhou, X. Pan, Z. Zhou, A. Shoberu, P. Zhang, J. Zou, *J. Org. Chem.* **2015**, *80*, 5348–5354. b) K. Sun, X. Wang, G. Li, Z. Zhu, Y. Jiang, B. Xiao, *Chem. Commun.* **2014**, *50*, 12880–12883. c) S. Chatani, T. Gong, B. A. Earle, M. Podgorski, C. N. Bowman, *ACS Macro Letters* **2014**, *3*, 315–318. d) S. Mani, P. Cassagnau, M. Bousmina, P. Chaumont, *Polymer* **2010**, *51*, 3918–3925. e) E. Font-Sanchis, C. Aliaga, E. V. Bejan, R. Cornejo, J. C. Scaiano, *J. Org. Chem.* **2003**, *68*, 3199–3204. f) J. B. Miller, J. R. Salvador, *J. Org. Chem.* **2002**, *67*, 435–442. g) M. R. Thompson, C. Tzoganakis, G. L. Rempel, *Polymer*, **1997**, *39*, 327–334.
- [15] D. J. Berrisford, C. Bolm, K. B. Sharpless, *Angew. Chem. Int. Ed.* **1995**, *34*, 1059–1070.
- [16] E. B. Kyba, K. Koga, L. R. Sousa, M. G. Siegel, D. J. Cram, *J. Am. Chem. Soc.* **1973**, *95*, 2692–2693.
- [17] P. Bako, E. Czinege, T. Bako, M. Czugler, L. Toke, *Tetrahedron: Asymmetry* **1999**, *10*, 4539–4551.

Table of Contents Entry



The efficient synthesis of TMSBr provides easy access to an alternative dichlorocyclopropanation reagent. Cyclopropanes from electron deficient alkenes have been synthesised and the mechanism has been probed with NMR and DFT calculations.

Keywords: Dichlorocyclopropane • Michael Addition • Cyclopropanation • Conjugate Addition • Cyclopropane