



Research in Progress: Medical Research Council United Kingdom Refractory Asthma Stratification Programme (RASP-UK)

Heaney, L. G., Djukanovi, R., Woodcock, A., Walker, S., Matthews, J. G., Pavord, I. D., ... RASP-UK Consortium (2015). Research in Progress: Medical Research Council United Kingdom Refractory Asthma Stratification Programme (RASP-UK). *Thorax*. DOI: 10.1136/thoraxjnl-2015-207326

Published in:
Thorax

Document Version:
Publisher's PDF, also known as Version of record

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

Copyright the authors 2015. This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See: <http://creativecommons.org/licenses/by/4.0/>

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.



OPEN ACCESS

AUDIT, RESEARCH AND GUIDELINE UPDATE

Research in progress: Medical Research Council United Kingdom Refractory Asthma Stratification Programme (RASP-UK)

Liam G Heaney,¹ Ratko Djukanovic,² Ashley Woodcock,³ Samantha Walker,^{4,5} John G Matthews,⁵ Ian D Pavord,⁶ Peter Bradding,⁷ Robert Niven,³ Chris E Brightling,⁷ Rekha Chaudhuri,⁸ Joseph R Arron,⁵ David F Choy,⁵ Douglas Cowan,⁸ Adel Mansur,⁹ Andrew Menzies-Gow,¹⁰ Ian Adcock,¹¹ Kian F Chung,¹¹ Chris Corrigan,¹² Peter Coyle,¹ Timothy Harrison,¹³ Sebastian Johnston,¹¹ Peter Howarth,² James Lordan,¹⁴ Ian Sabroe,¹⁵ Jeannette Bigler,¹⁶ Dirk Smith,¹⁶ Matthew Cattley,¹⁷ Richard May,¹⁷ Lisa Pierre,¹⁸ Chris Stevenson,¹⁸ Glenn Crater,¹⁹ Frank Keane,²⁰ Richard W Costello,²¹ Val Hudson,²² David Supple,²² Tim Hardman²³

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/thoraxjnl-2015-207326>).

For numbered affiliations see end of article.

Correspondence to

Professor Liam G Heaney, Centre for Infection and Immunity, Queen's University of Belfast, Level 8, Belfast City Hospital, Lisburn Road, Belfast, BT9 7AB, UK; l.heaney@qub.ac.uk

Received 28 May 2015

Revised 17 June 2015

Accepted 22 June 2015

ABSTRACT

The UK Refractory Asthma Stratification Programme (RASP-UK) will explore novel biomarker stratification strategies in severe asthma to improve clinical management and accelerate development of new therapies. Prior asthma mechanistic studies have not stratified on inflammatory phenotype and the understanding of pathophysiological mechanisms in asthma without Type 2 cytokine inflammation is limited. RASP-UK will objectively assess adherence to corticosteroids (CS) and examine a novel composite biomarker strategy to optimise CS dose; this will also address what proportion of patients with severe asthma have persistent symptoms without eosinophilic airways inflammation after progressive CS withdrawal. There will be interactive partnership with the pharmaceutical industry to facilitate access to stratified populations for novel therapeutic studies.

INTRODUCTION

The Medical Research Council has recently funded an ambitious stratified medicines programme in severe asthma (the UK Refractory Asthma Stratification Programme (RASP-UK <http://www.rasp.org.uk/>). In asthma, disease stratification has traditionally followed a 'one-size fits all' approach, based on level of asthma control and response to 'stepwise' increases in corticosteroid (CS) treatment. More recently, stratification has been based on inflammatory biomarkers, with sputum analysis and transcriptomic profiling demonstrating that ~50% of patients with asthma have typical eosinophilic airway inflammation driven by 'Type 2' cytokines (T2 cytokines: interleukin (IL)-4, IL-5, IL-13); this T2-high asthma has been shown to respond well to CS therapy. In the remainder of patients, there is limited evidence for a major role for T2 cytokines and this has been termed 'T2-low asthma', where patients show little or no response to CS therapy.¹ There is also evidence of significant

disease heterogeneity in severe asthma, with only 25%–50% of patients having the prototypic T2 gene signature consistent with T2-high disease. In many cases of severe asthma, it therefore seems likely that the CS dose has been escalated inappropriately to try and manage persistent symptoms that are not CS responsive. Given the clear evidence that CS responsiveness is confined to T2-high disease, the cornerstone of the RASP-UK stratification programme is around failure of response to CS therapy in severe asthma.

We propose that in patients with severe asthma, failure to respond to high-dose CS treatment occurs for three principal reasons:

1. Non-adherence to CS treatment.
2. Impaired CS responsiveness—in some patients, despite adherence with high-dose inhaled CS, there is persistent T2-high inflammation/eosinophilia and these patients frequently require systemic CS to achieve an acceptable level of disease control.
3. Non-responsiveness to CS in patients with T2-low asthma—we believe that CS will have been escalated inappropriately in this group in clinical practice.

NON-ADHERENCE TO CS TREATMENT

Non-adherence to inhaled CS and oral CS therapy is a major clinical problem, present in 30%–50% of patients with difficult-to-control asthma. Non-adherence can be intentional or non-intentional, is poorly detected by physicians and asthma nurses and is associated with poor asthma outcomes and increased healthcare costs.² Moreover, enrolment of non-adherent patients into clinical trials is believed to contribute to the frequently observed improvement in placebo arms of asthma clinical trials, complicating the interpretation of trial results and reducing statistical power. Importantly, expensive new biological therapies targeting T2-high patients can potentially be used inappropriately in

To cite: Heaney LG, Djukanovic R, Woodcock A, et al. *Thorax* Published Online First: [please include Day Month Year] doi:10.1136/thoraxjnl-2015-207326

patients who would often be controlled with improved adherence to inhaled CS. Patients with difficult-to-control asthma and fractional exhaled nitric oxide (FeNO) >45 ppb have a high risk of exacerbation³ and the key clinical question is whether failure to control their disease is because of non-adherence to inhaled CS treatment or whether they have relative CS-resistant T2-high asthma and, thus, require additional treatment with drugs that target T2 mechanisms (eg, anti-IL-5 and anti-IL-13 treatments). RASP-UK will therefore use biomarker-based assessments of inhaled CS exposure and response in biomarker-high patients ('FeNO suppression test'—reference 4 online supplementary data) using remote monitoring technologies to define CS responsiveness and thus optimise inhaled CS where appropriate.

IMPAIRED CS RESPONSIVENESS

In some patients with severe asthma, despite adherence with high-dose inhaled CS, there is persistent T2-high inflammation and airway eosinophilia and these patients frequently require systemic CS to achieve disease control. The availability of many new biological therapies targeting T2-high disease in the next 5 years will generate many interesting questions, including differential efficacy between monoclonal antibodies targeting IL-5/IL-5R (mepolizumab, bendralizumab, relizumab), IL-13 (lebrikizumab, tralokinumab) and IL-4R α (dupilumab). Other strategies targeting the T2-axis, including anti-CrTh2 and novel anti-IgE therapies (quilizumab, Medi-4212, QGE031), will also be targeting overlapping patient groups; identifying which patients respond better to different classes of drugs may require 'head-to-head' studies. Many of these new therapies are expected to come to market with a companion diagnostic or predictive biomarker(s) of clinical response.

NON-RESPONSIVENESS TO CS

Inhaled CS remain the mainstay of asthma treatment, controlling disease through suppression of T2/eosinophilic inflammation in the airway. However, not all symptoms in severe asthma respond to CS, and CS responsiveness is minimal in the absence of T2 inflammation.¹ Therefore, new algorithms for guiding CS treatment in patients with severe asthma are needed, which are not solely based on clinical symptoms and lung function and which can be delivered in the clinic. Titrating CS against sputum eosinophilia improves asthma control by reducing asthma exacerbations⁴ but this strategy is difficult to deliver in real-world clinical settings. FeNO is straightforward to measure, but when used as a single biomarker, it has been disappointing in titrating CS in patients with moderate asthma.⁵ Periostin is a secreted matricellular protein from a number of tissues including airway epithelial cells and is upregulated by T2 cytokines (IL-4 and IL-13) and is associated with airway eosinophilia in patients with severe asthma.¹ An assay for serum periostin is being developed to help identify a periostin-high responder population for the anti-IL-13 monoclonal antibody lebrikizumab.

We have previously examined the predictive value of FeNO, blood eosinophils and serum periostin as a composite biomarker to predict exacerbation risk in the placebo arms of clinical trials with lebrikizumab and omalizumab in patients taking at least 500 μ g fluticasone propionate and a second controller (see online supplementary data). This analysis has demonstrated that, individually, these biomarkers are all correlated with exacerbation risk, but using the three biomarkers in a 'composite' scoring system further differentiated subjects on the basis of exacerbation rate (see online supplementary figure E1). This predictive composite score is independent of asthma symptoms (measured by asthma control questionnaire) and lung function

(see online supplementary figures E2 and E3) which current asthma guidelines advocate for treatment adjustment.

RASP-UK will therefore use this scoring system to adjust CS treatment in a randomised, pragmatic, multicentre, parallel group trial in patients with severe asthma and baseline FeNO <45 ppb comparing standard care with treatment guided by composite biomarkers (as discussed above, patients with FeNO \geq 45 ppb have a high risk of exacerbation and are not candidates for CS reduction). Exploratory outcomes will examine the prognostic value of selected, novel blood and urine biomarkers from the Unbiased BIOmarkers in PREDiction of respiratory disease outcomes (U-BIOPRED) programme for exacerbation risk (<http://www.europeanlung.org/en/projects-and-research/projects/u-biopred/home>). The programme will also characterise the inflammatory and microbiomic profiles (viral, bacterial, fungal) in exacerbations in both study arms.

One of the anticipated outcomes of this biomarker-based stratification will be identification of a patient group with T2-low severe asthma who have persistent symptoms and exacerbations which are not CS responsive and in whom T2 inflammation biomarkers (eosinophils, FeNO, periostin) are adequately controlled with moderate doses of inhaled CS. We know that this patient group exists based on data from a comparable sputum stratification study where one-third of patients had a sputum eosinophil count of <1.9% when repeatedly measured over 12 months despite steroid withdrawal.⁴ There are currently few treatment options for these patients representing a significant unmet medical need where new targets need to be identified so drugs can be developed. One of the core aims of RASP-UK is to characterise this group in detail to define important structure/function/symptom relationships. This will include transcriptomic and microbiomic profiling compared with age and gender matched patients with refractory eosinophilic asthma and stable asthma controlled with low-dose inhaled CS. These groups will be followed up for 12 months to identify their exacerbation pattern and stability of the phenotype. Previous research has focused on eosinophilia as a prognostic for exacerbation, but 'non-eosinophilic cohorts' also experience exacerbations and potent anti-eosinophilic therapies have not eradicated severe exacerbations in clinical trials. A better understanding of the mechanism and phenotype of exacerbations in this T2-low cohort is fundamental to targeting exacerbations in this group.

CONCLUSION

RASP-UK is a logical, clinically focused, stratification programme which will develop a clinical model to assess and phenotype patients with severe asthma (figure 1). It will identify appropriate patients for novel biological treatments that target the T2-cytokine axis (T2 biomarker high when adherent to high dose inhaled CS), will optimise CS treatment so that excessive CS exposure does not cause unnecessary morbidity and will facilitate a research focus and better understanding in a major area of 'unmet need', specifically non-T2/non-CS responsive severe asthma. Harmonised protocols for patient phenotyping will also allow stratified patient groups to enter proof of concept clinical trials which target the biology of their disease rather than using the usual broad inclusion criteria, where many of the patients are unlikely to respond; this will streamline and reduce size for early drug development in severe asthma. The RASP-UK Consortium vision is to effect a paradigm shift in asthma care through a logical stratified approach to severe asthma and to establish a long-term collaborative infrastructure for translational research excellence in the UK.




Figure 1 Flow diagram for the RASP-UK Clinical Stratification Programme. ATS, American Thoracic Society; ERS, European Respiratory Society; FeNO, fractional exhaled nitric oxide; FP, Fluticasone Propionate; RASP-UK, UK Refractory Asthma Stratification Programme.

Author affiliations

- ¹Centre of Infection and Immunity, The Queen's University of Belfast, Belfast, UK
- ²Academic Unit of Clinical and Experimental Sciences, University of Southampton, Southampton, UK
- ³Institute of Inflammation and Repair, The University of Manchester, Manchester Academic Health Science Centre, Manchester, UK
- ⁴Asthma UK, London, UK
- ⁵Genentech Inc., South San Francisco, California, USA
- ⁶Nuffield Department of Medicine, The University of Oxford, Oxford, UK
- ⁷Department of Infection, Immunity and Inflammation, Institute for Lung Health, University of Leicester, Leicester, UK
- ⁸Greater Glasgow Health Board, Glasgow, UK
- ⁹Heart of England NHS Foundation Trust, Birmingham, UK
- ¹⁰Royal Brompton & Harefield NHS Foundation Trust, London, UK
- ¹¹Imperial College of Science, Technology and Medicine, London, UK
- ¹²Kings College London, London, UK
- ¹³University of Nottingham, Nottingham, UK
- ¹⁴The Newcastle upon Tyne NHS Foundation Trust, Newcastle upon Tyne, UK
- ¹⁵School of Medicine and Biomedical Sciences, University of Sheffield, Sheffield, UK
- ¹⁶Amgen Inc., Thousand Oaks, California, USA
- ¹⁷Astra Zeneca, London, UK
- ¹⁸Janssen Research & Development LLC, London, UK
- ¹⁹Aerocrine AB, Solna, Sweden
- ²⁰Vitalograph Inc, Ennis, Ireland
- ²¹Department of Medicine, Royal College of Surgeons Ireland, Dublin, Ireland
- ²²Public and Patient Representative, London, UK
- ²³Niche Science & Technology, Richmond, UK

Acknowledgements DFC and JRA for analysis of the clinical trial biomarker data and generation of the biomarker algorithm. Sofia Mosesova (Genentech Inc.) for statistical input to the programme.

Contributors All of the listed authors were involved in the Consortium meetings to develop, write and review the RASP-UK programme which was submitted and funded by the Medical Research Council in January 2015.

Funding Medical Research Council (MR/M016579/1) and Industrial Partner project contributions.

Competing interests None declared.

Provenance and peer review Not commissioned; internally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See: <http://creativecommons.org/licenses/by/4.0/>

REFERENCES

- 1 Arron JR, Choy DF, Scheerens H, *et al.* Non-invasive biomarkers that predict treatment benefit from biologic therapies in asthma. *Ann Am Thorac Soc* 2013;10 (Suppl):S206–13.
- 2 Heaney LG, Horne R. Non-adherence in difficult asthma; time to take it seriously. *Thorax* 2012;67:268–70.
- 3 Kupczyk M, ten Brinke A, Sterk PJ, *et al.* Frequent exacerbators—a distinct phenotype of severe asthma. *Clin Exp Allergy* 2014;44:212–21.
- 4 Green RH, Brightling CE, McKenna S, *et al.* Asthma exacerbations and sputum eosinophil counts: a randomised controlled trial. *Lancet* 2002;360:1715–21.
- 5 Petsky HL, Cates CJ, Lasserson TJ, *et al.* A systematic review and meta-analysis: tailoring asthma treatment on eosinophilic markers (exhaled nitric oxide or sputum eosinophils). *Thorax* 2012;67:199–208.

THORAX

Research in progress: Medical Research Council United Kingdom Refractory Asthma Stratification Programme (RASP-UK)

Liam G Heaney, Ratko Djukanovic, Ashley Woodcock, Samantha Walker, John G Matthews, Ian D Pavord, Peter Bradding, Robert Niven, Chris E Brightling, Rekha Chaudhuri, Joseph R Arron, David F Choy, Douglas Cowan, Adel Mansur, Andrew Menzies-Gow, Ian Adcock, Kian F Chung, Chris Corrigan, Peter Coyle, Timothy Harrison, Sebastian Johnston, Peter Howarth, James Lordan, Ian Sabroe, Jeannette Bigler, Dirk Smith, Matthew Cattley, Richard May, Lisa Pierre, Chris Stevenson, Glenn Crater, Frank Keane, Richard W Costello, Val Hudson, David Supple and Tim Hardman

Thorax published online July 23, 2015

Updated information and services can be found at:
<http://thorax.bmj.com/content/early/2015/07/23/thoraxjnl-2015-207326>

These include:

Supplementary Material

Supplementary material can be found at:
<http://thorax.bmj.com/content/suppl/2015/07/29/thoraxjnl-2015-207326.DC1.html>

References

This article cites 5 articles, 2 of which you can access for free at:
<http://thorax.bmj.com/content/early/2015/07/23/thoraxjnl-2015-207326#BIBL>

Open Access

This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See:
<http://creativecommons.org/licenses/by/4.0/>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Open access](#) (149)
[Asthma](#) (1650)
[Inflammation](#) (947)

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>