



**QUEEN'S
UNIVERSITY
BELFAST**

Balancing the (carbon) budget: Using linear inverse models to estimate carbon flows and mass-balance ^{13}C : ^{15}N labelling experiments in low oxygen sediments.

Hunter, W., van Oevelen, D., & Witte, U. (2013). Balancing the (carbon) budget: Using linear inverse models to estimate carbon flows and mass-balance ^{13}C : ^{15}N labelling experiments in low oxygen sediments.. Poster session presented at European Geoscience Union 2013, Vienna, Austria.

Queen's University Belfast - Research Portal:

[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

Copyright the authors 2013.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

William R Hunter^{1,3}, Dick Van Oevelen², Ursula, Witte¹
¹Oceanlab, University of Aberdeen AB41 6AA, UK.
²Royal Netherlands Institute for Sea Research, Yerseke, Netherlands.
³Present Address: Department for Limnology, Universität Wien, Austria. Email: william.hunter@univie.ac.at

Introduction

Approximately 6 % of the continental margin sea floor experiences persistent dysoxia within oxygen minimum zones (OMZs) (Helly & Levin, 2004). OMZs are predicted to grow as a consequence of climate change, with implications for marine biogeochemical cycles (Stramma et al., 2008). The Arabian Sea OMZ impinges upon the Indian continental margin at bathyal depths (150 – 1500 m) generating a depth-dependent oxygen gradient.

In 2008 a multi-national expedition led by Prof Hiroshi Kitazato (JAMSTEC, Japan) investigated the effects of oxygen availability, sediment geochemistry and community structure upon carbon & nitrogen cycling pathways at the Indian margin OMZ. *In situ* ¹³C-tracer experiments were conducted to quantify OM processing by sediment bacteria and fauna (Witte et al., 2011; Hunter et al., 2012 a; b). However, no empirical data on ¹³C fluxes from the sediment were available. We closed the carbon budget for each ¹³C-tracer experiment using linear inverse modelling (LIM) to reconstruct sediment carbon fluxes.

Experimental Design & Model Formulation




Fig 1. Bathymetric Map of the Indian Continental margin with experimental stations marked (left) & corresponding O₂ profiles (right).

Fig 2. Deployment of semi-enclosed mesocosms at the Indian margin seafloor.

Experimental Design & Data Acquisition

Semi-enclosed mesocosms deployed fixed doses of ¹³C-labelled diatoms (*Thalassiosira weissflogii*) at the seafloor across the Indian margin OMZ (540 – 1100 m, Figs 1, 2). The ¹³C label was traced into sediment OM, bacteria, foraminifera and metazoan fauna (A. Enge & P. Heinz, *unpub. data*; Hunter et al., 2012a; b; Witte et al., 2011).

Linear Inverse Models

Food web models were constructed following Van Oevelen et al. (2010). Food web components and flow linkages were fixed *a priori*. Flow magnitudes were constrained using data from the mesocosm experiments. Final model solutions were obtained by Bayesian sampling for best fit from 25,000 iterations of each model.

Linear Inverse Modelling Solutions




Fig 3. LIM solutions for carbon flows across the Indian margin OMZ. Barcharts reveal contributions of detrital, bacterial and faunal OM sources to organismal carbon budgets.

Abbreviations: lDet, labile detritus; rDet, refractory detritus; Bac, Bacteria; Nem, Nematodes, SDFPol / pSDF Surface Deposit Feeding polychaetes, SSDFPol / pSSDF, subsurface deposit feeding polychaetes; Othpol / pOth, other polychaetes Crust, macrofaunal crustacea; Moll, molluscs; Sip, sipunculans; Asc, ascidians; Oph, megafaunal ophiuroids; Dec, megafaunal decapods

Summary

1. Food-web model complexity increases concomitantly with oxygen availability.
2. Across the Indian margin OMZ labile phytodetritus was primarily processed by foraminifera and metazoan macrofauna.
3. LIM estimates community respiration to be greatest at the 800 m stations, driven by higher metazoan faunal contributions.
4. Bacterial contributions to sediment respiration increased concomitantly with the depth-dependent oxygen gradient.
5. At present, foraminiferal data are only available for station T1 540 m. Foraminifera must be integrated into all models to accurately reconstruct sediment carbon fluxes.




Fig 4. Sediment community respiration and the relative contributions of bacteria and major faunal groups estimated from LIM solutions.

Acknowledgements

Figure 2 images collected by the submersible *Shinkai 6500*, courtesy of JAMSTEC. This study was funded by The Carnegie Trust, Scotland. WRH was funded under a NERC doctoral training grant. The authors wish to thank Prof. H. Kitazato, the YK08-11 science team, and the captain and crew of the RV *Yokosuka*. Without their unwavering dedication and enthusiasm this work would not have been possible. Mass Spectroscopy was carried out at the Royal Netherlands Institute for Sea Research, UC Davis Stable Isotope Facility and Macaulay Land Use Research Institute.

References

Helly and Levin (2004). *Deep-Sea Res Part I* 51 (9): 1159-1168.
 Hunter et al. (2012a). *Biogeosciences* 9: 993-1006.
 Hunter et al. (2012b). *ISME Journal* 6 (11): 2140-2151.
 Stramma et al. (2008). *Science* 320 (5876): 655-658.
 Van Oevelen et al. (2010). *Ecosystems* 13: 32-45.
 Witte et al., (2011). *Geophys Res Abs.* Vol. 13, EGU2011-3089.