



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



ELSEVIER

Contents lists available at ScienceDirect

International Journal of Infectious Diseases

journal homepage: www.elsevier.com/locate/ijidINTERNATIONAL
SOCIETY
FOR INFECTIOUS
DISEASES

Editorial

COVID-19 and the scaled-down 2020 Hajj Pilgrimage—Decisive, logical and prudent decision making by Saudi authorities overcomes pre-Hajj public health concerns



Introduction

The abrupt appearance of SARS-CoV-2 as a novel lethal zoonotic pathogen causing COVID-19 disease in humans in late December 2019 (WHO, 2020), and its explosive global spread (Hui et al., 2020) caught health authorities worldwide by surprise and exposed the ill-preparedness of global public health systems to deal with the appearance of a new pathogen. Apart from generic prevention and control issues of public health and lockdown measures to limit epidemic spread, specific issues of Mass Gathering sporting and religious events came under specific spotlight (Alzahrani et al., 2020; Memish et al., 2020; Baloch et al., 2020; McCloskey et al., 2020; Petersen et al., 2020b).

Mass gathering events present important health challenges related to the public health services and health of the host country population, the attendees and their home countries (Memish et al., 2014; Memish et al., 2019). The 2009 Hajj was held during the 2009 H1N1 influenza pandemic and focused attention on developing Mass Gatherings Medicine as a formal discipline, resulting in the formation of a coalition of global academic and public health faculty and virtual WHO mass gathering collaborating centres to guide development of, and update, optimal public health and medical prevention and treatment guidelines at mass gathering events (Memish et al., 2014).

Mass gathering events and COVID-19

WHO with global mass gathering partners, developed comprehensive key recommendations for COVID-19, and since end of February 2020, there was a stepwise increase in cancellation, temporary suspension or postponement of international and national religious, sporting, musical, and other mass gatherings (WHO, 2020b; McCloskey et al., 2020; Petersen, McCloskey et al., 2020a). Apart from focusing on major sporting events, global attention has been on Saudi Arabia and the Umrah and Hajj pilgrimages. Approximately 10 million people from 182 countries travel to Saudi Arabia annually for the Hajj and Umrah pilgrimages (Memish et al., 2014, 2019). The Umrah pilgrimage can be performed anytime during the year, and thousands of pilgrims from all continents arrive in Saudi Arabia every month. Saudi Arabia with its extensive experience and commitment to pilgrim safety and wellbeing, was quick off the mark and on February 27th,

2020, restricted inbound flights and local and international pilgrims were prevented from travelling to Makkah and Madinah for the Umrah pilgrimage.

Social and economic impacts of the Hajj and COVID-19

For the July 2020 Hajj, approximately 2.4 million pilgrims were expected in Saudi Arabia (Saudi Ministry of Hajj and Umrah, 2020), but they were made aware that the growing COVID-19 pandemic might put their plans into disarray. The Saudi Ministry of Health made regular announcements that the format of the Hajj 2020 was being debated and evaluated based on COVID-19 situation globally and domestically. Several countries from where large numbers of Muslim pilgrims originate (Malaysia, Indonesia, India and Singapore) subsequently announced that they were barring their pilgrims from attending the 2020 Hajj. These cancellations were anticipated to have major social and economic impacts on national economies, individual livelihoods and public morale. The Umrah and Hajj pilgrimages together generate over \$12 billion annually for businesses and the economy, thus limiting or cancelling the Hajj would come at a huge cost.

Modelling of COVID-19 and public health impact at Hajj 2020

Alzahrani et al. (2020), in their Autoregressive Integrated Moving Average (ARIMA) model, had forecast the expected daily number of COVID-19 cases in Saudi Arabia over ensuing weeks before the 2020 Hajj. They forecasted that the number of COVID-19 cases in Saudi Arabia would continue growing and could reach up to 7,668 new cases per day, with over 127,129 cumulative daily cases in a matter of weeks if stringent precautionary and control measures were not implemented to limit the spread of COVID-19. They suggested that the Saudi authorities faced very difficult decisions regarding the 2020 Hajj Pilgrimage which was scheduled for the last week of July 2020 and they had two options: cancellation of the Hajj or implementation of extreme preventive and control measures in the cities of Mecca and Madinah required to avoid a COVID-19 catastrophe of widespread national and international spread (Alzahrani et al., 2020).

With the global spread of COVID-19 showing no signs of abating, the national and global public health impact, risk to livelihoods and economic costs, religious rulings and sensitivities,

pros and cons of holding the Hajj were discussed and debated leading to the final prudent decision by the Saudi government to hold a scaled-down 2020 Hajj (Atique and Jumallah, 2020; Al Saidu et al., 2020; Baloch et al., 2020; Ebrahim & Memish, 2020; McCloskey et al., 2020; Memish et al., 2020a). Of public health importance were the close proximity in which pilgrims interacted and lived during the Hajj; respiratory tract infections are known to transmit easily during the Hajj; an estimated 40% of pilgrims are elderly and many have comorbid conditions increasing the risk of COVID-19 associate mortality; two-thirds of pilgrims originate from countries with weak public health capabilities and COVID-19 spread in either direction was to be avoided at all costs. To introduce and maintain social distancing measures and a safe public health environment for the Hajj rites to be completed by all pilgrims had become a major public health challenge and seemed near impossible. Muslim religious leaders and ethicists were unified in concluding that the preservation of life during pandemics takes priority over practicing religious rituals, and that a small number of people nominally perform the Hajj under strict supervision and public health guidance. Islamic tradition relates to a prophetic saying ‘if you hear of an outbreak of a plague in a land, do not enter it; but if that epidemic breaks out in a place where people are residing, do not leave that place.’ The correlate being, preserving the lives of pilgrims and people of the host country and countries from which pilgrims originate, is more important than the pilgrimage itself.

Hajj 2020-decisive political and public health action

In early June 2020, after careful consideration, the Saudi government took decisive, logical, logistical and prudent decisions (Saudi Ministry of Hajj and Umrah, 2020) to overcome these pre-Hajj nightmares of public health, political, economic and religious concerns. For the first time since the Kingdom of Saudi Arabia was formed in 1932, the decision to bar pilgrims arriving in Saudi Arabia from foreign countries was made. The 2020 Hajj was scaled down considerably, and participation for Hajj rituals was restricted to only 1000 people with a negative COVID-19 test, residing within the Kingdom of which ‘foreign’ residents would comprise two-thirds of all selected pilgrims from a pool of local workers, health care workers and security personnel, – especially those who had recovered from COVID-19. Those aged 65 years and over and those with comorbid conditions would be barred. While all holy sites would remain open, adequate physical distancing and disinfection measures were put in place with oversight and assistance at regular intervals during the pilgrim’s journey. Wearing masks was mandatory, and pilgrims would be subject to temperature checks and placed in quarantine if required. All pilgrims were given well thought out kits that include disinfectants, masks, a prayer rug, the ihram (a seamless white garment required to be worn by pilgrims), sterilised pebbles for the stoning ritual at Jamaraat. Throughout the duration of the Hajj, the pilgrims would have to keep a social distance of one and a half meters and were guided by well laid out markers and Hajj coordinators. No pilgrims would be allowed to touch the Kaaba or kiss the black stone at its corner-both of which are regular customs during the Hajj. Pilgrims would also have to be quarantined for 14 days after the pilgrimage.

The Hajj 2020 was a public health success and ended on Monday 3rd August 2020. The decisive, logical and prudent decision making by Saudi authorities which enabled the pre-Hajj nightmare of public health, political, economic and religious concerns to be overcome. The successful completion of the 2020 Hajj is a major tribute to the leadership and commitment of the Saudi authorities, and it reflects their extensive experience of organising the annual Hajj pilgrimage, and continued commitment to improving public health issues related to mass gatherings events. The decisive actions, public health

preparedness and strict implementation of public health prevention and intervention measures, pre-Hajj, during Hajj and post-Hajj, serve as exemplars for other mass gathering religious and sporting events.

Previous scaled-down Hajj pilgrimages

The 2020 Hajj was not the first time the Hajj has been scaled down. Historically, the Hajj has been scaled down several times before due to infectious disease outbreaks. Between 1830 and 1930, there were at least 27 cholera outbreaks among pilgrims in Mecca (Peters, 1994). Massive cholera outbreaks throughout the 19th century resulted in the suspension of Hajj in 1837 and 1846. The cholera outbreak in 1865 in Saudi Arabia led to the establishment of quarantine ports to limit the spread of the disease during Hajj. Since Saudi Arabia’s foundation in 1932 the Hajj has never been cancelled and has not missed any year.

Conclusions

The ongoing SARS-CoV-2 pandemic, yet once again, highlights the continuing threat of new emerging infectious diseases with epidemic potential, including the persistent threat of the Middle East Respiratory Syndrome (MERS) (Periman et al., 2020; Memish et al., 2020b), to global health security. As of 2nd August 2020, there have been 17,660,523 confirmed cases of COVID-19 worldwide, including 275,905 cases from Saudi Arabia. reported to WHO (2020a). The WHO has emphasized the importance and need for collaborative and committed public health efforts over the longer term to achieve COVID-19 control globally. Regularly recurring mass gathering events should be used as important global sentinel surveillance platforms and opportunities for the conduct of research and evaluation of a range of global public health issues (Shafi et al., 2016; Memish et al. 2019). Increased investments in the fields of mass gathering medicine and ‘One-Human–Environmental-Animal-Health’ programs (Zumla et al., 2016a, b) are required for pro-active surveillance, and development of accurate prevention, management, and control guidelines to protect the health of attendees of mass gatherings and of host country populations.

Author declarations/Conflicts of interest

All authors are members and founders of the Global Centre for Mass Gatherings Medicine. Dr Shuja Shafi is director of Mass Gatherings and Global Health Network, London, UK. All authors declare no conflicts of interest.

Acknowledgements

Sir Zumla is co-Principal Investigator of the Pan African-European Network on Emerging and Re-Emerging Infections (PANDORA-ID-NET–<https://www.pandora-id.net/>) funded by the European and Developing Countries Clinical Trials Partnership the EU Horizon 2020 Framework Programme. Sir Zumla is in receipt of a National Institutes of Health Research senior investigator award.

References

- Al Saidu AMO, Nur FA, Al-Mandhari AS, El Rabbat M, Hafeez A, Abubakar A. Decisive leadership is a necessity in the COVID-19 response. *Lancet* 2020; doi:[http://dx.doi.org/10.1016/S0140-6736\(20\)31493-8](http://dx.doi.org/10.1016/S0140-6736(20)31493-8) [published online ahead of print, 2020 July 3] S0140-6736(20)31493-31498.
- Alzahrani SI, Aljamaan IA, Al-Fakih EA. Forecasting the spread of the COVID-19 pandemic in Saudi Arabia using ARIMA prediction model under current public health interventions. *J Infect Public Health* 2020;13(7):914–9. doi:<http://dx.doi.org/10.1016/j.jiph.2020.06.001>.

- Atique S, Itumalla R. Hajj in the Time of COVID-19. *Infect Dis Health* 2020;25(3):219–21, doi:http://dx.doi.org/10.1016/j.idh.2020.04.001 [published online ahead of print, 2020 April 15].
- Baloch Z, Ma Z, Ji Y, Ghanbari M, Pan Q, Aljabr W. Unique challenges to control the spread of COVID-19 in the Middle East. *J Infect Public Health* 2020; , doi:http://dx.doi.org/10.1016/j.jiph.2020.06.034 [published online ahead of print, 2020 July 13] S1876-0341(20)30564-30565.
- Ebrahim SH, Memish ZA. Saudi Arabia's drastic measures to curb the COVID-19 outbreak: temporary suspension of the Umrah pilgrimage. *J Travel Med* 2020;27(3), doi:http://dx.doi.org/10.1093/jtm/taaa029 taaa029.
- Hui DS, I Azhar E, Madani TA, et al. The continuing 2019-nCoV epidemic threat of novel coronaviruses to global health—the latest 2019 novel coronavirus outbreak in Wuhan, China. *Int J Infect Dis* 2020;91:264–6, doi:http://dx.doi.org/10.1016/j.ijid.2020.01.009.
- McCloskey B, Zumla A, Lim PL, et al. A risk-based approach is best for decision making on holding mass gathering events. *Lancet* 2020;395(10232):1256–7, doi:http://dx.doi.org/10.1016/S0140-6736(20)30794-7.
- Memish ZA, Zumla A, Alhakeem RF, et al. Hajj: infectious disease surveillance and control. *Lancet* 2014;383(9934):2073–82, doi:http://dx.doi.org/10.1016/S0140-6736(14)60381-0.
- Memish ZA, Steffen R, White P, et al. Mass gatherings medicine: public health issues arising from mass gathering religious and sporting events. *Lancet* 2019;393(10185):2073–84, doi:http://dx.doi.org/10.1016/S0140-6736(19)30501-X.
- Memish ZA, Ahmed Y, Alqahtani SA, Ebrahim SH. Pausing super spreader events for COVID-19 mitigation: International Hajj pilgrimage cancellation. *Travel Med Infect Dis* 2020a;36:101817, doi:http://dx.doi.org/10.1016/j.tmaid.2020.101817 [published online ahead of print, 2020 July 13].
- Memish ZA, Perlman S, Van Kerkhove MD, Zumla A. Middle East respiratory syndrome. *Lancet* 2020b;395(10229):1063–77, doi:http://dx.doi.org/10.1016/S0140-6736(19)33221-0.
- Perlman S, Azhar EI, Memish ZA, Hui DS, Zumla A. Confronting the persisting threat of the Middle East respiratory syndrome to global health security. *Lancet Infect Dis* 2020;20(2):158–60, doi:http://dx.doi.org/10.1016/S1473-3099(19)30347-0.
- Peters FE. Steamships and cholera: the Hajj in modern times. In: Peters FE, editor. *The Hajj: the Muslim Pilgrimage to Mecca and the Holy Places*. New Jersey, Princeton: Princeton University Press and Princeton Paperbacks; 1994. p. 266–315.
- Petersen E, McCloskey B, Hui DS, et al. COVID-19 travel restrictions and the International Health Regulations—Call for an open debate on easing of travel restrictions. *Int J Infect Dis* 2020a;94:88–90, doi:http://dx.doi.org/10.1016/j.ijid.2020.04.029.
- Petersen E, Memish ZA, Zumla A, Maani AA. Transmission of respiratory tract infections at mass gathering events. *Curr Opin Pulm Med* 2020b;26(3):197–202, doi:http://dx.doi.org/10.1097/MCP.0000000000000675.
- Saudi Ministry of Hajj and Umrah. Saudi Ministry of Hajj and Umrah. 2020. <https://www.haj.gov.sa/en#>.
- Shafi S, Dar O, Khan M, et al. The annual Hajj pilgrimage—minimizing the risk of ill health in pilgrims from Europe and opportunity for driving the best prevention and health promotion guidelines. *Int J Infect Dis* 2016;47:79–82, doi:http://dx.doi.org/10.1016/j.ijid.2016.06.01.
- WHO, 2020 <https://www.who.int/emergencies/diseases/novel-coronavirus-2019> [Accessed 2 August 2020].
- WHO. Key Planning Recommendations for Mass Gatherings in the Context of the Current COVID-19 Outbreak <https://www.who.int/publications/i/item/10665-332235> - accessed 1st August, 2020. 2020.
- Zumla A, Alagaili AN, Cotten M, Azhar EI. Infectious diseases epidemic threats and mass gatherings: refocusing global attention on the continuing spread of the Middle East Respiratory syndrome coronavirus (MERS-CoV). *BMC Med* 2016a;14(1)132, doi:http://dx.doi.org/10.1186/s12916-016-0686-3 Published 2016 Sep 7.
- Zumla A, Dar O, Kock R, et al. Taking forward a 'One Health' approach for turning the tide against the Middle East respiratory syndrome coronavirus and other zoonotic pathogens with epidemic potential. *Int J Infect Dis* 2016b;47:5–9 doi:10.

Alimuddin Zumla

Division of Infection and Immunity, University College London, and
NIHR Biomedical Research Centre, UCL Hospitals NHS Foundation
Trust, London, United Kingdom

Esam I. Azhar^{a,b}

^aSpecial Infectious Agents Unit, King Fahd Medical Research Center,
Jeddah, Saudi Arabia

^bMedical Laboratory Sciences Department, Faculty of Applied Medical
Sciences, King Abdulaziz University, Jeddah, Saudi Arabia

Saleh Alqahtani^{a,b}

^aDepartment of Medicine, King Faisal Specialist Hospital & Research
Center, Riyadh, Saudi Arabia

^bDivision of Gastroenterology and Hepatology, Johns Hopkins
University, Baltimore, MD, USA

Shuja Shafi

Mass Gatherings and Global Health Network, London, United
Kingdom

Ziad A. Memish^{a,b,*1}

^aResearch & Innovation Centre, King Saud Medical City, Ministry of
Health, and College of Medicine, Alfaisal University, Riyadh, Saudi
Arabia

^bHubert Department of Global Health, Emory University, Atlanta, USA

¹All authors contributed equally.

* Corresponding author at: Research & Innovation Centre, King
Saud Medical City, Ministry of Health, and College of Medicine,
Alfaisal University, Riyadh, Saudi Arabia.

E-mail addresses: a.i.zumla@gmail.com (A. Zumla),
eazhar@kau.edu.sa (E. Azhar),
Salqaht1@jhmi.edu (S. Alqahtani),
msshafi12@gmail.com (S. Shafi),
zmemish@yahoo.com (Z. Memish).