



Conditions of *Mytilus edulis* extracellular body fluids and shell composition in a pH-treatment experiment: Acid-base status, trace elements and $\delta^{11}\text{B}$

Agnes Heinemann, Jan Fietzke, Frank Melzner, Florian Böhm, and Jörn Thomsen

Leibniz Institute of Marine Sciences (IFM-GEOMAR), D-24148 Kiel, Germany (jfietzke@ifm-geomar.de)

Dieter Garbe-Schönberg

Institute of Geosciences, University of Kiel, D-24118 Kiel, Germany

Anton Eisenhauer

Leibniz Institute of Marine Sciences (IFM-GEOMAR), D-24148 Kiel, Germany

[1] *Mytilus edulis* were cultured for 3 months under six different seawater $p\text{CO}_2$ levels ranging from 380 to 4000 μatm . Specimen were taken from Kiel Fjord (Western Baltic Sea, Germany) which is a habitat with high and variable seawater $p\text{CO}_2$ and related shifts in carbonate system speciation (e.g., low pH and low CaCO_3 saturation state). Hemolymph (HL) and extrapallial fluid (EPF) samples were analyzed for pH and total dissolved inorganic carbon (C_T) to calculate $p\text{CO}_2$ and $[\text{HCO}_3^-]$. A second experiment was conducted for 2 months with three different $p\text{CO}_2$ levels (380, 1400 and 4000 μatm). Boron isotopes ($\delta^{11}\text{B}$) were investigated by LA-MC-ICP-MS (Laser Ablation–Multicollector–Inductively Coupled Plasma–Mass Spectrometry) in shell portions precipitated during experimental treatment time. Additionally, elemental ratios (B/Ca, Mg/Ca and Sr/Ca) in the EPF of specimen from the second experiment were measured via ICP-OES (Inductively Coupled Plasma–Optical Emission Spectrometry). Extracellular pH was not significantly different in HL and EPF but systematically lower than ambient water pH. This is due to high extracellular $p\text{CO}_2$ values, a prerequisite for metabolic CO_2 excretion. No accumulation of extracellular $[\text{HCO}_3^-]$ was measured. Elemental ratios (B/Ca, Mg/Ca and Sr/Ca) in the EPF increased slightly with pH which is in accordance with increasing growth and calcification rates at higher seawater pH values. Boron isotope ratios were highly variable between different individuals but also within single shells. This corresponds to a high individual variability in fluid B/Ca ratios and may be due to high boron concentrations in the organic parts of the shell. The mean $\delta^{11}\text{B}$ value shows no trend with pH but appears to represent internal pH (EPF) rather than ambient water pH.

Components: 10,800 words, 5 figures, 6 tables.

Keywords: *Mytilus edulis*; acid base; boron isotopes; extrapallial fluid; ocean acidification; trace metals.

Index Terms: 0419 Biogeosciences: Biomineralization; 0424 Biogeosciences: Biosignatures and proxies; 0454 Biogeosciences: Isotopic composition and chemistry (1041, 4870).

Received 8 July 2011; **Revised** 14 November 2011; **Accepted** 17 November 2011; **Published** 12 January 2012.

Heinemann, A., J. Fietzke, F. Melzner, F. Böhm, J. Thomsen, D. Garbe-Schönberg, and A. Eisenhauer (2012), Conditions of *Mytilus edulis* extracellular body fluids and shell composition in a pH-treatment experiment: Acid-base status, trace elements and $\delta^{11}\text{B}$, *Geochem. Geophys. Geosyst.*, 13, Q01005, doi:10.1029/2011GC003790.


1. Introduction

[2] About 50% of anthropogenic carbon dioxide (CO₂) released to the atmosphere is absorbed by the global oceans. It is predicted that the oceans approach a pH of ~7.3 within the next 300 years [e.g., *Caldeira and Wickett, 2003; Sabine et al., 2004*]. This leads to a shift in the inorganic carbon equilibrium towards higher CO₂ and lower CO₃²⁻ concentrations. Therefore, the calcium carbonate (CaCO₃) saturation state (Ω) will decrease by about 50% (for $p\text{CO}_2$ of 840 μatm) [e.g., *Feely et al., 2004; Fabry et al., 2008*] and higher-latitude oceans are predicted to become undersaturated with respect to aragonite by the year 2050 [*Orr et al., 2009; Cao and Caldeira, 2008*] which may have considerable consequences for marine calcifying organisms [*Orr et al., 2005*]. Many (mostly short-term studies, days to week duration) ocean acidification experiments have been conducted and different responses to high $p\text{CO}_2$ were found. The majority of the investigated species showed declining rates of calcification and reproduction [see *Doney et al., 2009; Ries et al., 2009*]. Most ocean acidification perturbation experiments are not able to properly account for the genetic adaptation potential of marine species, as time limitations usually prevent multi generation experiments. Thus, calcifying marine organisms from habitats with naturally high CO₂ concentrations can serve as analogues for future ocean conditions [e.g., *Hall-Spencer et al., 2008*].

[3] Kiel Fjord (Western Baltic Sea, Germany) provides ideal conditions for ocean acidification studies due to summer hypoxia in bottom waters and upwelling of CO₂ enriched waters [*Hansen et al., 1999; Lehmann et al., 2002*]. Kiel Fjord is already today frequently exposed to seawater $p\text{CO}_2$ values that are predicted for the future global ocean of the next 100–300 years [*Thomsen et al., 2010; Caldeira and Wickett, 2003*]. The habitat is characterized by low salinity (10–20), low alkalinity (1900–2150 $\mu\text{mol kg}^{-1}$), low pH (minimum value <7.5) during summer and autumn, and high $p\text{CO}_2$ (maximum value of 2340 μatm ; 1 $\mu\text{atm} = 0.101$ Pa) events. The CaCO₃ saturation state in the Kiel Fjord therefore can reach minimum values of $\Omega_{\text{arag}} = 0.34$ and $\Omega_{\text{calc}} = 0.58$ [*Thomsen et al., 2010*]. It can be assumed that calcifying communities in this habitat have already adapted to a fluctuating carbonate system speciation with frequent high $p\text{CO}_2$ events for multiple generations.

[4] Predictions of future ocean pH scenarios can be improved by studies of climate shifts in the past. In

this regard the elemental and isotopic composition of biogenic carbonates (e.g., bivalve shells, coral skeletons, foraminifera) serve as proxies for the reconstruction of past ocean chemistry. In particular, boron to calcium ratios (B/Ca) have been shown to decrease with declining pH in some biological and inorganic carbonates [*Hemming and Hanson, 1992; Sanyal et al., 2000*]. This results from the fact that in aqueous solutions, boric acid dissociates to B(OH)₄⁻ and H⁺ (equation (1)). Consequently, the proportion of boric acid to borate is pH dependent.



The borate ion is suggested to be preferentially incorporated into carbonates precipitated from seawater [*Vengosh et al., 1991; Hemming and Hanson, 1992; Hemming et al., 1995; Pagani et al., 2005*]. For this reason, the isotopic signature of B(OH)₄⁻ will be recorded in the carbonate. *Klochko et al.* [2009] and *Rollion-Bard et al.* [2011] proposed that both species can be incorporated. In contrast, *Foster* [2008] investigated B/Ca ratio in foraminifera and reported that the partition coefficient of borate between water and shell carbonate is primarily influenced by seawater [CO₃²⁻] and not directly by seawater pH. The results of *Yu et al.* [2007] showed the opposite trend of those shown by *Foster* [2008]. *Foster* [2008] suggested this indicates a species specific control on boron incorporation.

[5] Boron isotopes ($\delta^{11}\text{B}$) have been measured in skeletons of different organisms like corals [*Reynaud et al., 2004; Hönisch et al., 2004*] and foraminifera [*Hönisch et al., 2003; Foster, 2008; Rollion-Bard and Erez, 2010*] grown in laboratory experiments as a pH proxy. Therefore, the combination of these two proxies (B/Ca for [CO₃²⁻] and $\delta^{11}\text{B}$ for pH reconstruction) might offer the opportunity to reconstruct the marine paleo-carbonate system as it is necessary to know two of the six parameters (pH, [CO₂], [HCO₃⁻], [CO₃²⁻], total alkalinity (A_T) and dissolved inorganic carbon (C_T)) to constrain the system and to establish how CaCO₃ saturation state of the oceans changed in the past [*Foster, 2008*].

[6] Several studies showed marine calcifying organisms being sensitive to elevated $p\text{CO}_2$ and lower pH [e.g., see *Doney et al., 2009*]. However, blue mussels from Kiel Fjord showed the ability to settle, survive and even calcify under similar low pH conditions, when food concentrations are high [*Thomsen et al., 2010*]. Therefore mussels may be good candidate species to provide high resolution records of environmental conditions (like temperature, salinity, pH) and thus contribute to the

reconstruction of past climate. Additionally, the wide distribution and adaptation to a broad range of environments [Gosling, 1992] suggests *M. edulis* as a model organism for such studies.

[7] Since different studies using bivalve shells as proxy archives yielded contradictory results [Klein *et al.*, 1996a, 1996b; Vander Putten *et al.*, 2000; Lazareth *et al.*, 2003; Immenhauser *et al.*, 2005; Freitas *et al.*, 2008; Wanamaker *et al.*, 2008]. It is necessary to understand fundamental processes of bivalve biomineralization in order to understand differences between studies [e.g., Carré *et al.*, 2006; Heinemann *et al.*, 2008]. Extracellular body fluids (hemolymph and extrapallial fluid) are the connection between tissues and shell. Especially the extrapallial fluid (EPF) may influence calcification processes as it fills the extrapallial cavity enclosed by the shell, the periostracum and the outer mantle margin [Wilbur and Saleuddin, 1983].

[8] Thus, to contribute to the understanding of the ability of calcifying organisms to live under acidified conditions and of biomineralisation mechanisms of *M. edulis* we sampled hemolymph (HL) and extrapallial fluid (EPF) of mussels at the end of two CO₂ perturbation experiments. Fluids were analyzed for pH, C_T and elemental ratios. To consider the suitability of *M. edulis* shell as a proxy archive for ocean pH, boron isotopes ($\delta^{11}\text{B}$) were investigated by LA-MC-ICP-MS (method see Fietzke *et al.* [2010]) in shell portions precipitated during the experimental incubation (shells from the long-term growth study described by Thomsen *et al.* [2010]).

2. Material and Methods

2.1. Culture and Samples

2.1.1. General Setup

[9] Two experiments were conducted to investigate how increased water *p*CO₂ influences different parameters (pH, *p*CO₂, HCO₃⁻, CO₃²⁻ and B/Ca, Mg/Ca and Sr/Ca ratios) of *Mytilus edulis* extracellular fluids and to test whether shell boron isotopes can be used as a pH proxy of the surrounding seawater.

[10] Atmospheric *p*CO₂ averaged 386 μatm in 2009 and concentrations are predicted to reach values between 700 and 1000 μatm by the year 2100 [Intergovernmental Panel on Climate Change, 2007]. Particular marine habitats like Kiel Fjord (Western Baltic Sea, Germany) are already exposed

to values above 2000 μatm and might occasionally encounter *p*CO₂ values of >4000 μatm if water surface *p*CO₂ doubles [see Thomsen *et al.*, 2010]. Thus, *p*CO₂ levels of 380–4000 μatm were chosen for the experimental incubations. All specimen used in the experiments were collected from subtidal populations in Kiel Fjord (54° 19.8' N; 10° 9.0' E) and cultured at the culturing facilities of the Leibniz Institute of Marine Sciences (IFM-GEOMAR) in Kiel. Animals were acclimated under control conditions for 2 weeks prior to experimentation.

[11] The experiments were conducted in a flow through system (200 ml min⁻¹ per experimental unit) using water from Kiel Fjord which was cleaned (50–5 μm filters, UV-sterilized) and pumped into a storage tank, where it was aerated. The air saturated water was pumped to a header tank and then supplied to the aquaria via gravity feed. Different CO₂-air-mixtures were used to equilibrate the experimental seawater (for a more detailed description see the auxiliary material).¹

[12] Water pH was measured daily between Mondays and Fridays with a WTW pH meter (pH 340i, electrode: Sen Tix 81 (calibrated with Radiometric IUPAC precision pH buffer 7 and 10), WTW GmbH, Weilheim, Germany; $\Delta\text{pH} \pm 0.01$). All pH values measured with the pH meter were corrected with respect to the pH values calculated from weekly A_T and C_T measurements (n = 223, R² = 0.949) as the latter provide more accurate results:

$$\text{pH}_{\text{corrected}} = 0.9781 * \text{pH}_{\text{measured}} + 0.2123 \quad (2)$$

For all pH values reported in this paper, the NBS scale was used.

[13] Hemolymph (HL) and extrapallial fluid (EPF) samples were taken directly after removing the bivalves from the aquaria. Valves were opened carefully to not injure the mantle and held open with a 1000 μl pipette tip. Hemolymph was drawn anaerobically with a syringe from the posterior adductor muscle and EPF was taken from the extrapallial space by inserting a flexible syringe needle (\emptyset 0.6 × 80 mm) between shell and the pallial line.

[14] From all aquaria water temperature, salinity (with WTW cond 315i salinometer and WTW TETRACON 325 probe) and pH were measured daily. A_T and C_T samples were taken for carbonate system calculations (Table 1; for more details of exp. 2 see Thomsen *et al.* [2010]).

¹Auxiliary materials are available in the HTML. doi:10.1029/2011GC003790.

Table 1. Water Conditions During Experimental Trials^a

Treatment	pH _{NBS}	A _T (μmol kg ⁻¹)	C _T (μmol kg ⁻¹)	pCO ₂ (μatm)	Ω _{calc}	Ω _{arag}
<i>First Experiment^b</i>						
387 μatm	8.02 ± 0.08	2031 ± 74	1965 ± 63	577 ± 95	1.8 ± 0.3	1.1 ± 0.2
560 μatm	7.93 ± 0.07	2031 ± 74	1988 ± 62	722 ± 97	1.5 ± 0.2	0.9 ± 0.1
840 μatm	7.86 ± 0.05	2032 ± 72	2004 ± 65	839 ± 80	1.3 ± 0.2	0.8 ± 0.1
1120 μatm	7.77 ± 0.05	2033 ± 73	2026 ± 66	1041 ± 93	1.1 ± 0.1	0.5 ± 0.1
1400 μatm	7.71 ± 0.05	2033 ± 73	2040 ± 66	1195 ± 98	0.9 ± 0.1	0.6 ± 0.1
4000 μatm	7.28 ± 0.08	2040 ± 72	2172 ± 66	3352 ± 566	0.4 ± 0.1	0.2 ± 0.04
<i>Second Experiment^c</i>						
387 μatm	8.13 ± 0.02	1966 ± 3	1891 ± 5	500 ± 30	1.94 ± 0.04	1.14 ± 0.04
1400 μatm	7.72 ± 0.06	1968 ± 5	1984 ± 12	1350 ± 200	0.81 ± 0.09	0.48 ± 0.06
4000 μatm	7.26 ± 0.04	1970 ± 4	2126 ± 13	3950 ± 220	0.28 ± 0.02	0.17 ± 0.01

^aData for experiment 2 are from *Thomsen et al.* [2010]. Errors are given as SD. Values represent mean over whole experiment.

^bTemperature in °C, 12.1 ± 0.6; salinity, 18.8 ± 1.59.

2.1.2. Experiment 1

[15] The first experiment was conducted from September, 3rd to December, 18th 2008. We cultured *Mytilus edulis* (~140 individuals per tank) ranging in size from 12.4 mm to 46.7 mm at six different pCO₂ levels (380, 560, 840, 1120, 1400 and 4000 μatm). Shell length was measured at the longest axis (from umbo to edge at opposite site of the shell) using calliper with an accuracy of 0.1 mm. Fluids were sampled from the five largest bivalves (45.6 ± 0.6 mm) in each treatment. The experimental aquaria had a volume of 15 l and were exposed to a 14:10 hour light:dark cycle. Water pH ranged from 8.02 ± 0.08 to 7.28 ± 0.08 in the different treatments. Temperature was 12.2 ± 0.6 °C and salinity 18.8 ± 1.6.

[16] During the first 6 weeks mussels were fed with “DTs Live Marine Phytoplankton–Premium Reef Blend” (DT’s Plankton Farm, Sycamore, IL, USA), a live phytoplankton mixture of three marine algae species (*Phaeodactylum tricornotum* (40%), *Nannochloropsis oculata* (40%) and *Chlorella sp.* (20%)). Since mussel growth was unsatisfactory during that period the animals were additionally fed with *Artemia salina* after that period as *Wong and Levinton* [2004] reported best growth of *M. edulis* with a mixture of phyto- and zooplankton. Food was provided on 3 days a week (~75 mg algal biomass and ~45 mg *Artemia salina* (both dry weight) day⁻¹ aquarium⁻¹) by closing the flow through for 4 hours and adding food directly into the aquaria.

[17] Fluid samples were centrifuged (30 s) and pH was measured with a microelectrode (WTW Mic-D). Then total dissolved inorganic carbon (C_T) was determined immediately after sampling via a Corning 965 CO₂ analyzer, (Olympic Ana-

lytical Service Essex, UK; accuracy 0.1 mM) in two 100 μl subsamples. Carbonate parameters (pCO₂, [HCO₃⁻], and [CO₃²⁻]) of the fluids were calculated from measured pH and C_T with using the Henderson-Hasselbalch equation:

$$p\text{CO}_2 = C_T \left(10^{\text{pH} - \text{p}K'_1} \cdot \alpha_{\text{CO}_2} + \alpha_{\text{CO}_2} \right)^{-1} \quad (3)$$

$$[\text{HCO}_3^-] = 10^{\text{pH} - \text{p}K'_1} \cdot \alpha_{\text{CO}_2} \cdot p\text{CO}_2 \quad (4)$$

$$[\text{CO}_3^{2-}] = 10^{\text{pH} - \text{p}K'_2} \cdot [\text{HCO}_3^-] \quad (5)$$

$$\text{p}K'_1 = \text{pH} - \log \left(\frac{C_T}{p\text{CO}_{2\alpha\text{CO}_2}} - 1 \right) \quad (6)$$

where α is the CO₂ solubility coefficient. pK₁' and pK₂' are the first and second apparent dissociation constants of carbonic acid. α_{CO2} was calculated (0.045 mol/l*μatm) [*Weiss*, 1974] and pK₂' (9.321) [*Roy et al.*, 1993] was chosen according to experimental temperature and salinity. pK₁' was calculated from pH_{NBS}, C_T and pCO₂ (measured in *Mytilus edulis* body fluids by [*Thomsen et al.*, 2010]) using equation (6) [*Albers and Pleschka*, 1967]. A linear relationship was found for pK₁' in relation to pH_{NBS}. After adjusting temperature and salinity according to this experiment the regression for pK₁' for *M. edulis* internal fluids was pK₁' = -0.1323pH + 7.2371 (R² = 0.3027).

2.1.3. Experiment 2

[18] From 14th of May to 13th of July a second (long-term growth) experiment was conducted using three pCO₂ levels (380, 1400 and 4000 μatm). The level of replication was four. Each 15 l aquarium contained eight small (mean 5.5 mm), eight medium sized (mean 13 mm) and two big (mean

Table 2. Instrumental Parameters Laser Ablation

Parameter	Value
<i>AXIOM MC-ICP-MS</i>	
Cool gas	14 l/min
Auxiliary gas	1.8 l/min
Nebulizer gas	0.9 l/min (Ar)
RF power	1250 W
Reflected power	3 W
Accelerating voltage	4972 V
Cones	R.A. Chilton RAC19/RAC705
Resolution	500res
<i>UP193FX</i>	
Ablation cell gas	0.6 l/min (He)
Spot size	150 μm
Fluence	2.3 J/cm ²
Fluence Preablation	0.6 J/cm ²
Repetition rate	30 Hz
Scan mode	Line scan (1000 μm @ 12 $\mu\text{m/s}$)
Scan mode Preablation	Line scan (1000 μm @ 50 $\mu\text{m/s}$)
Pause between lines	90 s

40 mm) mussels. Shells from the medium specimen (13.3 ± 1.4 mm) and fluids from the big specimen were investigated in this study. The experiment was conducted under a mean temperature of 13.8 ± 0.6 °C and a salinity of 15 ± 0.6 . The light/dark cycle was 14:10 hours. Food (*Rhodomonas sp.* suspension, 2900 ± 1200 cells ml⁻¹) was provided continuously by adding it into the aquaria at a rate of 100 ml min⁻¹. Body fluids of the big specimen were frozen at -20 °C directly after sampling and used for elemental analyses by ICP-OES (SPECTRO *Ciros*^{CCD} SOP) at the Institute of Geosciences, University Kiel, Germany. The tissue of the medium sized mussels was removed and shells were dried at 20 °C for 5 days.

[19] EPF pH could not be measured directly in experiment 2 specimen as there was not enough material for sub sampling and therefore no possibility to measure pH directly without contaminating the fluid samples with respect to boron (due to the high boron content of the glass pH electrode). Therefore boron isotopes and elemental ratios (experiment 2) were compared to EPF pH values estimated from experiment 1 fluid values ($n = 12$) and HL pH values measured in large *M. edulis* of an additional experiment presented by *Thomsen et al.* [2010]. The regression for the fluids and the corresponding water was $\text{pH}_{\text{EPF}} = 0.383(37) \text{pH}_{\text{water}} + 4.493(288)$ ($R^2 = 0.914$). Additionally the mean offset between EPF and HL (0.037; experiment 1 this study; $n = 6$ each) was considered. Regression results for each treatment and the mean offset between HL and EPF pH have been subtracted from water pH of experiment 2.

2.2. Analytical Methods

[20] The fluids from the second experiment (5–7 samples per treatment) were thawed and diluted 50fold with ultrapure 2% HNO₃. They were analyzed for Mg/Ca, Sr/Ca and B/Ca elemental ratios by ICP-OES (Inductively Coupled Plasma–Optical Emission Spectrometry; SPECTRO *Ciros*^{CCD} SOP) at the Institute of Geosciences, University Kiel, Germany. An intensity-ratio calibration procedure was applied using matrix-matched calibration standards and IAPSO seawater as a consistency standard. External precision of the elemental ratios was ~0.1% RSD for Mg/Ca and Sr/Ca, and 1–4% RSD for B/Ca.

[21] Ten shells from each treatment of the second experiment were broken to obtain fragments that were grown under treatment conditions (defined by length growth). Subsequently, the fragments were bleached in 10% NaOCl (1% active chlorine) for two days to remove organic components (periostracum and other shell organics). After 24 hours samples were placed into an ultra sonic bath for 10 min and the bleaching solution was replaced. Afterwards shells were rinsed three times with ultrapure water (18.2 MOhm) and then dried at 20 °C. To prevent dissolution of the carbonates the water was adjusted to a pH of ~9 by adding NH₄OH. Boron isotopes were measured by LA-MC-ICP-MS (Laser Ablation–Multicollector–Inductively Coupled Plasma–Mass Spectrometry; Thermo Fisher MC-ICP-MS *AXIOM*, originally designed and manufactured by VG) connected to an ESI New Wave Research *UP193FX* excimer laser ablation system equipped with an ESI New Wave Research *LFC* (large format cell) via a standard sample standard bracketing procedure described by *Fietzke et al.* [2010]. Soda lime glass SRM (NIST610) was used as external standard as *Fietzke et al.* [2010] showed that there is no matrix effect between carbonate and glass. First, three lines (1000 μm x 150 μm) were measured on the NIST. Then, seven lines of sample and seven lines of standard were measured in alternation. Prior to each of these measuring procedures, a preablation was conducted to remove surface contamination. All instrumental parameters are given in Table 2. The location of the sample lines on *M. edulis* shells is shown in Figure 1.

3. Results and Discussion

3.1. Water Parameters

[22] Since the experimental incubation systems used in this study were flow through designs with con-




Figure 1. (a) Modified schematic anatomy of a bivalve (redrawn after McConnaughey and Gillikin [2008]). (b) Position of laser ablation lines on *M. edulis* shell (scheme of whole shell and picture of sampled Laser area).

tinuous supply of water from Kiel Fjord, all treatments were influenced by changes in Kiel Fjord water chemistry. Although the water was vigorously aerated prior to introduction into the experimental aquaria, mean seawater $p\text{CO}_2$ deviated from the nominal values. Mean seawater conditions during the two experiments are given in Table 1. Carbonate system speciation measured in the extracellular fluids are not compared to these mean values but to values from dates of sampling (Table 3).

3.2. Acid-Base Parameters of Fluid Samples (Experiment 1)

[23] Overviews of all mean extracellular acid-base parameters of the internal fluids (HL and EPF) are given in Table 3 and Figure 2 (for detailed data and regressions see the auxiliary material).

[24] High extracellular $p\text{CO}_2$ values between 1000 and 4000 μatm are found in all aquatic ectothermic metazoans, as diffusive excretion of metabolic CO_2 depends on a relatively steep gradient of CO_2 from the body fluids to the seawater (see Melzner *et al.* [2009] for a review). Additional increases in extracellular $p\text{CO}_2$ under hypercapnia [Pörtner *et al.*, 2004; Michaelidis *et al.*, 2005; Spicer *et al.*, 2007; Thomsen *et al.*, 2010] are necessary in order to maintain metabolic CO_2 flux. In this study a linear increase of fluid $p\text{CO}_2$ with increasing water $p\text{CO}_2$

was measured, with slightly lower values in the HL (1648–3122 μatm) than in the EPF (1938–3430 μatm). However, $p\text{CO}_2$ values in HL and EPF were not significantly different from each other. At the highest $p\text{CO}_2$ treatment a reduced $p\text{CO}_2$ diffusion gradient could be observed (only low offset between water and extracellular body fluid $p\text{CO}_2$), indicating reductions in metabolism in experiment 1. Similar reductions in metabolic rate have been observed in *M. galloprovincialis* under increased water $p\text{CO}_2$ by Michaelidis *et al.* [2005]. This was indicated by a significant decrease in oxygen consumption. However, no reductions in metabolism were witnessed in another experiment using Baltic Sea *M. edulis* at 4000 μatm under higher food concentrations, suggesting that the observed reduction in the CO_2 diffusion gradient in experiment 1 of this study might be due to food limitation [Thomsen and Melzner, 2010]. A recent study on the same mussel population could confirm the strong role of food supply on biomineralisation and shell corrosion intensity [Melzner *et al.*, 2011].

[25] High HL and EPF $p\text{CO}_2$ values are the primary driving force for comparatively low HL and EPF pH values: values range from 7.37 to 7.54 (HL) and 7.34 to 7.52 (EPF) respectively, which is significantly lower in all treatments than seawater pH with values of 8.10 to 7.39. Due to the progressive reductions in the $p\text{CO}_2$ gradient between extracellular fluids and seawater, the pH offset

Table 3. Acid-Base Status of Hemolymph and Extrapallial Fluid and of the Water at Date of Sampling (Experiment 1)

Treatment $p\text{CO}_2$ (μatm) at Day of Sampling	EPF (SD, n)	HL (SD, n)	Water at Day of Sampling
pH			
452	7.52 (0.04; 3)	7.54 (0.05; 3)	8.10
574	7.50 (0.09; 4)	7.53 (0.05; 3)	8.01
731	7.46 (0.05; 3)	7.52 (0.04; 3)	7.91
906	7.45 (0.02; 3)	7.47 (0.05; 4)	7.82
1123	7.41 (0.04; 4)	7.47 (0.07; 4)	7.73
2724	7.34 (0.09; 4)	7.37 (0.02; 4)	7.39
C_T (mM)			
452	1.79 (0.18; 3)	1.63 (0.31; 3)	1.96
574	1.41 (0.03; 4)	1.62 (0.24; 3)	1.98
731	1.71 (0.18; 3)	1.79 (0.21; 3)	2.01
906	1.55 (0.07; 3)	1.77 (0.14; 4)	2.03
1123	1.85 (0.05; 3)	1.61 (0.34; 4)	2.05
2724	1.91 (0.07; 4)	1.89 (0.31; 4)	2.15
$p\text{CO}_2$ (μatm)			
452	1938 (93; 3)	1648 (183; 3)	452
574	1717 (427; 4)	1623 (190; 3)	574
731	2297 (284; 3)	1987 (383; 3)	731
906	2033 (169; 3)	2226 (348; 4)	906
1123	2611 (181; 4)	2199 (744; 4)	1123
2724	3430 (696; 4)	3122 (578; 4)	2724
HCO_3^- (mM)			
452	1.70 (0.18; 3)	1.55 (0.30; 3)	1.85
574	1.33 (0.04; 4)	1.54 (0.23; 3)	1.89
731	1.61 (0.17; 3)	1.70 (0.19; 3)	1.92
906	1.46 (0.07; 3)	1.67 (0.13; 4)	1.94
1123	1.73 (0.06; 4)	1.52 (0.31; 4)	1.96
2724	1.77 (0.08; 4)	1.75 (0.28; 4)	2.02
CO_3^{2-} (mM)			
452	0.028 (0.005; 3)	0.027 (0.009; 3)	0.085
574	0.020 (0.005; 4)	0.027 (0.005; 3)	0.070
731	0.022 (0.004; 3)	0.027 (0.003; 3)	0.056
906	0.020 (0.001; 3)	0.024 (0.003; 4)	0.047
1123	0.023 (0.003; 4)	0.021 (0.003; 4)	0.038
2724	0.019 (0.004; 4)	0.019 (0.003; 4)	0.017

between fluids and water decreased as well. At the highest seawater $p\text{CO}_2$ level (2724 μatm) pH values of EPF/HL and water became nearly identical. For each treatment, pH values of HL were slightly lower than EPF pH with a mean offset of 0.04 units. However, these differences were not significantly different. Fluid pH values decreased by 0.17 units in the HL and by 0.18 units in the EPF compared to a drop of 0.71 units in the water. The change of hemolymph pH relative to the

change in seawater ($\Delta\text{pH}_{\text{HL}}/\Delta\text{pH}_{\text{SW}}$) in this study was consistent with that found by *Michaelidis et al.* [2005] (Table 4). The pH in *M. galloprovincialis* HL decreased from 7.55 to 7.36 (SW: 8.05 to 7.3). Although the control water $p\text{CO}_2$ in their study was more than double compared to the study of *Thomsen et al.* [2010] and this study (1079 μatm and 470/452 μatm , respectively) the water pH was nearly the same. This may be due to the difference in alkalinity but also to the observed accumulation of bicarbonate in that study. In general the Mediterranean Sea is characterized by a higher alkalinity (2500 $\mu\text{mol kg}^{-1}$) in comparison to the Baltic Sea ($\sim 1900 \mu\text{mol kg}^{-1}$). The resulting higher buffer capacity leads to a smaller drop in pH in the Mediterranean Sea at comparable seawater $p\text{CO}_2$ values. This explains the different results with respect to hemolymph pH between the studies of *Thomsen et al.* [2010] and of *Michaelidis et al.* [2005] although the $p\text{CO}_2$ was nearly similar. *Thomsen et al.* [2010] measured a drop in *M. edulis* HL pH relative to ambient water pH twice as high (8.05 to 7.08 (SW) related to 7.59 to 7.16 (HL)) as measured by *Michaelidis et al.* [2005] and in this study. *Thomsen et al.* [2010] conducted experiments at $p\text{CO}_2$ values ranging from 470 to 4250 μatm . However, total alkalinity in the present study was 130 to 150 $\mu\text{mol/kg}$ higher than in that of *Thomsen et al.* [2010], which is related to the lower salinity in the latter (11.8 compared to 18.8 in this study). This disparity in salinity/alkalinity was due to the high seasonal variability in Kiel Fjord and is likely to be responsible for the different drop in body fluid pH. These results show the high variability in extracellular pH that can be caused by the strong influence of alkalinity and of the reduced metabolism (after three months of experiment with low food conditions) on the pH decrease in response to increased $p\text{CO}_2$.

[26] However, no extracellular accumulation of bicarbonate (HCO_3^-) could be measured in HL or EPF, indicating that *M. edulis* does not actively buffer HL or EPF to avoid decreasing pH values. For both fluids, $[\text{HCO}_3^-]$ was below that of the seawater and no significant differences (non-linear regressions) were found between HL and EPF. Accumulation of $[\text{HCO}_3^-]$ (nearly equivalent to a net excretion of protons) is an efficient mechanism of extracellular pH stabilization that is primarily employed by active marine ectothermic organisms that are characterized by pH sensitive respiratory pigments (see *Melzner et al.* [2009] for a review). The magnitude of this $[\text{HCO}_3^-]$ accumulation response varies between taxa, but highest degrees of $[\text{HCO}_3^-]$




Figure 2. Acid-base status of hemolymph and extrapallial fluid and of the seawater at sampling days compared to treatment $p\text{CO}_2$ at the day of sampling (exp. 1).

related pH compensation are found in teleost fish, decapod crustaceans and cephalopod molluscs [Larsen *et al.*, 1997; Pane and Barry, 2007; Gutowska *et al.*, 2010]. Two studies also reported $[\text{HCO}_3^-]$ accumulation in mussels to buffer extracellular pH [Lindinger *et al.*, 1984; Michaelidis *et al.*, 2005].

However, both of these were conducted in closed/recirculating systems and it is evident [Lindinger *et al.*, 1984] or very possible [Michaelidis *et al.*, 2005] that this increase in extracellular $[\text{HCO}_3^-]$ was due to increases in ambient water $[\text{HCO}_3^-]$ due to external/internal shell dissolution [see Thomsen

Table 4. Comparison Between the Hemolymph Acid-Base of This Study and the Studies of *Michaelidis et al.* [2005] and *Thomsen et al.* [2010]

	pCO ₂ SW (μatm)	pCO ₂ HL (μatm)	Δ pCO ₂ (HL-SW)	pH _{SW}	pH _{HL}	ΔpH _{HL} /ΔpH _{SW}	Salinity	A _T
<i>M. edulis</i> , control	452	1648	1196	8.10	7.54	0.239	18.8	2031
This study, treatment	2724	3122	398	7.39	7.37		18.8	2040
<i>M. edulis</i> , control	470	1697	1227	8.05	7.55	0.402	15.0	1901
<i>Thomsen et al.</i> [2010], treatment	4310	4960	650	7.08	7.16		15.0	1891
<i>M. galloprovincialis</i> , control	1079	1513	434	8.05	7.59	0.267	32.0	
<i>Michaelidis et al.</i> [2005], treatment	5026	5724	697	7.30	7.36		32.0	

et al., 2010]. Consequently, carbonate concentrations are lower in HL and EPF than in seawater, decreasing with decreasing pH (increasing pCO₂). In the highest pCO₂ treatment (2724 μatm), HL and EPF [CO₃²⁻] were almost equal to seawater [CO₃²⁻] within error bars.

[27] Generally, it is important to note that the acid-base conditions in the body fluids, especially in the EPF, which is in direct contact with the inner shell surface, may impact calcification and existing shell structures. Reduced shell growth under high pCO₂ (4000 μatm) was reported by *Thomsen et al.* [2010] and *Thomsen and Melzner* [2010]. Yet experiment 2 specimens increased their shell mass by >150% during the 8 week experiment, indicating great biological control and an ability to precipitate a shell in fluids that are highly undersaturated with regard to calcium carbonate. However, results from *Thomsen et al.* [2010] demonstrate that external shell dissolution at the umbo region and thinner aragonite layers may be consequences of high seawater and EPF pCO₂ values.

3.3. Elemental Ratios of Extrapallial Fluid and Water (Experiment 2)

[28] Elemental ratios (B/Ca, Mg/Ca and Sr/Ca) were measured in the EPF of *M. edulis* cultured at

three different pCO₂ levels (experiment 2, Table 5). Figure 3 shows the elemental ratios compared to EPF pH. The pH values were extrapolated using a regression of HL data on water pH (see methods). An increase of all elements (B, Mg and Sr) relative to Ca could be observed with rising pH. Although a high variability of the element concentrations was found in the fluid, especially B/Ca and Mg/Ca showed a linear trend (significant at 95% confidence level), while there is no significant trend for Sr/Ca. The variability between different individuals may be due to a strong individual physiological influence on element partitioning. The magnitude of increase with pH was variable for the different elements. For B/Ca ratios (19.8%) it was slightly more than twice as high as for Mg/Ca ratios (7.9%) and lowest for Sr/Ca ratios (1.2%).

[29] While minor amounts of Mg²⁺ and Sr²⁺ substitute for Ca²⁺ during calcification boron is believed to be incorporated into carbonates as B(OH)₄ replacing the carbonate ion in the lattice [*Vengosh et al.*, 1991; *Hemming and Hanson*, 1992; *Hemming et al.*, 1995; *Pagani et al.*, 2005]. Several studies reported different abiotic factors influencing the composition of bivalve shells. The B/Ca ratio has been shown to be related to pH in different carbonates [*Hemming and Hanson*, 1992; *Hobbs and*

Table 5. pH and Me/Ca Ratios of Treatment Water and EPF^a

Treatment in μatm	Culture pH (SD; n) Experiment 2	EPF pH Calculated	Percent Shell Mass Growth
380	8.08 (0.01; 3)	7.55	242
1400	7.68 (0.01; 4)	7.40	236
4000	7.23 (0.03; 4)	7.22	158

Treatment in μatm	EPF B/Ca (SD; n) in mmol/mol	Water B/Ca (SD; n) in mmol/mol	EPF Sr/Ca (SD; n) in mmol/mol	Water Sr/Ca (SD; n) in mmol/mol	EPF Mg/Ca (SD; n) in mmol/mol	Water Mg/Ca (SD; n) in mmol/mol
380	56.2 (1.08; 5)	54.3 (1.54; 3)	8.10 (0.09; 6)	8.09 (0.05; 3)	4947 (54; 6)	4859 (22; 3)
1400	55.0 (3.56; 5)	53.9 (3.72; 4)	8.05 (0.12; 5)	8.13 (0.04; 4)	4878 (125; 5)	4974 (124; 4)
4000	52.7 (2.71; 6)	53.1 (2.09; 4)	8.06 (0.08; 6)	8.13 (0.12; 4)	4817 (62; 6)	4898 (116; 4)

^aWater pH values represent the day of sampling. SD represents the error of different individuals (EPF) and different aquaria, respectively. The precision based on repeated standard (IAPSO) measurements was 0.20–0.25% for Mg, Ca and Sr and 0.88% for boron. EPF pH was calculated from experiment 1 and *Thomsen et al.* [2010] (see methods).




Figure 3. Elemental ratios of the EPF vs. calculated internal (EPF) pH values. Error bars show 1SD. Linear regressions show an insignificant increase (19.8% for B/Ca, 7.9% for Mg/Ca and 1.2% for Sr/Ca).

Reardon, 1999; Sanyal *et al.*, 2000; Foster, 2008]. Ni *et al.* [2007] found a systematic increase of B/Ca in *G. ruber* and *G. sacculifer* with increasing test size and suggested growth rate variations influencing the incorporation of boron. The results of McCoy *et al.* [2011] in *Mytilus californianus* shell did not reflect a distinct year-to-year correlation of boron concentrations with pH. However, the annual boron concentration probably reflects seasonal pH changes. The authors concluded that biological control of pH and/or boron concentrations in the EPF are the reasons for these observations. Our results show low EPF pH to be a result of high extracellular $p\text{CO}_2$ (Figure 2). As discussed above, high EPF $p\text{CO}_2$ values are necessary to excrete metabolic CO_2 .

Mg/Ca ratios in carbonates have been shown to be temperature-related in inorganically precipitated calcite [e.g., Oomori *et al.*, 1987; Lopez *et al.*, 2009] as well as in biogenic carbonates precipitated by different species like foraminifera [Nürnberg *et al.*, 1996; Elderfield and Ganssen, 2000; Kısakürek *et al.*, 2008] or bivalves [Klein *et al.*, 1996a; Wanamaker *et al.*, 2008]. Hiebenthal [2009] also reported a positive Mg/Ca-temperature relation in *M. edulis* shells from Kiel Fjord. Salinity was shown to likewise have an effect on Mg/Ca ratios in *M. edulis* calcite [Dodd, 1965].

[30] Other studies suggested different biological influences being responsible for the variations of




Figure 4. Distribution coefficient of EPF and water elemental ratios compared to shell mass growth of bivalves during experimental time (exp. 2). Errors are given as 1SD.

Me/Ca ratios between individuals of one species but also within single shells. In addition to their results mentioned above, *McCoy et al.* [2011] found organic-rich winter growth bands containing elevated B/Ca ratios. A comparable effect was reported by *Schöne et al.* [2010] for trace elements in *Arctica islandica* shells. They found a strong influence of organic matrix on the determination of Mg, Sr and Ca. This confirms our results from qualitative laser ablation measurements that show boron concentrations in the periostracum being about one magnitude higher than in the carbonate. *Carré et al.* [2006] observed increasing Me/Ca ratios (for Mg, Ba, Mn and especially Sr) in aragonitic bivalve shells (*Mesodesma donacium*, *Chione subrugosa*) when crystal growth rates increase. Their model predicts decreasing Ca^{2+} -channel selectivity when rates increase. *Klein et al.* [1996b] found significantly higher Sr/Ca ratios in a young, rapidly grown *Mytilus trossulus* than in a slowly grown adult individual. They concluded that shell precipitation along lateral margins is dominantly controlled by mantle metabolic activity at the site of carbonate formation. This suggests that also the elemental composition is dependent on the mantle activity as it seems to be influenced by growth rates. *Carré et al.* [2006] suggested the Sr/Ca measurements from different sections of *M. trossulus* shells (measured by *Klein et al.* [1996b]) can also be explained by differences in crystal growth. The results from *Ford et al.* [2010] support that of this study and show Mg/Ca ratios in shells of *Mytilus californianus* to be a function of growth rate rather than temperature related. *Takesue et al.* [2008] also found growth rate dependent alterations in Sr/Ca,

B/Ca and Ba/Ca ratios in valves from *Corbula amurensis*. Sr/Ca seems to be influenced by both temperature and salinity [*Dodd, 1965; Wanamaker et al., 2008*] but also by precipitation rates [*Lorens, 1981; Lorrain et al., 2005; Freitas et al., 2006*].

[31] Experiment 2 consisted of four replicate aquaria (two big individuals per aquarium) for each treatment, and variability in the elemental ratios of the water between different aquaria was observed (single values see auxiliary material). To eliminate this problem Figure 4 shows the distribution coefficients between EPF and water. When plotted against growth rates (shell mass growth during experiment in % from [*Thomsen et al., 2010*]) elemental ratios are slightly lower in the EPF than in the ambient water at lower growth rates at 4000 μatm . For this treatment external shell dissolution at the umbo region as well as thinner aragonite layers were observed [*Thomsen et al., 2010*]. At high growth rates ($p\text{CO}_2$ of 385 and 1400 μatm), elemental ratios are higher in the EPF than in the ambient water. Thus, with increasing growth rates elements like B, Mg and Sr become enriched in the EPF with respect to Ca. This effect may be due to higher growth and calcification rates. Therefore the EPF is more depleted with respect to calcium when compared to other elements and the elemental ratios of the EPF are shifted. As elemental ratios in the EPF increased with growth rate it is also likely that the elemental ratios in the shells rose when precipitated from altered fluids. Therefore our results support findings from different studies showing increased Me/Ca ratios in shells grown at high rates [e.g., *Klein et al., 1996b; Carré et al., 2006; Ford et al., 2010*].

Table 6. pH and $\delta^{11}\text{B}$ of Treatment Water and EPF^a

Culturing Water pH (2SE; n) Experiment 2	EPF pH Calculated	Shell $\delta^{11}\text{B}$ (SD; n) (‰, Experiment 2)	Shell Length (SD; n) (mm, Experiment 2)
8.11 (0.008; 53)	7.546	17.4 (3.1; 10)	22.9 (2.4; 10)
7.73 (0.020; 53)	7.396	18.3 (4.8; 11)	24.0 (1.4; 11)
7.28 (0.035; 53)	7.221	17.9 (5.5; 11)	20.7 (1.9; 11)

[32] Food availability, temperature, salinity and elevated $p\text{CO}_2$ (decreased pH) are influencing growth rates (and accordingly the amount of organic matter in the shell) of *M. edulis* [e.g., Malone and Dodd, 1967; Kautsky, 1982; Kossak, 2006; Thomsen et al., 2010]. While the biological performance depends on these biotic and abiotic factors, seawater $p\text{CO}_2$ or pH does not seem to be directly reflected in Me/Ca ratios of bivalve carbonates, rather, they might influence them indirectly via decreased rates of growth and calcification due to a repartitioning of the energy budget of the mussel [Thomsen and Melzner, 2010]. Our results show that the dependence between growth and Me/Ca ratios incorporation is represented in the extrapallial fluid. Thus, elemental ratios (B/Ca, Mg/Ca, and Sr/Ca) in the EPF of *M. edulis* may primarily serve as an indicator for growth rates/shell dissolution as they become enriched with increasing length growth and depleted by dissolution. This indicates that *Mytilus edulis* shells cannot be directly used as archive for experienced environmental conditions. Extensive calibrations are necessary to see how parameters such as temperature or salinity influence growth rate. With such calibrations and one for the influence of growth rate on Me/Ca ratios in the shell it may be possible to reconstruct environmental conditions. However, the encountered variability is high due to individual physiology and a high amount of replicates must be measured.

3.4. Boron Isotope ($\delta^{11}\text{B}$) Data of *M. edulis* Shells (Experiment 2)

[33] Shell portions precipitated during the experimental duration (additional length growth) were investigated for their $\delta^{11}\text{B}$ composition (Table 6). While the analytical precision of the carbonate boron isotope ratio determination was in the order of 0.9 ‰ the variability between individuals from the same treatment was much higher (3.1–5.5 ‰). This analytical precision results from a very low boron content in the shells, still the inter-individual variability is the dominating factor compromising the use of $\delta^{11}\text{B}$. Using the empirical borate $\delta^{11}\text{B}/\text{pH}$ dependence [Klochko et al., 2006] and the differ-

ence in the ambient water pH we expect a $\delta^{11}\text{B}$ change of 4.3 ‰ for this pH range (7.28–8.11). Considering this we would expect to find a systematic change in the mean $\delta^{11}\text{B}$ values depending on the pH. The results showed a high variability not only between different individuals but also within single shells. The high individual variability was already observed for the fluid elemental ratios. Additionally the considerable amount of boron in the periostracum (see section 3.3) suggests the organic matrix surrounding every single carbonate crystal could be a reason for this variability. Bivalve shells typically contain up to 5% of organic matter in the prismatic and nacre layer, respectively [e.g., Wilbur and Bernhardt, 1984]. The variability may be caused by individual and physiological influences as well as (crystal) growth rate. It is obvious that one factor inducing the variability in shell $\delta^{11}\text{B}$ is the variability in body fluid pH (EPF), not only between individuals but also over time. Extracellular pH for example was shown to decrease when bivalves close their shell valves [e.g., Crenshaw and Neff, 1969; Crenshaw, 1972]. Crenshaw [1972] observed a drop in EPF pH of *Mercenaria mercenaria* from 7.41 to 7.25 while valves were closed for 15 min or longer and a maximum drop in *M. edulis* to 7.2. The pattern how often and how long they close their valves for instance can be different between individuals. Average values of $\delta^{11}\text{B}$ ranged from 17.4 in the lowest $p\text{CO}_2$ (highest pH) to 17.9 ‰ in the highest $p\text{CO}_2$ (lowest pH) treatment, respectively. However, the highest value (18.3 ‰) was found at 1400 μatm but within the high variability no significant difference could be observed. Moreover no clear trend could be observed with pH but values differed from those of other biominerals when plotted against ambient water pH (Figure 5). Inorganically precipitated calcite [Sanyal et al., 2000], foraminifera [Sanyal et al., 2001] and corals [Hönisch et al., 2004; Reynaud et al., 2004] are characterized by $\delta^{11}\text{B}$ values that fit above the measured $\delta^{11}\text{B}$ curve of $\text{B}(\text{OH})_4^-$ presented by Klochko et al. [2006] (Figure 5) which correlates well with ambient water pH. For the curves plotted the following parameters were used: $\text{pK}_\text{B} = 8.597$ [Dickson, 1990]; $\alpha = 1.0194$ [Kakihana et al., 1977]




Figure 5. $\delta^{11}\text{B}$ measured in *M. edulis* shells and precipitated under different pCO_2 conditions in relation to data of other carbonates. Blue diamonds are displayed versus mean water pH during time of experiment 2, pink diamonds display the same $\delta^{11}\text{B}$ data plotted against EPF pH from experiment 1. *Tridacna gigas* data from Kasemann *et al.* [2009] are shown as a function of the approximated EPF pH value of *Tridacna squamosa* (at 28°C, SAL 31) observed by Ip *et al.* [2006] as no other data could be found. Errors are given as 1SD and display biological variability. The analytical precision of the carbonate boron isotope ratio determination was in the order of 0.9‰.

and $\alpha = 1.0272$ [Klochko *et al.*, 2006], respectively. Our data, when plotted against ambient water pH, differ from this pattern. When depicted against EPF pH, the pH at the calcification site, they also fit into this range between theoretical and measured curve. As HL and EPF pH are relatively low (calculated 7.22–7.55 for this experiment), $\delta^{11}\text{B}$ plots in a section of the curve where pH dependent changes in $\delta^{11}\text{B}$ are less pronounced. It seems likely that $\delta^{11}\text{B}$ in *M. edulis* shell reflects the internal fluid pH (EPF) and not that of ambient water. Because pH changes of internal fluids are minor in comparison to ambient water pH changes, only small changes in boron isotope fractionation of the shell can be expected. This fact and the high individual variability render boron isotopes in *M. edulis* shells a less sensitive pH-proxy for seawater pH variations.

[34] However, boron isotopes may be beneficial in providing more information on the processes of biomineralization in bivalves. The extrapallial space is described as the area where calcification takes place. Recent studies show how complex the underlying biological mechanisms probably are. Various organic components, such as a hydrophobic silk gel, acidic proteins, chitin and amorphous calcium carbonate as precursor phases are probably involved in the formation of bivalve shells [e.g., Levi-Kalishman *et al.*, 2001; Weiss *et al.*, 2009; Suzuki *et al.*, 2009; Weiss, 2010]. Suzuki *et al.* [2009] recently identified acidic matrix proteins (Pif97, Pif80) that bind to aragonite crystals and to chitin and might be important regulators of nacre

formation. Suzuki *et al.* [2009] and Weiss [2010] proposed that crystals are formed in a microenvironment separated from the extrapallial fluid via a chitinous membrane. Nevertheless $\delta^{11}\text{B}$ in the shell indicate that shell formation proceeds at pH values typical of the EPF which indicates that calcification is influenced by EPF conditions independent whether calcification occurs separated in a microenvironment close to the shell.

4. Conclusions

[35] Body fluids of *Mytilus edulis* from Kiel Fjord kept under different seawater pCO_2 levels are characterized by low HL and EPF pH values. This is due to the requirements of metabolic CO_2 excretion. HL and EPF pH decreases with 0.024 units per 0.1 units in seawater pH. No significant bicarbonate accumulation to buffer internal pH could be observed. Elemental ratios (Me/Ca) in the extrapallial fluid correlate with growth rates due to the higher consumption of $[\text{Ca}^{2+}]$ during calcification compared to other elements. More data on the mechanisms of biomineralisation and the general composition of shells not only related to ambient water but also to the body fluids are needed. Also the impact of the EPF on calcification independent of the exact precipitation area (directly the EPF or a microenvironment) and the involvement of components like hydrophobic silk gel, acidic proteins, chitin and also maybe amorphous calcium carbonate should be subject of future

studies. It's necessary to better understand the underlying mechanisms of trace element partitioning and isotope fractionation in order to demonstrate a robust use of proxies in bivalve shells. Trace metal composition of the extrapallial fluid and hence of the shell seems to reflect abiotic conditions only indirectly via the impact on metabolism and growth rate. Boron isotopes in *M. edulis* shell do not reflect seawater pH, they seem to represent internal pH (EPF). In addition, boron isotopes show a high variability between individuals and also within single shells. This may reflect variable contributions of boron rich organic matrices present between single crystals. In the future, more attention should be paid to these organic shell parts also with respect to geochemistry as they often are not completely removed before measuring carbonates.

Acknowledgments

[36] The authors wish to kindly acknowledge Magdalena Gutowska for constantly supporting this study. We thank Ulrike Panknin for help with bivalve sampling and culturing algae, Karin Kißling for supporting ICP-OES measurements and Susann Grobe for carbonate chemistry analysis. AH is grateful for statistical help provided by Claas Hiebenthal and Martin Wahl. AH and JF gratefully acknowledge Hauke Vollstaedt for supporting figure corrections. This study was funded by DFG Excellence Cluster "Future ocean" grants to FM. This work is a contribution to the German Ministry of Education and Research (BMBF)-funded project "Biological Impacts of Ocean ACIDification" (BIOACID). The authors like to thank two anonymous reviewers for their constructive and detailed reports significantly improving the manuscript.

References

- Albers, C., and K. Pleschka (1967), Effect of temperature on CO₂ transport in elasmobranch blood, *Respir. Physiol.*, **2**, 261–273, doi:10.1016/0034-5687(67)90031-X.
- Caldeira, K., and M. E. Wickett (2003), Anthropogenic carbon and ocean pH, *Nature*, **425**(6956), 365, doi:10.1038/425365a.
- Cao, L., and K. Caldeira (2008), Atmospheric CO₂ stabilization and ocean acidification, *Geophys. Res. Lett.*, **35**, L19609, doi:10.1029/2008GL035072.
- Carré, M., I. Bentaleb, O. Bruguier, E. Ordinola, N. T. Barrett, and M. Fontugne (2006), Calcification rate influence on trace elements concentrations in aragonitic bivalve shells: Evidences and mechanisms, *Geochim. Cosmochim. Acta*, **70**(19), 4906–4920, doi:10.1016/j.gca.2006.07.019.
- Crenshaw, M. A. (1972), Inorganic composition of molluscan extrapallial fluid, *Biol. Bull.*, **143**, 506–512, doi:10.2307/1540180.
- Crenshaw, M. A., and J. M. Neff (1969), Decalcification at the mantle-shell interface in molluscs, *Am. Zool.*, **9**, 881–889.
- Dickson, A. G. (1990), Standard potential of the reaction: AgCl_s + 1/2 H₂ = Ag_s + HCl_{Aq} and the standard acidity constant of the ion HSO₄⁻ in synthetic sea-water from 273.15 K to 318.15 K, *J. Chem. Thermodyn.*, **22**, 113–127, doi:10.1016/0021-9614(90)90074-Z.
- Dodd, J. R. (1965), Environmental control of strontium and magnesium in *Mytilus edulis*, *Geochim. Cosmochim. Acta*, **29**(5), 385–398, doi:10.1016/0016-7037(65)90035-9.
- Doney, S. C., V. J. Fabry, R. A. Feely, and J. A. Kleypas (2009), Ocean acidification: The other CO₂ problem, *Annu. Rev. Mater. Sci.*, **1**, 169–192, doi:10.1146/annurev.marine.010908.163834.
- Elderfield, H., and G. Ganssen (2000), Past temperature and δ¹⁸O of surface ocean waters inferred from foraminiferal Mg/Ca ratios, *Nature*, **405**(6785), 442–445, doi:10.1038/35013033.
- Fabry, V. J., B. A. Seibel, R. A. Feely, and J. C. Orr (2008), Impacts of ocean acidification on marine fauna and ecosystem processes, *ICES J. Mar. Sci.*, **65**, 414–432, doi:10.1093/icesjms/fsn048.
- Feely, R. A., C. L. Sabine, K. Lee, W. Berelson, J. A. Kleypas, V. J. Fabry, and F. J. Millero (2004), Impact of anthropogenic CO₂ on the CaCO₃ system in the oceans, *Science*, **305**(5682), 362–366, doi:10.1126/science.1097329.
- Fietzke, J., A. Heinemann, I. Taubner, F. Böhm, J. Erez, and A. Eisenhauer (2010), Boron isotope ratio determination in carbonates via LA-MC-ICP-MS using soda-lime glass standards as reference material, *J. Anal. At. Spectrom.*, **25**, 1953–1957, doi:10.1039/c0ja00036a.
- Ford, H., S. A. Schellenberg, B. J. Becker, D. L. Deutschman, K. A. Dyck, and P. L. Koch (2010), Evaluating the skeletal chemistry of *Mytilus californianus* as a temperature proxy: Effects of microenvironment and ontogeny, *Paleoceanography*, **25**, PA1203, doi:10.1029/2008PA001677.
- Foster, G. L. (2008), Seawater pH, pCO₂ and [CO₃²⁻] variations in the Caribbean Sea over the last 130 kyr: A boron isotope and B/Ca study of planktic foraminifera, *Earth Planet. Sci. Lett.*, **271**, 254–266, doi:10.1016/j.epsl.2008.04.015.
- Freitas, P., L. Clarke, H. Kennedy, C. A. Richardson, and F. Abrantes (2006), Environmental and biological controls on elemental (Mg/Ca, Sr/Ca and Mn/Ca) ratios in shells of the king scallop *Pecten maximus*, *Geochim. Cosmochim. Acta*, **70**, 5119–5133, doi:10.1016/j.gca.2006.07.029.
- Freitas, P. S., L. J. Clarke, H. A. Kennedy, and C. A. Richardson (2008), Inter and intra-specimen variability masks reliable temperature control on shell Mg/Ca ratios in laboratory- and field-cultured *Mytilus edulis* and *Pecten maximus* (bivalvia), *Biogeosciences*, **5**(5), 1245–1258, doi:10.5194/bg-5-1245-2008.
- Gosling, E. (1992), Systematics and geographic distribution of *Mytilus*, in *The Mussel Mytilus: Ecology, Physiology, Genetics and Culture*, edited by E. Gosling, pp. 1–19, Elsevier, Amsterdam.
- Gutowska, M. A., F. Melzner, M. Langenbuch, C. Bock, G. Claireaux, and H. O. Pörtner (2010), Acid-base regulatory ability of the cephalopod (*Sepia officinalis*) in response to environmental hypercapnia, *J. Comp. Physiol. B*, **180**, 323–335, doi:10.1007/s00360-009-0412-y.
- Hall-Spencer, J. M., R. Rodolfo-Metalpa, S. Martin, E. Ransome, M. Fine, S. M. Turner, S. J. Rowley, D. Tedesco, and M. C. Buia (2008), Volcanic carbon dioxide vents show ecosystem effects of ocean acidification, *Nature*, **454**, 96–99, doi:10.1038/nature07051.
- Hansen, H., H. Giesenhausen, and G. Behrends (1999), Seasonal and long-term control of bottom-water oxygen deficiency in a stratified shallow-water coastal system, *ICES J. Mar. Sci.*, **56**, 65–71, doi:10.1006/jmsc.1999.0629.

- Heinemann, A., J. Fietzke, A. Eisenhauer, and K. Zumholz (2008), Modification of Ca isotope and trace metal composition of the major matrices involved in shell formation of *Mytilus edulis*, *Geochem. Geophys. Geosyst.*, *9*, Q01006, doi:10.1029/2007GC001777.
- Hemming, N. G., and G. N. Hanson (1992), Boron isotopic composition and concentration in modern marine carbonates, *Geochim. Cosmochim. Acta*, *56*, 537–543, doi:10.1016/0016-7037(92)90151-8.
- Hemming, N. G., R. J. Reeder, and S. R. Hart (1995), Mineral-fluid partitioning and isotopic fractionation of boron in synthetic calcium carbonate, *Geochim. Cosmochim. Acta*, *59*, 371–379, doi:10.1016/0016-7037(95)00288-B.
- Hiebenthal, C. (2009), Sensitivity of *A. islandica* and *M. edulis* towards environmental changes: A threat to the bivalves: An opportunity for palaeo-climatology?, PhD thesis, Christian-Albrechts-Univ., Kiel, Germany. [Available at http://eldiss.uni-kiel.de/macau/receive/dissertation_diss_00003647.]
- Hobbs, M. Y., and E. J. Reardon (1999), Effect of pH on boron coprecipitation by calcite: Further evidence for nonequilibrium partitioning of trace elements, *Geochim. Cosmochim. Acta*, *63*, 1013–1021, doi:10.1016/S0016-7037(98)00311-1.
- Hönisch, B., J. Bijma, A. D. Russell, H. J. Spero, M. R. Palmer, R. E. Zeebe, and A. Eisenhauer (2003), The influence of photosynthesis on the boron isotopic composition of foraminifera shells, *Mar. Micropaleontol.*, *49*, 87–96, doi:10.1016/S0377-8398(03)00030-6.
- Hönisch, B., N. G. Hemming, A. G. Grottoli, A. Amat, G. N. Hanson, and J. Bijma (2004), Assessing scleractinian corals as recorders for paleo-pH: empirical calibration and vital effects, *Geochim. Cosmochim. Acta*, *68*, 3675–3685, doi:10.1016/j.gca.2004.03.002.
- Immenhauser, A., T. F. Nägler, T. Steuber, and D. Hippler (2005), A critical assessment of mollusk $^{18}\text{O}/^{16}\text{O}$, Mg/Ca, and $^{44}\text{Ca}/^{40}\text{Ca}$ ratios as proxies for Cretaceous seawater temperature seasonality, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, *215*(3–4), 221–237, doi:10.1016/j.palaeo.2004.09.005.
- Ip, Y. K., A. M. Loong, K. C. Hiong, W. P. Wong, S. F. Chew, K. Reddy, B. Sivaloganathan, and J. S. Ballantyne (2006), Light induces an increase in the pH of and a decrease in the ammonia concentration in the extrapallial fluid of the giant clam *Tridacna squamosa*, *Physiol. Biochem. Zool.*, *79*(3), 656–664, doi:10.1086/501061.
- Intergovernmental Panel on Climate Change (2007), *Climate Change 2007: The Physical Science Basis: Working Group I Contribution to the Fourth Assessment Report of the IPCC*, edited by S. Solomon et al., Cambridge Univ. Press, New York.
- Kakihana, H., M. Kotaka, S. Satoh, M. Nomura, and M. Okamoto (1977), Fundamental studies on the ion-exchange separation of boron isotopes, *Bull. Chem. Soc. Jpn.*, *50*, 158–163.
- Kasemann, S. A., D. N. Schmidt, J. Bijma, and G. L. Foster (2009), In situ boron isotope analysis in marine carbonates and its application for foraminifera and palaeo-pH, *Chem. Geol.*, *260*, 138–147, doi:10.1016/j.chemgeo.2008.12.015.
- Kautsky, N. (1982), Growth and size structure in a baltic *Mytilus edulis* population, *Mar. Biol. Berlin*, *68*, 117–133, doi:10.1007/BF00397599.
- Kisakürek, B., A. Eisenhauer, F. Böhm, D. Garbe-Schönberg, and J. Erez (2008), Controls on shell Mg/Ca and Sr/Ca in cultured planktonic foraminiferan, *Globigerinoides ruber* (white), *Earth Planet. Sci. Lett.*, *273*(3–4), 260–269.
- Klein, R. T., K. C. Lohmann, and C. W. Thayer (1996a), Bivalve skeletons record sea-surface temperature and $\delta^{18}\text{O}$ via Mg/Ca and $^{18}\text{O}/^{16}\text{O}$ ratios, *Geology*, *24*(5), 415–418, doi:10.1130/0091-7613(1996)024<0415:BSRSST>2.3.CO;2.
- Klein, R. T., K. C. Lohmann, and C. W. Thayer (1996b), Sr/Ca and $^{13}\text{C}/^{12}\text{C}$ ratios in skeletal calcite of *Mytilus trossolus*: Covariation with metabolic rate, salinity and carbon isotopic composition of sea water, *Geochim. Cosmochim. Acta*, *60*, 4207–4221, doi:10.1016/S0016-7037(96)00232-3.
- Klochko, K., A. J. Kaufman, W. Yoa, R. H. Byrne, and J. A. Tossell (2006), Experimental measurement of boron isotope fractionation in seawater, *Earth Planet. Sci. Lett.*, *248*, 276–285, doi:10.1016/j.epsl.2006.05.034.
- Klochko, K., G. D. Cody, J. A. Tossell, P. Dera, and A. J. Kaufman (2009), Re-evaluating boron speciation in biogenic calcite and aragonite using ^{11}B MAS NMR, *Geochim. Cosmochim. Acta*, *73*, 1890–1900, doi:10.1016/j.gca.2009.01.002.
- Kossak, U. (2006), How climate change translates into ecological change: Impacts of warming and desalination on prey properties and predator-prey interactions in the Baltic Sea, PhD thesis, Christian-Albrechts-Univ., Kiel, Germany. [Available at http://eldiss.uni-kiel.de/macau/receive/dissertation_diss_00001910.]
- Larsen, B., H. O. Pörtner, and F. Jensen (1997), Extra- and intracellular acid-base balance and ionic regulation in cod (*Gadus morhua*) during combined and isolated exposures to hypercapnia and copper, *Mar. Biol. Berlin*, *128*(2), 337–346, doi:10.1007/s002270050099.
- Lazareth, C. E., E. Vander Putten, L. Andre, and F. Dehairs (2003), High resolution trace element profiles in shells of the mangrove bivalve *Isognomon ehippium*: a record of environmental spatio-temporal variations?, *Estuarine Coastal Shelf Sci.*, *57*(5–6), 1103–1114, doi:10.1016/S0272-7714(03)00013-1.
- Lehmann, A., W. Krauss, and H. H. Hinrichsen (2002), Effects of remote and local atmospheric forcing on circulation and upwelling in the Baltic Sea, *Tellus, Ser. A*, *54*(3), 299–316, doi:10.1034/j.1600-0870.2002.00289.x.
- Levi-Kalishman, Y., G. Falini, L. Addadi, and S. Weiner (2001), Structure of the nacreous organic matrix of a bivalve mollusk shell examined in the hydrated state using Cryo-TEM, *J. Struct. Biol.*, *135*, 8–17, doi:10.1006/jsbi.2001.4372.
- Lindinger, M. I., D. J. Lauren, and D. G. McDonald (1984), Acid-base-balance in the sea mussel, *Mytilus edulis*. 3. Effects of environmental hypercapnia on intracellular and extracellular acid-base-balance, *Mar. Biol. Lett.*, *5*, 371–381.
- Lopez, O., P. Zuddas, and D. Faivre (2009), The influence of temperature and seawater composition on calcite crystal growth mechanisms and kinetics: Implications for Mg incorporation in calcite lattice, *Geochim. Cosmochim. Acta*, *73*(2), 337–347, doi:10.1016/j.gca.2008.10.022.
- Lorens, R. (1981), Sr, Cd, Mn and Co distribution coefficients in calcite as a function of calcite precipitation rate, *Geochim. Cosmochim. Acta*, *45*, 553–561, doi:10.1016/0016-7037(81)90188-5.
- Lorrain, A., D. Gillikin, Y. M. Paulet, L. Chavaud, A. Lemerrier, J. Navez, and L. Andre (2005), Strong kinetic effects on Sr/Ca ratios in the calcitic bivalve *Pecten maximus*, *Geology*, *33*, 965–968, doi:10.1130/G22048.1.
- Malone, P., and J. Dodd (1967), Temperature and salinity effects on calcification rate in *Mytilus edulis* and its paleo-ecological implications, *Limnol. Oceanogr.*, *12*(3), 432–436, doi:10.4319/lo.1967.12.3.0432.
- McConnaughey, T. A., and D. P. Gillikin (2008), Carbon isotopes in mollusk shell carbonates, *Geo Mar. Lett.*, *28*(5–6), 287–299, doi:10.1007/s00367-008-0116-4.
- McCoy, S. J., L. F. Robinson, C. A. Pfister, J. Wootton, and N. Shimizu (2011), Exploring B/Ca as a pH proxy in

- bivalves: relationships between *Mytilus californianus* B/Ca and environmental data from the northeast Pacific, *Biogeosciences Discuss.*, 8, 5587–5616, doi:10.5194/bgd-8-5587-2011.
- Melzner, F., M. A. Gutowska, M. Langenbuch, S. Dupont, M. Lucassen, M. C. Thorndyke, M. Bleich, and H. O. Pörtner (2009), Physiological basis for high CO₂ tolerance in marine ectothermic animals: pre-adaptation through lifestyle and ontogeny?, *Biogeosciences*, 6, 2313–2331, doi:10.5194/bg-6-2313-2009.
- Melzner, F., P. Stange, K. Trübenbach, J. Thomsen, I. Casties, U. Panknin, and M. A. Gutowska (2011), Interactive effects of food supply and seawater pCO₂ on calcification and internal shell surface dissolution in the blue mussel *Mytilus edulis*, *PLoS ONE*, 6(9), e24223, doi:10.1371/journal.pone.0024223.
- Michaelidis, B., C. Ouzounis, A. Pleras, and H. O. Pörtner (2005), Effects of long-term moderate hypercapnia on acid-base balance and growth rate in marine mussels *Mytilus galloprovincialis*, *Mar. Ecol. Prog. Ser.*, 293, 109–118, doi:10.3354/meps293109.
- Ni, Y., G. L. Foster, B. Bailey, T. Elliott, D. N. Schmidt, P. Pearson, B. Haley, and C. Coath (2007), A core top assessment of proxies for the ocean carbonate system in surface-dwelling foraminifers, *Paleoceanography*, 22, PA3212, doi:10.1029/2006PA001337.
- Nürnberg, D., J. Bijma, and C. Hemleben (1996), Assessing the reliability of magnesium in foraminiferal calcite as a proxy for water mass temperatures, *Geochim. Cosmochim. Acta*, 60(5), 803–814, doi:10.1016/0016-7037(95)00446-7.
- Oomori, T., H. Kaneshima, Y. Maezato, and Y. Kitano (1987), Distribution coefficient of Mg²⁺ ions between calcite and solution at 10–50°C, *Mar. Chem.*, 20(4), 327–336, doi:10.1016/0304-4203(87)90066-1.
- Orr, J. C., et al. (2005), Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms, *Nature*, 437(7059), 681–686, doi:10.1038/nature04095.
- Orr, J. C., et al. (2009), Research priorities for understanding ocean acidification: Summary from the Second Symposium on the Ocean in a High-CO₂ World, 22(4), 182–189, doi:10.5670/oceanog.2009.107.
- Pagani, M., D. Lemarchand, A. Spivack, and J. Gaillardet (2005), A critical evaluation of the boron isotope-pH proxy: the accuracy of ancient pH estimates, *Geochim. Cosmochim. Acta*, 69, 953–961, doi:10.1016/j.gca.2004.07.029.
- Pane, E. F., and J. P. Barry (2007), Extracellular acid-base regulation during short-term hypercapnia is effective in a shallow-water crab but ineffective in a deep-sea crab, *Mar. Ecol. Prog. Ser.*, 334, 1–9, doi:10.3354/meps334001.
- Pörtner, H. O., M. Langenbuch, and A. Reipschläger (2004), Biological impact of elevated ocean CO₂ concentrations: Lessons from animal physiology and Earth history, *J. Oceanogr.*, 60, 705–718, doi:10.1007/s10872-004-5763-0.
- Reynaud, S., N. G. Hemming, A. J. Leclerc, and J.-P. Gattuso (2004), Effect of pCO₂ and temperature on the boron isotopic composition of the zooxanthellate coral *Acropora sp.*, *Coral Reefs*, 23, 539–546, doi:10.1007/s00338-004-0399-5.
- Ries, J. B., A. L. Cohen, and D. C. McCorkle (2009), Marine calcifiers exhibit mixed responses to CO₂-induced ocean acidification, *Geology*, 37, 1131–1134, doi:10.1130/G30210A.1.
- Rollion-Bard, C., and J. Erez (2010), Intra-shell boron isotope ratios in the symbiont-bearing benthic foraminiferan *Amphistegina lobifera*: Implications for δ¹¹B vital effects and paleo-pH reconstructions, *Geochim. Cosmochim. Acta*, 74, 1530–1536, doi:10.1016/j.gca.2009.11.017.
- Rollion-Bard, C., D. Blamart, J. Trebosc, G. Tricot, A. Mussi, and J.-P. Cuif (2011), Boron isotopes as pH proxy: A new look at boron speciation in deep-sea corals using ¹¹B MAS NMR and EELS, *Geochim. Cosmochim. Acta*, 75, 1003–1012, doi:10.1016/j.gca.2010.11.023.
- Roy, R. N., L. N. Roy, K. M. Vogel, C. Porter-Moore, T. Pearson, C. E. Good, F. J. Millero, and D. M. Campbell (1993), The dissociation constants of carbonic acid in seawater at salinities 5 to 45 and temperatures 0 to 45 °C, *Mar. Chem.*, 44, 249–267, doi:10.1016/0304-4203(93)90207-5.
- Sabine, C. L., et al. (2004), The oceanic sink for anthropogenic CO₂, *Science*, 305(5682), 367–371, doi:10.1126/science.1097403.
- Sanyal, A., M. Nugent, R. J. Reeder, and J. Bijma (2000), Seawater pH control on the boron isotopic composition of calcite: evidence from inorganic calcite precipitation experiments, *Geochim. Cosmochim. Acta*, 64, 1551–1555, doi:10.1016/S0016-7037(99)00437-8.
- Sanyal, A., J. Bijma, H. Spero, and D. W. Lea (2001), Empirical relationship between pH and the boron isotopic composition of *Globigerinoides sacculifer*: implications for the boron isotope paleo-pH proxy, *Paleoceanography*, 16, 515–519, doi:10.1029/2000PA000547.
- Schöne, B. R., Z. Zhang, J. Jacob, D. G. Gillikin, T. Tütken, D. Garbe-Schönberg, T. MacConnaughey, and A. Soldati (2010), Effect of organic matrices on the determination of the trace element chemistry (Mg, Sr, Mg/Ca, Sr/Ca) of aragonitic bivalve shells (*Arctica islandica*)—Comparison of ICP-OES and LA-ICP-MS data, *Geochem. J.*, 44, 23–37.
- Spicer, J. I., A. Raffo, and S. Widdicombe (2007), Influence of CO₂-related seawater acidification on extracellular acid-base balance in the velvet swimming crab *Necora puber*, *Mar. Biol. Berlin*, 151, 1117–1125, doi:10.1007/s00227-006-0551-6.
- Suzuki, M., K. Saruwatari, T. Kogure, Y. Yamamoto, T. Nishimura, T. Kato, and H. Nagasawa (2009), An acidic matrix protein, Pif, is a key macromolecule for nacre formation, *Science*, 325, 1388–1390, doi:10.1126/science.1173793.
- Takesue, R. K., C. R. Bacon, and J. K. Thompson (2008), Influences of organic matter and calcification rate on trace elements in aragonitic estuarine bivalve shells, *Geochim. Cosmochim. Acta*, 72(22), 5431–5445, doi:10.1016/j.gca.2008.09.003.
- Thomsen, J., and F. Melzner (2010), Moderate seawater acidification does not elicit long-term metabolic depression in the blue mussel *Mytilus edulis*, *Mar. Biol. Berlin*, 157(12), 2667–2676, doi:10.1007/s00227-010-1527-0.
- Thomsen, J., et al. (2010), Calcifying invertebrates succeed in a naturally CO₂ enriched coastal habitat but are threatened by high levels of future acidification, *Biogeosciences*, 7, 3879–3891, doi:10.5194/bg-7-3879-2010.
- Vander Putten, E., F. Dehairs, E. Keppens, and W. Baeyens (2000), High distribution of trace elements in the calcite shell layer of modern *Mytilus edulis*: Environmental and biological controls, *Geochim. Cosmochim. Acta*, 64(6), 997–1011, doi:10.1016/S0016-7037(99)00380-4.
- Vengosh, A., Y. Kolodny, A. Starinsky, A. R. Chivas, and M. T. McCulloch (1991), Coprecipitation and isotopic fractionation of boron in modern biogenic carbonates, *Geochim. Cosmochim. Acta*, 55, 2901–2910, doi:10.1016/0016-7037(91)90455-E.
- Wanamaker, A. D., K. J. Kreutz, T. Wilson, H. W. Borns, D. S. Introne, and S. Feindel (2008), Experimentally determined Mg/Ca and Sr/Ca ratios in juvenile bivalve calcite for *Mytilus edulis*: implications for paleotemperature reconstructions,

- Geo Mar. Lett.*, 28(5–6), 359–368, doi:10.1007/s00367-008-0112-8.
- Weiss, I. M. (2010), Jewels in the pearl, *ChemBioChem*, 11, 297–300, doi:10.1002/cbic.200900677.
- Weiss, I. M., S. Kaufmann, B. Heiland, and M. Tanaka (2009), Covalent modification of chitin with silk-derivatives acts as an amphiphilic self-organizing template in nacre biomineralisation, *J. Struct. Biol.*, 167, 68–75, doi:10.1016/j.jsb.2009.04.005.
- Weiss, R. F. (1974), Carbon dioxide in water and seawater: The solubility of a non-ideal gas, *Mar. Chem.*, 2, 203–215, doi:10.1016/0304-4203(74)90015-2.
- Wilbur, K. M., and A. M. Bernhardt (1984), Effects of amino acids, magnesium and molluscan extrapallial fluid on crystallization of calcium carbonate: In vitro experiments, *Biol. Bull.*, 166, 251–259, doi:10.2307/1541446.
- Wilbur, K. M., and A. Saleuddin (1983), Shell formation, in *The Mollusca*, vol. 4, *Physiology, Part 1*, edited by A. Saleuddin and K. M. Wilbur, pp. 235–287, Academic, New York.
- Wong, W. H., and J. S. Levinton (2004), Culture of the blue mussel *Mytilus edulis* (Linnaeus, 1758) fed both phytoplankton and zooplankton: a microcosm experiment, *Aquacult. Res.*, 35(10), 965–969, doi:10.1111/j.1365-2109.2004.01107.x.
- Yu, J., H. Elderfield, and B. Hoenisch (2007), B/Ca in planktonic foraminifera as a proxy for surface seawater pH, *Paleoceanography*, 22, PA2202, doi:10.1029/2006PA001347.