

# The observation of the thin-ice thickness distribution within the Laptev Sea polynya using MODIS data

S. Adams<sup>1</sup>, S. Willmes<sup>1</sup>, D. Schröder<sup>1,2</sup>, G. Heinemann<sup>1</sup>

<sup>1</sup>Environmental Meteorology, University of Trier, Trier, Germany;

<sup>2</sup>Centre for Polar Observation and Modelling, University College London, London, UK

\*corresponding author: Susanne Adams (susanne.adams@uni-trier.de)



## 1. Motivation

- Laptev Sea = Ice factory
- Variables essential for the ice-production calculation:
  - Polynya area
  - Thin-ice thickness distribution
  - Atmospheric Variables
- Thin-ice layer = Insulation layer that effectively reduces the heat loss to the atmosphere
- Remote sensing data is suitable to derive the thin-ice thickness distribution, however, the spatial resolution issue has to be taken into account
- We use high-resolution MODIS data to derive the thin-ice thickness distribution within a polynya

## 2. Thermal thin ice thickness retrieval




Figure 1: Thin-ice thickness retrieval scheme.  $L_{\downarrow}$  and  $L_{\uparrow}$  are the incoming and outgoing long-wave radiation components,  $H_0$  and  $E_0$  are the turbulent fluxes,  $Q_A$  is the net energy flux to the atmosphere,  $Q_i$  is the conductive heat flux through the ice,  $T_s$  is the ice-surface temperature,  $T_f$  is the freezing temperature of sea water and  $h_i$  is the ice thickness.

- Thin-ice thickness retrieval is based on the relation between ice-surface temperature and thin-ice thickness
- Calculation of TIT following Yu and Lindsay, 2003)
- Atmospheric heat flux to the atmosphere  $Q_A$  equals the conductive heat flux through the ice  $Q_i$  (Fig. 1)
- Modification of the algorithm at two calculation steps:
  - (1) Calculation of the turbulent heat fluxes (iterative bulk approach based on Launiainen & Vihma, 1990 instead of simple bulk equations)
  - (2) Calculation of the atmospheric emission coefficient required for the determination of the incoming long-wave radiation (new improved parameterization following Jin et al., 2006)

## 3. Example of the MODIS thin-ice thickness

### Example of MODIS single scene




Figure 2: MODIS  $T_s$  from 6 January 2009 0205 UTC; corresponding NCEP  $T_a$  from 6 January 2009 0000 UTC; ice-thickness distribution as calculated with MODIS  $T_s$  and NCEP atmospheric variables (TIT.MODIS+NCEP).

### Example of a daily MODIS TIT map




Figure 3: Daily MODIS TIT map from 6 January 2009 (gray = thick ice).

- Daily maps are assimilated into the sea ice model (see Section 5)

## Acknowledgements

This work is part of the German-Russian cooperation 'System Laptev Sea' funded by the BMBF under grant 03G0759D. MODIS data is provided by NASA, NCEP Reanalysis data by the U.S. National Centre for Environmental Prediction / Dept. of Energy (NCEP/DOE)

## 4. Sensitivity analysis of the MODIS thin-ice thicknesses

(a)	MODIS	NCEP
Ice-surface temp.	$\pm 1.6^\circ\text{C}$	
2-m air temp.		$\pm 4.5^\circ\text{C}$
10-m wind speed		$\pm 1.3\text{ m s}^{-1}$
Relative humidity		$\pm 20\%$

Tables: (a) Uncertainties of the input variables for the calculation of thin-ice thicknesses. Values from Hall et al. (2004), Ernstdorf et al. (2011), Renfrew et al. (2002). (b) Results of the Monte Carlo error estimation for winter 2007/08 and 2008/09.

(b)	Winter 2007/08	Winter 2008/09	Mean of both winters
Ice class (m)	TIT <sub>MODIS+NCEP</sub> (cm)	TIT <sub>MODIS+NCEP</sub> (cm)	TIT <sub>MODIS+NCEP</sub> (cm)
0.00 - 0.05	$\pm 1.0$	$\pm 1.0$	$\pm 1.0$
0.05 - 0.10	$\pm 2.0$	$\pm 2.2$	$\pm 2.1$
0.10 - 0.20	$\pm 5.2$	$\pm 5.3$	$\pm 5.3$
0.20 - 0.30	$\pm 16.8$	$\pm 12.0$	$\pm 14.4$
0.30 - 0.40	$\pm 34.2$	$\pm 28.4$	$\pm 31.3$
0.40 - 0.50	$\pm 36.7$	$\pm 60.2$	$\pm 48.5$
mean up to 0.20	$\pm 4.7$	$\pm 4.6$	$\pm 4.7$
mean up to 0.50	$\pm 26.1$	$\pm 36.0$	$\pm 31.1$

- $T_s$  and 2-m air temperature ( $T_a$ ) strongly influence the calculation of the ice thickness (not shown)
- Underestimation of  $T_a$  = strong underestimation of TIT; overestimation of  $T_a$  = moderate overestimation of TIT
- Uncertainties in the atmospheric variables have a smaller impact on very thin ice than on thicker ice
- The atmospheric data ( $T_a$ ) have a strong impact on the quality of the retrieved ice thickness (Fig. 4)




Figure 4: (a) Averaged sum of reference thin-ice pixels for ice classes cl1 0 - 0.05 m, cl2 0.05 - 0.1 m and cl3 0.1 - 0.2 m. (b)-(i) Anomaly to reference ice thickness for the input variables with their minimum and maximum uncertainty.

## 5. Combined remote sensing – sea ice model approach




Figure 5: Number of grid points for different ice-thickness classes for April 2008. (a) MODIS thin-ice thickness data. The data is interpolated to FESOM's 5-km grid. (b) Simulations of FESOM's available model data are used only. (c) Simulations of FESOM's assimilated model run. All grid cells are shown.

## References

- T. Ernstdorf, D. Schröder, S. Adams, G. Heinemann, R. Timmermann, and S. Danilov, "Impact of atmospheric forcing data on simulations of the Laptev Sea polynya dynamics using the sea-ice ocean model FESOM," *J. Geophys. Res.*, vol. 116, no. C12, 2011
- D. K. Hall, J. R. Key, K. A. Case, G. A. Riggs, and D. J. Cavalieri, "Sea ice surface temperature product from MODIS," *IEEE Trans. Geosci. Remote Sensing*, vol. 42, no. 5, pp. 1076–1087, 2004.
- X. Jin, D. Barber, and T. Papakyriakou, "A new clear-sky downward longwave radiative flux parameterization for Arctic areas based on rawinsonde data," *J. Geophys. Res.*, vol. 111, no. D24, 2006.
- J. Launiainen and T. Vihma, "Derivation of turbulent surface fluxes — An iterative flux-profile method allowing arbitrary observing heights," *Environmental Software*, vol. 5, no. 3, pp. 113–124, 1990.
- A. Renfrew, G. W. K. Moore, P. S. Guest, and K. Bumke, "A Comparison of Surface Layer and Surface Turbulent Flux Observations over the Labrador Sea with ECMWF Analyses and NCEP Reanalyses," *Journal of Physical Oceanography*, *J. Phys. Oceanogr.*, vol. 32, no. 2, pp. 383–400, [http://dx.doi.org/10.1175/1520-0485\(2002\)032<0383:ACOSLA>2.0.CO;2](http://dx.doi.org/10.1175/1520-0485(2002)032<0383:ACOSLA>2.0.CO;2), 2002.
- Y. Yu and R. W. Lindsay, "Comparison of thin ice thickness distributions derived from RADARSAT Geophysical Processor System and advanced very high resolution radiometer data sets," *J. Geophys. Res.*, vol. 108, no. C12, 2003.