

### 4.7.3 Shore face profiles in the area of Cape Mamontov Klyk: echo sounding, seawater and sea bottom deposits sampling

*Mikhail N. Grigoriev and Waldemar Schneider*

#### 4.7.3.1. Introduction

One of the tasks of the expedition "Lena-Anabar 2003" was a determination of shore face features in the area of Cape Mamontov Klyk at the adjacent near-shore shelf. In this region, a bathymetric survey, seawater and bottom sediment sampling were carried out in August 2003. The main reason of seabed research is a purpose to conduct a relatively deep drilling on the shallow shelf from the sea ice in the nearest future. Previous shore face investigations in this sector of the Laptev Sea (near Terpay-Tumsa Cape) have been carried out in 2000 (Are et al., 2001).


#### 4.7.3.2. Methods

Bathymetric studies were conducted with help the echo-sounding device and rubber boat with "Honda" engine. The field of bathymetric survey has occupied an area about 60 km<sup>2</sup> (4 x 15 km), between isobath 1 and 10 meters (Figure 4.7-10). The length of all bathymetric profiles exceeds 150 km. For bottom sediment and seawater sampling a standard sampling dredger and bathometer were used.

#### 4.7.3.3. Preliminary results

As a result of bathymetric survey the bathymetric scheme of studied seabed was created. The mean shore face inclination at that part of the shelf is extremely slightly, about 0.0007. At least two quite evident sub-sea terraces (or terrace-like surfaces) were discovered at the shore face. The first vast sub-sea terrace is over the range of depth about 5-6 meters and another terrace begins from 9.5 meters depth (see Fig. 4.7-10).

It is very interesting that a few years ago approximately the same sub-sea terraces were found near the Cape Terpay-Tumsa in 60 km east of the Cape Mamontov Klyk (Are et al., 2001). Probably, these forms are results of several stages of coastal erosion activity. Taking into account that an average coastal retreat rate for the whole studied coast is about 4.0 m/year, the shallow sub-sea terrace, placed from 4.5 km to 11.5 km from the shoreline, was formed during a period of about 2900-1000 years BP. It is very difficult to evaluate the age of formation of the deeper terrace, beginning in a distance of about 14.5 km from the coast. It is possible that this terrace was formed during first stage of sea level stabilization in Holocene (about 5000-3600 years BP).



**Figure 4.7-10.** Bathymetric scheme and the shore face relief adjacent to the Mamontov Klyk Cape area (August 2003). Bathymetric profiles - white dotted line. Vertical white line is a location of proposed drilling profile.

During summer fieldwork 30 bottom sediments samples were collected from 0 to 10 meters depth as well as ice and frozen sediment samples from the coastal outcrops for different types of analysis, which were transported to Germany (Tabs. 4.7-2, 4.7-3). At present time all samples are processed in the laboratories.

**Table 4.7-2:** List of bottom sediment samples, Profile MAK-3 (August 2003)

No.	Name	Water depth (m)	Number of samples
1	S. MAK-3 (0 m)	0	3
2	S. MAK-3 (1 m)	1	3
3	S. MAK-3 (2 m)	2	3
4	S. MAK-3 (3 m)	3	3
5	S. MAK-3 (4 m)	4	3
6	S. MAK-3 (5 m)	5	3
7	S. MAK-3 (6 m)	6	3
8	S. MAK-3 (7 m)	7	3
9	S. MAK-3 (8 m)	8	3
10	S. MAK-3 (9 m)	9	3
11	S. MAK-3 (10 m)	10	3

**Table 4.7-3:** Frozen water and frozen sediment samples „Lena – Anabar 2003“

No.	Name	Depth (Water depth / Depth of sampling)	Description	Number of samples
1	MAK-3	0m; 2,5m; 5m; 7,5m; 10m	Frozen sea water samples	5
2	MAK-3	7/0m; 7/3,5m; 7/7m; 10/0m;10/5m;10/10m 0m;2,5/0m;5/0m;5/5m;	Frozen sea water sample (Unfiltered)	10
3	MAK-3	0/0m; 2,5/0m; 5/0m; 5/5m; 7/0m; 7/3,5m; 7/7m; 10/0m; 10/5m; 10/10m)	Frozen sea water sample (Filtered)	10

#### 4.7.3.4. Further investigations

The study of shore face profile structure and dynamics in the area of Cape Mamontov Klyk will be continued in spring 2005. The new facts concerning development of the shore face could be very useful for study of sub-sea permafrost evolution and regional paleogeographical reconstructions on the whole.

#### 4.8. References

- Amaral, J.A., Ekins, A., Richards, S.R. and Knowles, R. (1998). Effects of selected monoterpenes on methane oxidation, denitrification and aerobic metabolism by bacteria in pure culture. *Appl. Environ. Microbiol.* 64: 520-525.
- Are F.E., Grigoriev M. N., Rachold V., Hubberten, H.-W., Razumov S. O., and W. Schneider (2001) Shoreface profiles of the central and western Laptev Sea coast: In: Russian-German Cooperation SYSTEM LAPTEV SEA-2000: The Expedition LENA 2000. Reports on Polar Research, 388: 60-64.
- Boyarsky O.G. and Mitt K.L. (1961). New data on relic ground ice in Olenek-Anabar tundra. *Merzlotnye Issledovania, MSU*, v. 1, p.154-162. (In Russian).
- Calhoun, A. and King, G.M. (1997). Regulation of root-associated methanotrophy by oxygen availability in the rhizosphere of two aquatic macrophytes. *Appl. Environ. Microbiol.* 63 (8): 3051-3058.
- Cao, M., Marshall, S. and Gregson, K. (1996). Global carbon exchange and methane emissions from natural wetlands: application of a process-based model. *J. Geophys. Res.* 101 (D9): 14399-14414.
- Chekanovsky N.L. (1896). The daybook of the expedition at rivers Nizhniaya Tunguska, Olenek and Lena in 1873-1875. *Transactions of Russian Geographical Society*, v. XX, ? 1, (In Russian).
- Christensen, T.R. (1993). Methane emission from arctic tundra. *Biogeochemistry* 21: 117-139.
- Drachev S.S., Savostin L.A., Groshev V.G., Bruni I.E. (1998). Structure and geology of the continental shelf of the Laptev Sea, Eastern Russia Arctic., *Tectonophysics* 298: 357-393.
- Galabala R.O. (1987). New data on delta Lena construction. In "Quaternary period of Northern-East Asia. Magadan, SVKNII DVO AN SSSR, p. 152-172. (In Russian).
- Grigoriev M. N., Rachold V., Are F. E., Hubberten H.-W., Razumov S. O., and W. Schneider (2001). Coastal erosion in the Western coast of the Laptev Sea (2001). In: *The Lena Delta expedition 2000. Reports on Polar Research, Bremerhaven, Germany*, 335, p. 54-60.
- Grigoriev M. N., Rachold V., Bolshiyarov D.Yu., Pfeiffer E.-M., Schirrmeister L., Wagner D., and H.-W. Hubberten (Eds.) (2003). *Russian-German Cooperation SYSTEM LAPTEV SEA 2000: The Expedition LENA 2002. Reports on Polar and Marine Research*, 466, Bremerhaven, Germany, 341 pp.
- Grigoriev M.N. (1993). Cryomorphogenesis of the Lena River mouth area. *Permafrost Institute SB AN USSR, Yakutsk, Russia*. 176 pp. (In Russian).
- Grigoriev N.F. (1966). *Permafrost in the Yakutian coastal Zone*. Nauka Press, Moscow, Russia, 180 pp. (In Russian)
- Grigoriev, M.N., and V.V. Kunitsky (2000). Ice Complex of Yakutian Arctic Coast as a Sediment Source on the Shelf.: *Hydro meteorological and Biogeochemical Studies in the Arctic*. Vladivostok: Arctic Regional Centre, Far-Eastern Branch of RAS, Vol. 2, pp. 109-116 (in Russian).
- Harris, R.C., Bartlett, K., Frolking, S. and Crill, P. (1993). Methane emissions from northern high-latitude wetlands. In: Oremland, R.S. (ed.) *Biogeochemistry of Global Change: Radioactively Active Trace Gases*. Chapman and Hall, New York, pp. 449-485.
- Kolpakov V.V. (1973). The paleogeographical meaning of Quaternary Aeolian deposits in the North of Eastern Siberia. In "Several problems of regional geology". M. p. 38-42. (In Russian).
- Krueger, M., Frenzel, P. and Conrad, R. (2001). Microbial processes influencing methane emission from rice fields. *Global Change Biology* 7: 49-63.
- Kutzbach, L., Wagner, D. and Pfeiffer, E.-M. (2004). Effect of microrelief and vegetation on methane emission from wet polygonal tundra, Lena Delta, Northern Siberia. *Biogeochemistry* 69: 341-362.

- Meyer, H. (2003). Studies on recent cryogenesis. – In: Russian-German Cooperation SYSTEM LAPTEV SEA - The Expeditions LENA 2002, Ed.: Grigoriev et al., Reports of Polar and Marine Research 466, p. 29-48.
- Meyer, H., A. Yu. Dereviagin, C. Siegert, and H.-W. Hubberten (2002a). Paleoclimate studies on Bykovsky Peninsula, North Siberia; hydrogen and oxygen isotopes in ground ice.- Polarforschung, 70, p. 37-51.
- Meyer, H., A. Yu. Dereviagin, C. Siegert, L. Schirrmeister, and H.-W. Hubberten (2002b). Palaeoclimate reconstruction on Big Lyakhovsky Island, North Siberia; hydrogen and oxygen isotopes in ice wedges.- Permafrost and Periglacial Processes, 13, p. 91-105.
- Moraes, F. and Khalil, M.A.K. (1993). Permafrost methane content: 2. Modeling theory and results. Chemosphere 26: 595-607.
- Pfeiffer E.-M. and M.N. Grigoriev (Eds.) (2002). Russian-German Cooperation SYSTEM LAPTEV SEA 2000: The Expedition LENA 2001. Reports on Polar Research, 426, Bremerhaven, Germany, 186 pp.
- Pfeiffer, E.-M., Akhmadeeva, I., Becker, H., Friedrich, K., Wagner, D., Quass, W., Zhurbenko, M., Zöllner, E., Boike, J. (1999). Modern Processes in Permafrost Affected Soils. In: Rachold, V. and Grigoriev, M.N. (ed.) Expeditions in Siberia 1998. Reports on Polar Research 315: pp. 19-80.
- Popp, T.J., Chanton, J.P., Whiting, G.J. and Grant, N. (2000). Evaluation of methane oxidation in the rhizosphere of a *Carex* dominated fen in north central Alberta, Canada. Biogeochemistry 51: 259-281.
- Puminov A.P. (1962). Covering Quaternary deposits of Anabar-Olenek interfluves. In "Quaternary geology and geomorphology of Siberia". Novosibirsk, Publications of IGiG SO AN SSSR, v. 27., p. 102-117. (In Russian).
- Rachold V., and M. N. Grigoriev (Eds.) (1999). Russian-German Cooperation SYSTEM LAPTEV SEA-2000: The Expedition LENA DELTA 1998. Reports on Polar Research, 315, Bremerhaven, Germany, 259 pp.
- Rachold V., and M. N. Grigoriev (Eds.) (2000). Russian-German Cooperation SYSTEM LAPTEV SEA-2000: The Expedition LENA 1999. Reports on Polar Research, 354, Bremerhaven, Germany, 269 pp.
- Rachold V., and M. N. Grigoriev (Eds.) (2001). Russian-German Cooperation SYSTEM LAPTEV SEA-2000: The Expedition LENA 2000. Reports on Polar Research, 388, Bremerhaven, Germany, 135 pp.
- Rivkina, E. and Gilichinsky, D. (1996). Methane as a paleoindicator of the dynamics of permafrost deposits. Lithology and Mineral Resources 31 (4), 396-399.
- Roulet, N.T., Jano, A., Kelly, C.A., Klinger, L.F., Moore, T.R., Protz, R., Ritter, J.A. and Rouse, W.R. (1994). Role of the Hudson Bay lowland as a source of atmospheric methane. J. Geophys. Res. 99 (D1). 1435-1454.
- Saks V.N. (1963). Quaternary period in Soviet Arctic. Publications of NIIGA, v. 77. (In Russian).
- Schirrmeister, L., Grosse, G., Schwamborn, G., Andreev, A.A., Meyer, Kunitsky, V.V., Kuznetsova, T.V., Dorozkina, M.V., Pavlova, Y.Y, Bobrov, A., Oezen, D. (2004). Late Quaternary history of the accumulation plain north of the Chekanovsky Ridge (Lena Delta, Russia). A multidisciplinary approach.- Polar geography, 27: 277-319.
- Sochava V.B. (1933). Tundra of Anabar River basin. Publications of Geographical Society, v. 65, ? 4. (In Russian).
- Soil Survey Staff, Soil Conservation Service, USDA (1998). Keys to Soil Taxonomy 8<sup>th</sup> edition. Pocahontas, Blacksburg, Virginia.
- Telepnev E.V. (1981). Sub-sea frozen zone of the near-shore part of the Big Lyakhovsky Island. In: Cryolithozone of the Arctic Shelf. Permafrost Institute SB AN USSR, Yakutsk, Russia, pp. 44-53 (In Russian).
- Toll E.V. (1894). The expedition of Academy of Science to New Siberian Islands and Coast of North Ocean. Izvestia Russkogo Geographicheskogo obshestva, v. 30, ? 4. (In Russian).

- Tolmachev I.P. (1904). New founding on Siberia's geology. Publications of St.-Petersburg's Community of Natural Scientists, v. XXXIV, ? 1. (In Russian).
- Vaikmäe, R. (1989). Oxygen Isotopes in Permafrost and Ground Ice - A new Tool for paleoclimatic Investigations.- Proceedings of the 5th Working Meeting Isotopes in Nature, Leipzig, September 1989, p. 543-553.
- Whalen, S.C., Reeburgh, W.S. (1992). Interannual variations in tundra methane emission: a 4-year time series at fixed sites. Global Biogeochem. Cycles 6: 139-159.
- Zhukov V.V., Gorina I.F., Pinchuk L.Ya. (1968). Cenozoic diamondiferous of Anabar-Olenek interfluves. Publications of NIIGA, v.156. (In Russian).