

# Constraints from Sr-Nd-Hf-Pb-Os isotopes for enriched and depleted Hawaiian plume components in the oldest Emperor Seamounts

KAJ HOERNLE<sup>1\*</sup>, BRUCE SCHAEFER<sup>2</sup>,  
MAXIM PORTNYAGIN<sup>1</sup>, FOLKMAR HAUFF<sup>1</sup> AND  
REINHARD WERNER<sup>1</sup>

<sup>1</sup>GEOMAR Helmholtz Centre for Ocean Research Kiel,  
Germany (\*correspondence: [khoernle@geomar.de](mailto:khoernle@geomar.de))

<sup>2</sup>Earth and Planetary Sciences, Macquarie University, NSW  
Australia

Depleted and enriched geochemical compositions have been found at the oldest (~85-70 Ma) end of the Hawaiian-Emperor hotspot chain. The origin of these components is controversial. During two R/V SONNE 201 cruise legs, fresh lavas were recovered from the NW Emperor seamounts (Suizei, Tenji and Meiji). Major and trace element and Sr-Nd-Hf-Pb-Os isotope investigations provide new insights into the origin and evolution of the Hawaiian plume volcanism in the Late Cretaceous. The recovered samples range from enriched tholeiites and alkali basalts (e.g., La/Yb = 1.5-6.5) to depleted tholeiites (La/Yb = 0.7-1.2). Initial Pb and Nd isotope ratios form a tight inverse linear array, consistent with mixing of 1) an enriched Kea-type component found in present day Hawaiian tholeiites (Abouchami et al., 2005, *Nature* 434: 851-856) and 2) a depleted component with less radiogenic Nd than Pacific MORB at a given Pb isotopic composition. Initial Pb and Os isotope ratios correlate inversely whereas Nd and Os isotope ratios correlate positively. The depleted component has radiogenic (superchondritic) Os and thus cannot be derived from a MORB source. Therefore, we believe that the Suizei depleted endmember was a component in the Hawaiian plume, possibly derived from recycled metasomatized oceanic lithospheric mantle (Regelous et al. 2003, *J. Petrol.* 44: 113-140; Huang et al. 2005, *G-cubed* 6: 1-52; Portnyagin et al., 2009, *EPSL* 287: 194-204) or ancient non-chondritic primitive mantle (Jackson et al., 2010, *Nature* 466: 853-856; Jackson and Carlson, 2011, *Nature* 476: 316-319). The enriched component, with similar Sr-Nd-Pb isotope ratios to the Hawaiian Kea component, however, has subchondritic Os isotope ratios, suggesting recycled ancient subcontinental lithospheric mantle in the source. The subchondritic Os contrasts with slightly superchondritic ratios found in lavas from Mauna Kea, posing the question as to whether the Kea component was really present in the early Hawaiian plume or if the differences in Os isotopic composition simply reflect temporal heterogeneity in the plume source.